

A finitely presented E_∞ -prop II: cellular context

Anibal M. Medina-Mardones^{ab}

^aDepartment of Mathematics, University of Notre Dame, Notre Dame, IN 46556, United States

^bMax Planck Institute for Mathematics, Bonn, 53111, Germany

Abstract

We construct, using finitely many generating cells and relations, props in the category of CW-complexes with the property that their associated operads are models for the E_∞ -operad. We use one of these to construct a cellular E_∞ -bialgebra structure on the interval and derive from it a natural cellular E_∞ -coalgebra structure on the geometric realization of a simplicial set which, passing to cellular chains, recovers up to signs the Barratt-Eccles and Surjection coalgebra structures introduced by Berger-Fresse and McClure-Smith. We use another prop, a quotient of the first, to relate our constructions to earlier work of Kaufmann and prove a conjecture of his. This is the second of two papers in a series, the first investigates analogous constructions in the category of chain complexes.

Received: 17th October, 2019. Accepted: 28th June, 2021.

MSC: 18M75, 55U10, 18M30.

Keywords: Operads and props, E_∞ -structures, simplicial sets, diagonal approximations, arc surfaces.

1. Introduction

This is the second of two papers that, with the exception of section 5.3., can be read independently. In the first [24], we work over the category of differential graded modules. In this one, we do so over the category of CW-complexes and CW-maps.

A purposeful construction of model for the E_∞ -operad is central in most contexts where commutativity up to coherent homotopies plays a role. No model for the E_∞ -operad can be described in terms of finitely many generating cells and relations. However, as demonstrated in this work, passing to a more general setting with multiple inputs and outputs allows to finitely present props whose associated operad is a model for the E_∞ -operad.

We introduce three such props related by quotient morphisms

$$\tilde{\mathcal{S}} \rightarrow \mathcal{S} \rightarrow \mathcal{MS}.$$

Email address: ammedmar@mpim-bonn.mpg.de (Anibal M. Medina-Mardones)

© Anibal M. Medina-Mardones, 2021, under a Creative Commons Attribution 4.0 International License.

The cellular chains on the props \mathcal{S} and \mathcal{MS} are isomorphic to the algebraic props introduced in [24], in particular, the cellular chains of the operad associated to \mathcal{MS} are isomorphic up to signs to the Surjection operad of [19, 1]. We use $\tilde{\mathcal{S}}$ to provide the interval with an E_∞ -bialgebra structure and derive from it a natural E_∞ -coalgebra structure on the geometric realization of a simplicial set extending a cellular approximation to the diagonal. We describe how the natural Barratt-Eccles and Surjection coalgebra structures defined in [1] and [19] on the normalized chains of simplicial sets are deduced, up to signs, from the E_∞ -bialgebra structure on the interval introduced here. Additionally, we prove that \mathcal{MS} is isomorphic to an Arc Surface prop [13] whose associated operad, introduced by Kaufmann in [11], was conjectured in section 4.4. loc. cit. to have cellular chains isomorphic to the Surjection operad.

Given the concise graphical language it provides, the combinatorial formulation we present in this paper is of independent interest. In a string topology interpretation [30, 9], we note that the isomorphism of our prop \mathcal{MS} and Kaufmann's stabilized Arc Surface prop induces an action of the cellular chains of \mathcal{MS} on the Hochschild cochains of a normalized semi-simple Frobenius algebra, see Theorem 8 in [10]. The coproduct and product in our presentation correspond, via the input-output duality, to the Chas-Sullivan product [5] and Goresky-Hingston coproduct [8] in the formalism of [12].

We present an overview of the content of this article. In the second section, we review the material on operads and props needed for the rest of the paper, in particular we define the notion of finite presentation of a cellular prop. In the third section, we finitely present the prop $\tilde{\mathcal{S}}$ and compute its homotopy type. In the fourth section, we construct a natural E_∞ -coalgebra structure on the geometric realization of simplicial sets from an $\tilde{\mathcal{S}}$ -bialgebra structure on the interval. In the final section, we introduce the props \mathcal{S} and \mathcal{MS} and study their relationships to Arc Surface props and the Surjection operad.

2. Preliminaries

We work in the symmetric monoidal category $(\text{CW}, \times, \mathbf{1})$ of CW-complexes and CW-maps. We denote the interval $[0, 1]$ endowed with its usual CW-structure by I .

2.1 E_∞ -operads and E_∞ -props. We say that an operad \mathcal{O} is Σ -free if the action of Σ_m on $\mathcal{O}(m)$ is free for every m . A Σ -free resolution of an operad \mathcal{O} is an operad morphism from a Σ -free operad to \mathcal{O} inducing a homotopy equivalence in each arity m .

For any $X \in \text{CW}$, there are two types of representations of an operad \mathcal{O} on X . They are referred to as \mathcal{O} -coalgebra and \mathcal{O} -algebra structures and are respectively given by collections of CW-maps

$$\{\mathcal{O}(m) \times X \rightarrow X^m\}_{m \geq 0} \quad \text{and} \quad \{\mathcal{O}(m) \times X^m \rightarrow X\}_{m \geq 0}$$

satisfying associativity, equivariance, and unitality relations.

The terminal operad $\mathbf{1} = \{\mathbf{1}\}_{m \geq 0}$ is of particular importance. Its (co)algebras define usual (co)commutative, (co)associative, and (co)unital (co)algebra structures.

Following May [18], an operad \mathcal{O} is called an E_∞ -operad if it is a Σ -free resolution of the terminal operad and $\mathcal{O}(0) = \mathbf{1}$.

A **prop** is a strict symmetric monoidal category $\mathcal{P} = (\mathcal{P}, \odot, 0)$ enriched in CW generated by a single object. For any prop \mathcal{P} with generator p denote the CW-complex $\text{Hom}_{\mathcal{P}}(p^{\odot n}, p^{\odot m})$ by $\mathcal{P}(n, m)$. The symmetry of the monoidal structure induces commuting right and left actions

of Σ_n and Σ_m on $\mathcal{P}(n, m)$. Therefore, we think of the data of a prop as a Σ -**biobject**, i.e., a collection $\mathcal{P} = \{\mathcal{P}(n, m)\}_{n, m \geq 0}$ of CW-complexes with commuting actions of Σ_n and Σ_m , together with three types of maps

$$\begin{aligned} \circ_h : \mathcal{P}(n_1, m_1) \times \cdots \times \mathcal{P}(n_s, m_s) &\rightarrow \mathcal{P}(n_1 + \cdots + n_s, m_1 + \cdots + m_s), \\ \circ_v : \mathcal{P}(n, k) \times \mathcal{P}(k, m) &\rightarrow \mathcal{P}(n, m), \\ \eta : \mathbf{1} &\rightarrow \mathcal{P}(n, n). \end{aligned}$$

These types of maps are referred to respectively as **horizontal compositions**, **vertical compositions**, and **units**. They are derived respectively from the monoidal product, the categorical composition, and the identity morphisms of \mathcal{P} .

For any CW-complex X there are two types of representations of a prop \mathcal{P} on X . They are referred to as \mathcal{P} -**bialgebra** and **opposite \mathcal{P} -bialgebra** structures and are respectively given by collections of CW-maps

$$\{\mathcal{P}(n, m) \times X^n \rightarrow X^m\}_{n, m \geq 0} \quad \text{and} \quad \{\mathcal{P}(n, m) \times X^m \rightarrow X^n\}_{n, m \geq 0}.$$

satisfying associativity, equivariance, and unitality relations.

Let U be the functor from the category of props to that of operads given by naturally inducing from a prop \mathcal{P} an operad structure on the Σ -module $U(\mathcal{P}) = \{\mathcal{P}(1, m)\}_{m \geq 0}$. Notice that a \mathcal{P} -bialgebra (resp. opposite \mathcal{P} -algebra) structure on X induces a $U(\mathcal{P})$ -coalgebra (resp. $U(\mathcal{P})$ -algebra) structure on X .

Following Boardman and Vogt [2], a prop \mathcal{P} is called an E_∞ -**prop** if $U(\mathcal{P})$ is an E_∞ -operad.

2.2 Free props and presentations. As described for example in [7], the **free prop** $F(\mathcal{B})$ generated by a Σ -bimodule \mathcal{B} is constructed using open directed graphs with no directed loops that are enriched with a labeling described next. We think of each directed edge as built from two compatibly directed half-edges. For each vertex v of a directed graph G , we have the sets $in(v)$ and $out(v)$ of half-edges that are respectively incoming to and outgoing from v . Half-edges that do not belong to $in(v)$ or $out(v)$ for any v are divided into the disjoint sets $in(G)$ and $out(G)$ of incoming and outgoing external half-edges. For any positive integer n , let $\bar{n} = \{1, \dots, n\}$ and $\bar{0} = \emptyset$. For any finite set S , denote the cardinality of S by $|S|$. The labeling is given by bijections


$$\overline{|in(G)|} \rightarrow in(G) \quad \overline{|out(G)|} \rightarrow out(G)$$

and

$$\overline{|in(v)|} \rightarrow in(v) \quad \overline{|out(v)|} \rightarrow out(v)$$


for every vertex v . We refer to the isomorphism classes of such labeled directed graphs with no directed loops as (n, m) -**graphs**. We consider the right action of Σ_n and the left action of Σ_m on a (n, m) -graph given respectively by permuting the labels of $in(G)$ and $out(G)$.

The free prop $F(\mathcal{B})$ is given by all (n, m) -graphs which are \mathcal{B} -decorated in the following way. To every vertex v of one such G , one assigns an element $p \in \mathcal{B}(|in(v)|, |out(v)|)$ and introduces the equivalence relations:



where $p \in \mathcal{B}(n, m)$, $\sigma \in \Sigma_n$, and $\tau \in \Sigma_m$.

For any Σ -bimodule \mathcal{B} , the above construction defines the free prop $F(\mathcal{B})$ associated to \mathcal{B} . It satisfies the following universal property: Let $\iota : \mathcal{B} \rightarrow F(\mathcal{B})$ be the morphism sending an element $p \in \mathcal{B}(n, m)$ to the labeled and decorated (n, m) -corolla



For any Σ -bimodule map $\phi : \mathcal{B} \rightarrow \mathcal{P}$ where \mathcal{P} is a prop, there exists a unique prop morphism

$$F(\phi) : F(\mathcal{B}) \rightarrow \mathcal{P}$$

such that

$$\phi = F(\phi) \circ \iota.$$

Furthermore, there is a canonical isomorphism $F(F(\mathcal{B})) \rightarrow F(\mathcal{B})$ given by regarding graphs containing graphs as graphs.

Given any bisequence of spaces $\{B(n, m)\}_{n, m \geq 0}$ the free Σ -bimodule B^Σ is defined by

$$B^\Sigma(n, m) = \Sigma_m \times B(n, m) \times \Sigma_n$$

and satisfies the following universal property: Let $\xi : B \rightarrow B^\Sigma$ be the bisequence map that crosses with the identity elements in the corresponding symmetric groups. For any bisequence map $\phi : B \rightarrow \mathcal{B}$, there exists a unique Σ -bimodule map $\phi^\Sigma : B^\Sigma \rightarrow \mathcal{B}$ such that $\phi = \phi^\Sigma \circ \xi$.

We will now describe what is meant by a **presentation** (G, Φ, R) of a prop.

The first piece of data is a collection $G = \{G_d\}$ of bisequences with each $G_d(n, m)$ a disjoint union of spaces isomorphic to I^d . Each such space is called a **generating d -cell in biarity** (n, m) . We denote the bisequence containing their boundaries by ∂G_d and notice that $(\partial G_d)^\Sigma = \partial G_d^\Sigma$.

The second piece of data Φ are the **generating attaching maps**. These are morphisms of Σ -bimodules

$$\varphi_d : \partial G_d^\Sigma \rightarrow F(G_{d-1}^\Sigma).$$


Let X_0 be equal to $F(G_0^\Sigma)$ and for $d > 0$ let X_d be equal to the pushout

$$\begin{array}{ccc} F(\partial G_d^\Sigma) & \xrightarrow{F(\varphi_d)} & F(F(G_{d-1}^\Sigma)) \cong F(G_{d-1}^\Sigma) & \longrightarrow & X_{d-1} \\ \downarrow & & & & \downarrow \\ F(G_d^\Sigma) & \longrightarrow & & & X_d \end{array}$$

The limit of this process X is endowed with the induced prop structure.

The third piece of data is a bisequence R of subcomplexes of X called the **relations**. Denote by $\langle R \rangle$ the smallest sub- Σ -bimodule in X containing R and closed under compositions. We say that the triple (G, Φ, R) is a presentation of the prop $X/\langle R \rangle$.

2.3 Immersion convention. Graphs immersed in the plane will be used to represent labeled directed graphs with no directed loops, the convention we will follow is that the direction is given from top to bottom and the labeling from left to right. For example,



3. The prop $\tilde{\mathcal{S}}$

In this section we define the prop $\tilde{\mathcal{S}}$ via a finite presentation and show it is an E_∞ -prop.

Definition 3.1. Let $\tilde{\mathcal{S}}$ be the prop generated by

$$\downarrow \in \tilde{\mathcal{S}}_0(1, 0) \quad \wedge \in \tilde{\mathcal{S}}_0(1, 2) \quad \Upsilon^{1-s} \in \tilde{\mathcal{S}}_1(2, 1) \quad \dagger^s \in \tilde{\mathcal{S}}_1(1, 1)$$

with generating attaching maps

$$\Upsilon^0 = \downarrow \downarrow \quad \Upsilon^1 = \downarrow \downarrow \quad \text{and} \quad \dagger^1 = \wedge \quad \dagger^0 = |$$

and restricted by the relations

$$\wedge = | \quad \Upsilon^{1-s} = \downarrow \downarrow \quad \dagger^s = |.$$


Recall that $\mathbf{1}$ stands for the terminal CW-complex, i.e., a single 0-cell.

Lemma 3.2. *Let*


$$\bar{\mathbf{1}}(n, m) = \begin{cases} \mathbf{1} & \text{if } n > 0 \\ \emptyset & \text{if } n = 0 \end{cases}$$

endowed with the trivial prop structure. The unique map $\tilde{\mathcal{S}} \rightarrow \bar{\mathbf{1}}$ is a homotopy equivalence.


Proof. For $n = 0$, we notice that $\tilde{\mathcal{S}}(0, m) = \emptyset = \bar{\mathbf{1}}(0, m)$. For $n > 0$ and $m \geq 0$ we start by showing that the CW-complexes $\tilde{\mathcal{S}}(n, m)$ and $\tilde{\mathcal{S}}(n, m + 1)$ are homotopy equivalent. Consider the collection of maps $\{i : \tilde{\mathcal{S}}(n, m) \rightarrow \tilde{\mathcal{S}}(n, m + 1)\}$ described by the following diagram




Consider also the collection of maps $\{r : \tilde{\mathcal{S}}(n, m + 1) \rightarrow \tilde{\mathcal{S}}(n, m)\}$ described by




The diagram below shows that $r \circ i$ is homotopic to the identity




Let us compute diagrammatically the composition $i \circ r$



The composition $i \circ r$ is homotopic to the identity since



These computations show that i and r are homotopy inverses, and the relations imposed on $\tilde{\mathcal{S}}$ imply that $\tilde{\mathcal{S}}(n, 0)$ contains only the class of




We conclude that each $\tilde{\mathcal{S}}(n, m)$ is contractible for $n > 0$. □

Theorem 3.3. *The prop $\tilde{\mathcal{S}}$ is an E_∞ -prop.*

Proof. Since by construction the action of Σ_m on $U(\tilde{\mathcal{S}})(m) = \tilde{\mathcal{S}}(1, m)$ is free, the theorem follows from the previous lemma. □

Remark 3.4. Notice that the operad obtained by restricting to $\{S(n, 1)\}$ is not Σ -free. For example,



in $\tilde{\mathcal{S}}(3, 1)$ is fixed by the transposition (1, 2).

4. Cellular E_∞ -coalgebra on simplicial sets

In this section we derive from an $\tilde{\mathcal{S}}$ -bialgebra structure on the interval a natural $U(\tilde{\mathcal{S}})$ -coalgebra structures on the geometric realization of simplicial sets.

Definition 4.1. Let us denote the singleton $\{0\}$ by Δ^0 and the interval $[0, 1] \subset \mathbb{R}$ by Δ^1 or I . For $d \geq 1$ let

$$\Delta^d = \{(x_1, \dots, x_d) \in I^d \mid x_1 \leq \dots \leq x_d\}.$$

For $i = 0, \dots, d+1$ the **coface maps** $\delta_i : \Delta^d \rightarrow \Delta^{d+1}$ and **codegeneracy maps** $\sigma_i : \Delta^{d+1} \rightarrow \Delta^d$ are respectively defined by

$$\delta_i(x_1, \dots, x_d) = \begin{cases} (0, x_1, \dots, x_d) & i = 0, \\ (x_1, \dots, x_i, x_i, \dots, x_d) & 0 < i < d + 1, \\ (x_1, \dots, x_d, 1) & i = d + 1, \end{cases}$$

and

$$\sigma_i(x_1, \dots, x_{d+1}) = (x_1, \dots, \hat{x}_i, \dots, x_{d+1}).$$

We give the spaces Δ^d the coarser CW-structure making coface and codegeneracy maps into CW-maps. With respect to this CW-structure an element (x_1, \dots, x_d) belongs to the k -skeleton of Δ^d if and only if the cardinality of $\{x_i \mid x_i \neq 0, 1\}$ is less than or equal to k .

The **simplex category** is the subcategory of CW-complexes with objects Δ^d and morphisms generated by coface and codegeneracy maps.




Figure 1: The simplex Δ^3 drawn with thick lines.

Definition 4.2. For arbitrary $x, y \in \Delta^1$ and $s \in I$ define:

1. the diagonal approximation $\Delta : \mathbf{\Delta}^1 \rightarrow \mathbf{\Delta}^1 \times \mathbf{\Delta}^1$ by

$$\Delta(x) = \begin{cases} (0, 2x) & \text{if } x \leq 1/2, \\ (2x - 1, 1) & \text{if } x \geq 1/2, \end{cases}$$

2. the join $\psi : \mathbf{I} \times \mathbf{\Delta}^1 \times \mathbf{\Delta}^1 \rightarrow \mathbf{\Delta}^1$ by

$$\psi_s(x, y) = sx + (1 - s)y,$$

3. the counit homotopy $\phi : \mathbf{I} \times \mathbf{\Delta}^1 \rightarrow \mathbf{\Delta}^1$

$$\phi_s(x) = \begin{cases} \frac{2x}{2-s} & \text{if } x \leq \frac{2-s}{2}, \\ 1 & \text{if } x \geq \frac{2-s}{2}, \end{cases}$$

4. and the terminal map $\varepsilon : \mathbf{\Delta}^1 \rightarrow \{0\}$.

Lemma 4.3. *The maps given by*

$$\begin{aligned} \Phi(\lrcorner, (x_1, \dots, x_d)) &= ((\pi_1 \Delta(x_1), \dots, \pi_1 \Delta(x_d)), (\pi_2 \Delta(x_1), \dots, \pi_2 \Delta(x_d))), \\ \Phi(\lfloor, (x_1, \dots, x_d)) &= (\varepsilon(x_1), \dots, \varepsilon(x_d)), \\ \Phi(\overset{1-s}{\Upsilon}, (x_1, \dots, x_d), (y_1, \dots, y_d)) &= (\psi_s(x_1, y_1), \dots, \psi_s(x_d, y_d)), \\ \Phi(\overset{s}{\dagger}, (x_1, \dots, x_d)) &= (\phi_s(x_1), \dots, \phi_s(x_d)), \end{aligned}$$

define a $\tilde{\mathcal{S}}$ -bialgebra structure on $\mathbf{\Delta}^d$.

Proof. For any $s \in \mathbf{I}$ and $x \in \mathbf{\Delta}^1$ each of the functions

$$\pi_1 \Delta, \pi_2 \Delta, \psi_s(x, -), \psi(-, x), \phi_s : \mathbf{\Delta}^1 \rightarrow \mathbf{\Delta}^1$$

is order preserving, so the maps above are well defined. By counting the number of distinct coordinates of (x_1, \dots, x_d) that are not equal to 0 or 1 before and after applying the maps above we can verify they are cellular. To check these maps define a $\tilde{\mathcal{S}}$ -structure we need to verify they satisfy the identities coming from the attaching maps and relations on the generating cells of $\tilde{\mathcal{S}}$. In what follows we use the isomorphisms $\mathbf{\Delta}^0 \times \mathbf{\Delta}^d \cong \mathbf{\Delta}^d \cong \mathbf{\Delta}^d \times \mathbf{\Delta}^0$ with no further comment. For $\mathbf{x} = (x_1, \dots, x_n)$ and $\mathbf{y} = (y_1, \dots, y_n)$ we have

Attaching maps:

$$\begin{aligned} \Phi(\overset{0}{\Upsilon}, \mathbf{x}, \mathbf{y}) &= \mathbf{y} \cong \Phi(\lfloor \lfloor, \mathbf{x}, \mathbf{y}) \\ \Phi(\overset{1}{\Upsilon}, \mathbf{x}, \mathbf{y}) &= \mathbf{x} \cong \Phi(\lfloor \lfloor, \mathbf{x}, \mathbf{y}) \\ \Phi(\overset{0}{\dagger}, \mathbf{x}) &= \mathbf{x} = \Phi(\lfloor, \mathbf{x}) \\ \Phi(\overset{s}{\dagger}, \mathbf{x}) &= \left(\left(\begin{cases} 2x_1 & \text{if } x_1 \leq \frac{1}{2} \\ 1 & \text{if } x_1 \geq \frac{1}{2} \end{cases}, \dots, \begin{cases} 2x_d & \text{if } x_d \leq \frac{1}{2} \\ 1 & \text{if } x_d \geq \frac{1}{2} \end{cases} \right) \right) \cong \Phi(\lrcorner, \mathbf{x}). \end{aligned}$$

Relations:

$$\begin{aligned} \Phi(\lrcorner, \mathbf{x}) &\cong 0 \cong \Phi(\lfloor, \mathbf{x}) \\ \Phi(\overset{1-s}{\Upsilon}, \mathbf{x}, \mathbf{y}) &\cong 0 \cong \Phi(\lfloor \lfloor, \mathbf{x}, \mathbf{y}) \\ \Phi(\overset{s}{\dagger}, \mathbf{x}) &\cong 0 \cong \Phi(\lfloor, \mathbf{x}). \end{aligned}$$

Finally we need to verify naturality. Given the coordinate-wise nature of the $\tilde{\mathcal{S}}$ -bialgebra structure, naturality with respect to codegeneracy maps and coface maps $\delta_i : \Delta^d \rightarrow \Delta^{d+1}$ for $0 < i < d$ is immediate. Using

$$\begin{aligned} \Phi(\lrcorner, \delta_0 x) &= ((\pi_1 \Delta(0), \pi_1 \Delta(x)), (\pi_2 \Delta(0), \pi_2 \Delta(x))) \\ &= ((0, \pi_1 \Delta(x)), (0, \pi_2 \Delta(x))) \\ &= \delta_0(\pi_1 \Delta(x), \pi_2 \Delta(x)) \\ &= \delta_0 \Phi(\lrcorner, x) \end{aligned}$$

and

$$\begin{aligned} \Phi(\overset{1-s}{\Upsilon}, \delta_0 x, \delta_0 y) &= (\psi_s(0, 0), \psi_s(x, y)) & \Phi(\overset{s}{\Upsilon}, \delta_0 x) &= (\phi_s(0), \phi_s(x)) \\ &= (0, \Phi(\overset{1-s}{\Upsilon}, x, y)) & &= (0, \Phi(\overset{s}{\Upsilon}, x)) \\ &= \delta_0 \Phi(\overset{1-s}{\Upsilon}, x, y) & &= \delta_0 \Phi(\overset{s}{\Upsilon}, x) \end{aligned}$$

we can deduce the naturality of δ_0 . The naturality of δ_d is derived analogously. □

Definition 4.4. A **simplicial set** Γ is a contravariant functor from the simplex category to the category of sets. Denote $\Gamma(\Delta^d)$ by Γ_d . Its **geometric realization** is the CW-complex

$$|\Gamma| = \coprod_{d \geq 0} \Gamma_d \times \Delta^d / \sim$$

where $\tau^*(\gamma) \times \mathbf{x} \sim a \times \tau(\mathbf{x})$ for any $\tau \in \text{Hom}(\Delta^d, \Delta^e)$, $\gamma \in \Gamma_e$ and $\mathbf{x} \in \Delta^d$.

Theorem 4.5. Let Γ be a simplicial set. A natural $U(\tilde{\mathcal{S}})$ -coalgebra structure is defined on its geometric realization by

$$U(\Phi)(g, (\gamma, \mathbf{x})) = ((\gamma, \pi_1 \Phi(g, \mathbf{x})), \dots, (\gamma, \pi_m \Phi(g, \mathbf{x})))$$

where $g \in U(\tilde{\mathcal{S}})(m)$, $\gamma \in \Gamma_d$ and $\mathbf{x} \in \Delta^d$.

Proof. To verify $U(\Phi) : U(\tilde{\mathcal{S}})(m) \times |\Gamma| \rightarrow |\Gamma|^m$ is a well defined map, consider two representatives $\tau^*(\gamma) \times \mathbf{x}$ and $\gamma \times \tau(\mathbf{x})$. We have

$$(\tau^*(\gamma), \pi_1 \Phi(g, \mathbf{x}), \dots, \tau^*(\gamma), \pi_m \Phi(g, \mathbf{x}))$$

is equivalent in $|\Gamma|^m$ to

$$(\gamma, \tau(\pi_1 \Phi(g, \mathbf{x})), \dots, \gamma, \tau(\pi_m \Phi(g, \mathbf{x})))$$

which equals

$$(\gamma, \pi_1 \Phi(g, \tau(\mathbf{x})), \dots, \gamma, \pi_m \Phi(g, \tau(\mathbf{x})))$$

since, by naturality,

$$\tau(\pi_i \Phi(g, \mathbf{x})) = \pi_i \tau^m \Phi(g, \mathbf{x}) = \pi_i \Phi(g, \tau(\mathbf{x})).$$

The equivariance of $U(\Phi)$ and the identity

$$U(\Phi)(h, U(\Phi)(g, (\gamma, \mathbf{x}))) = U(\Phi)(h \circ g, (\gamma, \mathbf{x}))$$

follow from those satisfied by Φ . □

Remark 4.6. It is not the case that $|\Gamma|$ carries an $\tilde{\mathcal{S}}$ -bialgebra structure. For example, if $|\Gamma| = \{x, y\}$ then $\Phi(\overset{1-s}{\underset{s}{\Upsilon}}, x, y)$ is not well defined.

Remark 4.7. For any $d \geq 0$ and $j \geq 0$ let $[j] = ([j]_1, \dots, [j]_d) \in \mathbf{\Delta}^d$ be given by

$$[j]_k = \begin{cases} 0 & k \leq d - j \\ 1 & k > d - j. \end{cases}$$

For any finite set of integers $\{v_0, \dots, v_k\}$, denote the convex closure of $[v_0], \dots, [v_k]$ by $[v_0, \dots, v_k]$. These subsets correspond to the cells of $\mathbf{\Delta}^d$ and we have

$$\begin{aligned} \Phi(\lrcorner, [v_0, \dots, v_k]) &= \prod_{i=0}^d [v_0, \dots, v_i] \times [v_i, \dots, v_k] \\ \Phi(\llcorner, [v_0, \dots, v_k]) &= [0] \\ \prod_{s \in \mathbf{I}} \Phi(\overset{1-s}{\underset{s}{\Upsilon}}, [v_0, \dots, v_j], [v_{j+1}, \dots, v_k]) &= [v_0, \dots, v_j, v_{j+1}, \dots, v_k] \\ \prod_{s \in \mathbf{I}} \Phi(\overset{s}{\Upsilon}, [v_0, \dots, v_k]) &= [v_0, \dots, v_k]. \end{aligned}$$

Remark 4.8. We conjecture that a natural cellular action of $U(\tilde{\mathcal{S}})$ can be defined on the geometric realization of cubical sets with or without connections. For cubical chains, an action of the prop obtained by applying cellular chains to $U(\tilde{\mathcal{S}})$ is defined in [15].

5. The prop \mathcal{MS}

In this section we introduce the finitely presented E_∞ -prop \mathcal{MS} . We provide a description of \mathcal{MS} in terms of oriented surfaces with a weighted 1-skeleton. We show that the operad associated to \mathcal{MS} is isomorphic to Kaufmann’s Arc Surface model for the E_∞ -operad [11], and that its cellular chains are isomorphic, up to signs, to the Surjection operad [19, 1]. These two identifications combine to verify a conjecture of Kaufmann.

5.1 Edge-weights and definition

Definition 5.1. Let \mathcal{S} be the prop generated by

$$\lrcorner \in \mathcal{S}(1, 0)_0 \quad \lrcorner \in \mathcal{S}(1, 2)_0 \quad \overset{1-s}{\underset{s}{\Upsilon}} \in \mathcal{S}(2, 1)_1$$

with generating attaching maps

$$\overset{1}{\underset{0}{\Upsilon}} = \lrcorner \lrcorner \quad \overset{0}{\underset{1}{\Upsilon}} = \lrcorner \lrcorner$$

and restricted by the relations


$$\lrcorner = \lrcorner = \lrcorner \quad \overset{1-s}{\underset{s}{\Upsilon}} = \lrcorner \lrcorner.$$

Remark 5.2. This prop is a strictly counital version of $\tilde{\mathcal{S}}$ and it receives a quotient map from it. We notice that a simplified version of the proof given for Lemma 3.2 shows this is an E_∞ -prop.


We define alternative coordinates on the prop \mathcal{S} . Consider a d -cell in $\mathcal{S}(n, m)$ and an (n, m) -graph G supporting it. The **edge-weight coordinates** of this cell are given by the assignment of a non-negative real number to each edge of G , referred to as its **weight**, satisfying the following conditions:¹

1. Edges of the form \downarrow have weight 0.
2. Edges in $out(G)$ have weight 1.
3. For every vertex v of G , the sum of the edge-weights in $in(v)$ and in $out(v)$ are the same.


Edge-weight coordinates are well defined as we can see from:




We can pass from the original coordinates induced from Υ to edge-weight coordinates via the following inductive procedure: Edges containing \downarrow are set to have weights 0. Edges in $out(G)$ are set to have weight 1. The other edges get their weights from




The passage from edge-weight coordinates to the original coordinates is induced by



Definition 5.3. Let \mathcal{MS} be the quotient of \mathcal{S} by the involutive, coassociative, associative, commutative and Leibniz relations




and




depending respectively on if $a_1 > b_1$, $a_1 = b_1$ or $a_1 < b_1$.

Remark 5.4. We can express the Leibniz relation in the following alternative way. Consider




¹We say an edge belongs to $in(G)$, $out(G)$, $in(v)$ or $out(v)$ if one of its two half-edges does.


with $a_1 + a_2 = b_1 + b_2$. In \mathbb{R}^2 consider the rectangle with opposite vertices at coordinates $(0, 0)$ and $(b_1 + b_2, 3)$. Cut along the lines joining $(b_1, 0)$ with $(b_1, 2)$ and $(a_1, 3)$ with $(a_1, 1)$. Deformation retract keeping the vertical coordinate invariant to a $(2, 2)$ -graph with labelings induced from the plane. Give this $(2, 2)$ -graph the edge-weight coordinates coming from the width of their corresponding sub-rectangle. For example, if $a_1 > b_1$ we have




Notation 5.5. We will utilize the following diagrammatic simplification



to represent labeled directed graphs resulting from iterated grafting of the product and coproduct in the left comb order



Definition 5.6. A canonical (n, m) -graph is an (n, m) -graph of the form



containing no internal vertices or copies of either \downarrow or \diamond and such that for each $i = 1, \dots, m$ the induced map

$$\{1, \dots, k_i\} \rightarrow \bigsqcup \{1, \dots, r_1\} < \dots < \{1, \dots, r_n\}$$

is order preserving.

Definition 5.7. Given an element γ in $\mathcal{MS}(n, m)$, thought of as an equivalence of weighted (n, m) -graphs, we say that an (n, m) -graph Γ **supports** it if Γ is equal to a representative of γ after forgetting its weights.

Lemma 5.8. For every element in $\mathcal{MS}(n, m)$ with $m > 0$ there exists a unique canonical graph supporting it.


Proof. Consider an element in $\mathcal{MS}(n, m)$ and an (n, m) -graph supporting it. We start by getting rid of all occurrences of \downarrow . Consider one such strand and follow it up until hitting a vertex, which

we must since $m > 0$. If the vertex we encounter is in a subgraph of the form Υ we can replace this with \Downarrow and continue the excursion up along one of the strands. If alternatively we encounter a vertex contained in a subgraph of one of the following forms \swarrow or \searrow we can replace this with \downarrow and choose another strand \downarrow to repeat the process.


We have constructed a (n, m) -graph with no copies of \downarrow supporting our element. We now use the Leibniz relation to ensure that with respect to the direction of the (n, m) -graph all vertices belonging to a subgraph of the form \swarrow appear before vertices belonging to subgraphs of the form Υ .

Now we now use coassociativity and associativity to enforce the left comb convention. Using commutativity we reorder the strands of each iterated graftings of Υ so that the order preserving condition is satisfied. We then scan the supporting graph and replace each copy of \diamond by a copy of \downarrow . This construction produces a canonical graph supporting our element.

In order to show the uniqueness of such canonical graph, we need to prove that the order in which we performed the replacements above does not matter, in the terminology of Gröbner bases [6, 17], that all critical monomials are confluent. For example,



and



The other compositions are verified similarly. □

5.2 Surface realization of \mathcal{MS} . For any element in $\mathcal{MS}(n, m)$ with $m > 0$ we faithfully associate an oriented surface equipped with a CW-structure having a weighted 1-skeleton.

Construction 5.9. Consider an element in $\mathcal{MS}(n, m)$ with $m > 0$ and the canonical (n, m) -graph supporting it as constructed in Lemma 5.8. By compactifying the open edges of the graph, we introduce $n + m$ new vertices. We glue to each of them both endpoints of an interval and call the resulting $n + m$ circles the incoming and outgoing boundary circles depending on the direction of the (n, m) -graph. We make this graph into a ribbon graph, i.e. give each vertex a




Figure 2: Illustrating Construction 5.9 with omitted weights and labelings

cyclic order of its incident edges as follows: For the new $n + m$ vertices choose any cyclic order and for all others chose the natural extension of the total order induced from the labeling.

Consider the surface associated to this ribbon graph.² Remove from it the disks attached to the boundary circles and collapse edges satisfying the following conditions: 1) one and only one of their endpoint is in a boundary circle and 2) no other edge incident to their interior endpoint has the same relative direction (towards or away from the vertex).

We refer to the directed and weighted 1-cells of the resulting CW-surface as **arcs** and notice the original element in \mathcal{MS} can be recovered from them.

Definition 5.10. The prop \mathcal{A} is defined by pushing forward the CW and prop structures from \mathcal{MS} to the image of Construction 5.9.

Remark 5.11. Notice that $\mathcal{A}(n, 0) = \emptyset$ for every n . Also, for a family of element in \mathcal{A} parametrized by the weight of an arc tending to zero, we see that the limit will remove the arc and the topology of the surface will possibly change.

5.3 Relations to earlier work. The reader familiar with [13] will recognize the elements of \mathcal{A} as examples of Arc Surfaces. We make the connection more precise with the following


Proposition 5.12. *The operad $U(\mathcal{A})$ is isomorphic to $St\mathcal{LGT}ree^1(m)$ as defined by Kaufmann in [11].*

Proof. Comparing with Definition 2.4 in [11] and section 4.1 in the same reference, we notice that any element in $U(\mathcal{A})(m)$ corresponds a to quasi-filling element in $\mathcal{LGT}ree(m)^1$ and that any such element arises this way. In the same reference, Corollary 2.2.7 states that any element in $St\mathcal{LGT}ree(m)^1$ corresponds to a unique quasi-filling element in $\mathcal{LGT}ree(m)^1$, so we have a bijection between $\mathcal{A}(m)$ and $St\mathcal{LGT}ree(m)^1$.

Comparing the stabilization process introduced in Definition 2.24 of [11] with Remark 5.11 makes this bijection into a cellular isomorphism.

We can use Remark 5.4 to describe the composition in $U(\mathcal{MS})$ and \mathcal{A} in terms of vertically invariant deformation retractions of cut rectangles:

²The surface associated to a ribbon graph is constructed by attaching a disk to each ribbon loop. A ribbon loop can be described as follows: Choose an edge of the ribbon graph and a direction for that edge. Select from the edges incident to the forward vertex v the one that follows directly after our original edge in the cyclic order associated to v . We provide this second edge with the direction that has v as its backward vertex and repeat this process until returning to our original edge in the original direction.



where $a_1 + \dots + a_q = 1$ and $b_1 + \dots + b_q = b$. This allows us to recognize the vertical composition in \mathcal{A} as that of the Arc Surface props [13]. See for example Section 1.2.2. in [11] for the definition of this composition. In particular, this shows the correspondence of the operadic compositions of $U(\mathcal{A})$ and $\{St\mathcal{LGT}ree^1(m)\}_{m \geq 1}$. \square


The prop $C_\bullet(\mathcal{MS})$ in the category of differential graded modules, resulting from applying the functor of cellular chains to the prop \mathcal{MS} , inherits a finite presentation with generators

$$\downarrow \in C_0(\mathcal{MS})(1, 0) \quad \wedge \in C_0(\mathcal{MS})(1, 2) \quad \Upsilon \in C_1(\mathcal{MS})(2, 1)$$

differential

$$\partial \downarrow = 0 \quad \partial \wedge = 0 \quad \partial \Upsilon = \downarrow \downarrow$$

and relations



Lemma 17 of [24] shows that the operad associated to $C_\bullet(\mathcal{MS})$ is isomorphic up to signs to the Surjection operad [19, 1]. We therefore have the following corollary to Proposition 5.12 which was conjectured by Kaufmann in 4.4 of [11].

Corollary 5.13. *The operad obtained by applying the cellular chains to $St\mathcal{LGT}ree^1$ is up to signs isomorphic to the Surjection operad.*

Remark 5.14. For the E_2 -suboperad, this was independently established in [16].

Let us now return to E_∞ -structures on simplicial sets. Consider the $\tilde{\mathcal{S}}$ -bialgebra structure on the standard simplices as described in Lemma 4.3. Applying the functor of cellular chains, we obtain a natural prop morphism

$$C_\bullet(\tilde{\mathcal{S}}) \rightarrow \text{End}(C_\bullet(\Delta^d)) \tag{1}$$

where $\text{End}(C_\bullet(\Delta^d))(n, m) = \text{Hom}(C_\bullet(\Delta^d)^{\otimes n}, C_\bullet(\Delta^d)^{\otimes m})$.

As can be seen from Remark 4.7, this map sends the generators of $C_\bullet(\tilde{\mathcal{S}})$ to the following functions, which we describe up to signs:

$$\begin{aligned} \wedge [v_0, \dots, v_q] &= \sum_{i=0}^q [v_0, \dots, v_i] \otimes [v_i, \dots, v_q], \\ \downarrow [v_0, \dots, v_q] &= \begin{cases} 1 & \text{if } q = 0 \\ 0 & \text{if } q > 0, \end{cases} \\ \Upsilon ([v_0, \dots, v_p] \otimes [v_{p+1}, \dots, v_q]) &= \begin{cases} [v_{\pi(0)}, \dots, v_{\pi(q)}] & \text{if } i \neq j \text{ implies } v_i \neq v_j \\ 0 & \text{if not,} \end{cases} \\ \dagger [v_0, \dots, v_q] &= 0. \end{aligned} \tag{2}$$

The fact that \dagger acts trivially serves as a motivation for considering the cellular chains on \mathcal{S} as introduced in Definition 5.1. We have an algebraic presentation of it with generators

$$\downarrow \in C_\bullet(\mathcal{S})(1, 0)_0 \quad \wedge \in C_\bullet(\mathcal{S})(1, 2)_0 \quad \Upsilon \in C_\bullet(\mathcal{S})(2, 1)_1$$

differential

$$\partial \downarrow = 0 \quad \partial \wedge = 0 \quad \partial \Upsilon = \downarrow | - | \downarrow$$

and relations

$$\Upsilon \quad \wedge - | \quad | - \wedge.$$

We can verify that these relations are satisfied by the assignments in (2), so we have a factorization

$$\begin{array}{ccc} C_\bullet(\tilde{\mathcal{S}}) & \longrightarrow & \text{End}(C_\bullet(\Delta^d)). \\ \downarrow & \nearrow & \\ C_\bullet(\mathcal{S}) & & \end{array}$$

The above morphism $C_\bullet(\mathcal{S}) \rightarrow \text{End}(C_\bullet(\Delta^d))$ was introduced and studied in [24] where it was related to the E_∞ -structure defined by McClure-Smith and Berger-Fresse [19, 1].

Remark 5.15. The restriction to biarity (1, 2) of this E_∞ -bialgebra induces Steenrod’s cup- i products [27, 26]. In [20], an axiomatic characterization of these products was given, and in [22], they were used to derive the nerve construction of higher categories as defined in [28]. In the present paper, we were able to obtain Steenrod’s cup- i products naturally from only four maps associated to the interval. We see this as further evidence of the fundamental nature of Steenrod’s cup- i products. See [21] for algorithms using them to incorporate cohomology operations into topological data analysis [4, 29], and [23, 3] for their use constructing cochains enforcing the Cartan and Adem relations at the cochain level. Higher arity products, analogous to the cup- i products, defining Steenrod operations at odd primes are defined in [14] and implemented in [25].

Acknowledgments. We would like to thank Dennis Sullivan, Bruno Vallette, Ralph Cohen, Stephan Stolz, Ralph Kaufmann, Kathryn Hess, Manuel Rivera and the anonymous referee for their insights, questions, and comments about this project. This work was partially supported by Innosuisse grant 32875.1 IP-ICT - 1.

References

- [1] Clemens Berger and Benoit Fresse. Combinatorial operad actions on cochains. *Mathematical Proceedings of the Cambridge Philosophical Society*, 137(1):135–174, 2004.
- [2] John Michael Boardman and Rainer M Vogt. *Homotopy invariant algebraic structures on topological spaces*, volume 347. Springer, 2006.
- [3] Greg Brumfiel, Anibal M. Medina-Mardones, and John Morgan. A cochain level proof of Adem relations in the mod 2 Steenrod algebra. [ArXiv:2006.09354](https://arxiv.org/abs/2006.09354), 2020.
- [4] Gunnar Carlsson. Topology and data. *Bull. Amer. Math. Soc. (N.S.)*, 46(2):255–308, 2009.
- [5] Moira Chas and Dennis Sullivan. String topology. [ArXiv:math/9911159](https://arxiv.org/abs/math/9911159), 1999.

- [6] Vladimir Dotsenko and Anton Khoroshkin. Gröbner bases for operads. *Duke Mathematical Journal*, 153(2):363–396, 2010.
- [7] Benoit Fresse. Props in model categories and homotopy invariance of structures. *Georgian Mathematical Journal*, 17(1):79–160, 2010.
- [8] Mark Goresky and Nancy Hingston. Loop products and closed geodesics. *Duke Math. J.*, 150(1):117–209, 2009.
- [9] Ralph M. Kaufmann. Moduli space actions on the Hochschild co-chains of a Frobenius algebra. II. Correlators. *J. Noncommut. Geom.*, 2(3):283–332, 2008.
- [10] Ralph M. Kaufmann. Noncommutative aspects of open/closed strings via foliations. *Rep. Math. Phys.*, 61(2):281–293, 2008.
- [11] Ralph M. Kaufmann. Dimension vs. genus: a surface realization of the little k -cubes and an E_∞ -operad. In *Algebraic topology—old and new*, volume 85 of *Banach Center Publ.*, pages 241–274. Polish Acad. Sci. Inst. Math., Warsaw, 2009.
- [12] Ralph M Kaufmann. A detailed look on actions on Hochschild complexes especially the degree 1 co-product and actions on loop spaces. [ArXiv:1807.10534](https://arxiv.org/abs/1807.10534), 2018.
- [13] Ralph M. Kaufmann, Muriel Livernet, and R. C. Penner. Arc operads and arc algebras. *Geom. Topol.*, 7:511–568, 2003.
- [14] Ralph M Kaufmann and Anibal M. Medina-Mardones. Cochain level May-Steenrod operations. [ArXiv:2010.02571](https://arxiv.org/abs/2010.02571), 2020.
- [15] Ralph M Kaufmann and Anibal M. Medina-Mardones. A combinatorial E_∞ algebra structure on cubical cochains. [ArXiv:2107.00669](https://arxiv.org/abs/2107.00669), 2021.
- [16] Ralph M. Kaufmann and Yongheng Zhang. Permutohedral structures on E_2 -operads. *Forum Math.*, 29(6):1371–1411, 2017.
- [17] Jean-Louis Loday and Bruno Vallette. *Algebraic operads*, volume 346 of *Grundlehren der Mathematischen Wissenschaften*. Springer, Heidelberg, 2012.
- [18] J Peter May. *The geometry of iterated loop spaces*, volume 271. Springer, 2006.
- [19] James McClure and Jeffrey Smith. Multivariable cochain operations and little n -cubes. *Journal of the American Mathematical Society*, 16(3):681–704, 2003.
- [20] Anibal M. Medina-Mardones. An axiomatic characterization of Steenrod’s cup- i products. [ArXiv:1810.06505](https://arxiv.org/abs/1810.06505), 2018.
- [21] Anibal M. Medina-Mardones. Persistence Steenrod modules. [ArXiv:1812.05031](https://arxiv.org/abs/1812.05031), 2018.
- [22] Anibal M. Medina-Mardones. An algebraic representation of globular sets. *Homology Homotopy Appl.*, 22(2):135–150, 2020.
- [23] Anibal M. Medina-Mardones. An effective proof of the Cartan formula: the even prime. *J. Pure Appl. Algebra*, 224(12):106444, 18, 2020.

- [24] Anibal M. Medina-Mardones. A finitely presented E_∞ -prop I: algebraic context. *High. Struct.*, 4(2):1–21, 2020.
- [25] Anibal M. Medina-Mardones. A computer algebra system for the study of commutativity up-to-coherent homotopies. [ArXiv:2102.07670](https://arxiv.org/abs/2102.07670), 2021.
- [26] Anibal M. Medina-Mardones. New formulas for cup- i products and fast computation of Steenrod squares. [ArXiv:2105.08025](https://arxiv.org/abs/2105.08025), 2021.
- [27] Norman E Steenrod. Products of cocycles and extensions of mappings. *Annals of Mathematics*, pages 290–320, 1947.
- [28] Ross Street. The algebra of oriented simplexes. *Journal of Pure and Applied Algebra*, 49(3):283–335, 1987.
- [29] Guillaume Tauzin, Umberto Lupo, Lewis Tunstall, Julian Burella Pérez, Matteo Caorsi, Anibal M. Medina-Mardones, Alberto Dassatti, and Kathryn Hess. giotto-tda: A topological data analysis toolkit for machine learning and data exploration, 2020.
- [30] Thomas Tradler and Mahmoud Zeinalian. Algebraic string operations. *K-Theory*, 38(1):59–82, 2007.