

U. PORTO

FMUP FACULDADE DE MEDICINA
UNIVERSIDADE DO PORTO

MESTRADO INTEGRADO EM MEDICINA

2020/2021

Miguel Leonardo Costa dos Santos

**Reverse shoulder Arthroplasty with (or
without) subscapularis repair: does it affect
range of motion and/or articular stability?**

Março, 2021

FMUP

Miguel Leonardo Costa dos Santos

**Reverse shoulder Arthroplasty with (or
without) subscapularis repair: does it affect
range of motion and/or articular stability?**

Mestrado Integrado em Medicina

Área: Ortopedia

Tipologia: Monografia

Trabalho efetuado sob a Orientação de:

Professor Doutor Manuel Gutierres

E sob a Coorientação de:

Doutor Bernardo Nunes e Professora Doutora Cláudia Dias

Trabalho organizado de acordo com as normas da revista:
EFORT Open Reviews - official journal of the European Federation
of National Associations of Orthopaedics and Traumatology

Março, 2021

FMUP

UC DISSERTAÇÃO/PROJETO (6º ANO) - DECLARAÇÃO DE INTEGRIDADE

Eu, **Miguel Leonardo Costa dos Santos**, abaixo-assinado, nº mecanográfico **201703011**, estudante do 6º ano do Ciclo de Estudos Integrado em Medicina, na Faculdade de Medicina da Universidade do Porto, declaro ter atuado com absoluta integridade na elaboração deste projeto de opção.

Neste sentido, confirmo que **NÃO** incorri em plágio (ato pelo qual um indivíduo, mesmo por omissão, assume a autoria de um determinado trabalho intelectual, ou partes dele). Mais declaro que todas as frases que retirei de trabalhos anteriores pertencentes a outros autores, foram referenciadas, ou redigidas com novas palavras, tendo colocado, neste caso, a citação da fonte bibliográfica.

Faculdade de Medicina da Universidade do Porto, 27/03/2021

Assinatura conforme cartão de identificação:

Miguel Leonardo Costa dos Santos

NOME

Miguel Leonardo Costa dos Santos

NÚMERO DE ESTUDANTE

201703011

E-MAIL

Lcosta_santos@hotmail.com

DESIGNAÇÃO DA ÁREA DO PROJECTO.

Ortopedia

TÍTULO MONOGRAFIA

Reverse Shoulder Arthroplasty with (or without) subscapularis repair: does it affect. range of motion and/or articular stability?

ORIENTADOR

Professor Doutor Manuel António Pereira Gutierres

COORIENTADORES

Doutor João Bernardo Matos Nunes e Professora Doutora. Cláudia Camila Dias

ASSINALE APENAS UMA DAS OPÇÕES:

É AUTORIZADA A REPRODUÇÃO INTEGRAL DESTA TRABALHO APENAS PARA EFEITOS DE INVESTIGAÇÃO, MEDIANTE DECLARAÇÃO ESCRITA DO INTERESSADO, QUE A TAL SE COMPROMETE.	<input checked="" type="checkbox"/>
É AUTORIZADA A REPRODUÇÃO PARCIAL DESTA TRABALHO (INDICAR, CASO TAL SEJA NECESSÁRIO, Nº MÁXIMO DE PÁGINAS, ILUSTRAÇÕES, GRÁFICOS, ETC.) APENAS PARA EFEITOS DE INVESTIGAÇÃO, MEDIANTE DECLARAÇÃO ESCRITA DO INTERESSADO, QUE A TAL SE COMPROMETE.	<input type="checkbox"/>
DE ACORDO COM A LEGISLAÇÃO EM VIGOR, (INDICAR, CASO TAL SEJA NECESSÁRIO, Nº MÁXIMO DE PÁGINAS, ILUSTRAÇÕES, GRÁFICOS, ETC.) NÃO É PERMITIDA A REPRODUÇÃO DE QUALQUER PARTE DESTA TRABALHO.	<input type="checkbox"/>

Faculdade de Medicina da Universidade do Porto, 27/03/2021

Assinatura conforme cartão de identificação: Miguel Leonardo Costa Santos

“A alegria está na luta, na tentativa, no sofrimento envolvido e não na vitória propriamente dita.”

Mahatma Gandhi

AGRADECIMENTOS

Ao professor Doutor Manuel Gutierres, por todo o incentivo e apoio na exploração desta temática, e por despertar ainda mais o gosto pela Ortopedia.

Ao Doutor Bernardo Nunes por todo o apoio e disponibilidade na elaboração desta monografia.

À professora Doutora Cláudia Dias pela disponibilidade demonstrada e apoio no estudo deste tema.

À minha esposa, Sara, por todo o apoio e disponibilidade que me permitiu ter tempo para a realização desta monografia e conclusão deste curso. A ela devo o incentivo, força e coragem para iniciar e terminar este percurso!

Aos meus dois filhos, Francisco e Henrique, por todo o carinho e amor que me transmitem nos momentos mais difíceis deste longo, mas enriquecedor percurso. A eles também peço desculpa por não ter estado mais presente.

Aos meus pais por todo o apoio e disponibilidade.

Aos meus amigos, por estarem presentes quando foi preciso.

A todos, muito obrigado!

Reverse Shoulder Arthroplasty with (or without) subscapularis repair: does it affect. range of motion and/or articular stability?

Artroplastia invertida do ombro com (ou sem) reparação do subescapular: Afeta a amplitude de movimento e/ou estabilidade articular?

Miguel Costa Santos¹, Cláudia Camila Dias^{1,2}, João Bernardo Nunes^{3,4}, Manuel Gutierrez^{1,3,4}

1 – Faculdade de Medicina da Universidade do Porto

2 - Departamento de Medicina da Comunidade, Informação e Decisão em Saúde da Faculdade de Medicina da Universidade do Porto.

3 – Centro Hospitalar Universitário de São João, EPE

4 – Departamento de Cirurgia da Faculdade de Medicina da Universidade do Porto

Correspondant Author: Miguel Leonardo Costa Santos, up201703011@edu.med.up.pt

Miguel Leonardo Costa dos Santos
Lcosta_santos@hotmail.com

Reverse Shoulder Arthroplasty with (or without) subscapularis repair: does it affect range of motion and/or articular stability?

ABSTRACT

Background: The aim of this bibliographic review is to assess the results between the repair and non-repair of the subscapularis tendon in reverse shoulder arthroplasty, regarding range of motion and joint stability.

Methods: Literature search performed in PUBMED using the QUERY “Reverse Shoulder arthroplasty” AND “subscapularis repair”, limited to January 1st, 2015 until 16 January 16th, 2021. The inclusion criteria were studies with adult patients submitted to reverse shoulder arthroplasty, comparing subscapularis repair or non-repair, and reporting range of motion and stability as an outcome.

Results: Most of the studies did not find significant differences between the repair and the non-repair of the subscapularis for most of the assessed outcomes. However, lateralization or medialization of the centre of rotation of the prosthesis was identified as an important variable when not repairing the subscapularis, where lateralized prostheses showed a decreased dislocation rate.

Conclusion: Most studies did not find significant advantages in repairing the subscapularis when considering range of motion. When the subscapularis is not repaired, lateralized prostheses may provide lower dislocations rates. There is not yet sufficient evidence to demonstrate the indication or contraindication for repair of the subscapular tendon.

Key words: Reverse Shoulder Arthroplasty, Subscapularis Repair, Stability, Range Of Motion.

INTRODUCTION

Reverse shoulder arthroplasty was initially proposed for the treatment of end-stage rotator cuff arthropathy(1,2). Expanding indications have gradually been suggested in primary and secondary arthritis(2,3), fractures of the proximal humerus and its sequelae(2–5), tumoral resection(2), massive rotator cuff tears(2–4,6,7), other painful cuff arthropathies(3,5,8) or arthroplasty revision(2–4). Good to excellent functional outcomes have been reported(2), especially regarding active elevation(2), external rotation(2), and abduction(2). Nevertheless, some complications can occur, such as neurologic injury, dislocation, periprosthetic fracture, hematoma, infection, scapular notching, mechanical baseplate failure, and acromial fracture(3,4,9–11).

Despite this, concerns have been raised regarding limited range of motion after reverse shoulder arthroplasty(12,13). Studies have demonstrated that possible causes may be related to impingement(14), prostheses design(5), medialization of the glenosphere or the humerus(5) and management of the subscapularis tendon(15,16). A recent study has demonstrated better outcomes regarding range of motion with the lateralization of the centre of rotation of the prosthesis. Inferior placement of the glenosphere and a varus humeral neck shaft were also shown to reduce the adduction deficit and impingement and thus improving range of motion(5,16,17).

The subscapularis is an active internal rotator of the humeral head(15,16,18) and its tendon is released during the deltopectoral approach (tenotomy or osteotomy)(8,15,16).

In the literature, systematic reviews have approached this subject focusing on different clinical outcomes, such as dislocation rates(19) or functional scores(19,20). Notwithstanding, there is still limited evidence regarding the effect of subscapularis management during reverse shoulder arthroplasty (repair or no-repair) in shoulder range of motion – specifically internal rotation.

The objective of this study is to perform a bibliographic review of studies comparing the repair of the subscapularis with the non-repair in reverse shoulder arthroplasty focusing on outcomes related to range of motion and stability.

METHODS

A literature search was performed in PUBMED using the QUERY “Reverse Shoulder arthroplasty” AND “subscapularis repair”, limited to January 1st, 2015 until January 16th, 2021. 140 articles were found, and a full-text reading was performed. Articles that fulfilled the following inclusion criteria were selected: adult patients submitted to reverse shoulder arthroplasty and comparison of subscapularis repair or non-repair and reporting range of motion as an outcome (active forward elevation, external rotation, internal rotation, abduction). After the application of the inclusion and exclusion criteria and reading of the articles, 26 articles were considered relevant to which reference will be made in the following paragraphs, throughout this review Joint stability was also analysed in this literature review.

SUBSCAPULARIS MUSCLE AND TENDON

The subscapularis muscle has important actions in the shoulder. First, it provides internal rotation(4,5,18,19,21) but it is also very important in abduction, adduction, extension, and flexion of the shoulder(4,5,18,21,22). It provides different actions according to the activated portion: the upper two thirds on the tendon – the tendinous portion – are responsible for internal rotation and abduction(18,22). This portion is innervated by the superior subscapular nerve, derived from the brachial plexus 8(receiving fibres from the fifth cervical nerve, C5)(18,21). When the lower third contracts – the muscular portion – adduction occurs(18,22). The lower portion is innervated by the inferior subscapular nerve, also derived from the brachial plexus (receiving fibres from the sixth cervical nerve, C6)(18,21). The stability of the

Miguel Leonardo Costa dos Santos
Lcosta_santos@hotmail.com

glenohumeral joint depends also on the subscapularis tendon forces, together with the posterior rotator forces(22).

“TO REPAIR OR NOT TO REPAIR? THAT IS THE QUESTION!”

Subscapularis tendon management is a source of controversy in reverse shoulder arthroplasty. Some authors suggested that the repair of the subscapularis may improve reverse shoulder arthroplasty stability(15,16), and range-of-motion(15,16) while decreasing joint dead space(15). Nevertheless, other studies did not find significant differences in these outcomes when the subscapularis was not repaired(22) and some authors even suggest that subscapularis repair may create a biomechanically unfavourable condition due to the change in the joint centre of rotation after reverse shoulder arthroplasty(15). Subscapularis repair has also been associated with limited external rotation and impairment in daily living activities(15).

EFFECT ON RANGE OF MOTION

Subscapularis repair can be important for some authors, and they suggest that it should be performed because of its role in shoulder internal rotation, one of the main movements in many activities of daily living, such as tucking in a shirt in the back of the pants or clothing attach on the back(20). A successful subscapularis tendon repair can enhance a better range of motion and strength on internal rotation(12,23,24). For these authors, a routine repair of the tendon is advised in every case of reverse shoulder arthroplasty(25) and may be performed by transosseous(26,27) or end-to-end repair(27).

On the other hand, repairing the subscapularis may antagonize deltoid and posterior cuff, limiting elevation and external rotation, increasing surgery time, and more post-surgery protection is required(23,25). It may also contribute to coracoid

impingement(23,25). Besides that, there is no well documented evidence of better outcomes or healing rates(23,25). Subscapularis repair was also associated with a higher rate of reoperations (52,9% vs 0%, with a P val. = 0,01)(4).

Vourazeris et al(15) have studied the outcomes in patients submitted or not to subscapularis repair(15). In this study, with 202 patients, no statistically significant differences in the outcome scores were found (Shoulder Pain and Disability Index (SPADI) Score(28), American Shoulder and Elbow Surgeons Shoulder (ASES) Score(28) and University of California – Los Angeles (UCLA) Shoulder Rating Score(29))(15). They also did not find any advantages regarding range of motion or strength whether the subscapularis was repaired or not(15). After these results they have concluded that repairing the subscapularis had no effect in any of the measured range of motion outcomes, (active external rotation, passive external rotation, active forward elevation, active internal rotation, active abduction, external rotation)(15). In lateral designs, repair might not be necessary because the lateralization of the centre of rotation increases the subscapularis tension, making it difficult for external rotation and abduction.(15).

Friedman et al(24), in their comparative study of subscapularis repairing or not, have shown that the subscapularis repaired cohort was associated with better outcome scores according to Simple Shoulder Test (SST)(28,30) and Constant Murley scores(31,32), and was significantly more effective at improving active internal and external rotation(24), but less active abduction score and passive external rotation score(24). These results show that positive outcomes, regarding both range of motion and outcome scores, are achievable both with and without subscapularis repair(24). These results are more associated with the lateralization of the centre of rotation of the prostheses, and Friedman et al showed that good outcomes are achieved with or without subscapularis repair, with a lateralized prosthesis(24). So, these authors report no differences between repairing or not, but better outcomes with a lateralized prosthesis instead of a medialized Grammont style prosthesis(24).

Rol et al(12) have also concluded that the subscapularis repair was not associated with better internal rotation outcomes(12). They associated limited internal rotation to the

Miguel Leonardo Costa dos Santos
Lcosta_santos@hotmail.com

impingement related to the reduced space around the glenosphere, to allow a normal rotation around the humerus(12).

Franceschetti et al(22) have developed a field study in Italy comparing the repair and non-repair of subscapularis(22). They achieved similar results as Vourazeris(15) and Friedman(24), regarding Constant Murley(31,32) and Visual Analogue Scale (VAS) scores (no differences between groups), but they have demonstrated that the abduction value was higher in the non-repairing group and the internal rotation value was significantly higher in the repairing group(22). Regarding external rotation, the non-repairing group have achieved higher values, but the difference was not statistically and functionally significant(22). No differences were found regarding forward flexion(22). They have also considered the hypothesis of a higher value of post-operative pain in the repairing group, due to tendon scarring and increased tension on the sutured tendon(22).

Werner et al(16) have demonstrated that the isolated repair of the subscapularis had no significant effect on American Shoulder and Elbow Surgeons Shoulder (ASES) score(28) in reverse shoulder arthroplasty(16). On the other hand, when combined with the glenosphere lateralization (instead of medialization, in the Grammont-style prostheses) resulted in a less improved American Shoulder and Elbow Surgeons Shoulder (ASES)(28) score, when comparing with non-repairing group or medialization group(16).

In the systematic review conducted by Malahias et al(33), regarding range of motion, three out of four studies (of which two were also analysed in this review(8,15)), reported that there were no significant differences between repair or not repair. Regarding internal rotation, one of the studies(24) demonstrated a significantly higher improvement in the repair group, but the other ones found no significant differences(33). Additionally, no differences were found regarding external rotation(33). Although subscapularis repair was proven safe and effective for the augmentation of reverse total shoulder arthroplasty(33), it did not offer any additional clinical or functional benefit in range of motion and joint strength of patients treated with lateralized reverse total shoulder arthroplasty(33). Therefore, it may not be routinely performed in patients who have a preoperatively sufficient subscapularis tendon(33).

In another systematic review with metanalysis developed by Matthewson et al (19), of seven studies, (three of them explored in this review(15,16,24)), it has been demonstrated that even though there is some weak evidence of some advantages in subscapularis repair, the functional outcomes are similar both in subscapularis repairing or not, in reverse shoulder arthroplasty(19). One of the studies have demonstrated better score in Constant Murley evaluation(31,32), improved internal and external rotation on the subscapularis repair group, which was supported by another study analysed in this review(24). Yet, in this study(24), there were only used lateralized centre of rotation prostheses.

EFFECT ON JOINT STABILITY

In the medialized centre of rotation prostheses, “Grammont-style”, the centre of rotation is moved to the glenoid face, reducing the load on the baseplate-glenoid interface, and increasing the moment arm of the deltoid(15). This improves active elevation(15) with better outcomes on de daily living activities(15). However, dislocation rates are a concern, so Vourazeris et al, in their study(15) investigated whether the subscapularis repair can be beneficial to increase the joint stability(15). Nevertheless, they did not find statistically relevant differences between subscapularis repairing or not concerning joint stability and dislocations rates(15). Vourazeris et al(15) advised that subscapularis repair should be effective in medial humeral–medial glenosphere designs (Grammont style). However, in lateral humeral or glenosphere designs, the subscapularis repair is exempt and can be left up to the surgeon(15).

In the study developed by Friedman et al(24), no difference was noted in the complication rates, scapular notching rates, or scapular notching grades between patients with subscapularis repair and those without repair(24). They have also concluded that a lateralized centre of rotation prostheses is associated with decreased deltoid wrapping and more tension on rotator cuff, contributing for better stability(24). In their study, there were no differences between both subscapularis repair and not

repair groups, but they advised that those results may not be the same in medial centre of rotation prostheses “Grammont-style”(24).

Another study developed last year by Oh(34) et al came across the identical results of those developed by Friedman(24) and Vourazeris(15). These authors concluded that the subscapularis repair is not influential in the joint stability, in cases of a lateralized shoulder arthroplasty(34). Lateralizing the centre of rotation extends the moment arms of both external and internal rotation, when compared with a medialized centre of rotation(34), restoring the conjugate force between the subscapularis and infraspinous muscle, regardless the reparation or not of subscapularis tendon(34). Beyond that, lateralization of centre of rotation decreases deltoid wrapping and gives more tension to the rotator cuff, increasing the joint stability(34). This way, joint stability can increase with the lateralization of the centre of rotation, but is independent of the subscapularis repair(34). It should be noted that these authors carried out a study only with prostheses with lateralized centre of rotation, varying only the lateralization degree(34).

Regarding joint stability, Franceschetti et al(22) did not find any significant differences between repair or not repair the subscapularis tendon. However, they can claim that the prosthesis model used in their study, with the lateralized centre of rotation can offer better stability outcomes than the traditional Grammont design, as it would increase the rotator cuff tendon tension, promoting better stability(22). Therefore, in the Grammont design, as there is less deltoid tension and greater risk of impingement, subscapularis repair can be advised to stabilize the prosthesis, preventing the dislocation(22).

When comparing dislocation and/or instability rates, Werner et al(16) could not find any differences between groups(16). They concluded, then, that subscapularis repair is beneficial when conjugated with glenosphere lateralization, otherwise they found no advantages in individualized subscapularis repair in medial glenosphere prostheses(16). In lateralized glenosphere prostheses, they have demonstrated that the repair of subscapularis can translate to substantially less clinical improvement(16). They have also concluded that, for these types of prostheses, the subscapularis repair does not improve stability. In fact this combination may be prejudicial to shoulder function(16).

Miguel Leonardo Costa dos Santos
Lcosta_santos@hotmail.com

In a systematic review of comparative trials, developed by Malahias et al(33), using five studies, of which four of them were also analysed in this literature review(8,15,16,24), these authors concluded that the subscapularis repair did not offer any clinical or functional advantages in lateralized reverse total shoulder arthroplasty, even though this procedure was suggested by some authors to improve the joint stability and decrease post-operative dislocations(33). Thereby, these authors do not support the subscapularis repair as routine use for patients under reverse shoulder arthroplasty with a preoperatively sufficient subscapularis tendon, because it does not decrease the dislocations rate(33).

In their systematic review, Matthewson et al(19) suggested that subscapularis repair significantly reduces the risk of dislocation in cases of medialized prostheses(19). In lateralized prostheses, there were no difference between groups regarding stability(19). This kind of prostheses appear to be protective against dislocation, when compared with medialized prostheses(19). However, the benefits of a lateralized centre of rotation should still be weighed against the potential increased risk of component loosening and revision surgery secondary to increased joint reaction forces(19).

Cheung et al(35), in their study, have identified lack of subscapularis repair as an independent factor of instability after reverse total shoulder arthroplasty(35). They have demonstrated that a successful subscapularis repair would decrease the probability of prosthesis dislocation(35). Therefore, they supported the repair of subscapularis tendon, whenever possible, to improve the prosthetic stability(35). Nevertheless, they have used medialized centre of rotation prostheses in their study, which may limit the extrapolation of their results to other prosthetic designs.

This bibliographic review has several limitations. This study is subject to selection bias, due to the non-systematic methodology that may have not identified all the relevant bibliography. The information retrieved from the included studies may have also been biased by the non-systematic data analysis. Nevertheless, an extensive review of the queried articles was performed to minimize these biases. The heterogeneity of the articles is also an important factor, regarding the groups of patients in each study, the types of prostheses used, and the outcomes reported. In this study, it was not possible

Miguel Leonardo Costa dos Santos
Lcosta_santos@hotmail.com

to compare data statistically because it is not a meta-analysis resulting from a systematic review. Another limitation of this review is that it was carried out based mostly on retrospective comparative studies. There is still not enough detail to be able to concretely analyse the intended outcomes (range of motion and joint stability).

CONCLUSION

Through the consulted and analysed bibliography, we can conclude that many authors did not find significant evidence of the superiority of subscapularis repair in reverse shoulder arthroplasty, both in terms of joint stability, functional level, and range of motion.

Although it was not the main aim of this study, through this review it was also possible to suggest that the repair of the subscapular tendon is more effective when used concomitantly with the use of prostheses with medial rotation centre, “Grammont-style”. With the lateralized rotation centre, some authors do not see benefit in repairing the subscapular tendon, and there are even some authors that advise against the reparation in this type of prostheses.

However, there is still insufficient data to conclude which cases can benefit from simple subscapularis tenotomy and the ones that should be repaired.

BIBLIOGRAPHY

1. GRAMMONT P, TROUILLOUD P, LAFFAY J, DERIES X. Etude et réalisation d'une nouvelle prothèse d'épaule. *Rhumatologie (Aix-les-Bains)*. 1987;
2. Vaz O. Artroplastia Invertida do Ombro [Internet]. 2018. Available from: <http://hdl.handle.net/10316/82407>
3. Familiari F, Rojas J, Doral MN, Huri G, McFarland EG. Reverse total shoulder arthroplasty. *EFORT Open Reviews*. 2018;3(2):58–69.
4. Entezari V, Henry T, Zmistowski B, Sheth M, Nicholson T, Namdari S. Clinically significant subscapularis failure after anatomic shoulder arthroplasty: is it worth repairing? *Journal of Shoulder and Elbow Surgery* [Internet]. 2020;29(9):1831–5. Available from: <https://doi.org/10.1016/j.jse.2020.01.070>
5. Nayak A, Hansen M, Worhacz K, Stowell R, Jacofsky M, Roche C, et al. Role of Subscapularis Repair on Muscle Force Requirements with Reverse Shoulder Arthroplasty. *Bulletin of the NYU hospital for joint diseases*. 2015 Jan;73:21–7.
6. Cavalier M, Jullion S, Kany J, Grimberg J, Lefebvre Y, Oudet D, et al. Management of Massive Rotator Cuff Tears: Prospective study in 218 patients. *Orthopaedics and Traumatology: Surgery and Research*. 2018 Dec;104(8):S193–7.
7. Cuff DJ, Pupello DR, Santoni BG. Partial rotator cuff repair and biceps tenotomy for the treatment of patients with massive cuff tears and retained overhead elevation: midterm outcomes with a minimum 5 years of follow-up. *Journal of Shoulder and Elbow Surgery* [Internet]. 2016;25(11):1803–9. Available from: <http://dx.doi.org/10.1016/j.jse.2016.04.001>
8. de Boer FA, van Kampen PM, Huijsmans PE. The influence of subscapularis tendon reattachment on range of motion in reversed shoulder arthroplasty: a clinical study. *Musculoskeletal Surgery*. 2016;100(2):121–6.
9. Boileau P, Watkinson DJ, Hatzidakis AM, Balg F. Grammont reverse prostheses: Design, rationale, and biomechanics. *Journal of Shoulder and Elbow Surgery*. 2005;14(1 SUPPL.):S147–61.
10. Berliner JL, Regalado-Magdos A, Ma CB, Feeley BT. Biomechanics of reverse total shoulder arthroplasty. *Journal of Shoulder and Elbow Surgery* [Internet]. 2015;24(1):150–60. Available from: <http://dx.doi.org/10.1016/j.jse.2014.08.003>
11. Jazayeri R, Kwon YW. Evolution of the reverse total shoulder prostheses. *Bulletin of the NYU Hospital for Joint Diseases*. 2011;69(1):50–5.

Miguel Leonardo Costa dos Santos
Lcosta_santos@hotmail.com

12. Rol M, Favard L, Berhouet J. Factors associated with internal rotation outcomes after reverse shoulder arthroplasty. *Orthopaedics and Traumatology: Surgery and Research*. 2019 Dec;105(8):1515–9.
13. Kim MS, Jeong HY, Kim JD, Ro KH, Rhee SM, Rhee YG. Difficulty in performing activities of daily living associated with internal rotation after reverse total shoulder arthroplasty. *Journal of Shoulder and Elbow Surgery*. 2020 Jan;29(1):86–94.
14. Krämer M, Bäunker A, Wellmann M, Hurschler C, Smith T. Implant impingement during internal rotation after reverse shoulder arthroplasty. the effect of implant configuration and scapula anatomy: A biomechanical study. *Clinical Biomechanics*. 2016 Mar;33:111–6.
15. Vourazeris JD, Wright TW, Struk AM, King JJ, Farmer KW. Primary reverse total shoulder arthroplasty outcomes in patients with subscapularis repair versus tenotomy. *Journal of Shoulder and Elbow Surgery*. 2017 Mar;26(3):450–7.
16. Werner BC, Wong AC, Mahony GT, Craig E v., Dines DM, Warren RF, et al. Clinical Outcomes After Reverse Shoulder Arthroplasty With and Without Subscapularis Repair: The Importance of Considering Glenosphere Lateralization. *The Journal of the American Academy of Orthopaedic Surgeons*. 2018;26(5):e114–9.
17. Roberson TA, Shanley E, Griscom JT, Granade M, Hunt Q, Adams KJ, et al. Subscapularis Repair Is Unnecessary After Lateralized Reverse Shoulder Arthroplasty. *JBJS Open Access*. 2018;3(3):e0056.
18. Moore, K. L.; Dalley, A. F.; Agur AMR. *Moore Clinically Oriented Anatomy EIGHTH EDITION*. Vol. 282, Wolters Kluwer. Wolters Kluwer Health; 2018. 1045–1059.
19. Matthewson G, Kooner S, Kwapisz A, Leiter J, Old J, MacDonald P. The effect of subscapularis repair on dislocation rates in reverse shoulder arthroplasty: a meta-analysis and systematic review. *Journal of Shoulder and Elbow Surgery*. 2019 May;28(5).
20. Simovitch RW, Friedman RJ, Cheung E v., Flurin PH, Wright T, Zuckerman JD, et al. Rate of Improvement in Clinical Outcomes with Anatomic and Reverse Total Shoulder Arthroplasty. *Journal of Bone and Joint Surgery - American Volume*. 2017 Nov;99(21):1801–11.
21. Vinet L, Zhedanov A. *Netter’s Atlas of Human Anatomy*. Vol. 44, Journal of Physics A: Mathematical and Theoretical. 2020. 51.
22. Franceschetti E, de Sanctis EG, Ranieri R, Palumbo A, Paciotti M, Franceschi F. The role of the subscapularis tendon in a lateralized reverse total shoulder arthroplasty: repair versus nonrepair. *International Orthopaedics*. 2019 Jan;43(11):2579–86.
23. Sanchez-Sotelo J, Athwal GS. How to Optimize Reverse Shoulder Arthroplasty for Irreparable Cuff Tears. *Current Reviews in Musculoskeletal Medicine*. 2020;13(5):553–60.

Miguel Leonardo Costa dos Santos
Lcosta_santos@hotmail.com

24. Friedman RJ, Flurin PH, Wright TW, Zuckerman JD, Roche CP. Comparison of reverse total shoulder arthroplasty outcomes with and without subscapularis repair. *Journal of Shoulder and Elbow Surgery* [Internet]. 2017;26(4):662–8. Available from: <http://dx.doi.org/10.1016/j.jse.2016.09.027>
25. Dedy NJ, Gouk CJ, Taylor FJ, Thomas M, Tan SLE. Sonographic assessment of the subscapularis after reverse shoulder arthroplasty: impact of tendon integrity on shoulder function. *Journal of Shoulder and Elbow Surgery* [Internet]. 2018;27(6):1051–6. Available from: <https://doi.org/10.1016/j.jse.2017.12.008>
26. Black EM, Lin A, Srikumaran U, Jain N, Freehill MT. Arthroscopic transosseous rotator cuff repair: Technical note, outcomes, and complications. *Orthopedics*. 2015;38(5):e352–8.
27. Hartline JT, Brolin TJ, Wan JY, Dibaba DT, Azar FM, Throckmorton TW. The effect of subscapularis management technique on outcomes and complication rates following reverse total shoulder arthroplasty. *Seminars in Arthroplasty* [Internet]. 2020;30(1):42–9. Available from: <https://doi.org/10.1053/j.sart.2020.04.005>
28. Angst F, Schwyzer HK, Aeschlimann A, Simmen BR, Goldhahn J. Measures of adult shoulder function: Disabilities of the Arm, Shoulder, and Hand Questionnaire (DASH) and Its Short Version (QuickDASH), Shoulder Pain and Disability Index (SPADI), American Shoulder and Elbow Surgeons (ASES) Society Standardized Shoulder . *Arthritis Care and Research*. 2011;63(SUPPL. 11):174–88.
29. Luz Yolanda Toro Suarez. UCLA Shoulder Score [Internet]. 2015. p. 1–27. Available from: <https://www.codetechnology.com/ucla-shoulder-score-tool/>
30. Gutiérrez S, Comiskey IV CA, Luo ZP, Pupello DR, Frankle MA. Range of impingement-free abduction and adduction deficit after reverse shoulder arthroplasty. Hierarchy of surgical and implant-design-related factors. *Journal of Bone and Joint Surgery - Series A*. 2008;90(12):2606–15.
31. Vrotsou K, Ávila M, Machón M, Mateo-Abad M, Pardo Y, Garin O, et al. Constant–Murley Score: systematic review and standardized evaluation in different shoulder pathologies. *Quality of Life Research* [Internet]. 2018;27(9):2217–26. Available from: <http://dx.doi.org/10.1007/s11136-018-1875-7>
32. Constant CR, Murley AHG. A clinical method of functional assessment of the shoulder. *Clinical Orthopaedics and Related Research*. 1987;No. 214(April):160–4.
33. Malahias MA, Gerogiannis D, Chronopoulos E, Kaseta MK, Brilakis E, Antonogiannakis E. Is subscapularis repair associated with better outcome compared to non-repair in reverse total shoulder arthroplasty? A systematic review of comparative trials. *Orthopedic Reviews*. 2019;11(3):134–41.
34. Oh JH, Sharma N, Rhee SM, Park JH. Do individualized humeral retroversion and subscapularis repair affect the clinical outcomes of reverse total shoulder arthroplasty?

Miguel Leonardo Costa dos Santos
Lcosta_santos@hotmail.com

Journal of Shoulder and Elbow Surgery [Internet]. 2020;29(4):821–9. Available from: <https://doi.org/10.1016/j.jse.2019.08.016>

35. Cheung E v., Sarkissian EJ, Sox-Harris A, Comer GC, Saleh JR, Diaz R, et al. Instability after reverse total shoulder arthroplasty. Journal of Shoulder and Elbow Surgery [Internet]. 2018;27(11):1946–52. Available from: <https://doi.org/10.1016/j.jse.2018.04.015>

ANEXES

MANUSCRIPT GUIDELINES - The official journal of the European Federation of National Associations of Orthopaedics and Traumatology (EFORT)

“Information for authors

EFORT Open Reviews is a peer-reviewed, ‘gold’ open access journal. The journal welcomes objective, comprehensive review articles across the whole field of orthopaedics and traumatology, from authors from all parts of the world.

Acceptance for publication is subject to rigorous double-blind peer review by our panel of expert reviewers, and all articles that are accepted for publication undergo rigorous scientific editing.

Google Scholar, PubMed, Scopus and Web of Science

EFORT Open Reviews is indexed in PubMed with free full text provided in PubMed Central (PMC). It is also indexed in the Web of Science Core Collection, Google Scholar, and Scopus. It is listed in the Directory of Open Access Journals (DOAJ).

Open Access

EFORT Open Reviews is an open access journal: authors pay an article processing charge (APC) for their accepted articles to be made freely and permanently accessible online immediately upon publication. For further information, please see Licensing & Article Publication Charges.

Online submission

All manuscript submissions should be made online at the EFORT Open Reviews ScholarOne Manuscripts site: <https://mc04.manuscriptcentral.com/eor>

If you have been invited to submit a paper, please click on the link within your invitation email to navigate directly to your ScholarOne Author Centre, where you can enter information about the paper and upload article and figure files.

Miguel Leonardo Costa dos Santos
Lcosta_santos@hotmail.com

If you are submitting an unsolicited manuscript, please read the manuscript guidelines carefully to check that your paper meets the criteria for submission, then click the 'Create Account' link and follow the on-screen instructions.

Manuscript guidelines

Articles are accepted for exclusive publication in EFORT Open Reviews and must not be previously published or under consideration elsewhere.

Review Articles

Review Articles should summarise the current state of understanding on a topic within the fields of orthopaedics and traumatology.

The journal's focus is on instructional review articles, which discuss and synthesize the key, recent, peer-reviewed literature on the topic. The aim is that readers may quickly gain an overview of current knowledge and practice in the field. Review articles should display sufficient originality in their analysis of the field and recommendations for practice, and should not merely repeat or paraphrase existing literature – all submitted articles are screened for similarity to published works, including the author's own previous publications. Your expert assessment of the most relevant literature may also cover current debates and advise on future directions. Illustrative figures, including annotated radiographs, images and diagrams should be included to aid understanding.

Systematic Reviews

In addition to instructional review articles, EOR publishes systematic reviews, which are more extensive analyses that critically assess all published studies on a topic, according to clearly defined search criteria.

Systematic Reviews must meet the requirements of the PRISMA guidelines for systematic reviews and meta-analyses. Authors are required to complete the PRISMA checklist and submit this with their manuscript. You can find the checklist [here](#). Authors should include a statement such as below in the abstract and the 'Materials & Methods' section of their manuscript:

'The review process was conducted according to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines.'

Although not an obligatory requirement, the journal encourages registration of Systematic Reviews with PROSPERO (International Prospective Register of Systematic Reviews). If registered, authors should include the registration number in the abstract and the 'Materials & Methods' section of their manuscript.

Miguel Leonardo Costa dos Santos
Lcosta_santos@hotmail.com

The abstract should be structured into Purpose, Methods, Results and Conclusions and be limited to a maximum of 250 words.

File type

The article text should be supplied as a Word DOC or RTF. Please use continuous line numbering in your main document file (in Word 2010, go to Page Layout>Line Numbers>Continuous). See additional guidelines on figure and table files below.

Word count

Review articles should be as concise as possible - we recommend a maximum word count of 4000 words.

Authorship

Each paper should have a maximum of 6 authors, and each author must have contributed substantially to the review. The corresponding author provides final approval for publication. Each author is required to complete an ICMJE conflict of interest form prior to publication.

Cover letter

You are required to submit a cover letter with your manuscript.

Key points to include:

- Editor's name (you can find this on the Editorial Board page)
- Your manuscript's title
- Statement that your paper in part or in whole has not been previously published and is not currently under consideration by another journal
- Brief description of the research you are reporting in your paper, why it is important, and why you think the readers of EFORT Open Reviews would be interested in it
- Your contact information
- Conflict of interest statement

If you need further help to write your cover letter, you can download and use our sample cover letter template here. This will download a Word document.

Title page

To ensure anonymity in peer review, the title page should be supplied separately from the body of the article, which should not contain any identifying information. The title page should contain the full title of the manuscript, author name(s) and affiliation(s), and the name, postal and email addresses of the corresponding author.

Abstract and keywords

For narrative, instructional Review Articles, the abstract should summarise the content and main conclusions of the article into 3–6 bullet points, and should be limited to a maximum of 250 words.

For Systematic Reviews, the abstract should be structured into Purpose, Methods, Results and Conclusions and be limited to a maximum of 250 words. Keywords should be supplied for every article, based upon 3–8 key terms that might be used when searching for the article online.

Figures and tables

Each review article should include a small number of figures or images to illustrate points (techniques, materials, procedures and so on) that are discussed in the text, in order to aid readers' understanding. For any figures that are reproduced from previously published material (including websites), permission must be requested from the copyright holder, and confirmation of permission must be provided to the EOR Editorial Office. It is the author's responsibility to request permission to reproduce figures.

Figures should be submitted as separate files, and should not be embedded in the Word doc. Tables should be embedded in the Word doc. All figures and tables must be clearly cited in the text, and figure legends should be listed at the bottom of the main text file.

Each figure should have a figure legend, and figures should be numbered consecutively.

Each table should have a short, descriptive heading, and tables should be numbered consecutively. Tables should be supplied using proper table formatting with columns and rows.

As far as possible, figures should be provided as original source files, and should not be embedded within another document. Images may be supplied as JPEG or TIFF files at 300dpi or higher; line art may be supplied as EPS files at 800 dpi or higher.

Conclusions

Articles should usually finish with a Conclusions section. This should be more than simply a repetition of the body text, and should provide clear recommendations based on the literature and the authors' experience (as appropriate). Bullet points may be used for succinct expression of recommendations. This section may also briefly discuss the need for ongoing study and future directions.

References

References should be kept to a minimum, comprising only works that have been studied in full by the authors, and should only cite published work. All references must be cited within the body of the article. The majority of references should refer to peer-reviewed literature published within the last 5 years. References should be presented using the Vancouver system, whereby references are numbered according to the order of citation in the text (not in alphabetical order). The reference list should appear at the end of the article. Vancouver style reference examples are as follows:

Journal article

Patel MS, Newey M, Sell P. A comparison of patient-reported outcome measures after spinal surgery. *Bone Joint J* 2015;97-B:366–371.

Book

Crawford Adams, J. *Standard orthopaedic operations*. Fourth ed. Edinburgh: Churchill Livingstone, 1992.

Chapter in a book

Winqvist RA, Frankel VH. Complications of implant use. In: Epps CH Jr, ed. *Complications in orthopaedic surgery*. Vol. 1. Philadelphia: JB Lippincott Company, 1978:99–129.

Web reference

Hazarika S, Baird E, Palan J. British Orthopaedics Trainees Association: BOTA positional statement on the use of simulation in surgical training, 2013. http://www.bota.org.uk/uploads/post_02291_BOTA_statement_simulation.pdf (date last accessed 25 March 2015).

Abstract Reference

Peterson L. Osteochondritis of the knee treated with autologous chondrocyte transplantation [abstract]. ISAKOS Congress, 2001.

Manuscript polishing

Miguel Leonardo Costa dos Santos
Lcosta_santos@hotmail.com

We recommend The Charlesworth Group, who provide academic editing services to help authors refine their language and clarify information in their texts, cover letters, and other materials needed to communicate clearly. If you would like to use this service please follow this link.

Audio, video and multimedia material

Audio and video clips and other complementary multimedia material are very much encouraged. Multimedia material should be of the highest possible quality. For further guidelines or advice on inclusion of multimedia material, please contact the Managing Editor.

Copyright

EFORT Open Reviews publishes manuscripts under a Creative Commons Attribution Non-Commercial license (CC-BY-NC) as standard, which allows others to re-use the work without permission so long as the work is properly referenced and the use is non-commercial. For more information, please visit <http://creativecommons.org/licenses/by-nc/4.0/>. If you are in receipt of funding and your funder requires publication under a CC-BY licence, please inform your Production Editor following acceptance.

Permissions

Authors are responsible for obtaining permission from copyright holders for reproducing any illustrations, tables, figures or lengthy quotations previously published elsewhere.

No patient identifying information should be published in written descriptions, photographs, or multimedia material unless essential, in which case written informed Consent must be provided.

Plagiarism screening

EFORT Open Reviews takes issues of plagiarism, copyright infringement and other breaches of best practice seriously. All submitted articles are screened using duplication-checking software, and we reserve the right to reject or retract any articles that are found to have plagiarised other work or reproduced copyright material without permission. This extends to reproducing sections of other works previously published by the submitting author(s). Further action may be taken as appropriate.”