

Atomic and Specificity Details of Mucin 1 O-Glycosylation Process by Multiple Polypeptide GalNAc-Transferase Isoforms Unveiled by NMR and Molecular Modeling

Helena Coelho, Matilde de las Rivas, Ana S. Grosso, Ana Diniz, Cátia O. Soares, Rodrigo A. Francisco, Jorge S. Dias, Ismael Compañón, Lingbo Sun, Yoshiki Narimatsu, Sergey Y. Vakhrushev, Henrik Clausen, Eurico J. Cabrita, Jesús Jiménez-Barbero, Francisco Corzana, Ramon Hurtado-Guerrero,* and Filipa Marcelo*



Cite This: <https://doi.org/10.1021/jacsau.1c00529>



Read Online

ACCESS |



Metrics & More




Article Recommendations



Supporting Information

ABSTRACT: The large family of polypeptide GalNAc-transferases (GalNAc-Ts) controls with precision how GalNAc O-glycans are added in the tandem repeat regions of mucins (e.g., MUC1). However, the structural features behind the creation of well-defined and clustered patterns of O-glycans in mucins are poorly understood. In this context, herein, we disclose the full process of MUC1 O-glycosylation by GalNAc-T2/T3/T4 isoforms by NMR spectroscopy assisted by molecular modeling protocols. By using MUC1, with four tandem repeat domains as a substrate, we confirmed the glycosylation preferences of different GalNAc-Ts isoforms and highlighted the importance of the lectin domain in the glycosylation site selection after the addition of the first GalNAc residue. In a glycosylated substrate, with yet multiple acceptor sites, the lectin domain contributes to orientate acceptor sites to the catalytic domain. Our experiments suggest that during this process, neighboring tandem repeats are critical for further glycosylation of acceptor sites by GalNAc-T2/T4 in a lectin-assisted manner. Our studies also show local conformational changes in the peptide backbone during incorporation of GalNAc residues, which might explain GalNAc-T2/T3/T4 fine specificities toward the MUC1 substrate. Interestingly, we postulate that a specific salt-bridge and the inverse γ -turn conformation of the PDTRP sequence in MUC1 are the main structural motifs behind the GalNAc-T4 specificity toward this region. In addition, in-cell analysis shows that the GalNAc-T4 isoform is the only isoform glycosylating the Thr of the immunogenic epitope PDTRP *in vivo*, which highlights the relevance of GalNAc-T4 in the glycosylation of this epitope. Finally, the NMR methodology established herein can be extended to other glycosyltransferases, such as C1GalT1 and ST6GalNAc-I, to determine the specificity toward complex mucin acceptor substrates.

KEYWORDS: O-GalNAc glycosylation, GalNAc-Ts, mucin-1, NMR, molecular dynamics, cell analysis



Mucin-type (GalNAc-type) O-glycosylation is one of the most abundant and diverse forms of posttranslational modifications (PTMs) and is differentially regulated by a large family of polypeptide GalNAc-transferases (GalNAc-Ts).¹ GalNAc-Ts are type-II membrane proteins that catalyze the transfer of N-acetylgalactosamine (GalNAc) from UDP-GalNAc to Ser/Thr residues (and possibly Tyr).¹ While 20 GalNAc-Ts isoforms in humans initiate the O-GalNAc-type glycosylation (hereafter, O-glycosylation) with distinct, although partly overlapping, kinetic properties and acceptor substrate specificities,² other types of protein O-glycosylation pathways are initiated by a less number of enzymes (ranging between one to four enzymes).^{3–5} No clear consensus motifs for O-glycosylation have been reported, making this PTM the most complex and differentially regulated type of protein

glycosylation. GalNAc-Ts are localized in the Golgi apparatus and distributed among all organs with different expression levels in each tissue.^{1,5,6} Thus, the available repertoire of GalNAc-Ts and their localization determine which proteins are O-glycosylated and where O-glycans are attached. In this context, some glycoproteins with fewer acceptor sites are glycosylated by specific GalNAc-Ts (nonredundant glycosyla-

Received: November 23, 2021

tion),^{7–11} while complex glycoproteins with multiple acceptor sites (e.g., mucins) are glycosylated by multiple GalNAc-Ts (redundant glycosylation).¹²

GalNAc-Ts contains a region in the Golgi lumen displaying an N-terminal catalytic that adopts a GT-A fold¹³ and belongs to CAZY family 27 and connected to a C-terminal lectin domain by a short flexible linker,^{14–17} which dictates the relative orientation between the catalytic and lectin domains. The GalNAc-Ts are unique among metazoan GTs because of the C-terminal GalNAc-binding lectin domain, which adopts a β -trefoil fold built from three homologous repeat subdomains, named α , β , and γ .¹⁷ However, only one or two specific subdomains (α , β , or γ) actively bind to the GalNAc moiety. For example, in GalNAc-T2, -T3, and -T4, only the α -subdomain recognizes the sugar moiety.^{2,14,18–22}

GalNAc-Ts can be classified into two classes based on their short-range (acceptor sites adjacent to prior glycosites) and long-range (acceptor sites far from prior glycosites) glycosylating capabilities. The short-range glycosylation preferences account for the glycosylation of glycopeptide substrates where a proximal glycosite to the acceptor site binds to the catalytic domain promoting glycosylation (one to three residues from the prior glycosite).^{23,24} In contrast, the long-range glycosylation preferences refer to the binding of a prior glycosite to the lectin domain, which subsequently facilitates the recognition of the distant acceptor site (6 to ~17 residues away from the acceptor site) by the catalytic domain, promoting the glycosylation event.¹² Both glycosylation preferences operate on N- or C-terminal directions, and specifically, for the long-range glycosylation preferences, this behavior is dictated by the GalNAc-T isoform-specific interdomain flexible linker motion.²⁵ GalNAc-T3/T4/T6/T12 preferentially glycosylate remote C-terminal sites from the prior N-terminal GalNAc site,^{2,14,18–22} whereas GalNAc-T1/T2/T14 exhibit the opposite preference.^{15,26,27}

Much of the knowledge about the GalNAc-Ts family is derived from *in vitro* enzyme studies with the characterization of products by mass spectrometry (MS) analysis.^{18,21,26,28–30} In addition, studies of isogenic cell lines, animal models, and individuals with impaired GalNAc-T2/T3 mutations have provided information on the nonredundant functions of several GalNAc-T isoforms.^{31–33} At the molecular level, insights into the catalytic mechanism of GalNAc-Ts have been revealed by X-ray crystallography, NMR, and molecular dynamics (MD) simulations^{23,25,34,35} by employing short (glyco)peptides. Specifically, our previous results suggest that GalNAc-Ts adopt a UDP-GalNAc-dependent induced-fit mechanism.³⁴

However, our understanding of the molecular basis of GalNAc-Ts substrate specificity and especially how these enzymes act in a coordinated manner on very complex substrates such as mucins are barely known. The best-studied model of coordinated *O*-glycosylation has been performed on the mucin 1 (MUC1) tandem repeats (TRs), which consist of repeating 20-mer units of GVT³S⁴APDT⁸RPAPGS¹⁴T¹⁵APPAH. *In vitro* and *in cell* studies have demonstrated that several GalNAc-Ts (GalNAc-T1/T2/T3/T11) initiate *O*-glycosylation in MUC1 and add GalNAc moieties to T³, T¹⁵, and S¹⁴ with different preferences and order of glycosylation.^{11,36} On the contrary, GalNAc-T4 is the only isoform requiring prior *O*-glycosylation of at least one site by the above isoforms to glycosylate *in vitro* the remaining acceptor sites, S⁴ and T⁸.^{10,22}

O-Glycosylation induces specific alterations in the chemical environments of protein residues, which can be readily monitored by high-resolution NMR spectroscopy,³⁷ allowing the determination of the glycosylation specificity for different acceptor sites. In this context, Gariépy and co-authors applied NMR to ascertain the order and kinetics of GalNAc-introduction in a MUC1-5TR substrate by GalNAc-T1 isoform.³⁸ NMR allows the determination of how glycosylation affects the protein structure conformation, and in cases of clustered modifications, how one glycosylation site affects the subsequent glycosylation reaction. Particularly, Ser and Thr *O*-GalNAc glycosylation was reported to elicit extended-like structures in short MUC1 glycopeptides.^{39–42}

On this basis, herein, we exploit NMR spectroscopy to characterize the full process of MUC1 *O*-glycosylation by GalNAc-T2/T3/T4 and to investigate the role of the lectin domain to guide catalysis (Figure 1).




Figure 1. MUC1 *O*-glycosylation by multiple GalNAc-Ts. (A) MUC1^{4TR} template used in this study. (B) 3D structure of the GalNAc-Ts isoforms used in this study and complexed with a glycopeptide (GalNAc-T2 PDB ID: 5AJP; GalNAc-T3 PDB ID: 6S24; GalNAc-T4 PDB ID: 6H0B). The lectin binding site is highlighted and displays GalNAc in yellow sticks and the essential Asp residue, conserved in all isoforms, in pink sticks. The Asp established H-bonds with OH-3 and OH-4 of GalNAc, which are identified as dashed blue lines.

The glycosylation process of a four TR domain of MUC1 (MUC1^{4TR}) (Figure 1A) was explored by using the wild-type GalNAc-T2, -T3, and -T4 isoforms and their corresponding impaired lectin mutants GalNAc-T2^{D458R}, GalNAc-T3^{D517H}, and GalNAc-T4^{D459H}, which contain a critical mutation in the conserved Asp residue located at the α -subdomain of the lectin domain (Figure 1B). Previous studies demonstrated that the mutation of this Asp to a positive charged residue (Arg or His) precludes the binding of GalNAc by the lectin domain.^{18,21,43}




Figure 2. Glycosylation of MUC1^{4TR} catalyzed by the GalNAc-T2 and its GalNAc-T2^{D458R} mutant. Table S5 details the experimental conditions. (A) Scheme of the MUC1^{4TR} glycosylation event in the presence of the GalNAc-T2 enzyme. The glycosylation sites are displayed in red in the amino acid sequence of MUC1^{4TR}; (B,C) ¹H,¹⁵N-HSQC spectra recorded during the glycosylation process of MUC1^{4TR} by the GalNAc-T2 (B) and its GalNAc-T2^{D458R} mutant (C) without UDP-GalNAc (black) and after 20 min (red) and 24 h (blue), using an excess of the donor UDP-GalNAc. Arrows indicate the observed CSP that occur as a result of the GalNAc additions at T¹⁵ (blue arrows), T³ (green arrows), and S¹⁴ (orange arrows). The * labeling in the amino acids on the spectra indicates that either Thr or Ser residues are glycosylated, and # labeling indicates shifts on glycosylated T¹⁵ after glycosylation of S¹⁴.

Through monitoring of the chemical shift perturbation (CSP) of the amide resonances in ¹H/¹⁵N-HSQC spectra of the ¹⁵N-isotopically labeled MUC1^{4TR}, our studies revealed a stepwise mechanism for each isoform, confirming the distinct glycosylation preferences employed by each isoform and highlighting the relevance of the lectin domain in these enzymes. The alterations in the conformation of MUC1^{4TR} upon GalNAc incorporation along the O-glycosylation process were also monitored, and their implication on the GalNAc-Ts activity was also inferred. Finally, cellular assays were performed to define the GalNAc-T isoforms involved in glycosylating S⁴ and T⁸ in the MUC1 structure, revealing

GalNAc-T4 isoform as the only isoform able of glycosylating the T⁸ of the immunogenic epitope PDTRP.

RESULTS AND DISCUSSION

NMR Analysis of ¹⁵N-MUC1^{4TR} Glycosylation by the GalNAc-T2 Isoform—Role of the Lectin Domain

¹⁵N-MUC1^{4TR} was used as a probe to explore MUC1 O-glycosylation by NMR (see material and methods for details, Figures S1 and S2 and Table S1 of the Supporting Information). The ¹⁵N-MUC1^{4TR} molecular weight was verified by MS, and its NMR spectrum was assigned by




Figure 3. Putative 3D models to explain the glycosylation of T³ and S¹⁴ of MUC1^{4TR} catalyzed by GalNAc-T2. (A) 3D view of complex GalNAc-T2/1 derived from a representative frame obtained from 0.5 μ s of MD simulation. A key CH- π interaction between the β -methyl group of T³ and the aromatic ring of F372 is shown in dashed lines (distance 4.8 ± 0.4 Å). (B) 3D view of the binding site of GalNAc-T2/2 derived from a representative frame obtained from 0.5 μ s of MD simulation. A relevant hydrogen bond, 38% populated through the whole trajectory, between the GalNAc at T⁶ of 2 (T¹⁵ in MUC1^{4TR}) and W265 is highlighted in dashed lines. In both structures, the catalytic domain is displayed in orange and the lectin domain is depicted in green. In the glycopeptides, the GalNAc units are shown in yellow sticks. UDP-GalNAc is displayed in gray sticks and Mn²⁺ is shown as a cyan sphere. Relevant residues of the glycopeptides and some residues of catalytic domain are also shown as sticks and in a different color. Schematic representation of the sequential glycosylation process at T³ and S¹⁴ of MUC1^{4TR} catalyzed by GalNAc-T2 is also displayed. These models highlight the N-terminal preference of GalNAc-T2.

standard procedures (Figure S2 and Table S1). The reaction with GalNAc-T2 and GalNAc-T2^{D458R} impaired lectin mutant was carried under an excess of UDP-GalNAc and was followed over time through the acquisition of a series of ¹H/¹⁵N-HSQC spectra. Figure 2 shows the selected ¹H/¹⁵N-HSQC spectra of MUC1^{4TR} before and after addition of UDP-GalNAc at 20 min and 24 h. In the ¹H/¹⁵N-HSQC spectra before the addition of UDP-GalNAc, apart from the N-terminals T3 and S4 and C-terminals A19 and H20 for which two peaks are visible, only one amide peak is visible for each of the residues in the same relative position on the MUC1^{4TR} (Figure 2B,C black spectra and Figure S2). The ¹H,¹⁵N combined CSPs observed on the ¹H/¹⁵N-HSQC spectra reflect the GalNAc transfer to MUC1^{4TR} (Figure S3A). In agreement with previously reported results,³⁶ the first glycosylation events in the MUC1^{4TR} take place at the T¹⁵ moieties (T^{15*}, where * denotes that the residue is covalently attached to GalNAc), as deduced by the strong CSPs observed at these residues (0.38 ppm) and their vicinal amino acids, with A¹⁶ experiencing a considerable CSP (0.28 ppm). Then, glycosylation at T³ is evidenced by its CSP (0.32 ppm) along with those observed for S⁴ (0.23 ppm), A⁵ (0.13 ppm), and V² (0.08 ppm). Finally, glycosylation at S¹⁴ is verified by its corresponding CSP (0.15 ppm) along with those observed for the glycosylated T^{15*} (0.23 ppm) (Figure 2B identified as T^{15#}). Furthermore, CSPs

are also detected for A¹⁶ (0.11 ppm) and G¹³ (0.10 ppm). The measurement of the cross-peak volumes in the ¹H/¹⁵N-HSQC spectra allowed the quantification of the % glycosylation of each acceptor site in all four TRs. Indeed, after 20 min of reaction (fast glycosylation), complete glycosylation at T¹⁵ (blue arrows) and T³ (green arrows) was observed (Figure 2B). In contrast, only ~70% of S¹⁴ residues (orange arrows) were glycosylated. Thus, one of the four TR domains is not glycosylated at S¹⁴ in MUC1^{4TR} and requires 24 h (exhaustive glycosylation) for completion (Figure 2B). MS analysis together with the full NMR assignment of the purified product confirmed that T¹⁵, T³, and S¹⁴ were fully glycosylated in all TRs, while S⁴ and T⁸ remained nonglycosylated (Figure S4 and Table S2).

The combined ¹H,¹⁵N-CSPs (Figures S3A) observed for the neighboring residues to the glycosylation site suggest local alterations of the chemical and structural environment of MUC1^{4TR} due to the incorporation of GalNAc units, which is consistent with previous observations with short MUC1 Tn-glycopeptides.^{39–42} The ¹H and ¹⁵N NMR chemical shifts for the nonglycosylated MUC1^{4TR} are indicative of a major random coil-like conformational ensemble. However, besides the multiple conformations in solution, local conformations, typically found in the extended region of the Ramachandran plot for the three potential glycosylation regions of MUC1,

VTSA (mixture of β -strand/inverse γ turn), DTR (inverse γ turn), and GSTA (polyproline II-like), were previously identified by Kinarsky *et al.*³⁹ Thus, according to the authors, nonglycosylated MUC1 has in these regions a natural tendency to adopt extended-like conformations. Herein, the chemical shift index (CSI)^{44,45} of the $^{13}\text{C}_\alpha$ / $^{13}\text{C}_\beta$ of the naked MUC1^{4TR} and the glycosylated (MUC1^{4TR}-T³*S¹⁴T¹⁵*) were calculated and compared (Figure S3B, gray and green, respectively), clearly indicating local deviations around VTS and GSTA from the random coil (small negative CSI values) toward an extended conformation (due to an increase of negative CSI values) after the introduction of GalNAc.

By using the GalNAc-T2^{D458R} lectin mutant and after 20 min of reaction (Figure 2C), CSPs were only detected for T¹⁵ and the residues in its vicinity, and only after 90 min, glycosylation at T³ (~20%) was detected (Figure S5). Even after 24 h, glycosylation at T³ was not completed (*ca.* 80%) (Figures 2C and S5), while only a very small fraction of S¹⁴ sites was glycosylated (5%). Thus, the glycosylation at T¹⁵ sites takes place prior to glycosylation at T³, and although the GalNAc-T2^{D458R} mutant can glycosylate T³ and S¹⁴ residues, its catalytic efficiency is affected by the mutation at the lectin domain. These results suggest that GalNAc-T2 uses its lectin domain to drive glycosylation at S¹⁴ but also to a certain degree at T³ in MUC1. This agrees with previous studies, showing that GalNAc-T2 glycosylation at S¹⁴ requires the lectin domain,¹⁸ and further suggests that also the glycosylation of T³ is assisted by the lectin domain.

The preference of GalNAc-T2 toward T¹⁵ versus T³ in MUC1^{4TR} was also supported through NMR experiments limiting the UDP-GalNAc concentration with respect to the number of GalNAc-T2 potential acceptor sites in MUC1^{4TR}. First, we used the corresponding molar equivalents of UDP-GalNAc to glycosylate 4 out of the 20 putative glycosylation sites available at MUC1^{4TR}. The quantification of the cross-peak volumes in the $^1\text{H}/^{15}\text{N}$ -HSQC allowed assessing that, while ~54% of the T¹⁵ residues were glycosylated (roughly two TR domains), only ~7% of GalNAc-incorporation was observed at T³ (Figure S6). When the concentration of the UDP-GalNAc donor is sufficient to glycosylate 10 acceptor sites, all T¹⁵ residues were fully glycosylated, while T³ was glycosylated in only ~56% and S¹⁴ did not show any trace of glycosylation (Figure S6). These results clearly demonstrate that GalNAc-T2 first glycosylates the T¹⁵ residues and initiates the addition of GalNAc at T³ before completing the glycosylation of all T¹⁵ residues. These observations are in agreement with the conclusions reported for MUC1 peptide fragments.^{18,36} Therefore, the reaction at the T³ site is assisted by the binding of the lectin domain of the enzyme to a prior glycosylated T¹⁵, evidencing that GalNAc-T2 displays a long-range N-terminal preference. To support this observation, MD simulations of 0.5 μs of GalNAc-T2 complexed with UDP-GalNAc/Mn²⁺ and MUC1 monoglycopeptide GVTSAPDTR-PAPGST*APPA (1) were performed (underlined is the acceptor site and * indicates the glycosylated amino acid). Initially, we run extensive unrestrained MD simulations on the complexes. However, under these conditions and due to the flexibility of the protein and the glycopeptides, the residue to be glycosylated was not properly oriented in the catalytic domain, observing that the peptide fragment close to this domain explored additional areas of the protein. Therefore, a distance restriction between the oxygen of the hydroxyl group of the reactive Ser/Thr residue and C1 of GalNAc in UDP-

GalNAc (distance O–C1 < 4.5 Å) was set in all MD simulations to maintain the correct orientation of the peptides in complex with the enzymes. The putative 3D structure of complex GalNAc-T2/1 was then generated using as starting coordinates the X-ray structure of a complex of GalNAc-T2 in an extended conformation (PDB entry 2FFU) and glycopeptide 1 (Figure 3A). The GalNAc at T¹⁵* of glycopeptide 1 was bound to the lectin domain. The complex GalNAc-T2/1 was stable throughout the MD trajectory (Figure S7A) suggesting that glycosylation at T³ can be lectin assisted, as inferred from the NMR experiments. According to the MD simulations, the methyl group of T³ is engaged in a CH– π interaction with the aromatic ring of F372 (distance 4.8 ± 0.4 Å, Figure 3A). Preference for glycosylation of Thr over Ser residues by GalNAc-T2 was previously observed.^{46,47} In this context, the stabilizing interaction between T³ and F372 is a structural feature that can explain why GalNAc-T2 prefers to glycosylate T³ rather than S⁴.

Finally, with UDP-GalNAc donor concentration enough to glycosylate all putative 20 acceptor sites, it was possible to reach full T¹⁵ and T³ glycosylation; however, only 45% of S¹⁴ was glycosylated (Figure S6). These data indicate that glycosylation at S¹⁴ is only initiated when all T¹⁵ and T³ residues have already been glycosylated. Moreover, the severe impairment of S¹⁴ glycosylation by the GalNAc^{D458R} mutation suggests that the S¹⁴ glycosylation might be promoted by a prior binding of the glycosylated T³ of the following TR to the lectin domain, a mechanism compatible with a long-range N-terminal preference. This lectin dependence to glycosylate S¹⁴ suggests that the glycosylation of the S¹⁴ located at TR4 of MUC1^{4TR} (TR4 S¹⁴) should have a considerably slower rate than the glycosylation of the other three S¹⁴ acceptor sites. The absence of a fifth tandem repeat in our construct precludes the glycosylation of TR4 S¹⁴ in a lectin-dependent manner. In fact, under an excess of UDP-GalNAc, GalNAc-T2 only takes 20 min (fast glycosylation) to glycosylate the three S¹⁴ residues (11 GalNAc units are attached to MUC1^{4TR}, providing a mass of 9831 *m/z*; Figure S3), when the last S¹⁴ residue requires 24 h (exhaustive glycosylation) to be glycosylated (note that at 16 h, the last S¹⁴ is still not glycosylated since only 85% of the S¹⁴ residues are glycosylated, data not shown). Due to the degeneracy of S¹⁴ cross peaks in MUC1^{4TR}, it was not possible to assure with absolute certainty that the last S¹⁴ to be glycosylated corresponds to TR4 S¹⁴. However, considering the lectin dependence of GalNAc-T2 to glycosylate S¹⁴, it is tempting to speculate that the last S¹⁴ residue to be glycosylated should be located at the TR4 of MUC1^{4TR}.

To understand the molecular basis of S¹⁴ glycosylation, we performed guided 0.5 μs MD simulations of GalNAc-T2 complexed with UDP-GalNAc/Mn²⁺ and a MUC1 diglycopeptide PAPGST*APPAHGVT*SA (2) (Figure 3B). In this diglycopeptide, the relative position of S at GST*AP and glycosylated T* at GVT*SA are equivalent to S¹⁴ in one tandem repeat and T³* in the following tandem repeat of MUC1^{4TR}. Therefore T* in GVT*SA (T³* in a TR+1 of MUC1^{4TR}) is bound to the lectin domain to perform glycosylation at S in the GST*AP motif (S¹⁴ in a TR of MUC1^{4TR}). The computational study shows that the distance of the hydroxyl group of S at GST*AP (equivalent to S¹⁴) to C1 of UDP-GalNAc (distance <4.5 Å) is stable through the whole trajectory (Figures 3B and S7B). A crucial hydrogen bond (38% populated along the MD trajectory) between the carbonyl group of GalNAc of the adjacent glycosylated T* in




Figure 4. Glycosylation of MUC1^{4TR-T³S¹⁴*T¹⁵*} by GalNAc-T4 and the GalNAc-T4^{D459H} mutant. Table S5 details the experimental conditions. (A) Scheme of the MUC1^{4TR-T³S¹⁴*T¹⁵*} glycosylation event catalyzed by GalNAc-T4. The glycosylation sites are displayed in red. (B,D) ¹H/¹⁵N-HSQC of the MUC1^{4TR-T³S¹⁴*T¹⁵*} product in the presence of GalNAc-T4 (B) and GalNAc-T4^{D459H} (D). In black, only with MnCl₂, and in magenta and blue, after addition of UDP-GalNAc (in excess) at 2 and 24 h, respectively. The arrows indicate CSP that occurs as result of GalNAc additions at S⁴ (gray arrows) and at T⁸ (yellow arrows). The * labeling in the amino acid on the spectra indicates glycosylation. (C,E) Time course of relative % of glycosylation at S⁴ and T⁸ residues by GalNAc-T4 (C) and GalNAc-T4^{D459H} (E) (see the Experimental Section for details).

GST*AP (T^{15*} in a TR of MUC1^{4TR}) and the NH of the sidechain of W265 could be behind the stable and short distance between the S residue at GST*AP (S^{14} in a TR of MUC1^{4TR}) and UDP-GalNAc and should be an essential feature for the reaction to occur. In conclusion, our data pinpoints the N-terminal preference of GalNAc-T2 to glycosylate T^3 and S^{14} in MUC1^{4TR} through a mechanism assisted by the lectin domain.

NMR Analysis of ^{15}N -MUC1^{4TR} Glycosylation by the GalNAc-T3 Isoform and Its Comparison to that of GalNAc-T2

Although GalNAc-T3 glycosylates the same three MUC1 glycosylation sites found for GalNAc-T2, the order of glycosylation is different.³⁶ To compare the glycosylation process of MUC1^{4TR} by GalNAc-T3 and its corresponding GalNAc-T3^{D517H} mutant, we used the same methodology as described above for GalNAc-T2. GalNAc-T3 glycosylates first T^3 , then T^{15} , and finally S^{14} . Interestingly, after 20 min of reaction time, three out of four T^3 and T^{15} residues were glycosylated, corresponding to the three TRs, while less than 5% of S^{14} was glycosylated (Figure S8). After 90 min, glycosylation at T^3 and T^{15} residues was completed (100%). In contrast, for GalNAc-T3^{D517H}, after 20 min (fast glycosylation), only glycosylation at T^3 residues occurred (65%, corresponding to ~3 TR domains), while after 90 min, glycosylation at T^3 and T^{15} residues reached 100 and ~25%, respectively. After 24 h (exhaustive glycosylation), the glycosylation of S^{14} residues is ~70% in the presence of GalNAc-T3, while it is not detected in the presence of GalNAc-T3^{D517H} (Figure S8). Moreover, even after 72 h, glycosylation at S^{14} is still not completed (*ca.* 88%) while remaining undetected with the mutant (Figure S8). Therefore, while GalNAc-T3-mediated glycosylation at T^3 is only dependent on the natural affinity of the catalytic domain toward the GVTS region of MUC1, remote glycosylation at T^{15} and S^{14} is clearly assisted by the lectin domain. The observed long-range glycosylation by GalNAc-T3 agrees with the contribution of the lectin domain to the recognition of the GalNAc moiety located at T^3 , which promotes the GalNAc transfer to the T^{15} and S^{14} residues. This result is in agreement with the preference of GalNAc-T3 to glycosylate C-terminal remote acceptor sites prior to N-terminal glycosites.^{28,48}

Interestingly, a comparison of the glycosylation rates on MUC1^{4TR} between both isoforms shows that GalNAc-T3 is slower to glycosylate S^{14} than GalNAc-T2. These differences in the ability to incorporate GalNAc at S^{14} agrees with a previous report using kinetics studies assisted by MS that demonstrate that GalNAc-T2 glycosylates S^{14} faster than GalNAc-T1/T3 isoforms.³⁶ These differences are likely, not due to the different directionality of the lectin-mediated long-range glycosylation preferences since GalNAc-T1 and T2 isoforms display the same type of long-range glycosylation preferences.^{15,26,27} More likely, the divergent rates for the GalNAc-T3 and T2 isoforms are due to how the prior glycosylated sites affect the conformation and/or presentation of each TR domain, which determine the recognition by the catalytic domain of these isoforms. To evaluate this hypothesis, the conformation of MUC1^{4TR}- T^3 * T^{15} * in solution was investigated. The preparation of MUC1^{4TR}- T^3 * T^{15} * and its NMR and MS characterization are described in SI and displayed in Figure S9 and Table S3, respectively. The ^1H , ^{15}N -CSP of MUC1^{4TR}- T^3 * T^{15} * using naked MUC1^{4TR} as reference (Figure S10A) and the

comparison of the ^{13}C -CSI of MUC1^{4TR}- T^3 * T^{15} * with the naked MUC1^{4TR} were scrutinized (Figure S10B), showing that glycosylation at T^3 and T^{15} modulates the conformation of the surrounding residues. Specifically, the ^{13}C -CSI difference for S^{14} before and after glycosylation at T^{15} is rather large (Figure S10B) and suggests a decrease in the flexibility of S^{14} and a shift toward an extended conformation. Previously, Kinarsky *et al.*³⁹ suggested that the GS¹⁴T¹⁵A region changes from polyproline II-like toward an inverse γ -turn conformation upon glycosylation of T^{15} in MUC1 short glycopeptides. In fact, similar behavior has been previously observed also by us.⁴⁹ Based on this, it is tempting to propose that GalNAc-T2 has more affinity to the inverse γ -turn conformation of GS¹⁴T¹⁵*A and preferentially glycosylates the S^{14} acceptor site in this region than GalNAc-T3 and T1.

NMR Analysis of ^{15}N -MUC1^{4TR} Glycosylation by the GalNAc-T4 Isoform—Differences in S^4 and T^8 Glycosylation

GalNAc-T4 isoform is the only isoform identified that is able to glycosylate *in vitro* S^4 and T^8 in the MUC1^{4TR} structure.^{10,22,50,51} To further explore this event by NMR, we used the GalNAc-glycopeptide MUC1^{4TR}- T^3 * S^{14} * T^{15} * as the acceptor substrate (Figure 4). The CSP analysis is now focused on the effects of glycosylation at S^4 and T^8 (Figures 4B,C and S11). Glycosylation at T^8 induces a strong CSP (0.51 ppm, yellow arrow) at this residue along with others at R^9 (0.29 ppm, yellow arrow) and D^7 (0.10 ppm, yellow arrow) (Figures 4B and S11). In contrast, glycosylation at S^4 only provides moderate CSP at the corresponding S^4 (0.18 ppm, gray arrow) and A^5 (0.07 ppm, gray arrow) (Figures 4B and S11).

After 5 h of reaction time, S^4 and T^8 residues were glycosylated 40 and 24%, respectively, while after 48 h, ~75–80% of the S^4 and T^8 residues were glycosylated (Figure 4C) indicating that one glycosylation is lacking for both S^4 and T^8 . Nevertheless, the mass analysis of the product purified by HPLC displayed two masses, ~11,255.6 *m/z* and ~11,458.6 *m/z*, which corresponded to two different glycosylated MUC1^{4TR} that account for 18 and 19 GalNAc units, respectively (Figure S12).

The NMR assignment of the major product with 19 glycosylated sites out of 20 allowed us to unequivocally identify that the TR1 S^4 is not glycosylated (Figures S12 and Table S4), suggesting that glycosylation of S^4 by GalNAc-T4 is lectin domain-assisted and requires the binding of the lectin domain to a GalNAc moiety present at the preceding MUC1 TR domain. This is compatible with a long-range C-terminal preference. In this scenario, the GalNAc-T4 lectin domain should bind the GalNAc moiety at either S^{14} or T^{15} at a given TR of MUC1^{4TR}- T^3 * S^{14} * T^{15} * product to drive the glycosylation process onto S^4 of the following TR domain. The analysis with the GalNAc-T4^{D459H} lectin mutant was also performed (Figures 4D,E and S13). While S^4 glycosylation was completely abolished in the presence of the mutant, a small fraction of T^8 glycosylation occurred very slowly and progressively with the time of incubation (after 24 h, 20% of the T^8 residues were glycosylated, reaching a maximum of 35% after 72 h). These results confirm the importance of the lectin domain in the glycosylation of S^4 and demonstrate that the T^8 glycosylation process is also severely affected in the presence of GalNAc-T4^{D459H} (Figure 4), strongly suggesting that the glycosylation of T^8 also takes place in a lectin domain-assisted manner. The results described above for MUC1^{4TR}-




Figure 5. Putative 3D models obtained by MD simulations to explain the glycosylation of S⁴ and T⁸ in MUC1^{4TR} catalyzed by GalNAc-T4. (A,B) Glycosylation of S⁴. A 3D view of two complexes of GalNAc-T4/3 derived from a representative frame obtained from 0.5 μ s of MD simulation in which GalNAc-T^{15*} (A) or GalNAc-S^{14*} (B) is accommodated in the lectin domain, respectively. A hydrogen bond between the carbonyl group of GalNAc at T^{3*} and the main chain of F284, populated in 30% during MD trajectory, is established in (A). (C) Glycosylation of T⁸. 3D view of GalNAc-T4/3 derived from a representative frame obtained from 0.5 μ s of MD simulation in which GalNAc-T^{15*} (C) is accommodated in the lectin domain. The methyl group of T⁸ is involved in CH- π interactions with the aromatic ring of F364 (6.3 ± 0.5 Å). In all structures, the catalytic domain is shown in light blue and the lectin domain in dark blue. In the glycopeptides, the GalNAc units are shown in yellow sticks and the glycopeptide backbone in green ribbon. UDP-GalNAc is displayed in gray sticks and Mn²⁺ is shown as a cyan sphere. Relevant residues of the glycopeptides and some residues of catalytic domain are also shown as sticks and in a different color. Schematic representation of the glycosylation process at S⁴ and T⁸ of MUC1^{4TR} catalyzed by GalNAc-T4 is also displayed. These models highlight the C-terminal preference of GalNAc-T4.

T^{3*}S^{14*}T^{15*} indicate that S⁴ and T⁸ are glycosylated by GalNAc-T4 in a lectin-dependent manner. However, it was not inferred whether the lectin domain binds at the previous glycosylated S¹⁴ or T¹⁵ to drive the glycosylation events at S⁴ or T⁸. Therefore, to uncover how the glycosylation takes place at those acceptor sites, 0.5 μ s MD simulations were performed using 3D models of GalNAc-T4 complexed with UDP-GalNAc/Mn²⁺ and MUC1 triglycopeptide PGS*T*AP-PAHGVT*SAPDTRPAPG (3). In these models, the lectin domain was bound to either GalNAc at S or T of the GSTA region (equivalent to S^{14*} or T^{15*} of a preceding TR of MUC1^{4TR}) to drive glycosylation at S in GVT*SA (equivalent to S⁴ in a TR of MUC1^{4TR}) or T in PDTRP (equivalent to T⁸ in a TR of MUC1^{4TR}) (Figures 5 and 6).

According to 0.5 μ s guided MD simulations (with a distance between the hydroxyl group of S⁴ and C1 of UDP-GalNAc <4.5 Å), glycosylation of S⁴ is feasible when the lectin domain accommodates either GalNAc at T^{15*} (Figures 5A and S14A) or S^{14*} (Figures 5B and S14B). In both possibilities, the GalNAc moiety at T^{3*} interacts through transient hydrogen bonds with several residues of the catalytic domain. Nevertheless, when T^{15*} is bound to the lectin domain, a hydrogen bond populated around 30% along the MD trajectory is established between the carbonyl group of GalNAc at T^{3*} and

the NH of F284 of the enzyme (Figure 5A), which could allow for the proper orientation of S⁴ for the glycosylation. The MD simulations (with a distance between the hydroxyl group of T⁸ and C1 of UDP-GalNAc <4.5 Å) indicate that T⁸ is not properly located in the active site when S^{14*} is bound to the lectin domain, suggesting that the glycosylation in T⁸ likely occurs with the assistance of the lectin domain bound preferentially at T^{15*} of the preceding TR domain of MUC1^{4TR} (Figures 5C and S14C). In this latest binding mode, GalNAc at T^{3*} forms transient hydrogen bonds with the enzyme, and the methyl group of T⁸ is involved in a CH- π interaction with the aromatic ring of F364 (6.3 ± 0.5 Å), which may favor the correct orientation of this residue for optimal glycosylation (Figure 5C). In the long-range glycosylation, the T⁸ residue in MUC1^{4TR} is located 13 residues apart in the sequence from the glycosylated T^{15*} of the preceding TR, while S⁴ is much closer (9 residues), being the shorter distance within the optimal range for GalNAc-T4 catalysis. Previous studies demonstrate that GalNAc-T4 prefers to glycosylate acceptor residues located 7–11 residues away from a prior glycosite,^{28,48} which might explain the slight preference of GalNAc-T4 to glycosylate S⁴ versus T⁸ residues in MUC1^{4TR}.




Figure 6. Putative 3D models obtained by MD simulations to explain the glycosylation of MUC1^{4TR} catalyzed by GalNAc-T4. (A) 3D view of two complexes of GalNAc-T4/4 derived from a representative frame obtained from 0.5 μ s of MD simulations. Hydrophobic interaction between the methyl groups of T³ and A368 (4.7 \pm 0.6 Å) is observed, which may favor the proper orientation of this threonine residue for efficient glycosylation. (B) Close view of STD–NMR experiments for short diglycopeptide (MUC1-T³*T¹⁵*). STD–NMR experiments were performed at 298 K in the presence of UDP (75 μ M) and MnCl₂ (150 μ M) with a molar ratio of 75:1 diglycopeptide:GalNAc-Ts (GalNAc-T2, -T3, and -T4). The reference spectrum (¹H NMR) is displayed in blue, while the STD spectrum (STD) is displayed in black for GalNAc-T4, in dark gray for GalNAc-T3, and in gray for GalNAc-T2. (C) Close view of GalNAc-T4/3 derived from a representative frame obtained from 0.5 μ s of MD simulation in which GalNAc-T¹⁵ is accommodated in the lectin domain. Interestingly, D⁷ is engaged in a high populated salt-bridge interaction with R372 (71 and 88%) (dashed line). In the structures, the catalytic domain is shown in light blue and the lectin domain in dark blue. In the glycopeptides, the GalNAc units are shown in yellow sticks and the glycopeptide backbone in green ribbon. UDP-GalNAc is displayed in gray sticks and Mn²⁺ is shown as a cyan sphere. Relevant residues of the glycopeptides and some residues of catalytic domain are also shown as sticks and in a different color. Relevant residues of the glycopeptide and the protein are also shown as sticks. Schematic representation of the glycosylation process at T³ and T⁸ of MUC1^{4TR} catalyzed by GalNAc-T4 is also displayed. These models highlight the C-terminal preference of GalNAc-T4.

Role of the GalNAc-T4 Catalytic Domain in Glycosylating MUC1^{4TR}

Glycosylation of S⁴ and T⁸ of MUC1 by GalNAc-T4 is a process dominated by its lectin long-range preference. However, other features may modulate the GalNAc-T4 activity: (1) conformational changes in MUC1 structure induced by the presence of GalNAc in adjacent sites of the glycosylation site and (2) the preference of the catalytic domain toward specific peptide regions and/or glycopeptides (catalytic-domain-dependent glycosylation). To evaluate how the prior glycosites may affect the subsequent GalNAc-T4 glycosylation reactions, additional experiments were performed using MUC1^{4TR}-T³*T¹⁵* (Figure S15) and MUC1^{4TR}-T¹⁵* (Figure S16) as acceptor substrates. In the presence of MUC1^{4TR}-T³*T¹⁵*, GalNAc-T4 prefers to glycosylate S⁴ (GVT*SA) over S¹⁴ (GST*AP) or T⁸ (PDTRP) (Figure S15). After 24 h, 60% of S⁴, ~15% in S¹⁴, and 8% of T⁸ residues were glycosylated. This result supports that GalNAc-T4 lectin domain likely binds to the GalNAc moiety at T¹⁵* of a preceding TR to drive glycosylation at S⁴ of the following TR. By comparing the GalNAc incorporation in MUC1^{4TR}-T³*T¹⁵* versus MUC1^{4TR}-T³*S¹⁴*T¹⁵* by GalNAc-T4

(Figures 4B and S15), differences in glycosylation of the two acceptor substrates were observed. The slower glycosylation of MUC1^{4TR}-T³*T¹⁵* versus MUC1^{4TR}-T³*S¹⁴*T¹⁵* indicates that S¹⁴ glycosylation must be priorly glycosylated by GalNAc-T2/-T3 to maximize the full glycosylation of MUC1 and exemplify the importance of the complementary and hierarchical functions of GalNAc-Ts during the O-glycosylation process of MUC1. Interestingly, when using the MUC1^{4TR}-T¹⁵* as the acceptor substrate, GalNAc-T4 first glycosylates T³ over S⁴, both located at the GVTSA motif (Figure S16). MD simulations performed on the complex between the enzyme and a glycopeptide as an acceptor substrate, PGST*APPAHGVTSAPDTRPAPG (4), show that a hydrophobic contact between the methyl group of T³ and A368 may help to align this residue for glycosylation, which could be the reason for the preference of T³ related to S⁴ (Figure 6A). These two residues are in the same region of MUC1 peptide (sharing a similar chemical environment), and in both cases, glycosylation is assisted by the lectin domain (T³ and S⁴ are eight and nine residues away from T¹⁵*, respectively). Thus, the less efficient capacity of GalNAc-T4 to glycosylate Ser than Thr residues is likely due to the poorer

binding to the catalytic domain, as shown by our calculations. In fact, as verified by other GalNAc-Ts (e.g., T2 and T3), also for GalNAc-T4, a preference was shown to glycosylate Thr over Ser in random peptides.⁴⁶ On the other hand, from the conformational perspective, the ¹³C-CSI analysis of MUC1^{4TR}-T³*S¹⁴*T¹⁵* versus the naked MUC1^{4TR} (Figure S4B) clearly indicates that glycosylation at T³ affects the conformation of the S⁴ neighboring residue. Our ¹³C-CSI analysis shows that the presence of GalNAc at T³ increases the population of β -strand-like conformation at the GVTSA motif and agrees with the conformational analysis of MUC1 short glycopeptides previously reported.^{39,52} This alteration in the local conformation of S⁴ might also play a role in the glycosylation of S⁴ by GalNAc-T4. Additionally, as previously mentioned, the presence of the neighboring GalNAc at T³* contributes to additional interactions with the catalytic domain favoring the glycosylation of the contiguous S⁴ (Figure 6A, h-bond between the carbonyl of GalNAc and NH of F284). The presence of GalNAc at T³* as a prerequisite for glycosylation of S⁴ at the GVTSA region of MUC1 agrees with the absence of glycosylation at S⁴ observed for the MUC1 glycopeptide mutant in which T³ was mutated by a valine residue.²² The existence of this short-range glycosylation preference by GalNAc-T4 was previously reported by us using short model glycopeptides.²³

In the case of T⁸ glycosylation (located at the PDTRP sequence), GalNAc-T4 glycosylates this residue in the same proportion as S⁴ at GVT*SA when the acceptor substrate is MUC1^{4TR}-T³*S¹⁴*T¹⁵* (Figure 4B). In this acceptor substrate, the glycosylation of T⁸ and S⁴ seems to occur independently. In addition, a small fraction of T⁸ glycosylation is detected in the case of the mutant GalNAc-T4^{D458H} (Figure 4D,E). Surprisingly, in the presence of MUC1^{4TR}-T³*T¹⁵*, T⁸ glycosylation occurs in the same proportion as that of S¹⁴ at GST*AP. CSP and CSI analyses between MUC1^{4TR}-T³*T¹⁵ and MUC1^{4TR}-T³*S¹⁴*T¹⁵* (Figure S17) illustrate that S¹⁴ displays negligible effects on the chemical environment and conformation of PDTRP MUC1^{4TR}, suggesting that GalNAc-S¹⁴ does not modulate the glycosylation of T⁸ at the PDTRP motif. Previous studies suggest that the DTR fragment at MUC1 falls in the inverse γ -turn geometry's area of the extended conformations in the Ramachandran plot.³⁹ This conformation is likely important for the recognition of the GalNAc-T4 catalytic domain according to our results, explaining why the GalNAc-T4^{D459H} still glycosylates T⁸. Overall, these data suggest that GalNAc-T4 has some natural preference to glycosylate the PDTRP region in the absence of a functional lectin domain. To further confirm this hypothesis, we performed saturation transfer difference (STD)-NMR experiments with GalNAc-T2, -T3, and -T4 using a short diglycopeptide containing one TR (MUC1-T³*T¹⁵*) (Figures 6B and S18).

STD signals corresponding to side chain protons of D⁷ and R⁹ at the PDTRP region were only detected in the case of GalNAc-T4. No STD signals were observed within the PDTRP sequence for either GalNAc-T2 or T3 isoform, demonstrating the specific recognition of the catalytic domain of GalNAc-T4 toward the inverse γ -turn PDTRP region around T⁸. Interestingly, the T⁸ at the PDTRP region is the only acceptor site at MUC1 flanked by two charged residues, which implies that these charge residues are likely specifically recognized by the GalNAc-T4 catalytic domain. Indeed, the analysis of the MD simulations on glycopeptide 3 in complex

with GalNAc-T4 (Figure 6C) points out a significant salt-bridge between D7 at the PDTRP region of 3 and the R372 residue of GalNAc-T4 (high populated salt-bridge around 70–90% along the MD trajectory). Remarkably, this residue of GalNAc-T4 is not conserved in GalNAc-T2 and T3, which could be a mechanism of this enzyme to recognize the most immunogenic domain of MUC1. In short, GalNAc-T4 needs its lectin domain to efficiently glycosylate both S⁴ and T⁸ residues. However, other structural features such as the conformation and composition of the peptide sequence together with GalNAc-T4 short-range glycosylation preference finely tune MUC1 O-glycosylation by GalNAc-T4.

Elucidation of the GalNAc-T Isoforms Glycosylating S4 and T8 Residues in Cells

It is well accepted that GalNAc-T1, -T2, and -T3 are the major isoforms glycosylating *in vitro* and *in vivo* T³, S¹⁴, and T¹⁵ in MUC1.^{36,51} Although it is well accepted that GalNAc-T4 is capable of glycosylation S⁴ and T⁸ *in vitro*,^{10,22} it is not clear whether GalNAc-T4 glycosylates these acceptor sites *in vivo*. One study showed that overexpression of GalNAc-T4 in CHO cells appears to enhance T⁸ glycosylation at the PDTRP sequence,⁵⁰ and more recently, knock out (KO) of the *GALNT4* gene in HEK293 cells revealed selective loss of the O-glycan at the Thr in the PDTR motif (T⁸) of the MUC1 tandem repeats when a reporter construct with seven TRs (MUC1^{7TRs}) was expressed.⁵³ Since loss of GalNAc-T4 in HEK293 cells did not appear to affect O-glycosylation at the Ser in VTSA (S⁴), we wanted to further confirm this finding using an alternative bottom-up MS approach. The traditional bottom-up MS strategy for MUC1 TRs is based on use of endo-Asp digestion with cleavage in the PDTR motif, and an O-glycan at the Thr site may bias the cleavage and result of the analysis. We therefore took advantage of the glycomucase BT4244 from *Bacteroides thetaiotaomicron* previously shown to cleave N-terminal to Ser or Thr residues with an α -GalNAc O-glycan attached.⁵⁴ We recently demonstrated that this enzyme efficiently cleaves the Tn glycoform of the MUC1^{7TRs} reporter with predominant cleavage in between the O-glycan diads at the VT³S⁴A and GS¹⁴T¹⁵A motifs.⁵⁵ The BT4244 glycomucase efficiently cleaves the MUC1 reporter expressed in HEK293^{KO} *COSMC/GALNT4* with O-glycosylation limited to α -GalNAc O-glycans,⁵⁶ and we could confirm that the T⁸ in PDT⁸R was not glycosylated in HEK293 cells with the KO of *GALNT4*, while the S⁴ in VT³S⁴A was efficiently O-glycosylated (Figure 7). This suggests that S⁴ may be O-glycosylated by other GalNAc-T isoforms in HEK293 cells despite those past studies with *in vitro* enzyme assays that have failed to identify such activities. Nevertheless, our results clearly confirm that GalNAc-T4 is the only isoform capable of glycosylating T⁸ in PDTRP *in vivo* and completing the O-glycosylation of the five glycosites in MUC1.

CONCLUSIONS

Nature has deployed a vast number of GalNAc-Ts isoforms to deal with very complex protein substrates, like mucins, ensuring their function in living organisms. Deciphering the molecular determinants behind the glycosylation of mucins by GalNAc-Ts is crucial to understand the mode of action and the fine specificity of GalNAc-Ts toward mucins and is essential to explain how the complex mucinome is created. Herein, we provide NMR-derived structural insights into the full process of MUC1 O-glycosylation involving consecutive steps




Figure 7. Exclusive contribution of GALNT4 in cells to MUC1 O-glycosylation at T8 in the PDTR site. Deconvoluted bottom-up analysis of the isolated MUC1^{7TR} reporter produced in HEK293^{KO} COSMC/GALNT4 using BT4244 glycomucase digestion. The BT4244 glycomucase digestion sites are indicated by arrows. All identified glycosites by MS/MS ETD analysis are illustrated with numbers assigned to each peak based on decreasing abundance. Only peaks with abundance above 10% were assigned.

catalyzed by GalNAc-T2/T3 and T4. Apart from confirming the order by which GalNAc-T2/T3 and T4 glycosylate the different acceptor sites in the MUC1 TRs, we also demonstrate that the lectin-mediated functions of these GalNAc-Ts are strongly involved in all subsequent glycosylation reactions after the first GalNAc-addition. The lectin domain strongly contributes to define the site, the order, and the orientation of glycosylation in a complex mucin template such as MUC1^{4TR} (Figure 8). Since GalNAc-T1 and T2 are rather ubiquitously expressed and are the workhorses to initiate the

O-glycoproteome in cells,³³ we predict that these together with GalNAc-T3 (and its close paralog GalNAc-T6) in cells will drive the initial O-glycosylation of MUC1, while GalNAc-T4 serves to complete glycosylation. This hierarchy in GalNAc-Ts seems to be fundamental to obtain a fully glycosylated MUC1 product. Furthermore, our NMR data show local conformational changes in TRs during the incorporation of GalNAc residues, which may correlate with the preferences of some of GalNAc-Ts toward specific (glyco)peptides conformations (e.g., GalNAc-T2 glycosylates Ser residues at the glycosylated GST*AP better than GalNAc-T3/T1).

We have also demonstrated that GalNAc-T4 is the unique isoform responsible to glycosylate the PDTRP region of MUC1 in cells, a process that depends significantly on its lectin domain and to a lesser extent on the exquisite affinity of its catalytic domain to the PDTRP motif. The inverse γ -turn conformation adopted by the DTR sequence in solution³⁹ concomitantly with a specific salt-bridge that can be established between the Asp of PDTRP and the Arg of the enzyme might be the main structural features behind GalNAc-T4 preference toward this region. The PDTRP motif of the MUC1 structure is highly immunogenic, and most of the anti-MUC1 antibodies specifically target this peptide region at MUC1.^{57,58} Thus, the evidence that only GalNAc-T4 isoform can glycosylate this region *in vivo* points to the importance of the regulation of GalNAc-T4 expression in the glycosylation of the PDTRP region of MUC1. In summary, this work highlights the relevance of the lectin domain in the MUC1 glycosylation process, which clearly pinpoints the lectin domain of GalNAc-Ts as an attractive target for the rational design of inhibitors toward GalNAc-Ts. Finally, the NMR methodology and protocols applied herein are robust and can be applied to determine the specificity of other glycosyltransferases, such as C1GalT1 and ST6GalNAc-I, toward MUC1 or other mucin acceptor substrates.




Figure 8. Overview of the proposed mechanism for MUC1 GalNAc O-glycosylation by GalNAc-T2, -T3, and -T4.

■ ASSOCIATED CONTENT**SI Supporting Information**

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/jacsau.1c00529>.

(PDF)

■ AUTHOR INFORMATION**Corresponding Authors**

Ramon Hurtado-Guerrero – Institute for Biocomputation and Physics of Complex Systems (BIFI), Laboratorio de Microscopias Avanzadas (LMA), University of Zaragoza, 50018 Zaragoza, Spain; Copenhagen Center for Glycomics, Department of Cellular and Molecular Medicine, University of Copenhagen, Copenhagen DK-2200, Denmark; Fundación ARAID, 50018 Zaragoza, Spain; orcid.org/0000-0002-3122-9401; Email: rhurtado@bifi.es

Filipa Marcelo – Associate Laboratory i4HB—Institute for Health and Bioeconomy, NOVA School of Science and Technology and UCIBIO, Department of Chemistry, Faculdade de Ciências e Tecnologia, Universidade NOVA de Lisboa, 2829-516 Caparica, Portugal; orcid.org/0000-0001-5049-8511; Email: filipa.marcelo@fc.unl.pt

Authors

Helena Coelho – Associate Laboratory i4HB—Institute for Health and Bioeconomy, NOVA School of Science and Technology and UCIBIO, Department of Chemistry, Faculdade de Ciências e Tecnologia, Universidade NOVA de Lisboa, 2829-516 Caparica, Portugal; CIC bioGUNE, Basque Research and Technology Alliance (BRTA), 48170 Derio, Spain; Department of Organic Chemistry II, Faculty of Science & Technology, University of the Basque Country, Leioa 48940 Bizkaia, Spain; orcid.org/0000-0003-1992-8557

Matilde de las Rivas – Institute for Biocomputation and Physics of Complex Systems (BIFI), Laboratorio de Microscopias Avanzadas (LMA), University of Zaragoza, 50018 Zaragoza, Spain

Ana S. Grosso – Associate Laboratory i4HB—Institute for Health and Bioeconomy, NOVA School of Science and Technology and UCIBIO, Department of Chemistry, Faculdade de Ciências e Tecnologia, Universidade NOVA de Lisboa, 2829-516 Caparica, Portugal

Ana Diniz – Associate Laboratory i4HB—Institute for Health and Bioeconomy, NOVA School of Science and Technology and UCIBIO, Department of Chemistry, Faculdade de Ciências e Tecnologia, Universidade NOVA de Lisboa, 2829-516 Caparica, Portugal

Cátia O. Soares – Associate Laboratory i4HB—Institute for Health and Bioeconomy, NOVA School of Science and Technology and UCIBIO, Department of Chemistry, Faculdade de Ciências e Tecnologia, Universidade NOVA de Lisboa, 2829-516 Caparica, Portugal

Rodrigo A. Francisco – Associate Laboratory i4HB—Institute for Health and Bioeconomy, NOVA School of Science and Technology and UCIBIO, Department of Chemistry, Faculdade de Ciências e Tecnologia, Universidade NOVA de Lisboa, 2829-516 Caparica, Portugal

Jorge S. Dias – Associate Laboratory i4HB—Institute for Health and Bioeconomy, NOVA School of Science and Technology and UCIBIO, Department of Chemistry,

Faculdade de Ciências e Tecnologia, Universidade NOVA de Lisboa, 2829-516 Caparica, Portugal

Ismael Compañón – Departamento de Química, Centro de Investigación en Síntesis Química, Universidad de La Rioja, E-26006 Logroño, Spain

Lingbo Sun – Copenhagen Center for Glycomics, Department of Cellular and Molecular Medicine, University of Copenhagen, Copenhagen DK-2200, Denmark

Yoshiki Narimatsu – Copenhagen Center for Glycomics, Department of Cellular and Molecular Medicine, University of Copenhagen, Copenhagen DK-2200, Denmark

Sergey Y. Vakhrushev – Copenhagen Center for Glycomics, Department of Cellular and Molecular Medicine, University of Copenhagen, Copenhagen DK-2200, Denmark; orcid.org/0000-0002-0418-5765

Henrik Clausen – Copenhagen Center for Glycomics, Department of Cellular and Molecular Medicine, University of Copenhagen, Copenhagen DK-2200, Denmark

Eurico J. Cabrita – Associate Laboratory i4HB—Institute for Health and Bioeconomy, NOVA School of Science and Technology and UCIBIO, Department of Chemistry, Faculdade de Ciências e Tecnologia, Universidade NOVA de Lisboa, 2829-516 Caparica, Portugal; orcid.org/0000-0002-0720-2751

Jesús Jiménez-Barbero – CIC bioGUNE, Basque Research and Technology Alliance (BRTA), 48170 Derio, Spain; Department of Organic Chemistry II, Faculty of Science & Technology, University of the Basque Country, Leioa 48940 Bizkaia, Spain; Ikerbasque, Basque Foundation for Science, 48009 Bilbao, Spain; Centro de Investigación Biomedica En Red de Enfermedades Respiratorias, 28029 Madrid, Spain; orcid.org/0000-0001-5421-8513

Francisco Corzana – Departamento de Química, Centro de Investigación en Síntesis Química, Universidad de La Rioja, E-26006 Logroño, Spain; orcid.org/0000-0001-5597-8127

Complete contact information is available at:

<https://pubs.acs.org/doi/10.1021/jacsau.1c00529>

Author Contributions

H.C. expressed and purified the MUC1^{4TR} and performed and analyzed the NMR experiments. A.D., A.S.G., C.O.S., R.F., and J.S.D. helped on the expression and purification of MUC1^{4TR} and the assignment of glycosylated MUC1^{4TR} products. M.R. expressed and purified the enzymes and mutants. I.C. synthesized the glycopeptide. L.S., Y.N., and S.Y.V. performed the cell assays and MS analysis. F.C. performed the MDs simulations. F.M. designed the research. F.M. and R.H.G. wrote the article together with the main contributions of F.C., E.J.C., J.J.-B., Y.N., H.C., and H.C. All authors read and approved the final manuscript.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

The authors acknowledge Fundação para a Ciência e a Tecnologia (FCT-Portugal) for funding the projects, IF/00780/2015, PTDC/BIA-MIB/31028/2017, and UCIBIO project (UIDP/04378/2020 and UIDB/04378/2020), and Associate Laboratory Institute for Health and Bioeconomy—i4HB project (LA/P/0140/2020). The authors also thank FCT-Portugal for the PhD grant attributed to ASG (SFRH/

BD/140394/2018) and A.D. (PD/BD/142847/2018) under the PTNMRPhD Program (PD00065/2013) and for the Norma transitória DL 57/2016 Program Contract to J.S.D. The NMR spectrometers are part of the National NMR Facility supported by FCT-Portugal (ROTEIRO/0031/2013-PINFRA/22161/2016, cofinanced by FEDER through COMPETE 2020, POCI and PORL and FCT through PIDDAC). RHG thanks ARAID, the Spanish Ministry of Science, Innovation and Universities (BFU2016-75633-P and PID2019-105451GB-I00), and Gobierno de Aragón (E34_R17 and LMP58_18 to R.H.-G.) with FEDER (2014–2020) funds for “Building Europe from Aragón” for the financial support. F.C. thanks Agencia Estatal Investigación (AEI, Spain) (grant RTI2018-099592-B-C21). This project has received funding from the European Union’s Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie Grant Agreement no. 956544. J.J.-B. and H.C. thank EU for funding (Tollerant ITN, GA-642157). F.M. and J.J.-B. acknowledge the COST Action GLYCONanoProbes (CA18132). J.J.-B. also thanks the European Research Council for the financial support (ERC-2017-AdG, project number 788143-RECGLYCANMR), AEI (Spain, grant RTI218-094751-B-C21), and CIBER, an initiative of Instituto de Salud Carlos III (ISCIII), Madrid, Spain. H.C. acknowledges the Lundbeck Foundation and the Danish National Research Foundation (DNRF107).

ABBREVIATIONS

GalNAc-Ts	polypeptide GalNAc-transferases
MUC1 ^{4TR}	mucin1 with four tandem repeats
MUC1 ^{7TR}	mucin1 with seven tandem repeats
NMR	nuclear magnetic resonance
CSP	combined chemical shift perturbations
STD	saturation transfer difference

REFERENCES

- Bennett, E. P.; Mandel, U.; Clausen, H.; Gerken, T. A.; Fritz, T. A.; Tabak, L. A. Control of Mucin-Type O-Glycosylation: A Classification of the Polypeptide GalNAc-Transferase Gene Family. *Glycobiology* **2012**, *22*, 736–756.
- Gill, D. J.; Clausen, H.; Bard, F. Location, Location, Location: New Insights into O-GalNAc Protein Glycosylation. *Trends Cell Biol.* **2011**, *21*, 149–158.
- Hurtado-Guerrero, R.; Davies, G. J. Recent Structural and Mechanistic Insights into Post-Translational Enzymatic Glycosylation. *Curr. Opin. Chem. Biol.* **2012**, *16*, 479–487.
- Narimatsu, Y.; Joshi, H. J.; Nason, R.; Van Coillie, J.; Karlsson, R.; Sun, L.; Ye, Z.; Chen, Y.-H.; Schjoldager, K. T.; Steentoft, C.; Furukawa, S.; Bensing, B. A.; Sullam, P. M.; Thompson, A. J.; Paulson, J. C.; Büll, C.; Adema, G. J.; Mandel, U.; Hansen, L.; Bennett, E. P.; Varki, A.; Vakhrushev, S. Y.; Yang, Z.; Clausen, H. An Atlas of Human Glycosylation Pathways Enables Display of the Human Glycome by Gene Engineered Cells. *Mol. Cell* **2019**, *75*, 394–407.
- Schjoldager, K. T.; Narimatsu, Y.; Joshi, H. J.; Clausen, H. Global View of Human Protein Glycosylation Pathways and Functions. *Nat. Rev. Mol. Cell Biol.* **2020**, *21*, 729–749.
- White, T.; Bennett, E. P.; Takio, K.; Sørensen, T.; Bonding, N.; Clausen, H. Purification and CDNA Cloning of a Human UDP-N-Acetyl-Alpha-D-Galactosamine:Polypeptide N-Acetylgalactosaminyltransferase. *J. Biol. Chem.* **1995**, *270*, 24156–24165.
- Chefetz, I.; Kohno, K.; Izumi, H.; Uitto, J.; Richard, G.; Sprecher, E. GALNT3, a Gene Associated with Hyperphosphatemic Familial Tumoral Calcinosis, Is Transcriptionally Regulated by Extracellular Phosphate and Modulates Matrix Metalloproteinase Activity. *Biochim. Biophys. Acta, Mol. Basis Dis.* **2009**, *1792*, 61–67.
- Kato, K.; Jeanneau, C.; Tarp, M. A.; Benet-Pagès, A.; Lorenz-Depiereux, B.; Bennett, E. P.; Mandel, U.; Strom, T. M.; Clausen, H. Polypeptide GalNAc-Transferase T3 and Familial Tumoral Calcinosis: Secretion of Fibroblast Growth Factor 23 Requires O-Glycosylation. *J. Biol. Chem.* **2006**, *281*, 18370–18377.
- Khetarpal, S. A.; Schjoldager, K. T.; Christoffersen, C.; Raghavan, A.; Edmondson, A. C. Loss of Function of GALNT2 Lowers High-Density Lipoproteins in Humans, Nonhuman Primates, and Rodents. *Cell Metab.* **2016**, *24*, 234–245.
- Bennett, E. P.; Hassan, H.; Mandel, U.; Mirgorodskaya, E.; Roepstorff, P.; Burchell, J.; Taylor-Papadimitriou, J.; Hollingsworth, M. A.; Merckx, G.; van Kessel, A. G.; Eiberg, H.; Steffensen, R.; Clausen, H. Cloning of a Human UDP-N-Acetyl- α -D-Galactosamine:Polypeptide N-Acetylgalactosaminyltransferase That Complements Other GalNAc-Transferases in Complete O-Glycosylation of the MUC1 Tandem Repeat. *J. Biol. Chem.* **1998**, *273*, 30472–30481.
- Hintze, J.; Ye, Z.; Narimatsu, Y.; Madsen, T. D.; Joshi, H. J.; Goth, C. K.; Linstedt, A.; Bachert, C.; Mandel, U.; Bennett, E. P.; Vakhrushev, S. Y.; Schjoldager, K. T. Probing the Contribution of Individual Polypeptide GalNAc-Transferase Isoforms to the O-Glycoproteome by Inducible Expression in Isogenic Cell Lines. *J. Biol. Chem.* **2018**, *293*, 19064–19077.
- de las Rivas, M.; Lira-Navarrete, E.; Gerken, T. A.; Hurtado-Guerrero, R. Polypeptide GalNAc-Ts: From Redundancy to Specificity. *Curr. Opin. Struct. Biol.* **2019**, *56*, 87–96.
- Hagen, F. K.; Hazes, B.; Raffo, R.; DeSa, D.; Tabak, L. A. Structure-Function Analysis of the UDP-N-Acetyl-D-Galactosamine: Polypeptide N-Acetylgalactosaminyltransferase. *J. Biol. Chem.* **1999**, *274*, 6797–6803.
- Fritz, T. A.; Raman, J.; Tabak, L. A. Dynamic Association between the Catalytic and Lectin Domains of Human UDP-GalNAc:Polypeptide α -N-Acetylgalactosaminyltransferase-2. *J. Biol. Chem.* **2006**, *281*, 8613–8619.
- Lira-Navarrete, E.; Iglesias-Fernández, J.; Zandberg, W. F.; Compañón, I.; Kong, Y.; Corzana, F.; Pinto, B. M.; Clausen, H.; Peregrina, J. M.; Vocadlo, D. J.; Rovira, C.; Hurtado-Guerrero, R. Substrate-Guided Front-Face Reaction Revealed by Combined Structural Snapshots and Metadynamics for the Polypeptide N-Acetylgalactosaminyltransferase 2. *Angew. Chem., Int. Ed.* **2014**, *53*, 8206–8210.
- Kubota, T.; Shiba, T.; Sugioka, S.; Furukawa, S.; Sawaki, H.; Kato, R.; Wakatsuki, S.; Narimatsu, H. Structural Basis of Carbohydrate Transfer Activity by Human UDP-GalNAc: Polypeptide α -N-Acetylgalactosaminyltransferase (Pp-GalNAc-T10). *J. Mol. Biol.* **2006**, *359*, 708–727.
- Hazes, B. The (QxW)₃ Domain: A Flexible Lectin Scaffold. *Protein Sci.* **1996**, *5*, 1490–1501.
- Wandall, H. H.; Irazoqui, F.; Tarp, M. A.; Bennett, E. P.; Mandel, U.; Takeuchi, H.; Kato, K.; Irimura, T.; Suryanarayanan, G.; Hollingsworth, M. A.; Clausen, H. The Lectin Domains of Polypeptide GalNAc-Transferases Exhibit Carbohydrate-Binding Specificity for GalNAc: Lectin Binding to GalNAc-Glycopeptide Substrates Is Required for High Density GalNAc-O-Glycosylation. *Glycobiology* **2007**, *17*, 374–387.
- Tenno, M.; Kézdy, F. J.; Elhammer, Å. P.; Kurosaka, A. Function of the Lectin Domain of Polypeptide N-Acetylgalactosaminyltransferase 1. *Biochem. Biophys. Res. Commun.* **2002**, *298*, 755–759.
- Fritz, T. A.; Hurley, J. H.; Trinh, L.-B.; Shiloach, J.; Tabak, L. A. The Beginnings of Mucin Biosynthesis: The Crystal Structure of UDP-GalNAc:Polypeptide α -N-Acetylgalactosaminyltransferase-T1. *Proc. Natl. Acad. Sci. U.S.A.* **2004**, *101*, 15307–15312.
- Pedersen, J. W.; Bennett, E. P.; Schjoldager, K. T. B. G.; Meldal, M.; Holmér, A. P.; Blixt, O.; Cló, E.; Levery, S. B.; Clausen, H.; Wandall, H. H. Lectin Domains of Polypeptide GalNAc Transferases Exhibit Glycopeptide Binding Specificity. *J. Biol. Chem.* **2011**, *286*, 32684–32696.
- Hassan, H.; Reis, C. A.; Bennett, E. P.; Mirgorodskaya, E.; Roepstorff, P.; Hollingsworth, M. A.; Burchell, J.; Taylor-Papadimitriou, J.; Clausen, H. The Lectin Domain of UDP-N-

Acetyl-D-Galactosamine: Polypeptide N-Acetylgalactosaminyltransferase-T4 Directs Its Glycopeptide Specificities. *J. Biol. Chem.* **2000**, *275*, 38197–38205.

(23) De Las Rivas, M.; Daniel, E. J. P.; Coelho, H.; Lira-Navarrete, E.; Raich, L.; Compañón, I.; Diniz, A.; Lagartera, L.; Jiménez-Barbero, J.; Clausen, H.; Rovira, C.; Marcelo, F.; Corzana, F.; Gerken, T. A.; Hurtado-Guerrero, R. Structural and Mechanistic Insights into the Catalytic-Domain-Mediated Short-Range Glycosylation Preferences of GalNAc-T4. *ACS Cent. Sci.* **2018**, *4*, 1274–1290.

(24) Fernandez, A. J.; Daniel, E. J. P.; Mahajan, S. P.; Gray, J. J.; Gerken, T. A.; Tabak, L. A.; Samara, N. L. The Structure of the Colorectal Cancer-Associated Enzyme GalNAc-T12 Reveals How Nonconserved Residues Dictate Its Function. *Proc. Natl. Acad. Sci. U.S.A.* **2019**, *116*, 20404–20410.

(25) De Las Rivas, M.; Lira-Navarrete, E.; Daniel, E. J. P.; Compañón, I.; Coelho, H.; Diniz, A.; Jiménez-Barbero, J.; Peregrina, J. M.; Clausen, H.; Corzana, F.; Marcelo, F.; Jiménez-Osés, G.; Gerken, T. A.; Hurtado-Guerrero, R. The Interdomain Flexible Linker of the Polypeptide GalNAc Transferases Dictates Their Long-Range Glycosylation Preferences. *Nat. Commun.* **2017**, *8*, 1959.

(26) Raman, J.; Fritz, T. A.; Gerken, T. A.; Jamison, O.; Live, D.; Liu, M.; Tabak, L. A. The Catalytic and Lectin Domains of UDP-GalNAc:Polypeptide α -N-Acetylgalactosaminyltransferase Function in Concert to Direct Glycosylation Site Selection. *J. Biol. Chem.* **2008**, *283*, 22942–22951.

(27) Lira-Navarrete, E.; De Las Rivas, M.; Compañón, I.; Pallarés, M. C.; Kong, Y.; Iglesias-Fernández, J.; Bernardes, G. J. L.; Peregrina, J. M.; Rovira, C.; Bernadó, P.; Bruscolini, P.; Clausen, H.; Lostao, A.; Corzana, F.; Hurtado-Guerrero, R. Dynamic Interplay between Catalytic and Lectin Domains of GalNAc-Transferases Modulates Protein O-Glycosylation. *Nat. Commun.* **2015**, *6*, 6937.

(28) Revoredo, L.; Wang, S.; Bennett, E. P.; Clausen, H.; Moremen, K. W.; Jarvis, D. L.; Ten Hagen, K. G.; Tabak, L. A.; Gerken, T. A. Mucin-Type α -Glycosylation Is Controlled by Short- And Long-Range Glycopeptide Substrate Recognition That Varies among Members of the Polypeptide GalNAc Transferase Family. *Glycobiology* **2016**, *26*, 360–376.

(29) Tetaert, D.; Richet, C.; Gagnon, J.; Boersma, A.; Degand, P. Studies of Acceptor Site Specificities for Three Members of UDP-GalNAc:N-Acetylgalactosaminyltransferases by Using a Synthetic Peptide Mimicking the Tandem Repeat of MUC5AC. *Carbohydr. Res.* **2001**, *333*, 165–171.

(30) Kong, Y.; Joshi, H. J.; Schjoldager, K. T. B. G.; Madsen, T. D.; Gerken, T. A.; Vester-Christensen, M. B.; Wandall, H. H.; Bennett, E. P.; Levery, S. B.; Vakhrushev, S. Y.; Clausen, H. Probing Polypeptide GalNAc-Transferase Isoform Substrate Specificities by in Vitro Analysis. *Glycobiology* **2015**, *25*, 55–65.

(31) Schjoldager, K. T.; Joshi, H. J.; Kong, Y.; Goth, C. K.; King, S. L.; Wandall, H. H.; Bennett, E. P.; Vakhrushev, S. Y.; Clausen, H. Deconstruction of O-glycosylation—Gal NA C-T Isoforms Direct Distinct Subsets of the O-glycoproteome. *EMBO Rep.* **2015**, *16*, 1713–1722.

(32) Bagdonaite, I.; Pallesen, E. M.; Ye, Z.; Vakhrushev, S. Y.; Marinova, I. N.; Nielsen, M. I.; Kramer, S. H.; Pedersen, S. F.; Joshi, H. J.; Bennett, E. P.; Dabelsteen, S.; Wandall, H. H. O-glycan Initiation Directs Distinct Biological Pathways and Controls Epithelial Differentiation. *EMBO Rep.* **2020**, *21*, 1–17.

(33) Narimatsu, Y.; Joshi, H. J.; Schjoldager, K. T.; Hintze, J.; Halim, A.; Steentoft, C.; Nason, R.; Mandel, U.; Bennett, E. P.; Clausen, H.; Vakhrushev, S. Y. Exploring Regulation of Protein O-Glycosylation in Isogenic Human HEK293 Cells by Differential O-Glycoproteomics. *Mol. Cell. Proteomics* **2019**, *18*, 1396–1409.

(34) De Las Rivas, M.; Coelho, H.; Diniz, A.; Lira-Navarrete, E.; Compañón, I.; Jiménez-Barbero, J.; Schjoldager, K. T.; Bennett, E. P.; Vakhrushev, S. Y.; Clausen, H.; Corzana, F.; Marcelo, F.; Hurtado-Guerrero, R. Structural Analysis of a GalNAc-T2 Mutant Reveals an Induced-Fit Catalytic Mechanism for GalNAc-Ts. *Chem.—Eur. J.* **2018**, *24*, 8382–8392.

(35) de las Rivas, M.; Paul Daniel, E. J.; Narimatsu, Y.; Compañón, I.; Kato, K.; Hermosilla, P.; Thureau, A.; Ceballos-Laita, L.; Coelho, H.; Bernadó, P.; Marcelo, F.; Hansen, L.; Maeda, R.; Lostao, A.; Corzana, F.; Clausen, H.; Gerken, T. A.; Hurtado-Guerrero, R. Molecular Basis for Fibroblast Growth Factor 23 O-Glycosylation by GalNAc-T3. *Nat. Chem. Biol.* **2020**, *16*, 351–360.

(36) Wandall, H. H.; Hassan, H.; Mirgorodskaya, E.; Kristensen, A. K.; Roepstorff, P.; Bennett, E. P.; Nielsen, P. A.; Hollingsworth, M. A.; Burchell, J.; Taylor-Papadimitriou, J.; Clausen, H. Substrate Specificities of Three Members of the Human UDP- N -Acetyl- α -d-Galactosamine:Polypeptide N -Acetylgalactosaminyltransferase Family, GalNAc-T1, -T2, and -T3. *J. Biol. Chem.* **1997**, *272*, 23503–23514.

(37) Smet-Nocca, C.; Broncel, M.; Wieruszkeski, J. M.; Tokarski, C.; Hanouelle, X.; Leroy, A.; Landrieu, I.; Rolando, C.; Lippens, G.; Hackenberger, C. P. Identification of O-GlcNAc sites within peptides of the Tau protein and their impact on phosphorylation. *Mol BioSyst.* **2011**, *7* (5), 1420–1429.

(38) Brokx, R. D.; Revers, L.; Zhang, Q.; Yang, S.; Mal, T. K.; Ikura, M.; Gariépy, J. Nuclear Magnetic Resonance-Based Dissection of a Glycosyltransferase Specificity for the Mucin MUC1 Tandem Repeat. *Biochemistry* **2003**, *42*, 13817–13825.

(39) Kinarsky, L.; Suryanarayanan, G.; Prakash, O.; Paulsen, H.; Clausen, H.; Hanisch, F. G.; Hollingsworth, M. A.; Sherman, S. Conformational Studies on the MUC1 Tandem Repeat Glycopeptides: Implication for the Enzymatic O-Glycosylation of the Mucin Protein Core. *Glycobiology* **2003**, *13*, 929–939.

(40) Corzana, F.; Busto, J. H.; Jiménez-Osés, G.; De Luis, M. G.; Asensio, J. L.; Jiménez-Barbero, J.; Peregrina, J. M.; Avenoza, A. Serine versus Threonine Glycosylation: The Methyl Group Causes a Drastic Alteration on the Carbohydrate Orientation and on the Surrounding Water Shell. *J. Am. Chem. Soc.* **2007**, *129*, 9458–9467.

(41) Madariaga, D.; Martínez-Sáez, N.; Somovilla, V. J.; García-García, L.; Berbis, M. A.; Valero-González, J.; Martín-Santamaría, S.; Hurtado-Guerrero, R.; Asensio, J. L.; Jiménez-Barbero, J.; Avenoza, A.; Busto, J. H.; Corzana, F.; Peregrina, J. M. Serine versus Threonine Glycosylation with α -o-GalNAc: Unexpected Selectivity in Their Molecular Recognition with Lectins. *Chem.—Eur. J.* **2014**, *20*, 12616–12627.

(42) Bermejo, I. A.; Usabiaga, I.; Compañón, I.; Castro-López, J.; Insausti, A.; Fernández, J. A.; Avenoza, A.; Busto, J. H.; Jiménez-Barbero, J.; Asensio, J. L.; Peregrina, J. M.; Jiménez-Osés, G.; Hurtado-Guerrero, R.; Cocinero, E. J.; Corzana, F. Water Sculptures the Distinctive Shapes and Dynamics of the Tumor-Associated Carbohydrate Tn Antigens: Implications for Their Molecular Recognition. *J. Am. Chem. Soc.* **2018**, *140*, 9952–9960.

(43) Yoshimura, Y.; Nudelman, A. S.; Levery, S. B.; Wandall, H. H.; Bennett, E. P.; Hindsgaul, O.; Clausen, H.; Nishimura, S. I. Elucidation of the Sugar Recognition Ability of the Lectin Domain of UDP-GalNAc:Polypeptide N-Acetylgalactosaminyltransferase 3 by Using Unnatural Glycopeptide Substrates. *Glycobiology* **2012**, *22*, 429–438.

(44) Abu-Abed, M.; Mal, T. K.; Kainosho, M.; MacLennan, D. H.; Ikura, M. Characterization of the ATP-Binding Domain of the Sarco(Endo)Plasmic Reticulum Ca²⁺-ATPase: Probing Nucleotide Binding by Multidimensional NMR. *Biochemistry* **2002**, *41*, 1156–1164.

(45) Wishart, D. S.; Sykes, B. D. The ¹³C Chemical-Shift Index: A Simple Method for the Identification of Protein Secondary Structure Using ¹³C Chemical-Shift Data. *J. Biomol. NMR* **1994**, *4*, 171–180.

(46) Daniel, E. J. P.; Las Rivas, M.; Lira-Navarrete, E.; García-García, A.; Hurtado-Guerrero, R.; Clausen, H.; Gerken, T. A. Ser and Thr Acceptor Preferences of the GalNAc-Ts Vary among Isoenzymes to Modulate Mucin-Type O-Glycosylation. *Glycobiology* **2020**, *30*, 910–922.

(47) Campos, D.; Freitas, D.; Gomes, J.; Magalhães, A.; Steentoft, C.; Gomes, C.; Vester-Christensen, M. B.; Ferreira, J. A.; Afonso, L. P.; Santos, L. L.; Pinto de Sousa, J.; Mandel, U.; Clausen, H.; Vakhrushev, S. Y.; Reis, C. A. Probing the O-Glycoproteome of

Gastric Cancer Cell Lines for Biomarker Discovery. *Mol. Cell. Proteomics* **2015**, *14*, 1616–1629.

(48) Gerken, T. A.; Revoredo, L.; Thome, J. J. C.; Tabak, L. A.; Vester-Christensen, M. B.; Clausen, H.; Gahlay, G. K.; Jarvis, D. L.; Johnson, R. W.; Moniz, H. A.; Moremen, K. The Lectin Domain of the Polypeptide GalNAc Transferase Family of Glycosyltransferases (PpGalNAc Ts) Acts as a Switch Directing Glycopeptide Substrate Glycosylation in an N- or C-Terminal Direction, Further Controlling Mucin Type O-Glycosylation. *J. Biol. Chem.* **2013**, *288*, 19900–19914.

(49) Macías-León, J.; Bermejo, I. A.; Asín, A.; García-García, A.; Compañón, I.; Jiménez-Moreno, E.; Coelho, H.; Mangini, V.; Albuquerque, I. S.; Marcelo, F.; Asensio, J. L.; Bernardes, G. J. L.; Joshi, H. J.; Fiammengo, R.; Blixt, O.; Hurtado-Guerrero, R.; Corzana, F. Structural Characterization of an Unprecedented Lectin-like Antitumoral Anti-MUC1 Antibody. *Chem. Commun.* **2020**, *56*, 15137–15140.

(50) Olson, F. J.; Bäckström, M.; Karlsson, H.; Burchell, J.; Hansson, G. C. A MUC1 Tandem Repeat Reporter Protein Produced in CHO-K1 Cells Has Sialylated Core 1 O-Glycans and Becomes More Densely Glycosylated If Coexpressed with Polypeptide-GalNAc-T4 Transferase. *Glycobiology* **2005**, *15*, 177–191.

(51) Hanisch, F. G.; Reis, C. A.; Clausen, H.; Paulsen, H. Evidence for Glycosylation-Dependent Activities of Polypeptide N-Acetylgalactosaminyltransferases RGalNAc-T2 and -T4 on Mucin Glycopeptides. *Glycobiology* **2001**, *11*, 731–740.

(52) Martínez-Sáez, N.; Peregrina, J. M.; Corzana, F. Principles of Mucin Structure: Implications for the Rational Design of Cancer Vaccines Derived from MUC1-Glycopeptides. *Chem. Soc. Rev.* **2017**, *46*, 7154–7175.

(53) Nason, R.; Büll, C.; Konstantinidi, A.; Sun, L.; Ye, Z.; Halim, A.; Du, W.; Sorensen, D. M.; Durbesson, F.; Furukawa, S.; Mandel, U.; Joshi, H. J.; Dworkin, L. A.; Hansen, L.; David, L.; Iverson, T. M.; Bensing, B. A.; Sullam, P. M.; Varki, A.; de Vries, E.; de Haan, C. A. M.; Vincentelli, R.; Henrissat, B.; Vakhrushev, S. Y.; Clausen, H.; Narimatsu, Y. Display of the Human Mucinome with Defined O-Glycans by Gene Engineered Cells. *Nat. Commun.* **2021**, *12*, 4070.


(54) Noach, I.; Ficko-Blean, E.; Pluinage, B.; Stuart, C.; Jenkins, M. L.; Brochu, D.; Buenbrazo, N.; Wakarchuk, W.; Burke, J. E.; Gilbert, M.; Boraston, A. B. Recognition of Protein-Linked Glycans as a Determinant of Peptidase Activity. *Proc. Natl. Acad. Sci. U.S.A.* **2017**, *114*, E679–E688.

(55) Konstantinidi, A.; Nason, R.; Čaval, T.; Sun, L.; Sorensen, D. M.; Furukawa, S.; Ye, Z.; Vincentelli, R.; Narimatsu, Y.; Vakhrushev, S. Y.; Clausen, H. Exploring Glycosylation of Mucins with O-Glycodomain Reporters Expressed in Glycoengineered HEK293 Cells. *J. Biol. Chem.* **2021**.

(56) Steentoft, C.; Vakhrushev, S. Y.; Vester-Christensen, M. B.; Schjoldager, K. T.-B. G.; Kong, Y.; Bennett, E. P.; Mandel, U.; Wandall, H.; Levery, S. B.; Clausen, H. Mining the O-Glycoproteome Using Zinc-Finger Nuclease-Glycoengineered SimpleCell Lines. *Nat. Methods* **2011**, *8*, 977–982.


(57) Coelho, H.; Matsushita, T.; Artigas, G.; Hinou, H.; Cañada, F. J.; Lo-Man, R.; Leclerc, C.; Cabrita, E. J.; Jiménez-Barbero, J.; Nishimura, S. I.; Garcia-Martín, F.; Marcelo, F. The Quest for Anticancer Vaccines: Deciphering the Fine-Epitope Specificity of Cancer-Related Monoclonal Antibodies by Combining Microarray Screening and Saturation Transfer Difference NMR. *J. Am. Chem. Soc.* **2015**, *137*, 12438–12441.


(58) Yoshimura, Y.; Denda-Nagai, K.; Takahashi, Y.; Nagashima, I.; Shimizu, H.; Kishimoto, T.; Noji, M.; Shichino, S.; Chiba, Y.; Irimura, T. Products of Chemoenzymatic Synthesis Representing MUC1 Tandem Repeat Unit with T-, ST- or STn-Antigen Revealed Distinct Specificities of Anti-MUC1 Antibodies. *Sci. Rep.* **2019**, *9*, 16641.



JACS Au
AN OPEN ACCESS JOURNAL OF THE AMERICAN CHEMICAL SOCIETY

Editor-in-Chief
Prof. Christopher W. Jones
Georgia Institute of Technology, USA

Open for Submissions 

pubs.acs.org/jacsau  ACS Publications
Most Trusted. Most Cited. Most Read.