

# P2Y1 receptor inhibition rescues impaired synaptic plasticity and astroglial Ca<sup>2+</sup>-dependent activity in the epileptic hippocampus

Andrés Martorell<sup>a,c</sup>, Mario Wellmann<sup>a,b</sup>, Felipe Guiffa<sup>a</sup>, Marco Fuenzalida<sup>a</sup>, Christian Bonansco<sup>a,\*</sup>

<sup>a</sup> Centro de Neurobiología y Fisiopatología Integrativa (CENFI), Instituto de Fisiología, Facultad de Ciencias, Universidad de Valparaíso, Valparaíso, Chile

<sup>b</sup> Escuela de Fonoaudiología, Facultad de Medicina, Universidad de Valparaíso, Valparaíso, Chile

<sup>c</sup> Escuela de Fonoaudiología, Facultad de Ciencias de la Rehabilitación, Universidad Andres Bello, Viña del Mar, Chile

## ARTICLE INFO

### Keywords:

Gliotransmission  
Synaptic plasticity  
Epilepsy  
Astrocytes  
Purinergic receptors  
Long-term potentiation  
Kindling

## ABSTRACT

Epilepsy is characterized by a progressive predisposition to suffer seizures due to neuronal hyperexcitability, and one of its most common co-morbidities is cognitive decline. In animal models of chronic epilepsy, such as kindling, electrically induced seizures impair long-term potentiation (LTP), deteriorating learning and memory performance. Astrocytes are known to actively modulate synaptic plasticity and neuronal excitability through Ca<sup>2+</sup>-dependent gliotransmitter release. It is unclear, however, if astroglial Ca<sup>2+</sup> signaling could contribute to the development of synaptic plasticity alterations in the epileptic hippocampus. By employing electrophysiological tools and Ca<sup>2+</sup> imaging, we found that glutamatergic CA3-CA1 synapses from kindled rats exhibit an impairment in theta burst (TBS) and high frequency stimulation (HFS)-induced LTP, which is accompanied by an increased probability of neurotransmitter release (Pr) and an abnormal pattern of astroglial Ca<sup>2+</sup>-dependent transients. Both the impairment in LTP and the Pr were reversed by inhibiting purinergic P2Y1 receptors (P2Y1R) with the specific antagonist MRS2179, which also restored the spontaneous and TBS-induced pattern of astroglial Ca<sup>2+</sup>-dependent signals. Two consecutive, spaced TBS protocols also failed to induce LTP in the kindled group, however, this impairment was reversed and a strong LTP was induced when the second TBS was applied in the presence of MRS2179, suggesting that the mechanisms underlying the alterations in TBS-induced LTP are likely associated with an aberrant modulation of the induction threshold for LTP. Altogether, these results indicate that P2Y1R inhibition rescues both the pattern of astroglial Ca<sup>2+</sup>-activity and the plastic properties of CA3-CA1 synapses in the epileptic hippocampus, suggesting that astrocytes might take part in the mechanisms that deteriorate synaptic plasticity and thus cause cognitive decline in epileptic patients.

## 1. Introduction

Epilepsy is characterized by a progressive predisposition to suffer seizures as a consequence of an imbalance in neuronal excitability, which has been associated behavioral and cognitive alterations commonly exhibited, affecting about 70 million people worldwide (Fisher et al., 2005; Halgren et al., 1991; Helmstaedter, 2002). The difficulties that patients with epilepsy face extend far beyond seizures and include severe and disabling comorbidities as cognitive impairment, which has profound consequences in their life quality (Holmes, 2015). Several lines of evidence indicate that an exacerbated glutamatergic activity, often referred to as the “glutamate hypothesis”, is one of the key alterations in the pathophysiology of epilepsy (Soukupova et al., 2015). However, little is known about the cellular and molecular mechanisms that underlie cognitive decline in epileptic patients – which include

deficits in attentional processes, working memory, and executive function –, and it is not fully understood how the aforementioned glutamatergic imbalance could contribute to its development (Campo et al., 2013; Celiker Uslu et al., 2019; Rai et al., 2015).

Kindling is the most employed experimental animal model of chronic epilepsy and closely resembles the development of epilepsy in humans (Löscher, 2011; McNamara et al., 1985; Morales et al., 2014; Morimoto et al., 2004). It consists in the presentation of repetitive, sub-convulsive electrical or chemical stimuli on a nervous structure (usually the amygdala or hippocampus) which elicits a gradual and progressive enhancement of electroencephalographic (EEG) activity and behavioral responses, culminating in generalized seizures. Numerous works suggest that the development of chronic epilepsy through electrical kindling and the induction of classical long-term potentiation (LTP) in glutamatergic synapses share a wide range of cellular and molecular

\* Corresponding author at: Instituto de Fisiología, Facultad de Ciencias, Universidad de Valparaíso, Avenida Gran Bretaña 1111, Valparaíso 2360102, Chile.

E-mail address: [christian.bonansco@uv.cl](mailto:christian.bonansco@uv.cl) (C. Bonansco).

<https://doi.org/10.1016/j.nbd.2020.105132>

Received 10 July 2020; Received in revised form 15 September 2020; Accepted 6 October 2020

Available online 10 October 2020

0969-9961/ © 2020 Centro de Neurobiología y Fisiopatología Integrativa (CENFI), Insto. de Fisiología, Universidad de Valparaíso. Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

mechanisms. Those mechanisms include the repetitive stimulation during the induction phase, the activity-dependent and input-specific strengthening of synaptic efficacy, the temporal coincidence between pre- and post-synaptic activity mediated by *N*-methyl-D-aspartate receptors (NMDARs; Larson and Munkácsy, 2015; Matsuzaki et al., 2004; Nicoll and Schmitz, 2005), among other mechanisms, all of which are also at the very basis of the plasticity processes that underlie learning and memory (Leung and Shen, 2006; Stasheff et al., 1989; Sutula et al., 1988). It has been suggested that repeated seizure activity causes an indiscriminate and widespread enhancement in synaptic efficacy, interfering with the synaptic plasticity (i.e.: LTP) required for information processing and learning (Reid and Stewart, 1997). In fact, previous research indicates that LTP is markedly reduced at the epileptic focus (i.e.: hippocampus) in humans with temporal lobe epilepsy (Beck et al., 2000). In line with those results, experiments with animal models suggests that kindling can suppress LTP (Leung and Wu, 2003), whereas electrically-induced seizures can block (Hu et al., 2005) or reduce it and impair learning and memory (Esmailpour et al., 2017; Ghotbeddin et al., 2018; Moore et al., 1993). However, the cellular mechanisms by which epilepsy – like kindling – can alter LTP remain unclear.

During the last 15 years, it has been shown that astrocytes modulate both short- and long-term synaptic plasticity as well as neuronal excitability through  $\text{Ca}^{2+}$ -dependent gliotransmitter release (Bonansco et al., 2011; Perea and Araque, 2005, 2007). For instance, it has been proposed that spontaneously released glutamate from astrocytes actively regulate the basal Pr of glutamatergic terminals by activating presynaptic metabotropic glutamate receptors of the mGluR1/5 subtypes, operating as a gain control mechanism that modulates the induction threshold for LTP (Bonansco et al., 2011; Panatier et al., 2011). In addition, growing evidence indicates that astrocytes directly participate in the pathophysiology of several brain conditions, including epilepsy (Nikolic et al., 2019; Pirttimaki et al., 2013; Shigetomi et al., 2019; Verkhratsky et al., 2017). In previous reports, we showed that reactive astrocytes from kindled rats exhibit an abnormal pattern of  $\text{Ca}^{2+}$  transients (i.e.:  $\text{Ca}^{2+}$ -mediated hyperexcitability) that enhances glutamate-mediated gliotransmission and increases neuronal glutamate release via presynaptic mGluR1/5 receptor activation (Álvarez-Ferradas et al., 2015). This abnormal pattern of astroglial  $\text{Ca}^{2+}$  activity requires purinergic P2Y1 receptor (P2Y1R) and Pannexin-1 hemichannel activation (Wellmann et al., 2018). Those mechanisms could represent a form of astrocyte-mediated metaplasticity and have been proposed as the basis for an astrocyte-to-neuron loop that chronically upregulate excitatory transmission, bringing epileptic circuits near seizure threshold (Álvarez-Ferradas et al., 2015; Cavus et al., 2005; Gómez-Gonzalo et al., 2010; Jones, 2015; Wellmann et al., 2018; Riquelme et al., 2020). Nonetheless, whether and how astroglial-mediated modulation of synaptic transmission contributes to the impairments in synaptic plasticity (i.e.: LTP) and thus to the development of cognitive decline in epileptic patients remains unknown.

Using hippocampal slices from fully kindled rats (Morales et al., 2014), we tested if different electrical stimulation protocols were able to induce LTP, assessing synaptic efficacy and astroglial  $\text{Ca}^{2+}$  activity before, during and after the induction of activity-dependent synaptic plasticity. We found that both theta-burst (TBS) and high frequency (HFS) stimulation-induced LTP are impaired in glutamatergic CA3-CA1 synapses from kindled animals. These impairments are associated with an anomalous pattern of astrocytic  $\text{Ca}^{2+}$  signals (i.e.: slow  $\text{Ca}^{2+}$ -transients; STs) that abnormally up-regulate the Pr of CA3-CA1 synapses. Both the impaired LTP and the increased Pr were reversed by inhibiting P2Y1 receptors with MRS2179, which also restored the pattern of astroglial  $\text{Ca}^{2+}$ -dependent activity. Altogether, our data suggest that hyper-excitable astrocytes are directly involved in the development of the glutamatergic dysfunction in the epileptic brain, likely impairing LTP by abnormally increasing the Pr in CA3-CA1 synapses. These results offer a new mechanism that explains the decreased plastic capacity of neural circuits and thus the cognitive decline in epileptic patients.

## 2. Methods

Animal care, surgery and recording procedures were in accordance with the guidelines laid down by the Institutional Animal Care and Ethics Committee from the Faculty of Sciences, Universidad de Valparaíso (DIUV Law 20,380, Chile) and NIH (USA).

### 2.1. Stereotaxic surgery and kindling protocol


Epileptogenesis was induced using a rapid kindling protocol (RK) previously described (Álvarez-Ferradas et al., 2015; Morales et al., 2014; Wellmann et al., 2018). Briefly explained, male Sprague-Dawley rats (35–40 postnatal days) were anesthetized and subjected to stereotaxic surgery procedures for electrode implantation. Following stereotaxic coordinates (Paxinos and Watson, 1998), two pairs of electrodes for cortical recording were bilaterally implanted in the primary motor cortex and the visual cortex, and one stimulation electrode in the right basolateral amygdala complex (BLA), plus an anchorage screw. Following surgery, rats had a recovery period of at least one week before the start of the kindling procedure.

The RK protocol consists of ten daily trains of biphasic rectangular current pulses at subthreshold after-discharge (ADs) intensity, applied for three consecutive days. Rats subjected to the RK protocol displayed progressive epileptic activity throughout the brain, accompanied by seizures that increased in severity as the protocol went on (Morales et al., 2014). ADs progression and seizure severity were assessed using EEG recordings and the Racine scale (De Jonge and Racine, 1987), respectively. Rats were considered fully kindled after having suffered at least five consecutive generalized epileptic seizures (i.e., Racine 4 and 5 states) accompanied by long-lasting repetitive ADs. Under these conditions, cortical and mesolimbic structures of fully kindled rats are considered epileptic tissue, including the hippocampal formation (Morales et al., 2014; Steward et al., 1991). The control group consisted of rats that were subjected to electrode implantation surgery but remained non-stimulated (i.e., sham), and naïve animals; results were pooled together because no significant differences were found in this nor in previous reports (Álvarez-Ferradas et al., 2015; Morales et al., 2014; Wellmann et al., 2018). All the experiments were performed at least one week after fully kindled states were reached (i.e.: 45–60 postnatal days). Rats were maintained ad-libitum before, during and after all procedures.

### 2.2. Electrophysiology and hippocampal slice preparation

Acute slices from the dorsal hippocampus were obtained from control and kindled rats as previously described (Bonansco and Buño, 2003; Bonansco et al., 2011). Briefly, rats were anesthetized and decapitated, the brain was rapidly removed through craniotomy and placed in ice-cold ( $< 4\text{ }^{\circ}\text{C}$ ) artificial cerebrospinal fluid (ACSF), gassed with a mixture of 95%  $\text{O}_2$  and 5%  $\text{CO}_2$  (pH 7.4). Transversal brain slices (300–350  $\mu\text{m}$  thick) were obtained with a Vibroslice microtome (VSL, WPI, USA) and incubated for 1 h at room temperature (21–24  $^{\circ}\text{C}$ ) in ACSF containing (in mM): 124.0 NaCl, 2.7 KCl, 1.25  $\text{KH}_2\text{PO}_4$ , 2.0  $\text{Mg}_2\text{SO}_4$ , 26.0  $\text{NaHCO}_3$ , 2.5  $\text{CaCl}_2$  and 10.0 glucose. Slices were then transferred to an immersion-recording chamber (2 ml), fixed to an upright microscope stage (FN100 IR; Nikon Inc.; Japan) equipped with infrared and differential interference contrast imaging devices and with a  $40\times$ -water immersion objective. Slices were perfused with carbogen-bubbled ACSF (5%  $\text{CO}_2$  and 95% oxygen; 2 ml/min) and maintained at room temperature (21–24  $^{\circ}\text{C}$ ). All recordings were made in presence of the GABA<sub>A</sub>R antagonist picrotoxin (PTX, 10  $\mu\text{M}$ ).

Two recording methods were employed: extracellular field potentials and whole-cell patch-clamp recordings (Bonansco et al., 2007). Field excitatory postsynaptic potentials (fEPSP) were performed with a glass pipette (2–4  $\text{M}\Omega$ , filled with perfusion medium) placed in the middle of the CA1 *stratum radiatum* and connected to an AC amplifier



**Fig. 1.** TBS fails to induce LTP at CA3-CA1 synapses and evokes an unresponsive pattern of astroglial  $\text{Ca}^{2+}$ -dependent activity in kindled hippocampal slices. (A) Average traces of excitatory postsynaptic currents (EPSC) before (black trace) and 40 min after TBS application (gray trace) in control ( $n = 5$  cells; 4 slices) and kindled group ( $n = 6$  cells; 5 slices). (B) (top) Time course of the changes in EPSC amplitude (%) before and after TBS and (bottom) normalized values of input resistance ( $R_i$ ) for each cell, measured throughout the experiments. (C) Superimposed averaged EPSCs evoked by the PPR protocol recorded before (black trace) and after (gray trace) TBS in control and kindled condition. (D) Summary graphs for average TBS-induced potentiation (left), baseline PPR (middle) and PPR change (%; right), measured before and 40 min after the protocol. (E-F) Fluorescence traces of spontaneous  $\text{Ca}^{2+}$  elevations from five representative astrocytes, recorded before, during and after the application of the TBS protocol in control (E) and kindled (F) groups. Blue boxes denote  $\text{Ca}^{2+}$ -dependent activity during TBS protocols. (Right) Summary graphs showing mean values of FT/ST ratio (fast  $\text{Ca}^{2+}$  transients/slow  $\text{Ca}^{2+}$  transients) and average number of slow  $\text{Ca}^{2+}$  transients (ST) and fast  $\text{Ca}^{2+}$  transients (FT) per astrocyte. Results expressed as mean  $\pm$  SEM. \*  $p < 0.05$ . (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

(P-5 series; Grass) with a gain of  $310.000\times$ , a 3.0-kHz low pass filter, and a 0.30 Hz high pass filter. To evoke fEPSPs, a bipolar concentric electrode (concentric platinum/iridium electrode, 125  $\mu\text{m}$  outer, FHC Inc., Bowdoin, ME, USA) was placed in the *stratum radiatum* within 100–200  $\mu\text{m}$  from the recording site to activate the Schaeffer collaterals fibers using bipolar cathodic stimulation (50  $\mu\text{sec}$ , 0.3 Hz, 20–100  $\mu\text{A}$ ) generated by a stimulator (A.M.P.I., Israel) connected to an isolation unit (Isoflex, AMPI), which elicited compound action potentials from the presynaptic axons (fiber volley; fV) followed by fEPSPs. Unless otherwise stated, whole-cell voltage-clamp recordings from CA1 pyramidal cells clamped at  $-60$  mV were made using a EPC-7 amplifier (Heka Instruments, Germany), and patch-type pipette electrodes ( $\sim 5$  M $\Omega$ ) containing (in mM): 130 KMeSO<sub>4</sub>, 10 HEPES, 4 ATP-Na<sub>2</sub> and 20 KCl, 290 mOsm, adjusted to pH 7.3. Excitatory postsynaptic currents (EPSCs) were elicited every 3 s, filtered at 3.0 kHz and acquired at 4.0 kHz, using an A/D converter (ITC-16; Instrutech, Germany) and

stored with Pulse FIT software (Heka instruments, Germany). Experiments started after a 5–10 min stabilization period following the establishment of the whole-cell configuration. Cells that exhibited a significant change in access resistance ( $> 20\%$ ) were excluded from the analysis. We used a paired-pulse protocol and calculated the paired-pulse ratio (PPR) to estimate putative presynaptic changes. PPR was defined as the ratio between the peak amplitude of the second and the first EPSCs (80 ms apart). Relative changes in PPT throughout the recordings were calculated as a percentage variation compared to baseline values. LTP was induced by using two well-known protocols delivered at the SCs: theta-burst stimulation (TBS) and high-frequency stimulation (HFS). TBS consisted in 10 trains of 5 pulses at 100 Hz, delivered at 200 ms intervals, containing 10–12 bursts 30 s apart; HFS consisted in 2 trains at 100 Hz (1-s duration; 30 s apart). Off-line recording analysis was performed using the software pClamp by Molecular Devices (USA).

### 2.3. $Ca^{2+}$ imaging in astrocytes

Intracellular  $Ca^{2+}$ -elevations from astrocytes were monitored by fluorescence microscopy using Fluo4-AM as cell-permeant  $Ca^{2+}$  indicator, as previously described (Wellmann et al., 2018). For  $Ca^{2+}$  imaging experiments, brain extraction and slice preparation were performed in sucrose enriched ACSF (215.0 mM). To confirm the specific recording of  $Ca^{2+}$  signals from astrocytes, slices were incubated first with the astroglial morphological marker sulforhodamine-101 (SR101; 0.5–1  $\mu$ M; Kafitz et al., 2008) for ~30 min in low  $Ca^{2+}$ /high  $Mg^{2+}$  ACSF at 32–34 °C. Slices were then transferred to a maintenance chamber with regular ACSF for 30 min and later incubated with FLUO4-AM (1–2  $\mu$ L of the dye dissolved in pluronic acid at 0.01% was dropped over the hippocampus, obtaining a final concentration of 5–10  $\mu$ M) for ~75 min in regular ACSF at room temperature. Astrocytes were imaged using a CCD camera (Andor DR328G; Andor Technologies PLC, Ireland) attached to a fluorescence microscope (Nikon, Japan). The camera was controlled and synchronized by the Niss-Elements AR 3.2 software (Nikon, Japan), which was also used for offline analyses. Cells were illuminated with a Xenon lamp at 490 nm (200–400 ms exposure; 36,700 mm<sup>2</sup> area), and images were acquired at 1 Hz for 5 min, regulated by a shutter (Lambda SC-Smart shutter, Sutter Instrument Company). Analyses of astroglial  $Ca^{2+}$  levels were restricted to the cell body and  $Ca^{2+}$ -transients were estimated as changes in the fluorescence signal over the baseline ( $\Delta F/F_0$ ) after background subtraction. For every individual cell, the baseline was obtained by extracting and averaging the fluorescence values of at least 30 consecutive frames where the astrocyte exhibited no spontaneous  $Ca^{2+}$ -dependent activity. Changes in fluorescence were considered as events when the  $\Delta F/F_0$  intensity exceeded the fluorescence of the baseline by at least two standard deviations for no less than five consecutive frames. For multi-peak astroglial  $Ca^{2+}$  activity, those events in which the  $\Delta F/F_0$  dropped to a half of the maximum fluorescence intensity in relation to the baseline were considered as independent transients. As we previously described (Wellmann et al., 2018), the duration of astroglial  $Ca^{2+}$  transients was distributed among two populations starting from the 75th percentile of the cumulative distribution – which was 17 s –, being that value employed as a cut-off criterion from which random events in the same astrocyte were classified as slow (STs > 17 s) or fast transients (FTs < 17 s; Fig. 1E–F). Also, we incorporated the FT/ST frequency ratio as an index of astroglial excitability (Wellmann et al., 2018), where a lower index would be associated to a greater  $Ca^{2+}$ -dependent excitability.

### 2.4. Statistical analysis

In all cases, a distribution analysis was performed first to determine if the data fitted a normal distribution (Shapiro-Wilk test, Kolmogorov-Smirnov test). According to that, a parametric (Student's two-tailed *t*-test) or a non-parametric test (Mann-Whitney test) was performed. For multiple comparisons, ANOVA or Kruskal-Wallis test was used as appropriate, both with post hoc Bonferroni correction. Unless otherwise stated, all data is expressed as mean  $\pm$  SEM. Differences were considered statistically significant at  $p < 0.05$ .

## 3. Results

### 3.1. TBS fails to induce LTP at CA3-CA1 synapses and causes an anomalous astroglial $Ca^{2+}$ -dependent response in kindled hippocampal slices


The effect of epileptogenesis over synaptic plasticity paradigms such as LTP are often controversial, having previous reports described increases (Krug et al., 1997; Ruethrich et al., 1996; Schneiderman, 1997), decreases (De Jonge and Racine, 1987; Gilbert and Mack, 1990; Maru and Goddard, 1987) or even complete suppressions of LTP (Leung and

Wu, 2003; Schubert et al., 2005; Suárez et al., 2012). Using hippocampal slices, we evaluated if the induction of LTP is altered in kindled rats (Morales et al., 2014). The application of a TBS protocol after a stable baseline (~10 min; Fig. 1A) induced a robust LTP in control rats (at 30–40 min;  $249.07 \pm 8.46\%$  of baseline,  $p = 0.016$ ;  $n = 6$ ), but failed to induce LTP in the kindled group (to  $105.23 \pm 7.3\%$  of baseline,  $p = 0.073$ ;  $n = 5$ ; Fig. 1A). In addition, kindled rats exhibited lower PPR values before and after the protocol compared to control animals (kindled:  $1.25 \pm 0.09$ ;  $n = 6$  vs control:  $1.75 \pm 0.05$ ;  $n = 5$ ;  $p = 0.03$ ; Fig. 1C, D). Since astrocytes from the epileptic hippocampus show an abnormal pattern of spontaneous  $Ca^{2+}$ -dependent activity (i.e.: increased frequency of slow transients, STs; Álvarez-Ferradas et al., 2015; Wellmann et al., 2018) that could alter astroglial-mediated mechanisms that regulate synaptic plasticity, we assessed if astrocytes in the kindled hippocampus respond with changes in their pattern of  $Ca^{2+}$ -dependent activity during and after the TBS protocol. In basal conditions, control slices exhibited both fast (i.e., FTs < 17 s) and slow (i.e., STs > 17 s) spontaneous astroglial  $Ca^{2+}$  transients, being FTs the most prevalent type of oscillations (Fig. 1E). During the TBS protocol both FTs and STs increased in number, but STs exhibited a greater relative increase compared to FTs (basal FT;  $6.1 \pm 0.5$  vs TBS FT;  $6.5 \pm 0.5$ ;  $p < 0.05$  and basal ST;  $1.4 \pm 0.1$  vs TBS ST;  $3.1 \pm 0.3$ ;  $p < 0.001$ ;  $n = 56$ ; Fig. 1E), which is reflected by the decrease in the FT/ST ratio (basal;  $4.5 \pm 0.60$  vs during TBS;  $2.3 \pm 0.37$ ,  $p = 0.021$ ; Fig. 1E). This shift in the overall  $Ca^{2+}$ -dependent activity pattern was quickly reversed after the protocol. In contrast, astrocytes from the kindling group showed a higher rate of basal activity (basal FT;  $5.1 \pm 0.4$  vs basal ST;  $4.5 \pm 0.1$ ;  $p < 0.01$ ;  $n = 43$ ), and a lower FT/ST ratio, which resembles the pattern of  $Ca^{2+}$ -dependent activity observed in the group control during the TBS protocol (Control TBS  $2.3 \pm 0.41$ ,  $n = 57$ , vs Kindling baseline,  $1.74 \pm 0.40$ ,  $n = 43$ ,  $p = 0.29$ ). During the protocol, STs augmented in number (Fig. 1F; basal;  $4.5 \pm 0.36$  vs during TBS;  $6.3 \pm 0.6$ ;  $p < 0.05$ ,  $n = 61$ ) but without fluctuations in the FT/ST ratio (Fig. 1F; basal  $1.74 \pm 0.40$ ,  $n = 43$ , vs TBS  $1.76 \pm 0.40$ ,  $n = 52$ ,  $p = 0.76$ ), indicating that both FTs and STs increased together without changes in the global pattern of astroglial  $Ca^{2+}$ -dependent activity in response to TBS, which suggests that the abnormal pattern of  $Ca^{2+}$ -dependent activity emerges in an uncoupled, synaptic activity-independent fashion. These findings show that both neurons and astrocytes in the epileptic hippocampus exhibit an abnormal response to the TBS protocol, which is characterized by an impairment in TBS-induced LTP in CA3-CA1 synapses and the maintenance of an abnormal pattern of astroglial  $Ca^{2+}$ -dependent activity.

### 3.2. Consecutive, spaced TBS protocols failed to induce LTP in the kindled hippocampus

Since a single TBS protocol did not induce LTP, we next tested if a stronger protocol consisting in two spaced (30 min inter-stimulus interval), consecutive TBS can induce a plastic change (Schubert et al., 2005). In control slices, both protocols (TBS1 and TBS2) induced a short period of post-tetanic facilitation followed by an increase in EPSC amplitude that reached a steady state after 30 min ( $159.19 \pm 10.11\%$  of baseline;  $n = 5$ ) for TBS1 and 50 min for TBS2 ( $208.88 \pm 25.50\%$  of baseline;  $n = 5$ ; Fig. 2D). In kindled slices, although post-tetanic facilitation was observed after both TBS1 and TBS2, neither protocols were able to induce LTP (Fig. 2D, kindled: TBS1;  $109.41 \pm 7.53\%$  versus TBS2;  $117.55 \pm 13.84\%$  respect to baseline;  $n = 6$ ;  $p > 0.05$ ). In addition, basal PPR differences between groups were maintained throughout the experiments (Fig. 2C; basal, control:  $1.55 \pm 0.06$  vs kindled:  $1.24 \pm 0.03$ ;  $p = 0.02$ ), without exhibiting any changes after TBS1 nor TBS2 (Fig. 2D; after 50 min, control:  $111.45 \pm 9.03\%$  of baseline vs kindled:  $103.18 \pm 3.18\%$  of baseline;  $p = 0.63$ ), which is consistent with the previous experiments. The fact that even repeated TBS protocols are not able to induce plastic changes in CA3-CA1 synapses from kindled rats further supports the previous results and





**Fig. 2.** Two consecutive, spaced TBS protocols were unable to induce LTP in kindled hippocampal slices. (A) Superimposed averaged EPSCs traces obtained in basal conditions (1), and after TBS1 (2) and TBS2 (3) in control ( $n = 5$ ) and kindled ( $n = 6$ ) groups. (B) (top) Time course of the changes in EPSC amplitude (%) and (bottom) normalized values of Ri for each cell, measured throughout the experiments. (C) Superimposed averaged EPSCs evoked by the PPR protocol recorded in basal condition (gray trace) and after TBS2 (black trace) from representative cells from control and kindled groups. (D) (left) Summary graphs for control and kindled condition for average TBS-induced potentiation after TBS1 and TBS2 (left), baseline PPR (middle) measured basal and after TBS2 and PPR change (%; right), measured before and 40 min after the protocol. Results expressed as mean  $\pm$  SEM. \*  $p < 0.05$ .

suggests that there is likely no room for these synapses to be potentiated by TBS-dependent mechanisms.


### 3.3. Impairment in HFS-induced LTP in hippocampal glutamatergic synapses from kindled rats

It has been proposed that astrocytes modulate synaptic plasticity and neuronal excitability not only at a single-synapse level but also at a neural circuit level (Ma et al., 2016; Ben Haim and Rowitch, 2017). If that is the case, the abnormal pattern of astroglial  $Ca^{2+}$ -dependent activity observed in kindled slices should affect the excitability and plasticity of large neuronal populations, which would be reflected in extracellular field recordings. In order to address that, we recorded field excitatory postsynaptic potentials (fEPSP) before and after the application of HFS – which is a stronger plasticity protocol compared to TBS – and population spikes. First, we compared fEPSP input-output relationships between control and kindled slices. Superimposed field responses showed that for a given fiber volley amplitude, the fEPSP is consistently larger in kindled rats, which is reflected by the input-output relationships (Fig. 3A). Second, we evaluated the population spike threshold, which has been reliably used as an indicator of post-synaptic excitability (Andersen et al., 1980). The stimulation intensity required to induce a population spike in the kindled group was consistently lower compared to the control group (Fig. 3B;  $0.29 \pm 0.17 \mu V$ ,  $n = 10$  in kindling vs  $0.51 \pm 0.01 \mu V$ ,  $n = 10$  in control;  $p = 0.032$ ). We next evaluated the relative changes in fEPSP amplitude after the application of the HFS protocol. In control rats HFS induced a short period of post-tetanic facilitation followed by a progressive enhancement in synaptic transmission, which reached its peak after 40–50 min ( $167.82 \pm 8.13\%$  of baseline,  $p < 0.001$ ;  $n = 8$ ; Fig. 3C–D). Kindled slices also exhibited a slight period of post-tetanic

facilitation, however, the subsequent LTP was of significantly lesser magnitude compared to the control group ( $125.51 \pm 7.81\%$  of baseline,  $p < 0.001$   $n = 6$ ; Fig. 3C–D). Again, in both groups the PPR remained unchanged after the protocol (Control basal;  $1.37 \pm 0.14$  and HFS;  $1.32 \pm 0.20$ ;  $n = 10$ ;  $P > 0.05$  vs Kindling basal;  $1.10 \pm 0.05$  and HFS;  $1.08 \pm 0.04$ ;  $n = 8$ ;  $P > 0.05$ ; Fig. 3D). Taken together, these findings suggest that the impairments in synaptic plasticity and excitability observed in the kindled group affects hippocampal neuronal circuits in a widespread manner, which could be associated with the abnormal pattern of  $Ca^{2+}$ -dependent activity in the astroglial network.

### 3.4. Rescuing the normal pattern of astrocytic $Ca^{2+}$ -dependent activity restores TBS-induced LTP in the epileptic hippocampus

Since we recently showed that P2Y1R inhibition restores the FTs/STs relationship and thus the normal pattern of astroglial  $Ca^{2+}$  activity in the kindled hippocampus (Álvarez-Ferradas et al., 2015; Wellmann et al., 2018), we assessed if MRS2179 can also rescue the impairment in TBS-induced LTP in CA3-CA1 synapses. Control experiments were also performed in the presence of MRS2179 to rule out other potential effects of the purinergic antagonist over synaptic plasticity. As predicted, P2Y1R blockade rescued TBS-induced LTP in the kindled group, reaching a magnitude comparable to the observed in the control group (Fig. 4D; kindled:  $160.42 \pm 12.11\%$  of baseline;  $n = 6$  vs control:  $149.70 \pm 23.51\%$  of baseline;  $n = 7$ ;  $p = 0.72$ ). MRS2179 also restored the basal PPR in the kindled group (Fig. 4D; kindled:  $1.47 \pm 0.05$ ;  $n = 4$  vs control:  $1.54 \pm 0.07$ ;  $n = 5$ ;  $p = 0.26$ ), which in both groups remained unchanged after the protocol (Fig. 4; kindling:  $106.33 \pm 23.51\%$  of baseline;  $n = 7$  vs control:  $98.25 \pm 6.31\%$  of baseline;  $n = 6$ ;  $p = 0.07$ ). Concurrently, P2Y1R blockade restored the



**Fig. 3.** Impairment in HFS-induced LTP in kindled hippocampal slices. (A) (left) Superimposed averaged fEPSPs traces with matching fiber volley amplitudes from control (thin trace) and kindled (thick trace) slices. (right) fEPSPs amplitudes plotted against the corresponding fiber volley amplitudes from control (open circles) and kindled (black circles) slices, showing the input-output relation. (B) (left) Superimposed, magnified representative fEPSPs traces showing the population spike threshold in control and kindled conditions. (right) Population spike threshold means values for both control and kindled groups. (C) (top) Superimposed representative fEPSP traces obtained before (1) and after (2) the HFS protocol in control and kindled condition. (bottom) Time course of the changes in fEPSPs amplitude (%) before (1) and after (2) HFS from control and kindled condition. (D). (left) Summary graphs for average HFS-induced LTP (%), (middle) PPR absolute and (right) PPR change (%), measured before (1) and after HFS (2). Results expressed as mean  $\pm$  SEM. \*  $p < 0.05$ .


pattern of astroglial  $Ca^{2+}$ -dependent activity in the kindled group (Fig. 4E), exhibiting a coupled, time-locked increase in STs oscillations (Fig. 4F;  $4.1 \pm 0.88$  before vs  $6.33 \pm 1.32\%$  after;  $n = 62$ ;  $p = 0.02$ ) in parallel to a decrease in the FT/ST ratio (Fig. 4F;  $3.91 \pm 0.53$  before vs  $1.34 \pm 0.25$  after;  $p = 0.01$ ), which was also observed in the control group with MRS2179 (Fig. 4F). These findings indicate that the impairment in TBS-induced LTP observed in the kindling group could be directly associated with the abnormal pattern of  $Ca^{2+}$ -dependent signaling from astrocytes. To confirm if the impairment in LTP is linked to P2Y1R activation, we employed the protocol consisting in two spaced (30 min inter-stimulus interval), consecutive TBSs again, but performed TBS2 in the presence of MRS2179. If P2Y1R blockade is sufficient condition to rescue TBS-induced LTP, pyramidal neurons from kindled slices should potentiate with TBS2 plus MRS2179 after TBS1 has failed to induce LTP. Indeed, while TBS1 alone had no effect, after 10 min incubation with MRS2179, TBS2 induced a strong LTP (Fig. 5A–B: TBS1:  $113.33 \pm 5.88\%$  of baseline vs TBS2 + MRS2179:  $185.42 \pm 16.19\%$  of baseline;  $n = 5$ ;  $p = 0.015$ ). As expected, P2Y1R blockade not only rescued LTP but also increased the PPR (Fig. 5C–D: baseline:  $1.21 \pm 0.09$ ; TBS1:  $1.22 \pm 0.06$ ; TBS2 plus MRS2179:  $1.54 \pm 0.11$ ;  $n = 5$ ;  $p = 0.007$ ). Altogether, these findings indicate that the inhibition of the purinergic signals (i.e.: P2Y1R) underlying astroglial  $Ca^{2+}$ -dependent hyper-excitability rescues TBS-induced LTP, suggesting that the overall pattern of spontaneous astroglial  $Ca^{2+}$  signals is crucial for the maintenance of the plastic properties of CA3-CA1 synapses.

#### 4. Discussion

Here we have followed up on our previous studies showing that astrocytes in the epileptic hippocampus exhibit an abnormal pattern of  $Ca^{2+}$  activity that requires P2Y1R (Wellmann et al., 2018), which causes an upregulation in glutamate gliotransmission and thus enhances the Pr of CA3-CA1 synapses through presynaptic mGluR1/5 activation (Álvarez-Ferradas et al., 2015). Our current findings indicate that this anomalous purinergic-mediated pattern of astroglial  $Ca^{2+}$  signaling is also associated with an impairment in LTP at CA3-CA1 synapses, and could operate as an aberrant mechanism of astrocyte-dependent metaplasticity that interferes with synaptic plasticity in the hippocampus. This dysregulation in astrocyte-neuron crosstalk – which can be reversed by controlling astrocyte-to-astrocyte purinergic signaling – could work as one of the mechanisms deteriorating synaptic plasticity in the epileptic brain and explain, at least partially, the development of cognitive symptoms in epileptic patients.

##### 4.1. Synaptic plasticity impairments are associated with an abnormal pattern of astroglial $Ca^{2+}$ -dependent signals in the epileptic hippocampus


Memory and learning alterations in epileptic patients have been associated with aberrant functional and structural neuroplasticity mechanisms, including changes in synaptic efficacy (i.e.: upregulated glutamatergic tone) as well as neuronal circuitries remodeling, both of which considered hallmarks of the epileptic brain (Chauvière, 2020;



**Fig. 4.** Inhibition of P2Y1R-mediated purinergic signals rescues the normal pattern of astrocytic  $Ca^{2+}$ -activity and recovers LTP in kindled hippocampal slices. (A) Average EPSC traces before (black trace) and 40 min after TBS application (gray trace) recorded in the presence of the specific P2Y1R antagonist MRS2179 (100  $\mu$ M) from control ( $n = 6$ ) and kindled ( $n = 7$ ) groups. (B) (top) Time course of the changes in EPSC amplitude (%) before and after TBS in the presence of MRS2179 and (bottom) normalized values of input resistance (Ri) for each cell, measured throughout the experiments. (C) Superimposed averaged EPSCs evoked by the PPR protocol recorded before (black trace) and after (gray trace) TBS performed with MRS2179 in control and kindled condition. (D) Summary graphs for average TBS-induced potentiation (left), baseline PPR (middle) and PPR change (%; right), measured before and 40 min after the protocol. (E) Fluorescence traces of spontaneous  $Ca^{2+}$  elevations from seven representative astrocytes recorded before, during and after the application of the TBS protocol in control perfused with MRS2179 (left), kindled in regular ASCF (middle) and kindled with MRS2179 (right). Blue boxes denote  $Ca^{2+}$ -dependent activity during TBS protocols. (F) Summary graphs showing mean values of FT/ST ratio (fast  $Ca^{2+}$  transients/slow  $Ca^{2+}$  transients) and average number of slow  $Ca^{2+}$  transients (ST) per astrocyte for control plus MRS 2179 (left), kindled in regular ASCF (middle) and kindled perfused with MRS2179 (right). Results expressed as mean  $\pm$  SEM. \*  $p < 0.05$ . (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

Mikkonen et al., 1998; Postnikova et al., 2019). Here we showed that neuroplasticity alterations in epilepsy extend beyond the neuronal domain and also affect  $Ca^{2+}$ -dependent activity in astrocytes, with neurons exhibiting an impairment in TBS-induced LTP and astrocytes showing an unresponsive, aberrant P2Y1R-mediated pattern of  $Ca^{2+}$  signals during the TBS protocol that is associated with the impairment

in synaptic plasticity. In fact, several studies have revealed that astrocytes are actively involved in the pathophysiology of epilepsy, showing not only structural changes (i.e.: reactive astrogliosis, Fedele et al., 2005; Shigetomi et al., 2019) but also functional abnormalities in  $Ca^{2+}$ -dependent signaling and gliotransmission that affect synaptic transmission (Álvarez-Ferradas et al., 2015; Alves et al., 2017; Nikolic et al.,



**Fig. 5.** The inhibition of P2Y1R recovers TBS-induced LTP in pyramidal neurons from kindled slices that were unable to express LTP with TBS alone. (A) Average EPSC traces recorded at baseline (1), 30 min after the first TBS (2), and 30 min after the second TBS (3) in presence of MRS2179 (100  $\mu$ M) from the kindled group ( $n = 5$ ). (B) (top) Time course of the changes in EPSC amplitude (%) before TBS1 (1), after TBS1 (2) and after TBS2 + MRS2179 in kindled slices, and (bottom) normalized values of input resistance (Ri) for each cell, measured throughout the experiments. (C) Superimposed averaged representative EPSC traces evoked by the PPR protocol recorded before (gray trace) and after TBS1 and TBS2 + MRS2179 (black trace) in the kindled group. (D) (left) Summary graphs for average TBS-induced potentiation and (right) PPR measured at baseline, after TBS1 and after TBS2 + MRS2179. Results expressed as mean  $\pm$  SEM. \*  $p < 0.05$ .

2019; Wellmann et al., 2018; see also Riquelme et al., 2020). Nonetheless, in spite that emerging evidence has shown that both spontaneous and evoked astroglial  $\text{Ca}^{2+}$ -dependent signals directly modulate several forms of short and long-term synaptic plasticity (Bonansco et al., 2011; Navarrete et al., 2012; Perea and Araque, 2007; Sherwood et al., 2017), it was not clear if and how astrocytes could contribute to the alterations in synaptic plasticity and thus to the cognitive decline observed in epileptic patients. Previous works have shown that the electrical stimulation of the Schaffer collaterals – the main glutamatergic afferents of CA1 pyramidal neurons – induces stimuli-coupled, transient astroglial  $\text{Ca}^{2+}$  signals (Fig. 1E; see also Perea and Araque, 2005; Honsek et al., 2012; Navarrete et al., 2012; Sibille et al., 2015) and gliotransmitter release, which in turn modulate synaptic plasticity in the healthy hippocampus (Covelo and Araque, 2018). Concordantly, our results indicate that astrocytes from both control and kindled slices respond to TBS with an increase in  $\text{Ca}^{2+}$ -dependent signals. However, astrocytes from control slices did not only exhibit an increase in  $\text{Ca}^{2+}$ -dependent activity but also a shift in the overall pattern of  $\text{Ca}^{2+}$ -signals, with STs showing a greater relative increment compared to FTs as revealed by the higher STs incidence accompanied by a decrease in the FT/ST ratio (Fig. 1E). In contrast, astrocytes from kindled slices showed an aberrant pattern of spontaneous astroglial  $\text{Ca}^{2+}$  transients in basal conditions, characterized by a higher incidence of long-lasting  $\text{Ca}^{2+}$  events (STs) that persisted during the TBS protocol, without any modifications in the overall  $\text{Ca}^{2+}$  pattern as it is indicated by the unchanged FT/ST ratio. Moreover, the pattern of  $\text{Ca}^{2+}$ -dependent activity observed in the kindled group in basal conditions is comparable to the one exhibited by control astrocytes during the TBS protocol, suggesting that, at rest, astrocytes from the kindled group are as active as if they were subjected to an intense synaptic input. These observations are consistent with the lower PPR values exhibited here by the kindling group, since it has been repeatedly shown that astroglial  $\text{Ca}^{2+}$ -

dependent activity regulate the Pr of C3-CA1 synapses through glutamate release and group I mGluRs activation at presynaptic terminals (Bonansco et al., 2011; Covelo and Araque, 2018; Panatier et al., 2011; Perea and Araque, 2007). In fact, astroglial-induced chronic modifications in the Pr and group I mGluRs signaling could account for the alterations in synaptic plasticity observed here. It has been shown that mGluRs activation in hyperexcitable conditions (zero  $\text{Mg}^{+2}$  or high concentrations of picrotoxin) can induce long-term depression by modulating voltage-dependent  $\text{Ca}^{2+}$  channels and  $\text{Ca}^{2+}$  release from intracellular stores in presynaptic terminals (Palmer et al., 1997; Schwartz and Alford, 2000). In addition, previous reports have shown that experimentally increasing the Pr by employing a high  $\text{Ca}^{2+}$  medium reduces the magnitude of LTP (Bolshakov and Siegelbaum, 1995; Schulz, 1997). Since astroglial  $\text{Ca}^{2+}$ -dependent hyperexcitability enhances glutamate gliotransmission and thus the Pr of CA3-CA1 synapses by tonically activating group I mGluRs (Álvarez-Ferradas et al., 2015), it is possible that stimulation protocols that normally induce LTP (TBS or HFS) trigger cellular and molecular mechanisms with opposing effects – depression vs potentiation – in the epileptic brain, canceling the induction of synaptic plasticity. Supporting this idea, our data shows that in kindled slices rescuing the normal pattern of astroglial  $\text{Ca}^{2+}$  activity restores the plastic properties of CA3-CA1. Although these results suggest that astroglial  $\text{Ca}^{2+}$ -dependent hyper-excitability could explain the impairment in synaptic plasticity by several mechanisms that include changes in glutamate gliotransmission, group I mGluRs activation, and Pr upregulation, we do not rule out other possible effects mediated by different gliotransmitters at the presynaptic or post-synaptic level.

Several works suggest that the inhibition of spontaneous astroglial  $\text{Ca}^{2+}$  transients with the gliotoxin fluorocitrate, intracellular BAPTA or the  $\text{IP}_3\text{R}$  blocker heparin increases the threshold for LTP, suggesting that the basal pattern of astroglial  $\text{Ca}^{2+}$  activity and gliotransmitter



release tonically tunes the threshold for synaptic plasticity induction (Bonansco et al., 2011; Panatier et al., 2011; Sherwood et al., 2017). Our experiments here allow to suggest that, in addition to this basal astroglial-mediated modulation of synaptic plasticity, the induction of LTP requires an activity-dependent, stimuli-coupled shift in the pattern of astroglial  $\text{Ca}^{2+}$ -dependent transients, which is absent in the hippocampus of epileptic rats. Supporting this idea, restoring the normal pattern of astroglial  $\text{Ca}^{2+}$ -dependent activity alone is enough for CA3-CA1 synapses to recover their plastic properties (Fig. 4). Since it has been suggested that astrocytes actively process synaptic information by exhibiting different patterns of  $\text{Ca}^{2+}$ -dependent activity (Covelo and Araque, 2018; Perea et al., 2016), the unresponsive  $\text{Ca}^{2+}$  activity pattern exhibited by astrocytes in the epileptic hippocampus during the TBS protocol allows to suggest that its capacity to decode and adequately respond to synaptic information is impaired, which could interfere with its modulatory function over neural circuits.

#### 4.2. The pattern of P2YRs-mediated astroglial $\text{Ca}^{2+}$ signals modulates synaptic plasticity in the epileptic hippocampus

Our experiments here and previous evidence suggest that the abnormal pattern of astroglial  $\text{Ca}^{2+}$  activity observed in the hippocampus of epileptic rats is driven by purinergic astrocyte-to-astrocyte signals that require the activation of P2Y1R (Fig. 4; Álvarez-Ferradas et al., 2015; Wellmann et al., 2018; see also Riquelme et al., 2020). It has been previously indicated that ATP-mediated purinergic signaling acts as the main pathway for the propagation of  $\text{Ca}^{2+}$ -dependent signals between astrocytes in several neuropathologies. In those conditions, ATP reaches the extracellular medium mostly through hemichannels and, by activating metabotropic P2Y1 and P2Y2 receptors (Bowser and Khakh, 2007; Cieřlak et al., 2017; Pascual et al., 2005), allows for the generation of aberrant astrocyte-to-astrocyte  $\text{Ca}^{2+}$  signals in a wide range of neuropathologies, including epilepsy (Shigetomi et al., 2019; Riquelme et al., 2020). These network-wide, purinergic-mediated alterations in astroglial  $\text{Ca}^{2+}$ -dependent activity could upregulate glutamate gliotransmission and not only abnormally modulate the Pr, as previously indicated, both also enhance synchronous activity and excitability of local neural circuits (Gómez-Gonzalo et al., 2010; Lee et al., 2013; Tian et al., 2005), which is one of the hallmarks of the epileptic brain.

As mentioned earlier, our results here showed that the synaptic activity triggered as prompted during TBS did not modify the global pattern of astrocytic  $\text{Ca}^{2+}$  transients in the kindled group. This specific finding suggests that  $\text{Ca}^{2+}$ -dependent gliotransmission could be increased and occur in a stimuli-uncoupled fashion in the epileptic hippocampus, meaning that is not elicited by activity-dependent neurotransmission. This could upregulate glutamate and ATP release from the astroglial network, modifying the induction threshold, degree, or direction of activity-dependent changes in synaptic efficacy (Jones, 2015; Notenboom et al., 2010; van Dam et al., 2004). In fact, in the kindled group, P2Y1Rs inhibition not only restored the basal pattern of astroglial  $\text{Ca}^{2+}$ -dependent activity and the PPR rate, but also the normal astroglial response to the TBS protocol and the plastic properties of CA3-CA1 synapses. Moreover, our results show that MRS2179 has an effect over the pattern of  $\text{Ca}^{2+}$ -dependent activity only in astrocytes from the kindled condition, suggesting that P2Y1R activation is a feature specifically associated with abnormal astroglial  $\text{Ca}^{2+}$  signals. In fact, recent evidence allows to suggest that the effects of MRS2179 are mostly mediated by the inhibition of astroglial and not neuronal P2Y1R: (i) ATP is released by astrocytes as a mechanism for autocrine and paracrine signaling, being P2Y1R activation one of the main pathways for astrocyte-to-astrocyte communication in pathological conditions (Riquelme et al., 2020; Shen et al., 2014); (ii) P2Y1R activation induces slow astroglial  $\text{Ca}^{2+}$  transients (Álvarez-Ferradas et al., 2015; Wellmann et al., 2018; Shigetomi et al., 2019); (iii) P2Y1R are preferentially and mostly expressed by astrocytes in the hippocampus

(Di Castro et al., 2011); (iv) P2Y1R are overexpressed in several animal models of experimental epilepsy and in brain tissue from epileptic patients (Alves et al., 2017); and (v) P2Y1R activation exacerbates epileptiform activity and seizure severity (Alves et al., 2020). Therefore, the dysregulation of the purinergic signals that mediate astrocyte-to-astrocyte communication alone could alter astroglial  $\text{Ca}^{2+}$ -dependent activity and thus profoundly disrupt not only the modulatory effect that astrocytes exert over basal synaptic transmission, but also modify the plastic properties of neighboring neural circuits.

Extracellular levels of adenosine, which acts as an endogenous anticonvulsant via adenosine type 1 receptor (A1Rs) activation, are also affected in the epileptic brain, exhibiting a marked decrease due to the overexpression of adenosine kinase (ADK) (Etherington and Frenguelli, 2004; Fedele et al., 2005). The decrease in A1Rs activation promotes glutamate release and postsynaptic depolarization, leading to an increase in neuronal excitability and epileptiform activity (Aronica and Crino, 2011; Shen et al., 2014). Since ATP/adenosine homeostasis is controlled exclusively by astrocytes, its alteration could chronically decrease adenosine availability and increase extracellular levels of ATP (Boison, 2012; Dossi et al., 2018), having a joint effect that could lower seizure threshold. These alterations in astroglial purinergic signaling and metabolism could also explain the reduced population spike threshold and larger fEPSPs exhibited by kindled rats (Fig. 2A-D), which is consistent with the increased firing frequency exhibited by CA1 pyramidal neurons in our kindling model (Morales et al., 2014). Altogether, these findings suggest that the impairment in LTP observed in the epileptic hippocampus is associated with an aberrant pattern of astroglial  $\text{Ca}^{2+}$ -dependent activity, which is reversed by the inhibition of P2Y1R mediated signals, restoring glutamatergic neurotransmission at rate that resembles physiological conditions. In this scenario, we propose that the increase in P2Y1R-mediated astrocyte-to-astrocyte purinergic signals – and probably the decrease in adenosinergic signals – causes an aberrant pattern of astroglial  $\text{Ca}^{2+}$ -dependent activity, which prevents astrocytes from decoding synaptic activity and normally modulate LTP (Covelo and Araque, 2018; Di Castro et al., 2011; Perea and Araque, 2005; Sibille et al., 2015). However, further studies are necessary to test whether and how different gliotransmitters, such as D-serine or ATP/adenosine (Henneberger et al., 2010), could alter astrocyte-mediated metaplasticity in the epileptic hippocampus.

Several evidence suggest that the repetitive activation of glutamatergic inputs can lead to LTP saturation (Morgan and Teyler, 2001; Schulz and Fitzgibbons, 1997), reaching the neural pathway a state in which no further potentiation is feasible and thus disrupting learning and memory encoding (Gruart, 2006; Leung and Shen, 2006; Moser, 1998). Chronic epilepsy induction – as in kindling models – requires the repeated activation of excitatory projections that arrive to the hippocampal formation (Morales et al., 2014; Morimoto et al., 2004), which resembles the process required to induce LTP saturation. The brief, low intensity trains of electrical stimulation employed in kindling protocols – delivered in a frequency similar to that used to induce LTP – could cause a permanent state of electroconvulsive seizures, which can also reduce the capacity to generate LTP (Reid and Stewart, 1997). We observed that the application of two consecutive TBS protocols failed to induce LTP in the kindled group, which could suggest that the impairment is indeed due to an on-demand state of synaptic saturation. However, after a first attempt that failed to induce LTP, the application of the second TBS in the presence of MRS2179 induced a robust LTP, suggesting that the mechanism that underlies the impairment in LTP is likely different from synaptic saturation.

In conclusion, our results here suggest that the abnormal pattern of astroglial  $\text{Ca}^{2+}$ -dependent signals in the epileptic hippocampus is associated with an impairment in TBS-induced LTP in CA3-CA1 synapses, and likely operates as an aberrant mechanism of astrocyte-dependent metaplasticity. The restoration of the normal pattern of astroglial  $\text{Ca}^{2+}$ -dependent activity by the blockade of P2Y1R-mediated signals rescued TBS-induced LTP, suggesting that the alterations in synaptic plasticity

in epileptic circuits could be largely associated to an aberrant crosstalk between astrocytes. By means of an aberrant pattern of  $\text{Ca}^{2+}$ -dependent activity and an upregulation in glutamate gliotransmission, astrocytes could not only abnormally increase the Pr, neuronal synchrony and excitability, but also interfere with the induction of synaptic plasticity in epileptic circuits, which constitutes a novel, key mechanism that could explain the cognitive decline in epileptic patients.

## Acknowledgments

The author would like to thank Dr. Gertrudis Perea and Dr. Andrés Chavez for their critical comments and suggestions for the present manuscript. This work was supported by Centro de Neurobiología y Fisiología Integrativa CENFI, UV (Grant DIUV-CI 01/2006) and Grants 1130491 from FONDECYT, CONICYT, Chile to Christian Bonansco, and 11171006 from FONDECYT, Conicyt, Chile, Millennium Nucleus NUMIND NC-130011 and Centro de Neurobiología y Fisiología Integrativa, CENFI, UV (Grant DIUV-CI 01/2006) to Marco Fuenzalida.

## References

- Álvarez-Ferradas, C., Morales, J.C., Wellmann, M., Nualart, F., Roncagliolo, M., Fuenzalida, M., Bonansco, C., 2015. Enhanced astroglial  $\text{Ca}^{2+}$  signaling increases excitatory synaptic strength in the epileptic brain. *Glia* 63 (1507), 1521.
- Alves, M., Gomez-Villafuertes, R., Delanty, N., Farrell, M.A., O'Brien, D.F., Miras-Portugal, M.T., Hernandez, M.D., Henshall, D.C., Engel, T., 2017. Expression and function of the metabotropic purinergic P2Y receptor family in experimental seizure models and patients with drug-refractory epilepsy. *Epilepsia* 58, 1603–1614.
- Alves, M., Smith, J., Engel, T., 2020. Differential expression of the metabotropic P2Y receptor family in the cortex following status epilepticus and neuroprotection via P2Y1 antagonism in mice. *Front. Pharmacol.* 10, 1558.
- Andersen, P., Silfvenius, H., Sundberg, S.H., Sveen, O., 1980. A comparison of distal and proximal dendritic synapses on CA1 pyramids in Guinea-pig hippocampal slices *in vitro*. *J. Physiol.* 307, 273–299.
- Aronica, E., Crino, P.B., 2011. Inflammation in epilepsy: clinical observations. *Epilepsia* 52, 26–32.
- Beck, H., Goussakov, I.V., Lie, A., Helmstaedter, C., Elger, C.E., 2000. Synaptic plasticity in the human dentate gyrus. *J. Neurosci.* 20, 7080–7086.
- Ben Haim, L., Rowitch, D.H., 2017. Functional diversity of astrocytes in neural circuit regulation. *Nat. Rev. Neurosci.* 18, 31–41.
- Boison, D., 2012. Adenosine dysfunction in epilepsy. *Glia* 60, 1234–1243.
- Bolshakov, V., Siegelbaum, S., 1995. Regulation of hippocampal transmitter release during development and long-term potentiation. *Science* 269, 1730–1734.
- Bonansco, C., Buño, W., 2003. Cellular mechanisms underlying the rhythmic bursts induced by NMDA microiontophoresis at the apical dendrites of CA1 pyramidal neurons. *Hippocampus* 13, 150–163.
- Bonansco, C., Fuenzalida, M., Olivares, V., Molina, C., Roncagliolo, M., 2007. Asynchronous transmission in the CA3-CA1 hippocampal synapses in the neurological mutant taiep rat. *J. Neurosci. Res.* 85, 223–229.
- Bonansco, C., Couve, A., Perea, G., Ferradas, C.A., Roncagliolo, M., Fuenzalida, M., 2011. Glutamate released spontaneously from astrocytes sets the threshold for synaptic plasticity. *Eur. J. Neurosci.* 33, 1483–1492.
- Bowser, D.N., Khakh, B.S., 2007. Vesicular ATP is the predominant cause of intercellular calcium waves in astrocytes. *J. Gen. Physiol.* 129, 485–491.
- Campo, P., Garrido, M.L., Moran, R.J., García-Morales, I., Poch, C., Toledano, R., Gil-Nagel, A., Dolan, R.J., Friston, K.J., 2013. Network reconfiguration and working memory impairment in mesial temporal lobe epilepsy. *NeuroImage* 72, 48–54.
- Cavus, I., Kasoff, W.S., Cassaday, M.P., Jacob, R., Gueorguieva, R., Sherwin, R.S., Abi-Saab, W.M., 2005. Extracellular metabolites in the cortex and hippocampus of epileptic patients. *Ann. Neurol.* 57 (2), 226–235.
- Celiker Uslu, S., Yuksel, B., Tekin, B., Sariahmetoglu, H., Atakli, D., 2019. Cognitive impairment and drug responsiveness in mesial temporal lobe epilepsy. *Epilepsy Behav.* 90, 162–167.
- Chauvière, L., 2020. Potential causes of cognitive alterations in temporal lobe epilepsy. *Behav. Brain Res.* 27 (378), 112310.
- Cieslak, M., Wojtczak, A., Komoszyński, M., 2017. Role of the purinergic signalling in epilepsy. *Pharmacol. Rep.* 69, 130–138.
- Covelo, A., Araque, A., 2018. Neuronal activity determines distinct gliotransmitter release from a single astrocyte. *Elife.* 30, 7.
- De Jonge, M., Racine, R.J., 1987. The development and decay of kindling-induced increases in paired-pulse depression in the dentate gyrus. *Brain Res.* 412, 318–328.
- Di Castro, M.A., Chuquet, J., Liaudet, N., Bhaukaurally, K., Santello, M., Bouvier, D., Tiret, P., Volterra, A., 2011. Local  $\text{Ca}^{2+}$  detection and modulation of synaptic release by astrocytes. *Nat. Neurosci.* 14, 1276–1284.
- Dossi, E., Blauwblomme, T., Moulard, J., Chever, O., Vasile, F., Guinard, E., Le Bert, M., Couillien, I., Pallud, J., Capelle, L., Huberfeld, G., Rouach, N., 2018. Pannexin-1 channels contribute to seizure generation in human epileptic brain tissue and in a mouse model of epilepsy. *Sci. Transl. Med.* 10, 443 pii: eaar3796.
- Esmaeilpour, K., Sheibani, V., Shabani, M., Mirnajafi-Zadeh, J., 2017. Effect of low frequency electrical stimulation on seizure-induced short- and long-term impairments in learning and memory in rats. *Physiol. Behav.* 168, 112–121.
- Etherington, L.A.V., Frenguelli, B.G., 2004. Endogenous adenosine modulates epileptiform activity in rat hippocampus in a receptor subtype-dependent manner. *Eur. J. Neurosci.* 19, 2539–2550.
- Fedele, D.E., Gouder, N., Güttinger, M., Gabernet, L., Scheurer, L., Rüllicke, T., Crestani, F., Boison, D., 2005. Astroglialosis in epilepsy leads to overexpression of adenosine kinase, resulting in seizure aggravation. *Brain* 128, 2383–2395.
- Fisher, R.S., van Boas, W.E., Blume, W., Elger, C., Genton, P., Lee, P., Engel, J., 2005. Epileptic seizures and epilepsy: definitions proposed by the international league against epilepsy (ILAE) and the International Bureau for Epilepsy (IBE). *Epilepsia* 46, 470–472.
- Ghotbeddin, Z., Moazedi, A.A., Yadollahpour, A., Rendi, F., Jalilifar, M., 2018. Improving cognitive task in kindled rats by using low frequency stimulation during epileptogenesis. *Metab. Brain Dis.* 33, 1525–1531.
- Gilbert, M.E., Mack, C.M., 1990. The NMDA antagonist, MK-801, suppresses long-term potentiation, kindling, and kindling-induced potentiation in the perforant path of the unanesthetized rat. *Brain Res.* 519, 89–96.
- Gómez-Gonzalo, M., Losi, G., Chiavegato, A., Zonta, M., Cammarota, M., Brondi, M., Vetri, F., Uva, L., Pozzan, T., de Curtis, M., Ratto, G.M., Carmignoto, G., 2010. An excitatory loop with astrocytes contributes to drive neurons to seizure threshold. *PLoS Biol.* 8, e1000352.
- Gruart, A., 2006. Involvement of the CA3-CA1 synapse in the acquisition of associative learning in behaving mice. *J. Neurosci.* 26, 1077–1087.
- Halgren, E., Stapleton, J., Domalski, P., Swartz, B.E., Delgado-Escueta, A.V., Walsh, G.O., Mandelkern, M., Bland, W., Ropchan, J., 1991. Memory dysfunction in epilepsy patients as a derangement of normal physiology. In: Smith, D.B., Treiman, D.M., Trimble, M.R. (Eds.), *Advances in Neurology*, Vol. 55. Neurobehavioral Problems in Epilepsy. Raven Press, Publishers, pp. 385–410.
- Helmstaedter, C., 2002. Effects of chronic epilepsy on declarative memory systems. *Prog. Brain Res.* 135, 439–453.
- Henneberger, C., Papouin, T., Oliet, S.H., Rusakov, D.A., 2010. Long-term potentiation depends on release of D-serine from astrocytes. *Nature.* 463, 232–236.
- Holmes, G.L., 2015. Cognitive impairment in epilepsy: the role of network abnormalities. *Epileptic Disord.* 2, 101–116.
- Honsek, S.D., Walz, C., Kafitz, K.W., Rose, C.R., 2012. Astrocyte calcium signals at Schaffer collateral to CA1 pyramidal cell synapses correlate with the number of activated synapses but not with synaptic strength. *Hippocampus* 22 (1), 29–42.
- Hu, B., Karnup, S., Zhou, L., Stelzer, A., 2005. Reversal of hippocampal LTP by spontaneous seizure-like activity: role of group I mGluR and cell depolarization. *J. Neurophysiol.* 93, 316–336.
- Jones, O.D., 2015. Astrocyte-mediated metaplasticity in the hippocampus: help or hindrance? *Neuroscience* 309, 113–124.
- Kafitz, K.W., Meier, S.D., Stephan, J., Rose, C.R., 2008. Developmental profile and properties of sulforhodamine 101-labeled glial cells in acute brain slices of rat hippocampus. *J. Neurosci. Methods* 169, 84–92.
- Krug, M., Koch, M., Grecksch, G., Schulzeck, K., 1997. Pentylentetrazol kindling changes the ability to induce potentiation phenomena in the hippocampal ca1 region. *Physiol. Behav.* 62, 721–727.
- Larson, J., Munkácsy, E., 2015. Theta-burst LTP. *Brain Res.* 1621, 38–50.
- Lee, H.U., Yamazaki, Y., Tanaka, K.F., Furuya, K., Sokabe, M., Hida, H., Takao, K., Miyakawa, T., Fujii, S., Ikenaka, K., 2013. Increased astrocytic ATP release results in enhanced excitability of the hippocampus. *Glia* 61, 210–224.
- Leung, L.S., Shen, B., 2006. Hippocampal CA1 kindling but not long-term potentiation disrupts spatial memory performance. *Learn. Mem.* 13, 18–26.
- Leung, L.S., Wu, C., 2003. Kindling suppresses primed-burst-induced long-term potentiation in hippocampal CA1. *Neuroreport* 14, 211–214.
- Löscher, W., 2011. Critical review of current animal models of seizures and epilepsy used in the discovery and development of new antiepileptic drugs. *Seizure* 20, 359–368.
- Ma, Z., Stork, T., Bergles, D.E., Freeman, M.R., 2016. Neuromodulators signal through astrocytes to alter neural circuit activity and behaviour. *Nature.* 539, 428–432.
- Maru, E., Goddard, G.V., 1987. Alteration in dentate neuronal activities associated with perforant path kindling. *Exp. Neurol.* 96, 19–32.
- Matsuzaki, M., Honkura, N., Ellis-Davies, G.C.R., Kasai, H., 2004. Structural basis of long-term potentiation in single dendritic spines. *Nature* 429, 761–766.
- McNamara, J.O., Bonhaus, D.W., Shin, C., Crain, B.J., Gellman, R.L., Giacchino, J.L., 1985. The kindling model of epilepsy: a critical review. *CRC Crit. Rev. Clin. Neurobiol.* 1, 341–391.
- Mikkonen, M., Soininen, H., Kälviäinen, R., Tapiola, T., Ylinen, A., Vapalahti, M., Paljärvi, L., Pitkänen, A., 1998. Remodeling of neuronal circuitries in human temporal lobe epilepsy: increased expression of highly polysialylated neural cell adhesion molecule in the hippocampus and the entorhinal cortex. *Ann. Neurol.* 44, 923–934.
- Moore, C.I., Browning, M.D., Rose, G.M., 1993. Hippocampal plasticity induced by primed burst, but not long-term potentiation, stimulation is impaired in area CA1 of aged Fischer 344 rats. *Hippocampus* 3, 57–66.
- Morales, J.C., Álvarez-Ferradas, C., Roncagliolo, M., Fuenzalida, M., Wellmann, M., Nualart, F.J., Bonansco, C., 2014. A new rapid kindling variant for induction of cortical epileptogenesis in freely moving rats. *Front. Cell. Neurosci.* 8, 200.
- Morgan, S.L., Teyler, T.J., 2001. Electrical stimuli patterned after the theta-rhythm induce multiple forms of LTP. *J. Neurophysiol.* 86, 1289–1296.
- Morimoto, K., Fahnestock, M., Racine, R.J., 2004. Kindling and status epilepticus models of epilepsy: rewiring the brain. *Prog. Neurobiol.* 73, 1–60.
- Moser, E.I., 1998. Impaired spatial learning after saturation of long-term potentiation. *Science* 281, 2038–2042.
- Navarrete, M., Perea, G., De Sevilla, D.F., Gómez-Gonzalo, M., Núñez, A., Martín, E.D., Araque, A., 2012. Astrocytes mediate *in vivo* cholinergic-induced synaptic plasticity.

- PLoS Biol. 10, e1001259.
- Nicoll, R.A., Schmitz, D., 2005. Synaptic plasticity at hippocampal mossy fiber synapses. *Nat. Rev. Neurosci.* 6, 863–876.
- Nikolic, L., Nobili, P., Shen, W., Audinat, E., 2019. Role of astrocyte purinergic signaling in epilepsy. *Glia*. <https://doi.org/10.1002/glia.23747>.
- Notenboom, R.G.E., Ramakers, G.M.J., Kamal, A., Spruijt, B.M., De Graan, P.N.E., 2010. Long-lasting modulation of synaptic plasticity in rat hippocampus after early-life complex febrile seizures. *Eur. J. Neurosci.* 32, 749–758.
- Palmer, M.J., Irving, A.J., Seabrook, G.R., Jane, D.E., Collingridge, G.L., 1997. The group I mGlu receptor agonist DHPG induces a novel form of LTD in the CA1 region of the hippocampus. *Neuropharmacology* 36, 1517–1532.
- Panatier, A., Vallée, J., Haber, M., Murai, K.K., Lacaille, J.C., Robitaille, R., 2011. Astrocytes are endogenous regulators of basal transmission at central synapses. *Cell* 146, 785–798.
- Pascual, O., Casper, K.B., Kubera, C., Zhang, J., Revilla-Sanchez, R., Sul, J.Y., Takano, H., Moss, S.J., McCarthy, K., Haydon, P.G., 2005. Astrocytic purinergic signaling coordinates synaptic networks. *Science* 310, 113–116.
- Paxinos, G., Watson, C., 1998. *The Rat Brain in Stereotaxic Coordinates*. Academic Press, San Diego.
- Perea, G., Araque, A., 2005. Properties of synaptically evoked astrocyte calcium signal reveal synaptic information processing by astrocytes. *J. Neurosci.* 25, 2192–2203.
- Perea, G., Araque, A., 2007. Astrocytes potentiate transmitter release at single hippocampal synapses. *Science* 317, 1083–1086.
- Perea, G., Gómez, R., Mederos, S., Covelo, A., Ballesteros, J.J., Schlosser, L., Hernández-Vivanco, A., Martín-Fernández, M., Quintana, R., Rayan, A., Díez, A., Fuenzalida, M., Agarwal, A., Bergles, D.E., Bettler, B., Manahan-Vaughan, D., Martín, E.D., Kirchhoff, F., Araque, A., 2016. Activity-dependent switch of GABAergic inhibition into glutamatergic excitation in astrocyte-neuron networks. *Elife* 5 pii: e20362.
- Pirttimäki, T., Parri, H.R., Crunelli, V., 2013. Astrocytic GABA transporter GAT-1 dysfunction in experimental absence seizures. *J. Physiol.* 591, 823–833.
- Postnikova, T.Y., Trofimova, A.M., Ergina, J.L., Zubareva, O.E., Kalemenev, S.V., Zaitsev, A.V., 2019. Transient switching of NMDA-dependent long-term synaptic potentiation in CA3-CA1 hippocampal synapses to mGluR1-dependent potentiation after pentylenetetrazole-induced acute seizures in young rats. *Cell. Mol. Neurobiol.* 39, 287–300.
- Rai, V.K., Shukla, G., Afsar, M., Poornima, S., Pandey, R.M., Rai, N., Goyal, V., Srivastava, A., Vibha, D., Behari, M., 2015. Memory, executive function and language function are similarly impaired in both temporal and extra temporal refractory epilepsy—A prospective study. *Epilepsy Res.* 109, 72–80.
- Reid, I.C., Stewart, C.A., 1997. Seizures, memory and synaptic plasticity. *Seizure* 6, 351–359.
- Riquelme, J., Wellmann, M., Sotomayor-Zárate, R., Bonansco, C., 2020. Gliotransmission: a novel target for the development of antiseizure drugs. *Neuroscientist*. <https://doi.org/10.1177/107385842090147>.
- Ruethrich, H., Grecksch, G., Becker, A., Krug, M., 1996. Potentiation effects in the dentate gyrus of pentylenetetrazol-kindled rats. *Physiol. Behav.* 60, 455–462.
- Schneiderman, J.H., 1997. The role of long-term potentiation in persistent epileptiform burst-induced hyperexcitability following GABAA receptor blockade. *Neuroscience* 1997 (81), 1111–1122.
- Schubert, M., Siegmund, H., Pape, H.C., Albrecht, D., 2005. Kindling-induced changes in plasticity of the rat amygdala and hippocampus. *Learn. Mem.* 12, 520–526.
- Schulz, P.E., 1997. Long-term potentiation involves increases in the probability of neurotransmitter release. *Proc. Natl. Acad. Sci. U. S. A.* 94, 5888–5893.
- Schulz, P.E., Fitzgibbons, J.C., 1997. Differing mechanisms of expression for short- and long-term potentiation. *J. Neurophysiol.* 78, 321–334.
- Schwartz, N.E., Alford, S., 2000. Physiological activation of presynaptic metabotropic glutamate receptors increases intracellular calcium and glutamate release. *J. Neurophysiol.* 84, 415–427.
- Shen, H.Y., Sun, H., Hanthorn, M.M., Zhi, Z., Lan, J.Q., Poulsen, D.J., Wang, R.K., Boison, D., 2014. Overexpression of adenosine kinase in cortical astrocytes and focal neocortical epilepsy in mice. *J. Neurosurg.* 120, 628–638.
- Sherwood, M.W., Arizono, M., Hisatsune, C., Bannai, H., Ebisui, E., Sherwood, J.L., Panatier, A., Oliet, S.H., Mikoshiba, K., 2017. Astrocytic IP<sub>3</sub> Rs: contribution to Ca<sup>2+</sup> signalling and hippocampal LTP. *Glia* 65, 502–513.
- Shigetomi, E., Saito, K., Sano, F., Koizumi, S., 2019. Aberrant calcium signals in reactive astrocytes: a key process in neurological disorders. *Int. J. Mol. Sci.* 20, 996.
- Sibille, J., Zapata, J., Teillon, J., Rouach, N., 2015. Astroglial calcium signaling displays short-term plasticity and adjusts synaptic efficacy. *Front. Cell. Neurosci.* 9, 189.
- Soukupova, M., Binaschi, A., Falcicchia, C., Palma, E., Roncon, P., Zucchini, S., Simonato, M., 2015. Increased extracellular levels of glutamate in the hippocampus of chronically epileptic rats. *Neuroscience* 301, 246–253.
- Stasheff, S., Anderson, W., Clark, S., Wilson, W., 1989. NMDA antagonists differentiate epileptogenesis from seizure expression in an in vitro model. *Science* 245, 648–651.
- Steward, O., Torre, E.R., Tomasulo, R., Lothman, E., 1991. Neuronal activity up-regulates astroglial gene expression. *Proc. Natl. Acad. Sci. U. S. A.* 88, 6819–6823.
- Suárez, L.M., Cid, E., Gal, B., Inostroza, M., Brotons-Mas, J.R., Gómez-Domínguez, D., de la Prida, L.M., Solís, J.M., 2012. Systemic injection of kainic acid differently affects LTP magnitude depending on its epileptogenic efficiency. *PLoS One* 7, e48128.
- Sutula, T., He, X., Cavazos, J., Scott, G., 1988. Synaptic reorganization in the hippocampus induced by abnormal functional activity. *Science* 239, 1147–1150.
- Tian, G.-F., Azmi, H., Takano, T., Xu, Q., Peng, W., Lin, J., Oberheim, N., Lou, N., Wang, X., Zielke, H.R., Kang, J., Nedergaard, M., 2005. An astrocytic basis of epilepsy. *Nat. Med.* 11, 973–981.
- van Dam, E.J., Kamal, A., Artola, A., de Graan, P.N., Gispen, W.H., Ramakers, G.M., 2004. Group I metabotropic glutamate receptors regulate the frequency-response function of hippocampal CA1 synapses for the induction of LTP and LTD. *Eur. J. Neurosci.* 19, 112–118.
- Verkhatsky, A., Zorec, R., Parpura, V., 2017. Stratification of astrocytes in healthy and diseased brain. *Brain Pathol.* 27, 629–644.
- Wellmann, M., Álvarez-Ferradas, C., Maturana, C.J., Sáez, J.C., Bonansco, C., 2018. Astroglial Ca<sup>2+</sup>-dependent hyperexcitability requires P2Y1 purinergic receptors and pannexin-1 channel activation in a chronic model of epilepsy. *Front. Cell. Neurosci.* 12, 446.