


Accepted Manuscript

Title: rainbow trout surviving infections of viral haemorrhagic septicemia virus (vhsv) show lasting antibodies to recombinant g protein fragments

Authors: P Encinas, E Gomez-Casado, Fregeneda Grandes, N.J Olesen, N Lorenzen, A Estepa, J.M Coll



PII: S1050-4648(11)00029-5

DOI: [10.1016/j.fsi.2011.01.021](https://doi.org/10.1016/j.fsi.2011.01.021)

Reference: YFSIM 1822

To appear in: *Fish and Shellfish Immunology*

Received Date: 27 October 2010

Revised Date: 22 January 2011

Accepted Date: 23 January 2011

Please cite this article as: Encinas P, Gomez-Casado E, Grandes F, Olesen NJ, Lorenzen N, Estepa A, Coll JM. rainbow trout surviving infections of viral haemorrhagic septicemia virus (vhsv) show lasting antibodies to recombinant g protein fragments, *Fish and Shellfish Immunology* (2011), doi: 10.1016/j.fsi.2011.01.021

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

RAINBOW TROUT SURVIVING INFECTIONS OF VIRAL HAEMORRHAGIC SEPTICEMIA VIRUS (VHSV) SHOW LASTING ANTIBODIES TO RECOMBINANT G PROTEIN FRAGMENTS

¹Encinas, P., ¹Gomez-Casado, E. ³Fregeneda-Grandes, ⁴Olesen, N.J., ⁴Lorenzen, N., ²Estepa, A., and
*¹Coll, J. M.

¹INIA, SGIT – Dept Biotecnología Crt. Coruña Km 7. 28040 Madrid, Spain.

²Universidad Miguel Hernández, IBMC. 03202 Elche, Spain

³Dept Sanidad Animal. Universidad León. 24071 León, Spain

⁴National Veterinary Institute, Technical University, Hangevej. 28200 Arhus N, Denmark

keywords,

trout antibodies, viral haemorrhagic septicemia, VHSV, G protein, fragments, survivors

* To whom correspondence should be sent

telph.34-1-3476850/fax 34-1-3572293/email juliocoll@inia.es

Rainbow trout antibodies (Abs) binding to recombinant fragments (frgs) derived from the protein G of the viral haemorrhagic septicemia virus (VHSV)-07.71 strain, could be detected by ELISA (frg-ELISA) in sera from trout surviving laboratory-controlled infections. Abs were detected not only by using sera from trout infected with the homologous VHSV isolate but also with the VHSV-DK-201433 heterologous isolate, which had 13 amino acid changes. Sera from healthy trout and/or from trout surviving infectious haematopoietic necrosis virus (IHNV) infection, were used to calculate cut-off absorbances to differentiate negative from positive sera. Specific anti-VHSV Abs could then be detected by using any of the following frgs: frg11 (56-110), frg15 (65-250), frg16 (252-450) or G21-465. While high correlations were found among the ELISA values obtained with the different frgs, no correlations between any frg-ELISA and complement-dependent 50 % plaque neutralization test (PNT) titres could be demonstrated. Between 4 to 10 weeks after VHSV-infection, more trout sera were detected as positives by using heterologous frg-ELISA rather than homologous PNT. Furthermore, the percentage of positive sera detected by frg11-ELISA increased with time after infection to reach 100 %, while those detected by complement-dependent PNT decreased to 29.4 %, thus confirming that the lack of neutralising Abs does not mean the lack of any anti-VHSV Abs in survivor trout sera. Preliminary results with sera from field samples suggest that further refinements of the frg-ELISA could allow detection of anti-VHSV trout Abs in natural outbreaks caused by different heterologous VHSV isolates. The homologous frg-ELISA method could be useful to follow G immunization attempts during vaccine development and/or to best understand the fish Ab response during VHSV infections. The viral frgs approach might also be used with other fish species and/or viruses.

The *in vitro* complement-dependent neutralisation tests (50 % plaque neutralization test, PNT) are usually employed to estimate protein G-specific antibodies (Abs) to viral haemorrhagic septicemia virus (VHSV)(Dorson and Torchy, 1979; Hattenberg-Baudoy et al., 1995; Jorgensen et al., 1991; Olesen and Jorgensen, 1986). In addition to complement-dependent PNT being highly specific and sensitive, detection of neutralizing Abs generally correlates well with protection to VHSV challenge (Lorenzen and Olesen, 1997; Lorenzen et al., 1993; Lorenzen et al., 1999). However, PNTs are also time-consuming, labour-intensive and require laboratories with cell culture facilities (Lorenzen and LaPatra, 1999; Olesen et al., 1991). Most important, because the VHSV isolate used for the complement-dependent PNT needs to be homologous or closely related to the isolate causing the infection, the assay is being prone to false negatives when applied to natural infections, since it is not always possible to match both isolates (Ahne et al., 1986; Fregeneda-Grandes and Olesen, 2007; Fregeneda-Grandes et al., 2009).

Binding to VHSV-captured enzyme linked immunosorbent assays (ELISA) has been alternatively used to estimate anti-VHSV Abs whether those be neutralising or not (Olesen et al., 1991). Other attempts to detect trout anti-VHSV protein G (G) Abs have used purified VHSV or yeast recombinant protein G4 (amino acid, 9-443) to coat solid-phases for indirect ELISAs (Estepa et al., 1994). However, some of these assays suffered from either high backgrounds and/or low sensitivities and the correlation of their results with protection has not been supported by any data (Lorenzen and Olesen, 1997; Lorenzen et al., 1993; Lorenzen et al., 1999).

To search for possible G short fragments (frgs) as alternative solid phases for indirect ELISAs, a 15-mer overlapping pepscan peptide library of the G protein of VHSV-07.71 was screened for trout Ab reactivity. However, it was not possible to map neutralizing monoclonal Abs by this method and very few peptides specifically reacted with homologous anti-VHSV trout sera (Fernandez-Alonso et al., 1998). Because, those results could have been due to the small size of the peptides used, trout Ab reactivity was then searched in larger G frgs. Thus the recombinant frg11 (amino acid 56-110) implicated in VHSV fusion (Estepa et al., 2001), was another alternative used to estimate anti-G binding Abs (Fernandez-Alonso et al., 1998; Rocha et al., 2002).

Here we have screened sera from trout surviving laboratory-controlled infections with homologous and heterologous VHSV isolates as a first step to explore the possibility of using recombinant G frgs to estimate the presence of anti-VHSV Abs in sera from trout surviving natural infections independently of their neutralizing or non-neutralizing activities. Thus, several recombinant frgs were derived from the VHSV-07.71 french isolate, while the sera were obtained from trout infected with either homologous (VHSV-07.71) or heterologous isolates (VHSV-DK-201433 from Denmark, differing in the protein G sequence by 13 amino acids from VHSV-07.71). In contrast to the Abs detected by complement-dependent PNT, Abs detected by frg-ELISA were found in more sera from trout infected not only with the homologous but also with the heterologous isolate under laboratory-controlled conditions. The results showed that more positive sera could be detected by frg-ELISA than by complement-dependent PNT at any time after infection. Furthermore, while the number of sera with positive ELISA increased, those with positive PNT decreased with time after infection, thus showing that the lack of neutralising Abs does not mean there are no other types of anti-VHSV Abs present in survivor trout sera. Preliminary results obtained with a limited amount of field samples suggest that further refinements could allow detection of anti-VHSV trout Abs in natural outbreaks caused by different heterologous VHSV isolates. This method will be useful to follow immunization attempts during vaccine development and/or the kinetic of fish Ab response following homologous VHSV infections. The antigenic viral frgs approach might also be used with other fish species and/or viruses.

Construction of recombinant fragments (frgs) of the glycoprotein G of viral haemorrhagic septicaemia virus (VHSV). The constructs, expression and purification of frgs of the protein G of VHSV were performed as described (Rocha et al., 2002). Briefly, the G protein sequence (Gene Bank accession number X59148) of the VHSV-07.71 strain, isolated in France from rainbow trout (*Oncorhynchus mykiss*) (LeBerre et al., 1977; Rocha et al., 2005) was used to design the frgs (G21-465, frg11, frg15 and frg16). Their sequences were then chemically synthesized and cloned into pRSET expression vectors that add poly histidine tails. The G21-465 was expressed in *Trichoplusia ni* (cabbage looper) insect larvae (Medin et al., 1995; Perez-Filgueira et al., 2007) while the other fragments frg11 (56-110), frg15 (65-250) and frg16 (252-450) were expressed in *E.coli*. Purification of the recombinant frgs was made by Ni²⁺ affinity chromatography followed by Sephadex chromatography as described (Rocha et al., 2002).

Confirmation of recognition of G frgs by western blotting with trout sera. Confirmation of fragment recognition by trout Abs was performed by western-blot on SDS-15 % polyacrylamide gels in buffer containing 2-β mercaptoethanol. The proteins in the gel were transferred to nitro-cellulose membranes (BioRad, Richmond, Vi) and blocked with dilution buffer (0.5 % bovine serum albumin, 0.1 % Tween-20, 0.01 % merthiolate, 0.005 % phenol red in phosphate buffered saline pH 6.7). Three mm wide strips were incubated with the corresponding trout Ab and with the peroxidase-conjugated rabbit anti-mouse (RAM-PO) Ab and developed with diaminobenzidine (DAB) staining.

Rainbow trout sera used for the screening. Rainbow trout *Oncorhynchus mykiss* (Walbaum) of ~ 50 g obtained from a certified VHS- and IHN-free trout farm were infected in Denmark under laboratory controlled conditions ($12 \pm 2^\circ\text{C}$ with constant aeration) by immersion for 2 h with 2×10^5 TCID₅₀ per ml of water of VHSV-201433-40 isolated from rainbow trout originating from an infected farm in Denmark (Fregeneda-Grandes and Olesen, 2007; Fregeneda-Grandes et al., 2009). A total of 77 sera samples were collected from survivor trout after 4 (n=23), 6 (n=21), 8 (n=16) and 10 (n=17) weeks. Five sera sampled from rainbow trout 12 weeks after infection with VHSV-07.71 (LeBerre et al., 1977) by immersion and 14 sera sampled from rainbow trout 4 weeks after intramuscular injection, were kindly provided by Dr. Jeannette Castric (AFSSA, now ANSES, France). In order to estimate cut-off values,

non-infected trout sera were collected from 15 healthy trout (500 g of body weight) maintained in a closed-circuit farm with no previous history of VHSV infections for more than 30 years (Escuela de Montes, Madrid, Spain). The non-infected sera cut-off values were used to study both the sera from VHSV-infected rainbow trout and 45 sera collected from rainbow trout surviving infection with IHNV strain 039-82 in USA with neutralization titres >200 kindly provided by Dr. Scott LaPatra (Clear Spring Foods, Inc, Idaho USA). Rainbow trout sera (n = 20) were collected from two farms in different Denmark locations with high (~80 %) and medium (~50 %) mortalities produced by VHSV outbreaks in 2006 (farm A 206277) and in 2008 (farm B 2008-50-362), respectively.

frg-ELISA assays of rainbow trout sera. Polystyrene plates of Maxisorb 96-wells (Nunc Immunoplates) were used as ELISA plates. The wells were coated with 5 µg per well of purified recombinant frg11 (56-110), frg15 (65-250), frg16 (252-450) or G21-465 in 100 µl 0.1 M carbonate buffer pH 9.6 and incubated overnight at 4 °C. To reduce the background, the coated wells were then blocked with 100 µl per well of dilution buffer (0.5 % bovine serum albumin, 0.1 % Tween-20, 0.01 % merthiolate, 0.005 % phenol red in phosphate buffered saline pH 6.7) containing 0.025 % of skimmed milk, and washed with 0.5 % Tween 20 in water before performing the ELISA. The wells were incubated during 60 min at room temperature with 100 µl/well of 200-fold diluted trout sera in dilution buffer. After washing, they were incubated during 30 min with anti-trout immunoglobulin MAb 1G7 mouse ascites diluted 1000-fold (Sanchez et al., 1991). After washing, 5000-fold diluted horseradish peroxidase-conjugated rabbit-anti-mouse antibody RAM-PO (Sigma Chem.Co, St.Louis, Mo) was added. After one more wash, a final incubation of 30 min with 0.5 M NaCl 0.01 % SDS was required when using frg15, frg16 and G21-465 to reduce their backgrounds. The colour reaction was developed by adding 1 mg/ml o-phenylenediamine in citrate buffer containing 3 mM H₂O₂. Optical density was measured by using dual wave length at 492-620 nm in a Labsystems ELISA reader. The optical density at 620 nm was used to correct for individual differences between wells as recommended by the manufacturer. Absorbances obtained in fragment coated wells that received no trout sera were subtracted from all the values (background absorbances varied from 0.05-0.15, depending on the fragment). To estimate frg-ELISA titres in large numbers of trout sera, the expression of the results as the absorbance obtained at a unique

200-fold dilution (fixed dilution) rather than the first 2-fold dilution that reached background absorbances (limiting dilution titre) was chosen (Figure 1). To correct for possible unspecific binding of rainbow trout sera obtained from field VHSV outbreaks, frg11-ELISA was carried out in parallel with plates coated with 5 µg per well of either frg11 and bovine serum albumin (BSA). The absorbance values obtained in plates coated with BSA were subtracted to the absorbances values obtained with plates coated with frg11.

VHSV complement-dependent 50 % plaque neutralization test (PNT). The complement-dependent 50 % PNT was performed in 96-well plates as previously described (Fregeneda-Grandes and Olesen, 2007). Briefly, 50 µl of serial 2-fold dilutions of previously heat inactivated trout sera in dilution medium (Eagles MEM with Tris buffer and 5 % fetal bovine serum), were mixed in round bottom 96-wells with 50 µl of fresh 30-fold diluted trout sera as a source of complement and incubated for 30 min at 15 °C on a rocker platform. Then, 100 µl of 8×10^3 pfu of the corresponding homologous VHSV isolate (either 07.71 or 201433) per ml were added to each well (the homologous isolate was chosen to maximize *in vitro* neutralization) and the plate incubated overnight at 15 °C on a rocker platform. Each serum-complement-virus mixture was then adsorbed to 2 replicate wells (10µl per well) with a monolayer of EPC cells for 1 h at 15 °C. The cell cultures were then overloaded with cell culture medium containing 1 % methyl cellulose and incubated for 5 days at 15 °C. Finally, the cells were fixed with 10 % phosphate buffered formaldehyde, washed and stained with 0.5 % crystal violet. After washing, the plates were air-dried and the lysis plaques counted. The complement-dependent 50 % PNT titre was calculated as the reciprocal value of the highest trout serum dilution causing a 50 % reduction of the average number of plaques in control cultures inoculated with non-infected trout serum. Because some trout sera at low dilutions show cellular toxicity, a cut-off value of 80 was chosen to discriminate negative from positive neutralization titers.

RESULTS

Preliminary experiments showed that rainbow trout antibodies (Abs) against the G protein of the french VHSV strain 07.71 (VHSV-07.71) could be estimated in sera from trout repeatedly injected intraperitoneally with both purified VHSV and G protein (hyperimmunized trout) by using the homologous recombinant fragments (frgs) as solid-phases for ELISA (frg-ELISA) (data not shown). Experiments using a few sera from trout surviving laboratory-controlled infections with the DK-201433 Danish VHSV heterologous isolate (differing with 13 amino acid mutations from the VHSV-07.71 isolate) suggested that the corresponding frg-ELISA could also be used to estimate the Ab levels in those sera. In both of the above mentioned cases, Western blotting confirmed that the absorbances obtained by frg-ELISA corresponded to true reactivity with the frgs. Thus, pooled sera from trout infected either with VHSV-07.71 having >10.000 ($n = 4$) and <80 ($n = 3$) PNT titres or with VHSV-201433 having >10.000 ($n = 4$) and <80 ($n = 4$) PNT titres, reacted equally well with frg11, frg15, frg16 or G21-465 by western blotting. In contrast, pooled non-infected trout sera ($n=15$) did not recognized any of the frgs (data not shown).

To semiquantitatively estimate frg-ELISA titres in large numbers of individual trout sera, the results obtained by using the limiting dilution (the first dilution to reach background absorbances) and fixed dilution (the absorbance at 200-fold dilution) methods were compared. Because the comparison of 11 trout sera selected within a wide range of frg11-ELISA titres (Figure 1A), showed a linear correlation between the results obtained with both methods (Figure 1B), the much simpler estimation of absorbances at the 200-fold dilution method was chosen to continue the experimentation.

To select the most appropriated recombinant frg of the protein G of VHSV to assess the presence of anti-VHSV Abs, trout sera were analysed by frg-ELISA by using solid phases coated with frg11, frg15, frg16 or G21-465 from VHSV-07.71. First, to distinguish negative from positive sera (cut-off values), sera from non-infected and IHNV-infected trout were compared with sera from VHSV-201433-infected rainbow trout. Sera from non-infected trout ($n=15$) showed a calculated cut-off absorbance value of 0.24. On the other hand, the distribution of absorbance values of sera from IHNV-infected trout ($n=45$) contained 6 sera (13.3 %) with absorbance values >0.5 . Whether those were due to crossreactivity

ACCEPTED MANUSCRIPT

between the G proteins of IHNV and VHSV or to Abs to VHSV was not investigated (there were considered as outliers). The inclusion of the remaining 39 sera from IHNV-infected trout to the 15 sera from non-infected trout to re-calculate cut-offs, resulted in a mean absorbance of 0.06 which corresponded also to a calculated cut-off value of 0.24 (mean + 2 standard deviations)(not shown). In contrast, Figure 2 shows the distribution to higher absorbance values of most of the sera from VHSV-201433-infected trout (0.46 ± 0.24 , $n=77$). Similar profiles but differing in their cut-off values were obtained when using frg15, frg16 or G21-465 as solid phases (data not shown).

Once the cut-off values for each frg were estimated, individual sera from trout harvested at different times after VHSV-201433 infection were assayed. The resulting absorbance values of trout sera infected with VHSV-201433 estimated with frg15, frg16 or G21-465 showed linear correlation coefficients of 0.91, 0.82 and 0.85, respectively when compared with those obtained with frg11 (Figure 3). A correlation coefficient of 0.84 was obtained when comparing 2 different experiments with frg11 (not shown). Similar results were obtained when using sera from trout infected with VHSV-07.71 (not shown). In contrast, the frg-ELISA absorbances of the sera did not correlate with their harvest time after VHSV infection. Thus, as one example, the distribution of values obtained from sera harvested 10 weeks after VHSV infection showed values ranging from the lowest to the largest frg-ELISA absorbances (Figure 3, black circles). Similar results were obtained with the sera obtained at other times after infection (Figure 3, data not differentially labelled). Because frg11 was the smallest G frg located at a region with the lowest amino acid variability (Benmansour et al., 1997; Einer-Jensen et al., 2004; Rocha et al., 2004) and showed the lowest background values, it was chosen to carry out the rest of the analysis with this frg.

Figure 4 shows the comparison between the titres obtained from frg11-ELISA and complement-dependent PNT. There is no correlation between frg11-ELISA and PNT values for the sera from trout infected either with VHSV-07.71 (Figure 3, open symbols), nor with sera from VHSV-201433 infected trout (Figure 3, black symbols). While a total of 90.1 % of the sera analysed from VHSV-infected trout were >0.24 by frg11-ELISA, only 45 % had PNT titres >80 .

The percentages of sera from VHSV-201433 infected trout at different times after infection were then classified in positive/negative classes by their complement-dependent PNT or frg11-ELISA ($n=77$)

titres. From 4 to 10 weeks after VHSV infection, the percentage of the positive sera estimated by frg11-ELISA increased from 86.9 to 100 % (Figure 5 black circles) while by PNT decreased from 65.2 to 29.4 % (Figure 5 black squares). Furthermore, 10 weeks after infection 100 % of the sera which were positive had absorbance values >0.4 . A total of 38.9 % of the sera that were negative by PNT were positive by ELISA, while a total of 3.3 % of the sera that were negative by ELISA were positive by PNT. Only 7.8 % of the VHSV-201433 infected sera were negative for both ELISA and PNT and all those sera were found at the earliest times after infection (4-6 weeks).

To preliminarily evaluate the potential use of frg11-ELISA to detect previous exposure to different VHSV isolates in different locations after natural infections, rainbow trout sera samples from 2 different farms were collected after their corresponding VHSV outbreaks caused either high or medium mortalities (~ 80 % farm A and ~50 % farm B, respectively) and assayed by PNT, VHSV-ELISA and frg11-ELISA. To increase the sensitivity of the PNT, the cut-off was established at a <40 dilution and several VHSV isolates were used for the test. To increase specificity of the frg11-ELISA, the absorbance values obtained by coating the plates with BSA were subtracted from the absorbance values obtained with plates coated with frg11. Results in farm A confirmed that best sensitivity for PNT was obtained when using the homologous isolate for the test (70 % positives) while only 50 or 15 % positives could be detected when using a related isolate or the F25 strain, respectively. Similar (65 %) or lower (40 %) percentages of positive sera were obtained when using VHSV or frg11-ELISA, respectively, for the sera from this farm (Table 2). In contrast, in farm B only 15 % of the sera could be detected as positives by either homologous PNT or VHSV-ELISA while up to 80 % of the sera resulted positive by frg11-ELISA.

DISCUSSION

The highest percentage of positive sera by frg-ELISA from trout surviving an VHSV infection under laboratory-controlled conditions were found 10 weeks after infection (100 %) in contrast to the positive sera by complement-dependent PNT (29.4 %). Those results show that the absence of neutralising Abs does not mean that other anti-VHSV binding Ab were absent in sera from surviving rainbow trout, therefore confirming earlier observations that showed non-neutralising Abs tend to persist longer than neutralising Abs in VHSV survivors (Enzmann and Konrad, 1993; Lorenzo et al., 1995; Olesen et al., 1991). However, since frg-ELISA is an *in vitro* method, it remains unknown whether or not the newly detected binding Abs are protective *in vivo*. Previous estimations of binding Abs by using VHSV-captured ELISA did not correlate with protection (Lorenzen and Olesen, 1997; Lorenzen et al., 1993; Lorenzen et al., 1999), however it might also be that the new binding Abs detected by frg-ELISA might recognize other additional epitopes not yet studied. On the other hand, because at least one *in vitro* non-neutralising monoclonal Ab was capable of providing *in vivo* passive protection (Lorenzen et al., 1990), it might also be possible that some of the binding Abs which were detected only by frg-ELISA, might also be important for *in vivo* protection. However that was not demonstrated in the present work. The hypothetical existence of protective frg-ELISA binding Abs could also explain why *in vitro* complement-dependent PNT are only detected in about 50 % of VHSV survivors (LeFrancois, 1984; Lorenzen and LaPatra, 1999; Lorenzen et al., 1990). Fractionation of the sera by affinity columns using immobilized frgs and assay of the corresponding fractions by frg-ELISA, PNT and *in vivo* passive transfer, might be used in the future to investigate the possible functional significance of frg11-binding Abs.

Complement-dependent PNT Abs could be detected with the highest sensitivity when the homologous VHSV or very similar isolates were used for testing the sera from DK-211443 infected trout (Fregeneda-Grandes and Olesen, 2007), despite their G protein sequences differing only by 24 amino acids with the DK-F1 VHSV isolate used for the PNT (Fregeneda-Grandes and Olesen, 2007). Similar results were obtained with some of the data reported here for farm A (Table 1). In comparison, frg-ELISA Abs could be detected in the sera from trout infected not only with the homologous VHSV-07.71 isolate

but also with the heterologous VHSV-201433 isolate. The 13 amino acid difference between the VHSV 07-71 and DK-201443 isolates (2 mutations in frg11, 3 in frg15, 7 in frg16 and 11 in G21-465) had no apparent effect on the frg-ELISA performances. However, whether a similar effect might be present in frg-ELISA with other VHSV isolates is not known. Because the highest sequence variability on the protein G of VHSV was localized in amino acid 245 to 300 (Benmansour et al., 1997; Einer-Jensen et al., 2004), new frg15 or frg16-like recombinants which will avoid the 245-300 hypervariable region, might be required to increase the scope of VHSV isolates which could be detected by those frgs. On the other hand, most of the variability found in frg11 was only restricted to 2 short amino acid stretches (Rocha et al., 2004) and therefore frg11 might detect more isolates than frg15 or frg16. An extensive work with more VHSV isolates and/or with the corresponding different sequence variations in frgs, will have to be performed to demonstrate their ability to detect Abs in trout infected with any VHSV isolate.

A high correlation existed among the results obtained by frg-ELISA whether frg11, frg15, frg16 or G21-465 were used for the assays. Among the frgs used, frg11 (amino acids 56-110) had the lowest backgrounds and therefore did not required additional washing steps. Previous work showed that frg11 from VHSV-07.71, could be used in ELISA to detect trout Abs to VHSV-07.71 (Rocha et al., 2002). Frg11 was initially selected because ~ 40 % of sera from trout hyperimmunised by repeated injections with purified VHSV-07.71 (Fernandez-Alonso et al., 1998), recognised the pepscan 15-mer peptides defining frg11, an stretch related to VHSV fusion (Nuñez et al., 1998). However, under the conditions reported before (Rocha et al., 2002), a serum from a hyperimmunized rainbow trout could only be ~ 100-200-fold diluted before the ELISA absorbances reached background absorbances obtained in the absence of any trout sera. In contrast, under the optimized conditions described in this work, hyperimmunized trout sera could be diluted up more than 10.000 fold to reach background absorbances (Figure 1). Because anti-VHSV trout Abs of the IgM class are usually of low affinity and bind non-specifically to almost any surface, they often produce false positives (Olesen et al., 1991). That phenomena have required the use of solid-phase Ab captured VHSV in the presence and in the absence of VHSV to distinguish true positives (Fregeneda-Grandes and Olesen, 2007; Jorgensen et al., 1991; Olesen et al., 1991) and caused also difficulties when using lectin-affinity purified protein G (Perez et al., 1998) and/or recombinant yeast G4

(Estepa et al., 1994). Frg-dependent backgrounds could be controlled in this work by blocking the frg-coated solid phases with low amounts of milk and by adding a final washing step of 0.5M NaCl, but those additives had to be carefully adjusted in concentration to avoid reducing true signals. Further studies might be required to control the higher backgrounds expected with most of the frgs with trout sera from farms, since trout sera from controlled-laboratory infections, like those described above, tend to show lower crossreactivities (data not shown). To preliminarily evaluate the potential use of frg11-ELISA to detect previous exposure to VHSV in natural infections, rainbow trout sera samples from 2 farms were obtained after having suffered their corresponding outbreaks and were analysed by several of the presently available methods (Table 2). To increase the specificity of the frg11-ELISA, the absorbance values obtained by coating the plates with BSA were subtracted from the absorbance values obtained with plates coated with frg11. Frg-ELISA resulted in higher prevalence of anti-VHSV Abs in sera from farm B (medium mortalities) than in the sera from farm A (higher mortalities). However, because of the small amount of sera (n=20 for each farm) and the large amount of variables of natural VHSV outbreaks, many more data should be gathered and studied before we can interpret those data and reach definitive conclusions.

ELISA might be preferred for massive screenings of trout sera for anti-VHSV Abs, because any kind of ELISA seems to be a simpler method to perform than complement-dependent PNT for that purpose. Further advantages of frg-ELISA could be that only the percentage of positive sera by frg-ELISA increased with time after VHSV infection and that frg-ELISA might be less sensitive to VHSV isolate sequence variations than PNT. However, it will be very important to know what the presence of those frg-binding Abs could mean from a functional standpoint (ie: whether those new frg-binding Abs are or are not protective) and whether or not more VHSV isolates could be detected by using G frgs than PNT. A practical method to detect survivors of VHSV natural outbreaks will be most welcome to detect asymptomatic carriers. However, because the number of sera studied here were limited and mostly came from laboratory controlled-conditions, further studies including many more sera from survivors of natural VHSV outbreaks with different VHSV isolates would need to be studied until such applications could be evaluated for frg-ELISA. Nevertheless, an ELISA method to detect trout serum Abs to VHSV, such as the

one described here, could be useful to follow immunization attempts during vaccine development and/or to follow the kinetics of fish Ab response during VHSV infections (Kurath et al., 2007).

ACCEPTED MANUSCRIPT

Table 1

Comparison among PNT, VHSV-ELISA and frg11-ELISA titres in rainbow trout sera obtained from 2 different located farms after VHSV outbreaks with high or medium mortalities

Rainbow trout farm	Serum No	50% PNT titre			ELISA titre	
		VHSV isolate used in test:			Solid-phase	
		F25	201433 -40	homologous	VHSV F25	frg11 07.71
A	1	<40	<40	<40	<40	0.08
	2	1280	2560	5120	640	-0.02
	3	<40	1280	320	<40	0.27
	4	<40	640	160	2560	-0.01
	5	80	<40	80	2560	0.00
	6	<40	<40	80	<40	0.92
	7	<40	<40	2560	640	0.01
	8	5120	2560	5120	640	0.19
	9	<40	5120	5120	2560	0.05
	10	<40	<40	640	<40	0.20
	11	<40	<40	320	<40	0.03
	12	<40	2560	<40	640	0.23
	13	<40	<40	<40	<40	0.02
	14	<40	<40	<40	2560	0.17
	15	<40	5120	5120	2560	0.08
	16	<40	<40	80	<40	0.28
	17	40	<40	320	640	0.19
	18	<40	640	2560	160	0.05
	19	<40	640	1280	160	-0.31
	20	<40	2560	5120	640	0.13
	positives, %	15	50	70	65	40
B	1	--	--	<40	<40	0.03
	2	--	--	<40	<40	0.26
	3	--	--	<40	<40	0.44
	4	--	--	<40	<40	0.28
	5	--	--	<40	160	0.35
	6	--	--	<40	160	0.47
	7	--	--	<40	<40	0.16
	8	--	--	<40	<40	0.20
	9	--	--	<40	<40	0.49
	10	--	--	320	640	0.24
	11	--	--	<40	<40	-0.07
	12	--	--	<40	<40	0.35
	13	--	--	<40	<40	0.63
	14	--	--	<40	40	0.28
	15	--	--	<40	<40	0.77
	16	--	--	<40	<40	0.49
	17	--	--	40	<40	0.10
	18	--	--	80	<40	0.37
	19	--	--	<40	<40	0.45
	20	--	--	<40	<40	0.18
	positives, %			15	15	80

Rainbow trout sera were collected from two farms in Denmark different locations after VHSV outbreaks with high (~80 %) and medium (~50 %) mortalities in farm A in 2006 (206277) and in farm B in 2008 (2008-50-362), respectively. ELISA were performed by either using captured-VHSV according to previously reported methods (Fregeneda-Grandes and Olesen, 2007), frg11 or bovine serum albumin (BSA) as solid-phases. --, not estimated. Homologous, the VHSV isolate used for the test was the one collected from the outbreak. The absorbance values obtained in plates coated with BSA were subtracted from the absorbances values obtained with plates coated with frg11 and cut-offs calculated accordingly. To increase their sensitivity, cut-offs were chosen as <40 for PNT and VHSV-ELISA. For frg11-ELISA, mean plus 2 standard deviations of non-infected sera (n=15) gave a cut-off value of 0.17. Numbers in bold, rainbow trout sera used as positives in each method to calculate percentage of positives (%).

Figure 1. Variation of frg11-ELISA absorbance values with dilution of VHSV-infected trout sera (A) and comparison of titres estimated by limiting and fixed dilution methods (B). Trout sera representing a whole range of different titres were selected for ELISA on plates coated with frg11 (frg11-ELISA) derived from the protein G of VHSV-07.71. Trout sera were 2-fold serially diluted from 200 to 25600-fold and ELISA performed as indicated in methods. A, absorbances at 492-620 nm of serial dilutions of 11 trout sera. B, comparison of ELISA titres defined either as the dilution required to obtain the cut-off of 0.24 units of absorbance (titre) or the absorbance value at a 200-fold dilution.

Figure 2. Distribution of frg11-ELISA values from non-infected + IHNV-infected and VHSV-infected trout sera. Sera from non-infected (n=15) and IHNV-infected rainbow trout (n=39) were used as negative specific controls for anti-G VHSV Abs. A total of 77 rainbow trout sera were obtained from VHSV-201433 infected rainbow trout and assayed. All the sera were 200-fold diluted and frg11-ELISA performed as indicated in methods. Absorbance values were classified in 0.1 absorbance classes and the resulting grouping represented. Open bars, sera from non-infected and IHNV-infected rainbow trout. Black bars, sera from VHSV-201433 -infected rainbow trout.

Figure 3. Comparison between frg-ELISAs absorbance values obtained with different G fragments by using VHSV-201433 survivor sera. Sera from rainbow trout surviving VHSV-201433 infection were diluted 200-fold and assayed by ELISA by using solid phases coated with purified frgs derived from the protein G of VHSV-07.71. The values obtained with frg15, frg16 and G21-465 were compared with the values obtained with frg11. Two experiments were performed with similar results, one of them is shown in the figure. -----, cut-off values were calculated by the following formula, mean absorbance of sera from IHNV-infected (n=39) and non-infected (n=15) rainbow trout + 2 x standard deviations. Straight line, linear fitting of the absorbance values obtained for the VHSV-201433 sera. ○, sera from rainbow trout harvested between 4 and 10 weeks after VHSV-201433 infection (n=77). ●, sera from rainbow trout harvested 10 weeks after VHSV-201433 infection (n= 17).

Figure 4. Lack of correlation between frg11-ELISA and complement-dependent 50 % PNT titres of VHSV-infected rainbow trout sera. Plates used for frg-ELISA were coated with VHSV-07.71 derived frg11. Rainbow trout were infected with VHSV-07.71 (open symbols) or VHSV-201433 (black symbols)

and their sera harvested at different times (4 to 10 weeks) after infection. Sera were 200-fold diluted and ELISA performed as indicated in methods and their means represented (n=2). Complement-dependent 50 % PNT were performed as indicated in methods. To show all the PNT values <80 in the figure, they have been arbitrarily modified in ± 10 increments. ▼, 4 weeks after infection with VHSV-201433. ▲, 6 weeks after infection with VHSV-201433. ■, 8 weeks after infection with VHSV-201433. ●, 10 weeks after infection with VHSV-201433. □, 4 weeks after intraperitoneal injection of VHSV-07.71. ○, 12 weeks after infection with VHSV-07.71. -----, cut-off values of 80 for PNT and 0.24 for frg11-ELISA.

Figure 5. Time course of the percentage of positive/negative sera from rainbow trout infected with VHSV-201433 by frg-ELISA and complement-dependent 50 % PNT of data from Figure 4.

Rainbow trout were infected with VHSV-201433 and sera from survivors harvested 4 (n=23), 6 (n=21), 8 (n=16) and 10 (n=17) weeks after infection. A 0.24 absorbance value and a dilution of 80-fold were used as the cut-offs to differentiate and group negative from positive sera by frg11-ELISA and complement-dependent PNT, respectively. Percentages were calculated by the formula, 100 x number of sera according to the specified criteria (see symbol explanations) / total number of sera analysed at any given time point after infection. ●, percentage of sera that were frg11-ELISA positive. ■, percentage of sera that were PNT positive. ○, percentage of sera that were both PNT negative, frg11-ELISA positive. □, percentage of sera that were both PNT positive, frg11-ELISA negative. ◇, percentage of sera that were both PNT negative, frg11-ELISA negative.

Acknowledgements.

Thanks are due to Dr. Jeannette Castric (AFSSA, France) and to Dr. Scott LaPatra (Clear Spring Foods, Inc, Idaho USA) by providing sera from rainbow trout infected by VHSV-07.71 and IHNV-039-82, respectively. We also thank Dr. Katja Einer-Jensen (NVI of Denmark) by their sequence data of VHSV isolate DK-201433. This work was supported by Spanish CICYT projects AGL08-03519-CO4-ACU, and CONSOLIDER INGENIO 2010: 2007-00002.

References

- Ahne, W.V., Jorgensen, P.E.V., Olesen, N.J., Scafer, W. and Steinhagen, P., 1986. Egtved virus: Occurrence of strains not clearly identifiable by means of virus neutralization tests. *Applied Ichthyology* 4, 187-189.
- Benmansour, A., Basurco, B., Monnier, A.F., Vende, P., Winton, J.R. and deKinkelin, P., 1997. Sequence variation of the glycoprotein gene identifies three distinct lineages within field isolates of viral haemorrhagic septicemia virus, a fish rhabdovirus. *Journal General Virology* 78, 2837-2846.
- Dorson, M. and Torchy, C., 1979. Complement Dependent Neutralization of Egtved Virus by Trout Antibodies. *Journal of Fish Diseases* 2, 345-347.
- Einer-Jensen, K., Ahrens, P., Forsberg, R. and Lorenzen, N., 2004. Evolution of the fish rhabdovirus viral haemorrhagic septicemia virus. *Journal General Virology* 85, 1167-1179.
- Enzmann, P.J. and Konrad, M., 1993. Longevity of antibodies in brown trout and rainbow trout following experimental infection with VHS virus. *Bulletin European Association Fish Pathologists* 13, 193-194.
- Estepa, A., Thiry, M. and Coll, J.M., 1994. Recombinant protein fragments from haemorrhagic septicaemia rhabdovirus stimulate trout leucocyte anamnestic in vitro responses. *Journal General Virology* 75, 1329-1338.
- Estepa, A.M., Rocha, A.I., Mas, V., Perez, L., Encinar, J.A., Nunez, E., Fernandez, A., Ros, J.M.G., Gavilanes, F. and Coll, J.M., 2001. A protein G fragment from the Salmonid viral hemorrhagic septicemia rhabdovirus induces cell-to-cell fusion and membrane phosphatidylserine translocation at low pH. *Journal of Biological Chemistry* 276, 46268-46275.
- Fernandez-Alonso, M., Lorenzo, G., Perez, L., Bullido, R., Estepa, A., Lorenzen, N. and Coll, J.M., 1998. Mapping of the lineal antibody epitopes of the glycoprotein of VHSV, a salmonid rhabdovirus. *Diseases Aquatic Organisms* 34, 167-176.
- Fregeneda-Grandes, J.M. and Olesen, N.J., 2007. Detection of rainbow trout antibodies against viral haemorrhagic septicaemia virus (VHSV) by neutralisation test is highly dependent on the virus isolate used. *Dis Aquat Organ* 74, 151-8.
- Fregeneda-Grandes, J.M., Skall, H.F. and Olesen, N.J., 2009. Antibody response of rainbow trout with single or double infections involving viral haemorrhagic septicaemia virus and infectious haematopoietic necrosis virus. *Dis Aquat Organ* 83, 23-9.
- Hattenberg-Baudoy, A.M., Danton, M., Merle, G. and DeKinkelin, P., 1995. Serum neutralization test for epidemiological studies of salmonid rhabdoviruses in France. *Veterinary Research* 26, 512-520.
- Jorgensen, P.E.V., Olesen, N.J., Lorenzen, N., Winton, J.R. and Ristow, S., 1991. Infectious hematopoietic necrosis (IHN) and viral haemorrhagic septicaemia (VHS): detection of trout antibodies to the causative viruses by means of plaque neutralization, immunofluorescence, and enzyme-linked immunosorbent assay. *Journal Aquatic Animal Health* 3, 100-108.
- Kurath, G., Purcell, M.K. and Garver, K.A., 2007. Fish rhabdovirus models for understanding host response to DNA vaccines. *CAB reviews* 2, 1-12.
- LeBerre, M., De Kinkelin, P. and Metzger, A., 1977. Identification sérologique des rhabdovirus des salmonidés. *Bulletin Office International Epizooties* 87, 391-393.
- LeFrancois, L., 1984. Protection against lethal viral infection by neutralizing and nonneutralizing monoclonal antibodies: a distinct mechanisms action in vivo. *Journal Virology* 51, 208-214.
- Lorenzen, N. and LaPatra, S.E., 1999. Immunity to rhabdoviruses in rainbow trout: the antibody response. *Fish & Shellfish Immunology* 9, 345-360.
- Lorenzen, N. and Olesen, N.J., 1997. Immunization with viral antigens: viral haemorrhagic septicemia. *Developmental Biological Standards* 90, 201-209.
- Lorenzen, N., Olesen, N.J. and Jorgensen, P.E.V., 1990. Neutralization of Egtved virus pathogenicity to cell cultures and fish by monoclonal antibodies to the viral G protein. *Journal General Virology* 71, 561-567.
- Lorenzen, N., Olesen, N.J. and Jorgensen, P.E.V., 1993. Antibody response in rainbow trout to VHS virus proteins. *Fish & Shellfish Immunology* 3, 461-473.
- Lorenzen, N., Olesen, N.J. and Koch, C., 1999. Immunity to VHS virus in rainbow trout. *Aquaculture* 172, 41-61.

- Lorenzo, G.A., Estepa, A., Chilmonczyk, S. and Coll, J.M., 1995. Mapping of the G-Regions and N-Regions of Viral Hemorrhagic Septicemia Virus (VHSV) Inducing Lymphoproliferation by Pepscan. *Veterinary Research* 26, 521-525.
- Medin, J.A., Gathy, K. and Coleman, M.S., 1995. Expression of foreign proteins in *Trichoplusia ni* larvae. *Methods Mol Biol* 39, 265-75.
- Nuñez, E., Fernandez, A.M., Estepa, A., Gonzalez-Ros, J.M., Gavilanes, F. and Coll, J.M., 1998. Phospholipid interactions of a peptide from the fusion-related domain of the glycoprotein of VHSV, a fish rhabdovirus. *Virology* 243, 322-330.
- Olesen, N., J., Lorenzen, N. and Jorgensen, P.E.V., 1991. Detection of rainbow trout antibody to Egtved virus by enzyme-linked immunosorbent assay (ELISA), immunofluorescence (IF), and plaque neutralization tests (50% PNT). *Diseases Aquatic Organisms* 10, 31-38.
- Olesen, N.J. and Jorgensen, P.E.V., 1986. Detection of neutralizing antibody to Egtved virus in Rainbow trout (*Salmo gairdneri*) by plaque neutralization test with complement addition. *Journal Applied Ichthyology* 2, 33-41.
- Perez-Filgueira, D.M., Resino-Talavan, P., Cubillos, C., Angulo, I., Barderas, M.G., Barcena, J. and Escribano, J.M., 2007. Development of a low-cost, insect larvae-derived recombinant subunit vaccine against RHDV. *Virology* 364, 422-30.
- Perez, L., Estepa, A. and Coll, J.M., 1998. Purification of the glycoprotein G from VHSV, a fish rhabdovirus, by lectin affinity chromatography. *Journal Virological Methods* 76, 1-8.
- Rocha, A., Fernandez-Alonso, M., Mas, V., Perez, L., Estepa, A. and Coll, J.M., 2002. Antibody response to a linear epitope of the protein G of VHS rhabdovirus in immunized trout. *Veterinary Immunology Immunopathology* 86, 89-99.
- Rocha, A., Ruiz, S. and Coll, J.M., 2005. Improvement of transfection efficiency of epithelioma papulosum cyprini carp cells by modification of their cell cycle and using an optimal promoter. *Marine Biotechnology* 6, 401-410.
- Rocha, A., Ruiz, S., Tafalla, C. and Coll, J.M., 2004. Conformation and fusion defective mutants in the hypothetical phospholipid-binding and fusion peptides of the protein G of viral haemorrhagic septicemia salmonid rhabdovirus. *Journal Virology* 78, 9115-9122.
- Sanchez, C., Coll, J.M. and Dominguez, J., 1991. One step purification of rainbow trout immunoglobulin. *Veterinary Immunology Immunopathology* 27, 383-392.

Table 1

Comparison among PNT, VHSV-ELISA and frg11-ELISA titres in rainbow trout sera obtained from 2 different located farms after VHSV outbreaks with high or medium mortalities

Rainbow trout farm	Serum No	50% PNT titre			ELISA titre	
		VHSV isolate used in test:			Solid-phase	
		F25	201433 -40	homologous	VHSV F25	frg11 07.71
A	1	<40	<40	<40	<40	0.08
	2	1280	2560	5120	640	-0.02
	3	<40	1280	320	<40	0.27
	4	<40	640	160	2560	-0.01
	5	80	<40	80	2560	0.00
	6	<40	<40	80	<40	0.92
	7	<40	<40	2560	640	0.01
	8	5120	2560	5120	640	0.19
	9	<40	5120	5120	2560	0.05
	10	<40	<40	640	<40	0.20
	11	<40	<40	320	<40	0.03
	12	<40	2560	<40	640	0.23
	13	<40	<40	<40	<40	0.02
	14	<40	<40	<40	2560	0.17
	15	<40	5120	5120	2560	0.08
	16	<40	<40	80	<40	0.28
	17	40	<40	320	640	0.19
	18	<40	640	2560	160	0.05
	19	<40	640	1280	160	-0.31
	20	<40	2560	5120	640	0.13
positives, %		15	50	70	65	40
B	1	--	--	<40	<40	0.03
	2	--	--	<40	<40	0.26
	3	--	--	<40	<40	0.44
	4	--	--	<40	<40	0.28
	5	--	--	<40	160	0.35
	6	--	--	<40	160	0.47
	7	--	--	<40	<40	0.16
	8	--	--	<40	<40	0.20
	9	--	--	<40	<40	0.49
	10	--	--	320	640	0.24
	11	--	--	<40	<40	-0.07
	12	--	--	<40	<40	0.35
	13	--	--	<40	<40	0.63
	14	--	--	<40	40	0.28
	15	--	--	<40	<40	0.77
	16	--	--	<40	<40	0.49
	17	--	--	40	<40	0.10
	18	--	--	80	<40	0.37
	19	--	--	<40	<40	0.45
	20	--	--	<40	<40	0.18
positives, %				15	15	80




Figure 1




Figure 1




Figure 2






Figure 2


frg15



frg16



G21-465



frg11

Figure 3




Figure 3

frg11-ELISA, A 492-620nm

1.5

1.0

0.5

0.0


100

1000

10000

PNT

Figure 4






Figure 5




Figure 5