



# Genome Sequences of Three Vaccine Strains and Two Wild-Type Canine Distemper Virus Strains from a Recent Disease Outbreak in South Africa

Angelika K. Loots,<sup>a,b</sup> Morné Du Plessis,<sup>a,c</sup>  Desiré Lee Dalton,<sup>a,d</sup> Emily Mitchell,<sup>a</sup> Estelle H. Venter<sup>b,e</sup>

National Zoological Gardens of South Africa, Gauteng, South Africa<sup>a</sup>; Department of Veterinary Tropical Diseases, Faculty of Veterinary Science, University of Pretoria, Onderstepoort, South Africa<sup>b</sup>; Department of Biotechnology, University of the Western Cape, Cape Town, South Africa<sup>c</sup>; Department of Zoology, University of Venda, Thohoyandou, South Africa<sup>d</sup>; College of Public Health, Medical and Veterinary Sciences, James Cook University, Townsville, Queensland, Australia<sup>e</sup>

**ABSTRACT** Canine distemper virus causes global multihost infectious disease. This report details complete genome sequences of three vaccine and two new wild-type strains. The wild-type strains belong to the South African lineage, and all three vaccine strains to the America 1 lineage. This constitutes the first genomic sequences of this virus from South Africa.

Canine distemper virus (CDV) (family *Paramyxoviridae*) is an enveloped, nonsegmented, single-stranded, negative-sense RNA virus responsible for the disease canine distemper (CD), which has a high mortality rate in domestic and wild carnivore hosts (1–3). Vaccination has proven to be an effective intervention strategy against CDV in domestic hosts (3, 4); however, there is still a lack of quantitative data on the effects of a CDV vaccine in wildlife (5).

The RNA genome is 15,690 nucleotides (nt) long and consists of 6 genes that encode for a single envelope-associated protein [matrix (M)], 2 glycoproteins [the hemagglutinin (H) and fusion (F) proteins], 2 transcriptase-associated proteins [phosphoprotein (P) and large (L) protein], and a nucleocapsid (N) protein (6). The organization of the major genes was 3′-N-P-M-F-H-L-5′, each separated by untranslated regions (7). Only one serotype of CDV has been recognized, with several cocirculating genotypes based on H-gene variation clustering into the lineages America 1 and 2; Asia 1 and 2; Europe–South America 1; Europe wildlife; South America 2 and 3; Arctic; Rockborn-like; and Africa and Africa 2 (8–11).

Here, we report the complete genome of the three CDV strains (CDV\_Buc, CDV\_Nobi, and CDV\_OVI) commonly found in vaccines and two wild-type strains (WT01 and WT02) from a recent CDV outbreak in South Africa. Both WT01 and WT02 were obtained from lung tissue collected from an infected African wild dog (*Lycaon pictus*), in Northern Cape Province, South Africa, and spotted hyena (*Crocuta crocuta*), in Limpopo Province, South Africa, respectively. Samples were cultured and passaged between one and three times on VerodogSLAM cells grown in 25 cm<sup>2</sup> tissue culture flasks (12). Viral RNA was extracted using Trizol (Invitrogen). Sequence-independent whole-genome reverse transcription-PCR amplification was used to prepare templates, which were sequenced on an Illumina MiSeq sequencer using the TruSeq sample preparation kit (Illumina). Data quality was assessed using FastQC version 0.11.2 software (<http://www.bioinformatics.babraham.ac.uk>). Paired sequence reads were analyzed with CLC Genomics Workbench version 6 (CLC bio, Aarhus, Denmark). Full-length genome sequences were assembled using a combination of mapping and *de novo* assembly.

Received 9 May 2017 Accepted 16 May 2017 Published 6 July 2017

**Citation** Loots AK, Du Plessis M, Dalton DL, Mitchell E, Venter EH. 2017. Genome sequences of three vaccine strains and two wild-type canine distemper virus strains from a recent disease outbreak in South Africa. *Genome Announc* 5:e00603-17. <https://doi.org/10.1128/genomeA.00603-17>.

**Copyright** © 2017 Loots et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Desiré Lee Dalton, [desire@nzc.ac.za](mailto:desire@nzc.ac.za).

The genomes of WT01, WT02, CDV\_Buc, CDV\_OVI, and CDV\_Nobi are 15,690 nt, 15,649 nt, 15,673 nt, 15,670 nt, and 15,649 nt, respectively. Amino acid lengths of the six proteins encoded by each of the four genomes were 522 (N), 506 (P), 334 (M), 662 (F), 604 (H), and 2,183 (L). Multiple sequence alignments and Bayesian phylogenetic analysis according to the H-gene revealed that WT01 and WT02 belong to the South African lineage. The three vaccine strains were grouped in the America 1 lineage, which is consistent with other known vaccine strain groupings.

The two wild-type CDV strains constitute the first report of genomic sequences in South Africa. Data suggest the formulation of new and updated vaccines, considering the level of genetic variability obtained. These data contribute to the knowledge of CDV, which may be beneficial in determining effective preventative, diagnostic, and control measures for CD in South Africa.

**Accession number(s).** The genome sequences of WT01, WT02, CDV\_Buc, CDV\_Nobi, and CDV\_OVI have been deposited in GenBank under the accession numbers [KY971528](#), [KY971532](#), [KY971529](#), [KY971530](#), and [KY971531](#), respectively.

## ACKNOWLEDGMENTS

This study was supported by the National Zoological Gardens of South Africa and funded by the National Research Foundation (NRF) Professional Development Program. We acknowledge wildlife veterinarian Peter Caldwell (Old Chapel Veterinary Clinic, Tshwane) for his invaluable contribution in collecting samples, as well as the Tswalu Kalahari Reserve for their permission to collect samples and data for this study. Karen Ebersohn and Peter Coetzee are recognized for their assistance in the laboratory and the Agricultural Research Council's Biotechnology Platform for sequencing on the MiSeq.

## REFERENCES

1. Appel MJG. 1987. Canine distemper virus, p 133–159. *In* Appel MJG (ed), *Virus infections of carnivores*. Elsevier Science, New York, NY.
2. Beineke A, Baumgärtner W, Wohlsein P. 2015. Cross-species transmission of canine distemper virus—an update. *One Health* 1:49–59. <https://doi.org/10.1016/j.onehlt.2015.09.002>.
3. Bronson E, Deem SL, Sanchez C, Murray S. 2007. Serologic response to a canarypox-vectored canine distemper virus vaccine in the giant panda (*Ailuropoda melanoleuca*). *J Zoo Wildl Med* 38:363–366. [https://doi.org/10.1638/1042-7260\(2007\)038\[0363:SRTACC\]2.0.CO;2](https://doi.org/10.1638/1042-7260(2007)038[0363:SRTACC]2.0.CO;2).
4. Sadler RA, Ramsay E, Mcaloose D, Rush RCV, Wilkes RP. 2016. Evaluation of two canine distemper virus vaccines in captive tigers (*Panthera tigris*). *J Zoo Wildl Med* 47:558–563. <https://doi.org/10.1638/2015-0223.1>.
5. Loots AK, Mitchell E, Dalton DL, Kotzé A, Venter EH. 2017. Advances in canine distemper virus pathogenesis research: a wildlife perspective. *J Gen Virol* 98:311–321. <https://doi.org/10.1099/jgv.0.000666>.
6. Diallo A. 1990. Morbillivirus group: genome organisation and proteins. *Vet Microbiol* 23:155–163. [https://doi.org/10.1016/0378-1135\(90\)90145-L](https://doi.org/10.1016/0378-1135(90)90145-L).
7. Sidhu MS, Husar W, Cook SD, Dowling PC, Udem SA. 1993. Canine distemper terminal and intergenic non-protein coding nucleotide sequences: completion of the entire CDV genome sequence. *Virology* 193:66–72. <https://doi.org/10.1006/viro.1993.1103>.
8. Ke GM, Ho CH, Chiang MJ, Sanno-Duanda B, Chung CS, Lin MY, Shi YY, Yang MH, Tyan YC, Liao PC, Chu PY. 2015. Phylodynamic analysis of the canine distemper virus hemagglutinin gene. *BMC Vet Res* 11:164. <https://doi.org/10.1186/s12917-015-0491-9>.
9. Panzera Y, Sarute N, Iraola G, Hernández M, Pérez R. 2015. Molecular phylogeography of canine distemper virus: geographic origin and global spreading. *Mol Phylogenet Evol* 92:147–154. <https://doi.org/10.1016/j.ympev.2015.06.015>.
10. Martinez-Gutierrez M, Ruiz-Saenz J. 2016. Diversity of susceptible hosts in canine distemper virus infection: a systematic review and data synthesis. *BMC Vet Res* 12:78. <https://doi.org/10.1186/s12917-016-0702-z>.
11. Harder TC, Osterhaus AD. 1997. Canine distemper virus—a morbillivirus in search of new hosts? *Trends Microbiol* 5:120–124. [https://doi.org/10.1016/S0966-842X\(97\)01010-X](https://doi.org/10.1016/S0966-842X(97)01010-X).
12. von Messling V, Svitek N, Cattaneo R. 2006. Receptor (SLAM [CD150]) recognition and the V protein sustain swift lymphocyte-based invasion of mucosal tissue and lymphatic organs by a morbillivirus. *J Virol* 80: 6084–6092. <https://doi.org/10.1128/JVI.00357-06>.