



# HHS Public Access

Author manuscript

*Prev Sci.* Author manuscript; available in PMC 2020 July 01.

Published in final edited form as:

*Prev Sci.* 2019 July ; 20(5): 665–673. doi:10.1007/s11121-019-0982-1.

## Power of the peer and parent: Gender differences, norms, and nonmedical prescription opioid use among adolescents in south central Kentucky

Kathleen L. Egan, PhD, MS<sup>1</sup>, Eric Gregory, EdD, CPS<sup>2</sup>, Vicki L. Osborne, PhD<sup>3,4</sup>, and Linda B. Cottler, PhD, MPH,<sup>4</sup> [FACE]

<sup>1</sup>Department of Health Education and Promotion, East Carolina University, Greenville, NC, USA

<sup>2</sup>Community Survey Solutions, LLC, Bowling Green, KY, USA

<sup>3</sup>Drug Safety Research Unit, Southampton, UK

<sup>4</sup>Department of Epidemiology, University of Florida, Gainesville, FL, USA

### Abstract

This study examined risk factors of nonmedical prescription opioid use (NMPOU) among adolescents and how risk factors differ by gender. In the fall of 2017, adolescents attending 6<sup>th</sup> through 12<sup>th</sup> grades across 44 schools in 10 south central Kentucky counties were invited to participate in an anonymous, school-based survey. A total of 11,761 adolescents completed the survey. Logistic regression was conducted to examine the association between NMPOU and constructs of the Theory of Reasoned Action (i.e., attitudes and subjective norms), descriptive norms (i.e., peer use), and parental control of prescription medications in the home. There were 297 (2.7%) adolescents who reported NMPOU in the past 12 months. In the adjusted multivariate logistic regression model, for both males and females, adolescents who perceived that more of their peers engaged in NMPOU were significantly more likely to endorse NMPOU. Whereas, male and female adolescents who perceived their peers disapproved of use were significantly less likely to report NMPOU. Parent disapproval was significantly associated with decreased NMPOU for females only. Moderated regression analyses revealed that gender moderated the relationship between parental disapproval and NMPOU. We found that during adolescence, NMPOU is influenced by peer norms for both genders and parental norms for females. These results indicate that prevention efforts should focus on changing adolescents' peer and parental norms related to NMPOU.

---

Correspondence concerning this article should be addressed to Kathleen Egan, Ph.D., M.S., Department of Health Education and Promotion, East Carolina University, 3105 Carol G. Belk Building, Greenville, NC 27858. Telephone: 252-328-2719., eganka18@ecu.edu.

Compliance with Ethical Standards

Disclosure of potential conflicts of interest

The authors declare that they have no conflict of interest.

Research involving human participants

All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards.

Informed consent

For this type of study formal consent was not required.

## Keywords

prescription drug; opioid; adolescent; norm; availability

---

## Introduction

The prevalence of nonmedical prescription opioid use (NMPOU), the use without a prescription or in ways other than prescribed, is second to marijuana use among adolescents (Center for Behavioral Health Statistics and Quality, 2017). The most recent National Survey on Drug Use and Health (NSDUH) found that 3.5% of adolescents ages 12 to 17 reported NMPOU in the past 12 months and 1.6% in the past 30 days (Center for Behavioral Health Statistics and Quality, 2017). In addition, 0.6% of adolescents met the DSM-IV criteria for either dependence or abuse of prescription pain relievers within the previous 12 months; only 0.2% less than 18–25 year olds and the same prevalence as individuals ages 26 and older (Center for Behavioral Health Statistics and Quality, 2017). Peak age of NMPOU onset is 16 years of age (Austic, McCabe, Stoddard, Ngo, & Boyd, 2015). Early age of NMPOU initiation has been associated with adverse consequences (Cerdá et al., 2013; McCabe, West, Morales, Cranford, & Boyd, 2007), including transition to heroin use in young adulthood (Cerdá et al., 2013). It is important to identify risk factors associated with NMPOU during adolescence in order to develop age appropriate intervention strategies to prevent and delay the onset of NMPOU.

Previous research has found the Theory of Reasoned Action (TRA) (Fishbein, 1979) and Theory of Planned Behavior (TPB) (Ajzen, 1991) to be useful in the explanation of substance use behaviors among adolescents and young adults (Gallucci, Martin, Beaujean, & Usdan, 2015; Maahs, Weidner, & Smith, 2016; Malmberg et al., 2012; Ponnet, Wouters, Walrave, Heirman, & Van Hal, 2014; Unger, Rohrbach, Howard-Pitney, Ritt-Olson, & Mouttapa, 2001). The TRA posits that behavior is guided by perceived consequences of the behavior (e.g., perceived risk) and perceived normative expectations of others (e.g., perceived approval). These perceptions result in the formation of a behavioral intention and ultimately the behavior of interest (Fishbein, 1979). The TPB expands upon the TRA to include perceived behavioral control of performing the behavior of interest; perceived behavioral control can either impact behavior directly or indirectly through intention (Ajzen, 1991).

Although the TRA and TPB have been used to understand substance use behavior, there are other factors that also contribute to substance use, including descriptive norms and availability of substances of abuse. Descriptive norms is a type of norm included in Social Norms Theory (Perkins & Berkowitz, 1986) that refers to individuals' perceptions of the prevalence of a specific behavior among others (e.g., peer use of prescription opioids). Studies have found that individuals tend to overestimate the use of substances by others and overestimation is related to personal use (Aas & Klepp, 1992; Campo et al., 2003; Egan, Erausquin, Milroy, & Wyrick, 2016; Kilmer, Geisner, Gasser, & Lindgren, 2015; Larimer et al., 2011; Martens et al., 2006; McCabe, 2008; Vidourek, King, & Burbage, 2014).

Availability refers to the ease and convenience of obtaining substances for use or abuse (Babor, Caetano, et al., 2010, p. 201). Prevention and intervention strategies that address the availability of substances are based on the hypothesis that limiting availability results in decreases of abuse and ultimately decreases of associated substance-related problems (Babor, Caulkins, et al., 2010; Maddahian, Newcomb, & Bentler, 1986; Paschall, Grube, & Kypri, 2009; Popova, Giesbrecht, Bekmuradov, & Patra, 2009). Prescription opioids have been reported to be commonly available to adolescents via friends or family members (Center for Behavioral Health Statistics and Quality, 2015; McCabe & Boyd, 2005). Additionally, previous research has found that adolescents report easy and unrestricted access to prescription medications in their own homes (Friese, Moore, Grube, & Jennings, 2013; McCabe, West, & Boyd, 2013; Ross-Durow, McCabe, & Boyd, 2013). These findings suggest that tracking and restricting access of prescription opioids may result in a decrease of NMPOU among adolescents.

Gender differences have been observed when examining NMPOU among both adolescents (Osborne, Serdarevic, Crooke, Striley, & Cottler, 2017) and adults (Back, Payne, Simpson, & Brady, 2010; Serdarevic, Striley, & Cottler, 2017). One study found that females, early initiators, and opiate users had a shorter length of time between onset of abuse and dependence which emphasized the importance of examining gender differences among adolescents who use NMPOs (Ridenour, Maldonado-Molina, Compton, Spitznagel, & Cottler, 2005). Among adolescents, these may arise because behavior is influenced by external factors which differ by gender. For example, parental monitoring and guidance can differ by gender, with different expectations directed towards males and females (Hyde, 2014; Witt, 1997). Substance use is often influenced by social networks and factors such as peer pressure (Chan, Kelly, Carroll, & Williams, 2017; Karakos, 2014; Kristjansson, Sigfusdottir, & Allegrante, 2013; McDonough, Jose, & Stuart, 2016; Song, Smiler, Wagoner, & Wolfson, 2012), which may vary by gender specific norms. There are mixed findings as to whether females are more heavily influenced by parental support and guidance compared to males (Dunn, Kitts, Lewis, Goodrow, & Scherzer, 2011) or if males are more susceptible to parental injunctive norms than females (Elek, Miller-Day, & Hecht, 2006). Yet, these gender differences in the context of NMPOU have yet to be explored.

While the TRA and TPB have been used to identify risk factors associated with the misuse of prescription stimulants among college students (Gallucci et al., 2015; Ponnet et al., 2014), to our knowledge, they have not been used to examine the misuse of prescription opioids among adolescents. Additionally, previous research has not accounted for the impact of descriptive norms, the availability of prescription drugs in the home, and gender when examining the relationship between these theoretical constructs and NMPOU. The aim of this study was to examine nonmedical prescription opioid use (NMPOU) among adolescents using the Theory of Reasoned Action and Social Norms Theory while also examining the impact of availability. We also sought to explore how these potential risk factors differed by gender. In accordance with theory and previous research, we hypothesized that perceived risk of NMPOU, parent and peer disapproval of NMPOU, peer NMPOU, and parental restriction of access to prescription opioids would be related to adolescent NMPOU. We hypothesized that these findings would be consistent for both genders but parental norms would have a greater impact for females compared to males.

## Methods

In the fall of 2017, adolescents attending 6<sup>th</sup> through 12<sup>th</sup> grades in 10 south central Kentucky counties were invited to participate in an anonymous, school-based survey. The survey was a collaborative project among regional community agencies. It assessed demographics, health behaviors (including substance use), mental health, parenting practices, and school experiences. The median population size of the 10 participating counties was 17,738 (range: 10,099 – 113,792), and on average the counties were 78% rural (range: 31.2% - 100%). All schools with 6<sup>th</sup> - 12<sup>th</sup> grades that were served by the regional community agencies were invited to participate in the survey and the majority participated (n=44 schools). Paper and pencil surveys were conducted within the classroom during the school day. Participation in the survey was voluntary and students were given the option to opt out of the survey or single questions within the survey. A total of 11,761 adolescents completed the survey (~64% response rate). The Wake Forest School of Medicine Institutional Review Board approved the study protocol for secondary analysis of existing data.

## Measures

**NMPOU.**—NMPOU was assessed with the following question: “Think back over the past 12 months. How often did you use prescription pain killers (i.e., Hydrocodone) to get high?.” Participants who endorsed “a few times a year,” “at least once a month,” “at least once a week,” or “several times a week” were considered past 12-month users. Students who stated “never” or “not within the past 12 months” were considered non-past 12 month users.

**Demographics.**—Gender (male or female), race (White, Black/African American American Indian/Native American, Asian/Pacific Islander, Multiracial, or Other), and grade (6–12<sup>th</sup>) were self-reported by participants.

**Covariates.**—Perceived risk of NMPOU was assessed by asking “how much do you think people risk harming themselves physically or in other ways if they use prescription pain killers (i.e., Hydrocodone) to get high?.” Response options were “no risk,” “slight risk,” “moderate risk,” or “great risk.” Perceived parent disapproval of nonmedical prescription drug use was queried with the item “how wrong would your parents feel it would be for you to use prescription drugs not prescribed to you.” Response options were “not wrong at all,” “a little bit wrong,” “wrong,” or “very wrong.” Perceived peer disapproval of NMPOU was queried with the item “how wrong would your friends feel it would be for you to use prescription pain killers (i.e., Hydrocodone) to get high.” Response options were “not wrong at all,” “a little bit wrong,” “wrong,” or “very wrong.” Perceived peer NMPOU was assessed with the question “think about the people about your age who you consider to be your best friends. How many of them used prescription pain killers (i.e., Hydrocodone) to get high?.” Response options were “none of them used,” “less than half of them used,” “about half of them used,” “more than half of them used,” and “all of them used.” Perceptions of parent(s) control of prescription opioids was captured with the question “do your parents take steps to ensure you do not have access to prescription medications?.”

## Statistical Analyses

Analyses consisted of bivariate and multivariate logistic regression. Bivariate logistic regression models were conducted to examine the unadjusted associations between NMPOU and demographics, perceived risk, perceived peer NMPOU, perceived peer and parent disapproval of NMPOU, and parental restriction of access to prescription medications in the home (Table 1). We examined the adjusted relationship between all variables listed above and NMPOU using a multivariate logistic regression model (Table 2). The impact of gender was assessed two ways: (1) gender stratified multivariate logistic regression models (Table 2) and (2) moderated multivariate logistic regression models (Table 3). For the five moderated models, in order to avoid potentially problematic high collinearity, the continuous variables were mean centered prior to creating the interaction terms. All models were adjusted for within-school clustering by treating school as a random effect using GENLINUX with a logit link function in SPSS version 24. P-values, adjusted odds ratios, and 95% confidence intervals are presented.

## Results

### Sample Characteristics

As seen in Table 1, an equal number of males and females participated in the survey (49.3% and 50.6%, respectively). Many of the participants were White (75.5%) followed by Black/African American (7.1%), Other (6.3%), Multiracial (6.2%), Asian/Pacific Islander (2.6%), and American Indian/Native American (2.4). The sample was slightly more diverse than the Census data for the participating counties which were on average 92% (range: 82.1% - 96.5%) white. As the grade levels increased from 6<sup>th</sup> through 12<sup>th</sup>, the percentage of participants per grade decreased slightly from 18.0% in 6<sup>th</sup> grade to 9.5% in 12<sup>th</sup> grade. There were 297 (2.7%) adolescents who reported NMPOU in the past 12 months. Of the 297 who reported NMPOU in the past 12 months, 45.5% (n=135) reported use a few times a year, 31.3% (n=93) at least once a month, 13.4% (n=40) at least once a week, and 9.8% (n=29) several times a week.

### Risk Factors of Nonmedical Prescription Opioid Use

In the unadjusted bivariate regression models (Table 1), neither gender nor race were statistically associated with NMPOU. Grade, perceived risk of NMPOU, perceived parent and peer disapproval of NMPOU, and perceived parent restriction of access to prescription opioids were significantly associated with NMPOU at  $p < 0.05$ .

Among the overall sample, after controlling for all variables, higher grade-level (AOR=1.07; CI=1.00–1.13) and perceived peer NMPOU (AOR=1.74; CI=1.57–1.95) were positively related to NMPOU, and perceived peer disapproval of NMPOU was negatively associated with NMPOU (AOR=0.74; CI=0.65–0.83) (shown in Table 2). These findings were consistent among males; higher grade-level (AOR=1.09; CI=1.00–1.19) and perceived peer NMPOU (AOR=1.75; CI=1.51–2.04) were positively associated with NMPOU, and perceived peer disapproval of NMPOU was negatively associated with NMPOU (AOR=0.78; CI=0.66–0.93). Among females, both perceived peer disapproval of NMPOU (AOR=0.69; CI=0.58–0.82) and perceived peer NMPOU (AOR=1.72; CI=1.47–2.02) were

significantly related to NMPOU in the same direction as they were for males. However, perceived parent disapproval of NMPOU was negatively related to NMPOU (AOR=0.78; CI=0.66, 0.93) but grade-level was not associated with NMPOU (Table 2). As shown in Table 3, the statistically significant interaction term of gender\*parental disapproval (AOR=0.74; CI=0.56, 0.97) suggests that gender moderates some of the relationship between parent disapproval of NMPOU and NMPOU. Gender did not appear to moderate the relationship between other constructs and NMPOU (Table 3).

## Discussion

In our theory-driven study of 11,761 middle and high schoolers in south central Kentucky, 2.7% of adolescents reported NMPOU in the past 12 months. In accordance with the Theory of Reasoned Action, Social Norms Theory, and literature on physical availability, perceived risk, parent and peer disapproval, peer use, and parental monitoring of prescription drugs were significantly related to NMPOU in bivariate analyses. However, among all adolescents, only perceived peer subjective and descriptive norms remained statistically significant in the multivariate model. Further examination of these NMPOU risk factors by gender revealed that while peer norms were statistically significant for both genders, parental disapproval was significant for females only and grade-level was significant for males only. Together, these findings suggest that, even after accounting for perceived risk and parental monitoring of prescription drugs at home, peer norms are the most powerful correlates of NMPOU among adolescents.

Among our sample, 2.7% reported NMPOU within the past 12 months which was similar to, but slightly less than, a national sample of adolescents ages 12 to 17 (3.5%) (Center for Behavioral Health Statistics and Quality, 2017). Our indicator of prevalence solely assessed recreational use of prescription opioids (“to get high”) which may contribute to the slight difference between the prevalence of NMPOU in our sample compared to the national sample. The adolescents who reported NMPOU in our sample may be at higher risk of adverse consequences compared to adolescents who use prescription opioids for medical reasons only given their recreational motivations for NMPOU (McCabe, Cranford, Boyd, & Teter, 2007).

Both adolescent males and females who perceived that their peers disapproved of NMPOU were significantly less likely to report NMPOU. This is consistent with the literature on the relationship between peer subjective norms and substance use (Malmberg et al., 2012; McCabe, 2008; Perkins, 2002; Ponnet et al., 2014). Additionally, we found that adolescents who perceived that their peers used NMPOU were statistically more likely to report past year NMPOU themselves. This finding is consistent with research on descriptive norms and other substance use (Neighbors, Lee, Lewis, Fossos, & Larimer, 2007), including general prescription drug use (McCabe, 2008). Given the cross-sectional nature of this study, we are not able to ascertain if perceptions of peer approval and NMPOU led to personal NMPOU to conform to perceived social norms or if NMPOU preceded the perceptions. However, Lewis et al (Lewis, Litt, & Neighbors, 2015) conducted a longitudinal study of alcohol consumption among college students and found a reciprocal relationship between norms and alcohol consumption; norms predicted alcohol use which subsequently predicted both norms



and alcohol use. Thus, a similar pattern may exist for NMPOU. Prevention efforts should focus on changing subjective and descriptive peer norms pertaining to NMPOU. There are several strategies and interventions that can be implemented to achieve this goal, including personalized normative feedback (Boyle, Earle, LaBrie, & Smith, 2017; Dotson, Dunn, & Bowers, 2015), social norms campaigns (Perkins, Linkenbach, Lewis, & Neighbors, 2010), and peer led interventions (MacArthur, Harrison, Caldwell, Hickman, & Campbell, 2016).

While parental disapproval did not impact NMPOU among males in the multivariate model, adolescent females who perceived that their parents disapproved of NMPOU were significantly less likely to report use. Our findings are similar to Dunn et al's (2011) who found that parental approval impacted adolescent females' marijuana use but not males' use. Our findings suggest that prevention strategies that include a parental norm component to address NMPOU may be more beneficial for female than male adolescents. That being said, parental norms were statistically significant prior to controlling for other variables, such as peer descriptive and injunctive norms, which suggests that they still impact males' NMPOU even if they are not as strong as other variables. Additional research is needed to better understand effective strategies that parents can use to dissuade NMPOU among their male sons. For example, parent-child communication strategies (Luk, Farhat, Iannotti, & Simons-Morton, 2010) and parenting styles (ablová, Pazderková, & Miovský, 2014) may be promising strategies to prevent NMPOU.

There are several limitations to our study. Our study was conducted in 10 counties located in south central Kentucky and may not be generalizable to other communities. However, our findings may apply to similar rural, southern contexts. According to the Theory of Planned Behavior and the Theory of Reasoned Action, intentions to perform a behavior occur between perceptions about the behavior and actually performing the behavior. While, we did not measure the construct of intention, previous research suggests that substance use behaviors may bypass intention and that measuring the intention of substance use is not necessary (Ajzen, 1991). We only examined the use of prescription opioids for recreational motivations. Thus, our findings should not be extrapolated to NMPOU for other motivations (e.g., self-treatment). While perceived risk, peer disapproval, and peer use were specific to prescription opioids, the measures of parent disapproval and access to medications in the home were not specific to opioids. The lack of specificity may have partially contributed to the decreased strength of the relationship between these two constructs and NMPOU in the multivariate analysis. The items that assessed opioid use or perceptions used "i.e.," a limiting statement, followed by "hydrocodone" which could result in an underestimate if the participants took this literally. In addition, substance use was self-reported and is subject to social desirability bias. Given the similarity of our findings to national estimates, it is unlikely that these two limitations significantly impacted our findings. As mentioned previously, given the cross-sectional nature of the data, we are not able to ascertain if the perceptions preceded NMPOU which is important since behavioral feedback may influence or reinforce perceptions.

## Conclusion

After accounting for perceived risk and parental monitoring of prescription drugs at home, peer norms are the most powerful correlates of NMPOU among adolescents in our sample. Specifically, both male and female adolescent NMPOU was related to peer norms, whereas, parental norms were only significantly associated with NMPOU for females. Prevention efforts that focus on peer norms related to NMPOU may result in a reduction of adolescent NMPOU.

## Acknowledgments

### Funding

Research reported in this article was supported by the NIDA T32 training grant at the UF Substance Abuse Training Center in Public Health from the National Institutes of Health (T32DA035167) and Substance Abuse and Mental Health Services Administration under SP019436. The content is solely the responsibility of the authors and does not represent the official views of the National Institute of Health or Substance Abuse and Mental Health Services Administration.

## References

- Aas H, & Klepp K-I (1992). Adolescents' alcohol use related to perceived norms. *Scandinavian Journal of Psychology*, 33(4), 315–325. 10.1111/j.1467-9450.1992.tb00920.x [PubMed: 1287824]
- Ajzen I (1991). The theory of planned behavior. *Organizational Behavior and Human Decision Processes*, 50(2), 179–211. 10.1016/0749-5978(91)90020-T
- Austic E, McCabe SE, Stoddard SA, Ngo QE, & Boyd C (2015). Age and Cohort Patterns of Medical and Nonmedical Use of Controlled Medication Among Adolescents: *Journal of Addiction Medicine*, 9(5), 376–382. 10.1097/ADM.000000000000142 [PubMed: 26291544]
- Babor T, Caetano R, Casswell S, Edwards G, Giesbrecht N, Graham K, ... Rossow I. (2010). *Alcohol: No Ordinary Commodity* (2nd ed.). Oxford: Oxford University Press Retrieved from [https://books.google.com/books/about/Alcohol\\_No\\_Ordinary\\_Commodity.html?id=bbcdAAAAQBAJ](https://books.google.com/books/about/Alcohol_No_Ordinary_Commodity.html?id=bbcdAAAAQBAJ)
- Babor T, Caulkins J, Edwards G, Fischer B, Foxcroft D, Humphreys K, ... Strang J. (2010). *Drug Policy and the Public Good* New York: Oxford University Press Retrieved from <http://www.oxfordscholarship.com/view/10.1093/acprof:oso/9780199557127.001.0001/acprof-9780199557127>
- Back SE, Payne RL, Simpson AN, & Brady KT (2010). Gender and Prescription Opioids: Findings from the National Survey on Drug Use and Health. *Addictive Behaviors*, 35(11), 1001–1007. 10.1016/j.addbeh.2010.06.018 [PubMed: 20598809]
- Boyle SC, Earle AM, LaBrie JW, & Smith DJ (2017). PNF 2.0? Initial evidence that gamification can increase the efficacy of brief, web-based personalized normative feedback alcohol interventions. *Addictive Behaviors*, 67, 8–17. 10.1016/j.addbeh.2016.11.024 [PubMed: 27978426]
- ablová L, Pazderková K, & Mioviský M (2014). Parenting styles and alcohol use among children and adolescents: A systematic review. *Drugs: Education, Prevention and Policy*, 21(1), 1–13. 10.3109/09687637.2013.817536
- Campo S, Brossard D, Frazer MS, Marchell T, Lewis D, & Talbot J (2003). Are social norms campaigns really magic bullets? Assessing the effects of students' misperceptions on drinking behavior. *Health Communication*, 15(4), 481–497. 10.1207/S15327027HC1504\_06 [PubMed: 14527868]
- Center for Behavioral Health Statistics and Quality. (2015). *Behavioral Health Trends in the United States: Results from the 2014 National Survey on Drug Use and Health (NSDUH Series H-50 No. HHS Publication No. SMA 15–4927)* Retrieved from Retrieved from <http://www.samhsa.gov/data/>
- Center for Behavioral Health Statistics and Quality. (2017). *Key substance use and mental health indicators in the United States: Results from the 2016 National Drug Use and Health (No. HHS*



- Publication No. SMA 17–5044, NSDUH Series H-52) Rockville, MD Retrieved from <https://www.samhsa.gov/data>
- Cerdá M, Ransome Y, Keyes KM, Koenen KC, Tardiff K, Vlahov D, & Galea S (2013). Revisiting the role of the urban environment in substance use: the case of analgesic overdose fatalities. *American Journal of Public Health*, 103(12), 2252–2260. 10.2105/AJPH.2013.301347 [PubMed: 24134362]
- Chan GCK, Kelly AB, Carroll A, & Williams JW (2017). Peer drug use and adolescent polysubstance use: Do parenting and school factors moderate this association? *Addictive Behaviors*, 64, 78–81. 10.1016/j.addbeh.2016.08.004 [PubMed: 27572180]
- Dotson KB, Dunn ME, & Bowers CA (2015). Stand-Alone Personalized Normative Feedback for College Student Drinkers: A Meta-Analytic Review, 2004 to 2014. *PLoS ONE*, 10(10). 10.1371/journal.pone.0139518
- Dunn MS, Kitts C, Lewis S, Goodrow B, & Scherzer GD (2011). Effects of Youth Assets on Adolescent Alcohol, Tobacco, Marijuana Use, and Sexual Behavior. *Journal of Alcohol and Drug Education; Lansing*, 55(3), 23–40.
- Egan KL, Erausquin JT, Milroy JJ, & Wyrick DL (2016). Synthetic Cannabinoid Use and Descriptive Norms among Collegiate Student-Athletes. *Journal of Psychoactive Drugs*, 48(3), 166–172. 10.1080/02791072.2016.1186305 [PubMed: 27224428]
- Elek E, Miller-Day M, & Hecht ML (2006). Influences of Personal, Injunctive, and Descriptive Norms on Early Adolescent Substance use. *Journal of Drug Issues*, 36(1), 147–172. 10.1177/002204260603600107
- Fishbein M (1979). A theory of reasoned action: Some applications and implications. *Nebraska Symposium on Motivation*, 27, 65–116.
- Friese B, Moore RS, Grube JW, & Jennings VK (2013). How Parents of Teens Store and Monitor Prescription Drugs in the Home. *Journal of Drug Education*, 43(3), 223–233. 10.2190/DE.43.3.b [PubMed: 25429166]
- Gallucci A, Martin R, Beaujean A, & Usdan S (2015). An examination of the misuse of prescription stimulants among college students using the theory of planned behavior. *Psychology, Health & Medicine*, 20(2), 217–226. 10.1080/13548506.2014.913800
- Ghandour LA, El Sayed DS, & Martins SS (2013). Alcohol and Illegal Drug Use Behaviors and Prescription Opioids Use: How Do Nonmedical and Medical Users Compare, and Does Motive to Use Really Matter? *European Addiction Research*, 19(4), 202–210. 10.1159/000345445 [PubMed: 23391856]
- Hyde JS (2014). Gender Similarities and Differences. *Annual Review of Psychology*, 65(1), 373–398. 10.1146/annurev-psych-010213-115057
- Karakos H (2014). Positive Peer Support or Negative Peer Influence? The Role of Peers among Adolescents in Recovery High Schools. *PJE. Peabody Journal of Education*, 89(2), 214–228. 10.1080/0161956X.2014.897094 [PubMed: 24839335]
- Kilmer JR, Geisner IM, Gasser ML, & Lindgren KP (2015). Normative perceptions of non-medical stimulant use: Associations with actual use and hazardous drinking. *Addictive Behaviors*, 42, 51–56. 10.1016/j.addbeh.2014.11.005 [PubMed: 25462654]
- Kristjansson AL, Sigfusdottir ID, & Allegrante JP (2013). Adolescent substance use and peer use: a multilevel analysis of cross-sectional population data. *Substance Abuse Treatment, Prevention, and Policy*, 8, 27. 10.1186/1747-597X-8-27
- Larimer ME, Neighbors C, LaBrie JW, Atkins DC, Lewis MA, Lee CM, ... Walter T. (2011). Descriptive drinking norms: For whom does reference group matter? *Journal of Studies on Alcohol and Drugs*, 72(5), 833–843. [PubMed: 21906510]
- Lewis MA, Litt DM, & Neighbors C (2015). The Chicken or the Egg: Examining Temporal Precedence Among Attitudes, Injunctive Norms, and College Student Drinking. *Journal of Studies on Alcohol and Drugs*, 76(4), 594–601. 10.15288/jsad.2015.76.594 [PubMed: 26098035]
- Luk JW, Farhat T, Iannotti RJ, & Simons-Morton BG (2010). Parent–child communication and substance use among adolescents: Do father and mother communication play a different role for sons and daughters? *Addictive Behaviors*, 35(5), 426–431. 10.1016/j.addbeh.2009.12.009 [PubMed: 20060651]

- Maahs JR, Weidner RR, & Smith R (2016). Prescribing Some Criminological Theory: An Examination of the Illicit Use of Prescription Stimulants Among College Students. *International Journal of Offender Therapy and Comparative Criminology*, 60(2), 146–164. 10.1177/0306624X14548530 [PubMed: 25156423]
- MacArthur G, Harrison S, Caldwell D, Hickman M, & Campbell R (2016). Peer- led interventions to prevent tobacco, alcohol and/or drug use among young people aged 11–21 years: a systematic review and meta- analysis. *Addiction (Abingdon, England)*, 111(3), 391–407. 10.1111/add.13224
- Maddahian E, Newcomb MD, & Bentler PM (1986). Adolescents' substance use: impact of ethnicity, income, and availability. *Advances in Alcohol & Substance Abuse*, 5(3), 63–78. 10.1300/J251v05n03\_05 [PubMed: 3487931]
- Malmberg M, Overbeek G, Vermulst AA, Monshouwer K, Vollebergh WAM, & Engels RCME (2012). The theory of planned behavior: Precursors of marijuana use in early adolescence? *Drug and Alcohol Dependence*, 123(1–3), 22–28. 10.1016/j.drugalcdep.2011.10.011 [PubMed: 22056217]
- Martens MP, Page JC, Mowry ES, Damann KM, Taylor KK, & Cimini MD (2006). Differences Between Actual and Perceived Student Norms: An Examination of Alcohol Use, Drug Use, and Sexual Behavior. *Journal of American College Health*, 54(5), 295–300. 10.3200/JACH.54.5.295-300 [PubMed: 16539222]
- McCabe, Sean E, West BT, Morales M, Cranford JA, & Boyd CJ. (2007). Does early onset of non-medical use of prescription drugs predict subsequent prescription drug abuse and dependence? Results from a national study. *Addiction (Abingdon, England)*, 102(12), 1920–1930. 10.1111/j.1360-0443.2007.02015.x
- McCabe, Sean Esteban. (2008). Misperceptions of non-medical prescription drug use: A web survey of college students. *Addictive Behaviors*, 33(5), 713–724. 10.1016/j.addbeh.2007.12.008 [PubMed: 18242002]
- McCabe, Sean Esteban, & Boyd CJ (2005). Sources of prescription drugs for illicit use. *Addictive Behaviors*, 30(7), 1342–1350. 10.1016/j.addbeh.2005.01.012 [PubMed: 16022931]
- McCabe, Sean Esteban, Cranford JA, Boyd CJ, & Teter CJ (2007). Motives, diversion and routes of administration associated with nonmedical use of prescription opioids. *Addictive Behaviors*, 32(3), 562–575. 10.1016/j.addbeh.2006.05.022 [PubMed: 16843611]
- McCabe, Sean Esteban, West BT, & Boyd CJ (2013). Leftover prescription opioids and nonmedical use among high school seniors: a multi-cohort national study. *The Journal of Adolescent Health: Official Publication of the Society for Adolescent Medicine*, 52(4), 480–485. 10.1016/j.jadohealth.2012.08.007 [PubMed: 23298996]
- McDonough MH, Jose PE, & Stuart J (2016). Bi-directional Effects of Peer Relationships and Adolescent Substance Use: A Longitudinal Study. *Journal of Youth and Adolescence*, 45(8), 1652–1663. 10.1007/s10964-015-0355-4 [PubMed: 26391360]
- Neighbors C, Lee CM, Lewis MA, Fossos N, & Larimer ME (2007). Are Social Norms the Best Predictor of Outcomes Among Heavy-Drinking College Students? *Journal of Studies on Alcohol and Drugs*, 68(4), 556–565. [PubMed: 17568961]
- Osborne V, Serdarevic M, Crooke H, Striley C, & Cottler LB (2017). Non-medical opioid use in youth: Gender differences in risk factors and prevalence. *Addictive Behaviors*, 72, 114–119. 10.1016/j.addbeh.2017.03.024 [PubMed: 28391071]
- Paschall MJ, Grube JW, & Kypri K (2009). Alcohol control policies and alcohol consumption by youth: a multi-national study. *Addiction*, 104(11), 1849–1855. 10.1111/j.1360-0443.2009.02698.x [PubMed: 19832785]
- Perkins H (2002). Social norms and the prevention of alcohol misuse in collegiate contexts. *Journal of Studies on Alcohol*, 14(Supplement), 164–172.
- Perkins HW, & Berkowitz AD (1986). Perceiving the community norms of alcohol use among students: some research implications for campus alcohol education programming. *The International Journal of the Addictions*, 21(9–10), 961–976. [PubMed: 3793315]
- Perkins H Wesley, Linkenbach JW, Lewis MA, & Neighbors C (2010). Effectiveness of social norms media marketing in reducing drinking and driving: A statewide campaign. *Addictive Behaviors*, 35(10), 866–874. 10.1016/j.addbeh.2010.05.004 [PubMed: 20619177]

- Ponnet K, Wouters E, Walrave M, Heirman W, & Van Hal G (2014). Predicting Students' Intention to use Stimulants for Academic Performance Enhancement. *Substance Use & Misuse* 10.3109/10826084.2014.952446
- Popova S, Giesbrecht N, Bekmuradov D, & Patra J (2009). Hours and Days of Sale and Density of Alcohol Outlets: Impacts on Alcohol Consumption and Damage: A Systematic Review. *Alcohol and Alcoholism*, 44(5), 500–516. 10.1093/alcalc/agg054 [PubMed: 19734159]
- Ridenour TA, Maldonado-Molina M, Compton WM, Spitznagel EL, & Cottler LB (2005). Factors associated with the transition from abuse to dependence among substance abusers: Implications for a measure of addictive liability. *Drug & Alcohol Dependence*, 80(1), 1–14. 10.1016/j.drugalcdep.2005.02.005 [PubMed: 16157227]
- Ross-Durow PL, McCabe SE, & Boyd CJ (2013). Adolescents' Access to Their Own Prescription Medications in the Home. *The Journal of Adolescent Health: Official Publication of the Society for Adolescent Medicine* 10.1016/j.jadohealth.2013.02.012
- Serdarevic M, Striley CW, & Cottler LB (2017). Sex differences in prescription opioid use. *Current Opinion in Psychiatry*, 30(4), 238–246. 10.1097/YCO.0000000000000337 [PubMed: 28426545]
- Song E-Y, Smiler AP, Wagoner KG, & Wolfson M (2012). Everyone Says It's OK: Adolescents' Perceptions of Peer, Parent, and Community Alcohol Norms, Alcohol Consumption, and Alcohol-Related Consequences. *Substance Use & Misuse*, 47(1), 86–98. 10.3109/10826084.2011.629704 [PubMed: 22216994]
- Unger JB, Rohrbach LA, Howard-Pitney B, Ritt-Olson A, & Mouttapa M (2001). Peer influences and susceptibility to smoking among California adolescents. *Substance Use & Misuse*, 36(5), 551–571. [PubMed: 11419487]
- Vidourek RA, King KA, & Burbage ML (2014). Synthetic Thc Use Among University Students: An Exploration of Perceived Harm, Ease of Access, and Friends' Use. *American Journal of Health Studies*, 29(1), 7–12.
- Witt SD (1997). Parental influence on children's socialization to gender roles. *Adolescence*, 32(126), 253–259. [PubMed: 9179321]

**Table 1.**

Sample characteristics of the overall sample and by nonmedical prescription opioid use

	Overall Sample	NMPOU		p-value
	(n=11,761) N(%)	No (n=10,718) N(%)	Yes (n=297) N(%)	
<b>Gender</b>			139 (46.8)	0.916
Male	5,497 (46.7)	5,070 (47.3)	143 (48.1)	
Female	5,644 (48.0)	5,283 (49.3)		
<b>Race</b>				0.158
White	8,346 (75.5)	7,823 (75.9)	209 (73.3)	
Black/African American	780 (7.1)	702 (6.8)	25 (8.4)	
Multiracial	681 (6.2)	263 (2.6)	26 (8.8)	
American Indian/Native American	260 (2.4)	242 (2.3)	4 (1.4)	
Asian/Pacific Islander	288 (2.6)	626 (6.1)	5 (1.8)	
Other	702 (6.3)	652 (6.9)	16 (5.6)	
<b>Grade</b>				<0.001
6th	2,114 (18.0)	1,953 (18.2)	21 (7.1)	
7th	1,982 (16.9)	1,884 (17.6)	16 (5.4)	
8th	2,089 (17.8)	1,982 (18.5)	36 (12.1)	
9th	1,476 (12.5)	1,364 (12.7)	37 (12.5)	
10th	1,214 (10.3)	1,113 (10.4)	50 (16.8)	
11th	1,267 (10.8)	1,165 (10.9)	53 (17.8)	
12th	1,117 (9.5)	985 (9.2)	77 (25.9)	
<b>Perceived risk of NMPOU</b>				<0.001
No Risk	1,185 (10.9)	1,081 (10.6)	31 (11.1)	
Slight Risk	757 (7.0)	672 (6.6)	52 (18.6)	
Moderate Risk	1,945 (17.9)	1,760 (17.3)	107 (38.4)	
Great Risk	6,973 (64.2)	6,648 (65.4)	89 (31.9)	
<b>Perceived parent disapproval of NMPOU</b>				<0.001
Not wrong at all	383 (3.7)	325 (3.4)	19 (7.2)	
A little bit wrong	137 (1.3)	106 (1.1)	28 (10.6)	
Wrong	599 (5.8)	537 (5.6)	43 (16.3)	
Very wrong	9,145 (89.1)	8,679 (90.0)	173 (65.8)	
<b>Perceived peer disapproval of NMPOU</b>				<0.001
Not wrong at all	805 (7.7)	662 (6.7)	89 (33.1)	
A little bit wrong	573 (5.5)	494 (5.0)	58 (21.6)	
Wrong	1,494 (14.2)	1,392 (14.1)	43 (16.3)	
Very wrong	7,618 (72.6)	7,295 (74.1)	173 (65.8)	
<b>Perceived peer NMPOU</b>				<0.001
None of them used	9,834 (90.2)	9,401 (92.2)	97 (34.0)	
Less than half of them used	611 (5.6)	499 (4.9)	89 (31.2)	
About half of them used	184 (1.7)	124 (1.2)	44 (15.4)	
More than half of them used	110 (1.0)	73 (0.7)	32 (11.2)	

	Overall Sample	NMPOU		p-value
	(n=11,761) N(%)	No (n=10,718) N(%)	Yes (n=297) N(%)	
All of them used	161 (1.5)	103 (1.0)	23 (8.1)	
<b>Perceived parent(s) restrict access to prescription opioids</b>	6,002 (64.3)	5,721 (65.2)	101 (42.1)	<0.001

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

**Table 2.**

Adjusted odds ratios for past year nonmedical prescription opioid use among the overall sample, males and females

	Overall (n=11,761)	Males (n=5,497)			Females (n=5,644)		
	AOR (95%CI)	NMPOU		AOR (95%CI)	NMPOU		AOR (95%CI)
		No N(%)	Yes N(%)		No N(%)	Yes N(%)	
<b>Gender</b>							
Male	-						
Female	1.07 (0.86, 1.34)						
<b>Race</b>							
White (referent)	-	3,691	93	-	3,899	109	-
Black/African American	0.93 (0.60, 1.43)	(75.3) 329 (6.7)	(68.9) 14	0.95 (0.51, 1.78)	(76.6) 351 (6.9)	(79.0) 8 (5.8)	0.92 (0.50, 1.67)
Multiracial	1.02 (0.65, 1.59)	290 (5.9)	(10.4) 14 (10.4)	0.91 (0.46, 1.80)	315 (6.2)	11 (8.0)	1.11 (0.61, 2.00)
Other	0.96 (0.66, 1.40)	330 (6.7)	9 (6.7)	0.92 (0.54, 1.57)	298 (5.9)	7 (5.1)	1.03 (0.60, 1.76)
<b>Grade</b>	1.07 (1.00, 1.13)*	3.50 (1.9)	4.90 (1.9)	1.09 (1.00, 1.19)	3.50 (1.9)	4.94 (1.8)	1.05 (0.96, 1.14)
<b>Risk of NMPOU</b>	0.97 (0.87, 1.10)	3.33 (1.0)	2.87 (1.9)	0.95 (0.81, 1.12)	3.42 (1.0)	2.96 (0.9)	0.99 (0.84, 1.17)
<b>Parent disapproval of NMPU</b>	0.91 (0.78, 1.07)	3.79 (0.7)	3.40 (1.0)	1.02 (0.81, 1.28)	3.86 (0.5)	3.42 (0.9)	0.79 (0.63, 0.98)*
<b>Peer disapproval of NMPU</b>	0.74 (0.65, 0.83)***	3.47 (0.9)	2.32 (1.2)	0.78 (0.66, 0.93)**	3.65 (0.8)	2.39 (1.1)	0.69 (0.58, 0.82)***
<b>Peer NMPOU</b>	1.74 (1.57, 1.95)***	1.13 (0.6)	2.30 (1.3)	1.75 (1.51, 2.04)***	1.13 (0.5)	2.24 (1.2)	1.72 (1.47, 2.02)***
<b>Parent(s) restrict access to PO</b>	0.87 (0.69, 1.10)	2,577 (63.4)	42 (38.9)	0.87 (0.62, 1.22)	3,010 (67.0)	51 (42.5)	0.89 (0.64, 1.22)

\* p&lt;0.05

\*\* p&lt;0.01

\*\*\* p&lt;0.001



**Table 3.**

Gender moderated regression analyses of past year nonmedical prescription opioid use among the overall sample (n=11,761)

	Model 1 AOR (95% CI)	Model 2 AOR (95% CI)	Model 3 AOR (95% CI)	Model 4 AOR (95% CI)	Model 5 AOR (95% CI)
<b>Gender</b>					
Male	1.07 (0.85, 1.34)	1.06 (0.85, 1.33)	1.04 (0.82, 1.32)	1.07 (0.86, 1.34)	1.11 (0.79, 1.55)
Female					
<b>Race</b>					
White (referent)					
Black/African	0.93 (0.60, 1.43)	0.93 (0.61, 1.44)	0.93 (0.60, 1.44)	0.93 (0.60, 1.43)	0.93 (0.60, 1.43)
American	1.02 (0.65, 1.59)	1.02 (0.65, 1.59)	1.02 (0.65, 1.59)	1.02 (0.65, 1.59)	1.02 (0.65, 1.59)
Multiracial	0.96 (0.66, 1.40)	0.70 (0.66, 1.42)	0.96 (0.66, 1.40)	0.96 (0.66, 1.40)	0.96 (0.66, 1.40)
Other					
<b>Grade</b>	1.07 (1.00, 1.13) <sup>*</sup>	1.07 (1.00, 1.13) <sup>*</sup>	1.07 (1.00, 1.13) <sup>*</sup>	1.07 (1.00, 1.13) <sup>*</sup>	1.07 (1.00, 1.13) <sup>*</sup>
<b>Risk of NMPOU</b>	1.00 (0.86, 1.17)	0.97 (0.87, 1.09)	0.97 (0.87, 1.10)	0.97 (0.87, 1.10)	0.97 (0.87, 1.09)
<b>Parent disapproval of NMPU</b>	0.91 (0.78, 1.07)	1.05 (0.85, 1.30)	0.90 (0.77, 1.06)	0.91 (0.78, 1.07)	0.91 (0.78, 1.07)
<b>Peer disapproval of NMPU</b>	0.74 (0.65, 0.83) <sup>***</sup>	0.73 (0.65, 0.83) <sup>***</sup>	0.80 (0.68, 0.93) <sup>***</sup>	0.74 (0.65, 0.83) <sup>***</sup>	0.74 (0.66, 0.83) <sup>***</sup>
<b>Peer NMPOU</b>	1.75 (1.57, 1.95) <sup>***</sup>	1.74 (1.56, 1.94) <sup>***</sup>	1.74 (1.56, 1.93) <sup>***</sup>	1.74 (1.57, 1.95) <sup>***</sup>	1.74 (1.57, 1.94) <sup>***</sup>
<b>Parent(s) restrict access to PO</b>	0.87 (0.69, 1.10)	0.88 (0.70, 1.11)	0.87 (0.69, 1.10)	0.87 (0.69, 1.10)	0.90 (0.65, 1.26)
<b>Gender*Risk</b>	0.95 (0.77, 1.17)	-	-	-	-
<b>Gender*Parent disapproval</b>	-	0.74 (0.56, 0.97) <sup>*</sup>	-	-	-
<b>Gender*Peer disapproval</b>	-	-	0.86 (0.71, 1.04)	-	-
<b>Gender*Peer NMPOU</b>	-	-	-	1.04 (0.85, 1.28)	-
<b>Gender*Restrict access</b>	-	-	-	-	0.94 (0.60, 1.48)

\*  
p<0.05

\*\*  
p<0.01

\*\*\*  
p<0.001