

RICHARD W. WILLY, PT, PhD, OCS¹ • LISA HALSEY, DPT¹ • ANDREW HAYEK, DPT¹
 HOLLY JOHNSON, BS¹ • JOHN D. WILLSON, PT, PhD¹

Patellofemoral Joint and Achilles Tendon Loads During Overground and Treadmill Running

The patellofemoral joint and Achilles tendon are among the most common sites of injuries sustained by runners. More specifically, patellofemoral pain and Achilles tendinopathy represent up to 25% and 9.5% of all running injuries, respectively.^{31,46} As a result of the high prevalence associated with these injuries, it is not surprising that individuals with these injuries make up a large proportion of patients in sports medicine clinics.^{14,35}

Factors previously related to patellofemoral pain and Achilles tendinopathy

in runners include injury history, age, strength deficits, training errors, structural issues, sex, and biomechanical overloading.^{11,18,32,33,37,39,54} Biomechanical loading of anatomical structures during

running is complex and multifaceted. Specifically, large biomechanical loads (peak load) are generally applied at a rapid rate (loading rate) and in a highly repetitive manner (cumulative load) to articular structures and tendons through the course of a run.^{8,11,20} Thus, measures of peak load, loading rate, and cumulative load of the patellofemoral joint cartilage and Achilles tendon should all be considered in biomechanical investigations of these structures.

Treadmills are commonplace in training and rehabilitation settings. Treadmills are convenient, particularly during inclement weather or when options for outdoor running are restricted. Treadmills are also routinely used in clinical gait analysis and gait-retraining programs due to the ability to evaluate and retrain running mechanics in a controlled environment.^{3,12,43} Further, treadmills are often a fixture in training programs and return-to-running programs after injury to the patellofemoral joint or Achilles tendon. Instrumented treadmills are now commonly used in biomechanical studies of ankle and knee mechanics during running.^{7,29,30,40,52} In particular, instrumented treadmills enable the study of repetitive gait cycles and facilitate more in-depth analyses, such as exertion and gait-modification studies.^{23,51} Despite their common use in both of these applications, little is known regarding the potential

● **STUDY DESIGN:** Level 4, controlled laboratory study.

● **BACKGROUND:** Little is known regarding how the potential differences between treadmill and overground running may affect patellofemoral joint and Achilles tendon loading characteristics.

● **OBJECTIVES:** To compare measures of loading of the patellofemoral joint and Achilles tendon across treadmill and overground running in healthy, uninjured runners.

● **METHODS:** Eighteen healthy runners ran at their self-selected speed on an instrumented treadmill and overground, while 3-D running mechanics were sampled. A musculoskeletal model derived peak load, rate of loading, and estimated cumulative load per 1 km of continuous running for the patellofemoral joint and Achilles tendon for each condition. Data were analyzed via paired *t* tests and Pearson correlations to detect differences and assess relationships, respectively, between the 2 running mediums.

● **RESULTS:** No differences ($P > .05$) were found between treadmill and overground running for peak load, rate of loading, or estimated cumulative

patellofemoral joint stress per 1 km of continuous running. However, treadmill running resulted in 12.5% greater peak Achilles tendon force ($P < .001$), 15.6% greater loading rate of Achilles tendon force ($P < .001$), and 14.2% greater estimated cumulative Achilles tendon force per 1 km of continuous running ($P < .001$) compared with overground running. There were strong ($r > 0.70$) and moderate agreements ($r > 0.50$) for most patellofemoral joint and Achilles measures, respectively, between treadmill and overground running.

● **CONCLUSION:** No differences were observed in loading characteristics to the patellofemoral joint between running mediums; however, treadmill running resulted in greater Achilles tendon loading compared with overground running. Future investigations should examine whether sudden bouts of treadmill running may increase the risk of mechanical overload of the Achilles tendon in runners who habitually train overground. *J Orthop Sports Phys Ther* 2016;46(8):664-672. Epub 12 May 2016. doi:10.2519/jospt.2016.6494

● **KEY WORDS:** ankle, biomechanics, knee, musculoskeletal model

¹Department of Physical Therapy, East Carolina University, Greenville, NC. Prior to initiation of this study, the research protocol was approved by the East Carolina University Human Subjects Research Board. The authors certify that they have no affiliations with or financial involvement in any organization or entity with a direct financial interest in the subject matter or materials discussed in the article. Address correspondence to Dr Richard Willy, Department of Physical Therapy, College of Allied Health Sciences, East Carolina University, Greenville, NC 27834. E-mail: willyr@ecu.edu • Copyright ©2016 *Journal of Orthopaedic & Sports Physical Therapy*[®]

differences of loading to the patellofemoral joint and the Achilles tendon during overground and treadmill running.

Seminal biomechanical comparisons between treadmill and overground running suggest that these running mediums have largely similar knee and ankle kinematics, particularly in the sagittal plane.^{19,40} However, potential differences in joint kinetics exist, suggesting that there are differences in loading characteristics of the patellofemoral joint and Achilles tendon between overground and treadmill running. For instance, treadmill running has been reported to result in an approximately 27% lower peak internal knee extensor moment compared with overground running.⁴⁰ The peak knee extensor moment likely closely relates to peak quadriceps force,¹ which in turn greatly influences patellofemoral joint reaction force.⁵² However, as knee flexion may also be less during treadmill running,^{19,40} a corresponding reduction in patellofemoral contact area may also occur.⁴ Therefore, it is unclear whether there are differences in patellofemoral joint stress (patellofemoral joint reaction force/patellofemoral contact area) between treadmill and overground running. Conversely, the peak plantar flexor moment and eccentric ankle joint power may be as much as 14% and 16% higher, respectively, during treadmill running,⁴⁰ suggesting greater demands on the Achilles tendon.

Previous work has also investigated the temporospatial differences between treadmill and overground running that may significantly affect cumulative loading of the patellofemoral joint and Achilles tendon. Compared with overground running, runners tend to adopt a 1% to 5% shorter step length during treadmill running.^{17,40} This potentially important temporospatial difference may have consequences for patellofemoral joint and Achilles tendon loading. First, a shorter step length during treadmill running may indicate a shorter stance phase, which may, in turn, result in a greater loading rate of the patellofemoral joint and Achil-

| TABLE 1 | | DEMOGRAPHICS FOR PARTICIPANTS (N = 18) | |
|----------------------------------|--|--|--|
| Characteristic | | Value* | |
| Age, y | | 23.6 ± 3.5 | |
| BMI, kg/m ² | | 22.2 ± 2.6 | |
| Running volume, km/wk | | 36.7 ± 26.5 | |
| Running experience, y | | 7.4 ± 3.6 | |
| Self-paced running velocity, m/s | | 2.9 ± 0.3 | |
| Treadmill comfort score (0-10) | | 9.6 ± 0.5 | |
| Tegner score (0-10) | | 6.9 ± 0.6 | |

Abbreviation: BMI, body mass index.
**Values are mean ± SD.*

les tendon if peak loads are of the same or greater magnitude as those during overground running. Second, the shorter step length associated with treadmill running may result in a greater number of steps (loading cycles) to cover a given distance, which may in turn increase cumulative loading on the patellofemoral joint and Achilles tendon during a sustained run.

The purpose of this study was to assess peak loads, rate of loading, and cumulative loading of the patellofemoral joint and the Achilles tendon during treadmill and overground running. Due to a reduced knee extensor moment, we hypothesized that treadmill running would result in reduced peak patellofemoral joint stress and patellofemoral joint stress loading rate. Conversely, we hypothesized that there would be greater Achilles tendon loading and loading rate during treadmill running. Finally, we hypothesized that greater cumulative patellofemoral joint stress and Achilles tendon loading would result due to a reduced step length during treadmill running.

METHODS

PRIOR TO CONDUCTING THE STUDY, the research protocol was approved by the East Carolina University Human Subjects Research Board. A sample-size estimate was calculated to determine the number of participants necessary to detect differences between conditions. Using an alpha of .05, a beta of .2, and the

means and variability of the peak knee extensor and plantar flexor moments between overground and treadmill running established by Riley and colleagues,⁴⁰ it was determined that 18 participants would be necessary to adequately power this study. Accordingly, 18 recreational runners (9 male, 9 female) were recruited from a large university and area running clubs.

All participants provided written and verbal consent prior to enrollment. In order to qualify, all participants were required to be habitual runners (defined as at least 10 km/wk for at least the previous 6 months), free of any lower extremity surgeries, and injury free for at least the previous 3 months. Participants had to be 18 to 35 years of age to limit the heterogeneity of biomechanics and Achilles tendon properties that may be introduced by those over 35 years of age.^{15,41} Comfort with treadmill running can affect running mechanics³⁸; therefore, only volunteers who were comfortable with treadmill running, defined as a minimum score of 8 on a visual analog scale (0 as completely uncomfortable and 10 as completely comfortable) were enrolled. While not an inclusion/exclusion criterion, involvement in endurance running (years of running experience) was also collected. **TABLE 1** displays demographics of the cohort of runners in this investigation.

Fifty-six retroreflective markers were affixed to the bilateral lower extremities,

pelvis, and trunk of each participant. Static calibration and dynamic hip trials²⁸ were collected. The pelvis coordinate system was defined by markers placed on the midline of the iliac crests and the greater trochanters. The thigh coordinate system was defined proximally by the hip joint center, calculated from the dynamic hip trial, and distally by the femoral condyles. The shank coordinate system was defined proximally by the tibial condyles and distally by the malleoli. Finally, the foot was defined proximally by the malleoli and distally by the first and fifth metatarsal heads and the distal aspect of the shoe. Tracking markers consisted of markers placed on the anterior superior iliac spines and shell-mounted clusters on the sacrum, the posterolateral aspect of the thigh and shank, and a cluster of 3 markers on the rearfoot. This is a common marker configuration and was similar to the marker set used by Fellin et al,¹⁹ a study of comparison for the present investigation.

After a 6-minute treadmill accommodation period,³⁴ 3-D running mechanics were sampled for 10 seconds at each participant's self-selected running speed. Participants were cued to choose this speed based on perception of their running pace during the middle of a standard training run. The self-selected running speed was based on the participant's feedback during the final 4 minutes of the treadmill accommodation period. Ground reaction forces and marker trajectories were sampled at 1000 Hz by the instrumented treadmill (Bertec Corporation, Columbus, OH) and 200 Hz by a 10-camera motion-capture system (Qualisys AB, Gothenburg, Sweden), respectively. Prior to study initiation, treadmill-speed calibration during running was performed using a digital tachometer every 0.2 m/s up to 4.0 m/s (HT-5500; Ono Sokki Co Ltd, Yokohama, Japan). The treadmill-running trial consisted of sustained running for no longer than 5 minutes, with approximately 10 minutes of rest provided to each runner between the end of the treadmill trial and

the start of the overground trial to minimize fatigue.

Next, 3-D overground running mechanics were sampled as runners traversed a 25-m runway at the same self-selected running speed ($\pm 3\%$) used during the treadmill-running trial. Each runner practiced the overground trial for several minutes to acclimate to the overground collection procedures, including establishment of running speed and runway starting position. Displacement of a single marker attached to the sacrum has previously been demonstrated to correspond to the displacement of a runner's estimated center of mass.^{21,22} Therefore, we tracked the anterior velocity of a sacral marker in real time to measure running speed as the runner traversed force plates flush with the runway floor (Advanced Mechanical Technology, Inc, Watertown, MA). In postprocessing, this method for tracking overground running velocity was highly correlated to the anterior velocity of the runner's estimated center of mass (correlation between anterior velocity of the sacral marker and estimated center of mass: Pearson $r = 0.96$, $P < .001$, with a root-mean-square error of 0.1 m/s). Any trials that fell outside the velocity range, in which the participant was visibly changing velocity in the capture volume or when the force plates were targeted by the participant, were discarded. The rationale for excluding trials in this manner was that different gait velocities and force-plate targeting can have marked effects on the magnitudes of segmental velocities, joint moments, and powers.^{2,6} Marker trajectories (Qualisys AB) and ground reaction forces were sampled with the exact same parameters as those utilized during the treadmill trial (200 Hz and 1000 Hz for kinematics and kinetics, respectively).

The order of testing (treadmill first, followed by overground testing) was chosen to determine each participant's safe self-selected running speed for the treadmill trials. In testing during protocol development, pilot subjects tended to self-select a running speed for over-




FIGURE 1. Patellofemoral joint reaction forces from both overground and treadmill running in the present study (hash marks correspond to ± 1 SD), contrasted with other published values of patellofemoral joint reaction forces during running. Chen and Powers⁹ utilized faster running velocity (present investigation, 2.9 m/s; Chen and Powers,⁹ 3.33 m/s), which may partly explain the higher values. In contrast, Lenhart et al²⁹ utilized nearly identical running velocities as those in the present investigation (2.8 m/s). Both the Chen and Powers⁹ and the Lenhart et al²⁹ models accounted for cocontraction of the knee musculature, as did the model utilized in the present investigation. In contrast, the model used by Sinclair and Selfe⁴² did not account for cocontraction of the knee musculature, which may have contributed to their lower patellofemoral joint reaction force values.

ground trials that was faster and did not represent a running speed that could be maintained on the treadmill. This mismatch between overground and treadmill running speeds might have been due to the fact that sustained running is not tested in overground trials, whereas treadmill running requires sustained running.

Data Processing and Musculoskeletal Model

Using a sagittal-frontal-transverse plane Euler angle sequence, joint coordinates were calculated with a 6-degree-of-freedom model (The MotionMonitor; Innovative Sports Training, Inc, Chicago, IL). Marker and ground reaction forces were filtered with a 15-Hz cutoff frequency via a low-pass, fourth-order Butterworth recursive filter. Matched cutoff filter frequencies are recommended to minimize nonphysiological signal artifacts during inverse dynamic routines that may occur in high-impact activities such as running.^{5,26} Internal joint moments




FIGURE 2. Time-series data for group mean data for sagittal plane knee kinematics and kinetics and patellofemoral joint stress during treadmill and overground running. (A) Knee flexion, (B) knee extension moment, and (C) patellofemoral joint stress.




FIGURE 3. Time-series data for group mean data for sagittal plane ankle kinematics, ankle kinetics, and Achilles tendon loading during treadmill and overground running. (A) Sagittal ankle power, (B) Achilles tendon force, (C) sagittal ankle moment, and (D) sagittal ankle kinematics. *Significant ($P < .05$). †Significant ($P < .005$).

were then derived using an inverse dynamic routine with published segmental inertial parameters¹³ and reported in the coordinate system of the distal segment. The dominant limb was used for all subsequent analyses. Separate, time-synchronized files of the vertical ground reaction force data were digitally filtered at 50 Hz using a low-pass, fourth-order Butterworth recursive filter and used to identify stance. Initial contact during the running trials was defined as the time when the vertical ground reaction force exceeded 20 N. Five stance phases of the dominant lower extremity (limb used to kick a ball) were analyzed from both the

treadmill and overground running trials. We retained the first 5 complete stance phases from the 10-second treadmill trial for analysis. For the overground trials, we chose the 5 trials with gait velocities that were closest to the treadmill gait speed to minimize the potential error that may be introduced by differing speeds between the 2 testing modes.

To calculate patellofemoral joint stress and Achilles tendon forces, we utilized a musculoskeletal model that has been fully described elsewhere.^{16,52,53} This model uses an inverse-dynamics approach to calculate hamstrings, quadriceps, gastrocnemius, and soleus muscle forces. As such,

this procedure accounts for knee joint cocontraction from the hamstrings and gastrocnemius.⁵² From the net hip extensor moment, hamstring force was calculated utilizing published hamstring and gluteus maximus cross-sectional areas and muscle moment arms as a function of hip angle.^{36,50} The net plantar flexor moment and the Achilles tendon muscle moment arm were then used to derive the Achilles tendon force.^{25,45} Achilles tendon force was further proportioned to the gastrocnemius and the soleus, based on the physiological cross-sectional area of each muscle.⁵⁰ To account for cocontraction about the knee, hamstring and gastrocnemius torques were calculated, using their respective moment arms at the knee, and then summed with the internal knee extension moment.^{24,44,45,49} Quadriceps force was then derived as the quotient of the adjusted quadriceps moment and the quadriceps moment arm.^{24,48} Patellofemoral joint reaction force was then calculated utilizing the quadriceps force as a function of knee joint angle.⁴⁷ See **FIGURE 1** for a comparison of patellofemoral joint reaction force output for our model compared with published values from other musculoskeletal models of varying complexity.^{9,29,42} Finally, patellofemoral joint stress was estimated as the quotient of the patellofemoral joint reaction force and sex-specific patellofemoral contact areas.⁴

A custom-written code in LabVIEW (National Instruments Corporation, Austin, TX) was used to calculate discrete variables. First, step length (meters) was

calculated. For patellofemoral joint stress and Achilles tendon force, the peak load, loading rate, and impulse (time integral) were recorded for each stance phase. Loading rates were calculated as the middle 60% of the rising curve between initial contact and the respective peaks of patellofemoral joint stress and Achilles tendon force (FIGURES 2 and 3) for each stance. Cumulative patellofemoral joint stress and cumulative Achilles tendon force were estimated as the load per 1 km of continuous running as the product of impulse per stance and number of strides to complete 1 km of continuous running (500 m/step length). To assist with interpreting our results, we also included peak knee extensor moment and peak plantar flexor moment in our analysis. Additionally, we calculated eccentric and concentric power for the ankle plantar flexors (joint power = sagittal plane angular velocity × joint moment), as these measures likely relate closely to energy storage and release of the plantar flexors.

All statistical analyses were performed with SPSS Version 20 (IBM Corporation, Armonk, NY). To detect differences between the 2 running modes, motion data were analyzed with a series of paired, 2-tailed *t* tests ($\alpha = .05$). Effect sizes (*d*) were also calculated to assess the magnitude of any differences, with a small effect corresponding to $d = 0.2$ to 0.4 , a moderate effect corresponding to $d = 0.4$ to 0.8 , and a large effect corresponding to $d \geq 0.8$.¹⁰ To assess the relationship between 2 running modes, discrete variables of interest were analyzed with Pearson's *r* ($\alpha = .05$).

RESULTS

WE FOUND NO DIFFERENCES AND excellent correlation for gait speed between overground and treadmill running for our participants (TABLE 2). All overground trials utilized in the analysis were within $\pm 2.6\%$ of the treadmill running speed. However, step length was significantly shorter ($P < .001$) during treadmill running compared with

| TABLE 2 | | GROUP DATA DURING TREADMILL AND OVERGROUND RUNNING FOR TEMPOROSPATIAL AND KNEE MEASURES* | | | |
|--|---------------|--|---------------------|-------------|-------------------|
| Variable | Treadmill | Overground | P Value | Effect Size | Pearson <i>r</i> |
| Gait speed, m/s | 2.88 ± 0.26 | 2.89 ± 0.27 | .50 | -0.04 | 0.97 [†] |
| Step length, m | 1.04 ± 0.10 | 1.10 ± 0.12 | <.0001 [†] | -0.62 | 0.86 [†] |
| Stance duration, ms | 273.1 ± 30.6 | 277.3 ± 26.1 | .23 | -0.15 | 0.88 [†] |
| Peak knee flexion angle, deg | -34.2 ± 3.5 | -34.3 ± 3.8 | .96 | 0.01 | 0.58 [‡] |
| Peak knee extension moment, Nm/m·kg | 1.18 ± 0.20 | 1.14 ± 0.27 | .28 | 0.19 | 0.77 [†] |
| Peak PFJ reaction force, BW | 4.0 ± 1.0 | 4.0 ± 0.8 | .99 | 0.00 | 0.81 [†] |
| Peak PFJ stress, mPA | 6.2 ± 1.4 | 6.1 ± 1.5 | .73 | 0.04 | 0.86 [†] |
| PFJ stress average loading rate, mPA/s | 131.5 ± 26.9 | 155.6 ± 61.3 | .11 | -0.55 | 0.39 |
| PFJ stress impulse, mPA·s | 0.71 ± 0.22 | 0.71 ± 0.16 | .84 | -0.03 | 0.85 [†] |
| Cumulative PFJ stress, mPA·s/km [§] | 344.5 ± 118.5 | 324.7 ± 73.3 | .21 | 0.21 | 0.88 [†] |

Abbreviations: BW, body weight; PFJ, patellofemoral joint.
**Values are mean ± SD unless otherwise indicated.*
[†]Significant ($P < .005$).
[‡]Significant ($P < .05$).
[§]Estimated PFJ stress to run 1 km continuously.

overground running. This difference was associated with a moderate effect size ($d = -0.62$), yet had an excellent correlation ($P < .001$, $r = 0.86$) between the 2 running modes. Interestingly, stance duration was not different and was highly correlated between the 2 running conditions.

Regarding all knee and patellofemoral joint measures, we found no differences between overground and treadmill running (TABLE 2, FIGURES 1 and 2). We also found moderate to excellent correlations for all knee measures, except for patellofemoral joint stress loading rate, which was not correlated. Specifically, peak knee flexion ($P = .96$, $d = 0.01$; $r = 0.58$, $P = .01$) and peak knee extension moment ($P = .28$, $d = 0.19$; $r = 0.77$, $P < .001$) were not different between the 2 running modes. Peak patellofemoral joint reaction force ($P = .99$, $d = 0.00$; $r = 0.81$, $P < .001$), peak patellofemoral joint stress ($P = .73$, $d = 0.04$; $r = 0.86$, $P < .001$), and the loading rate of patellofemoral joint stress ($P = .09$, $d = -0.55$) were also not different between conditions. However, there was a nonsignificant correlation between the running modes for the loading rate of patellofemoral joint stress ($r = 0.39$, $P = .11$). Despite the additional 23 steps estimated to run 1 km continuously

during treadmill running, estimated cumulative patellofemoral joint stress per 1 km of continuous running ($P = .21$, $d = 0.21$; $r = 0.88$, $P < .001$) during treadmill running was not different from that of the overground condition.

In contrast, we found moderate to large differences at the ankle between overground and treadmill running (TABLE 3, FIGURE 3). With the exception of peak plantar flexor moment and estimated cumulative Achilles tendon force per 1 km of continuous running, all ankle and Achilles values were moderately to strongly correlated between the 2 running modes. Though we found no difference in peak dorsiflexion angle ($P = .32$, $d = -0.15$; $r = 0.81$, $P < .001$), the peak plantar flexor moment ($P = .001$, $d = 1.17$) was significantly greater and not correlated ($r = 0.36$, $P = .14$) during treadmill running compared with overground running. Additionally, peak Achilles tendon force ($P < .001$, $d = 1.01$; $r = 0.52$, $P = .03$), Achilles tendon loading rate ($P < .001$, $d = 0.61$; $r = 0.62$, $P = .006$), Achilles tendon force impulse per stance ($P = .02$, $d = 0.63$; $r = 0.53$, $P = .02$), and estimated cumulative Achilles tendon force per 1 km of continuous running ($P < .001$, $d = 1.04$; $r = 0.39$, $P = .12$) were all significantly greater during

TABLE 3

GROUP DATA DURING TREADMILL AND OVERGROUND RUNNING FOR ANKLE AND ACHILLES TENDON DISCRETE VARIABLES*

| Discrete Variable | Treadmill | Overground | P Value | Effect Size | Pearson <i>r</i> |
|---|--------------|--------------|--------------------|-------------|-------------------|
| Peak dorsiflexion angle, deg | 22.4 ± 3.0 | 22.8 ± 3.0 | .32 | -0.15 | 0.81 [†] |
| Peak plantar flexor moment, Nm/m·kg | -1.52 ± 0.20 | -1.33 ± 0.12 | .001 [†] | 1.17 | 0.36 |
| Peak Achilles force, BW | 5.35 ± 0.782 | 4.68 ± 0.533 | <.001 [†] | 1.01 | 0.52 [‡] |
| Achilles loading rate, BW/s | 65.1 ± 10.8 | 54.7 ± 10.5 | <.001 [†] | 0.61 | 0.62 [‡] |
| Achilles impulse, BW·s | 0.66 ± 0.13 | 0.59 ± 0.08 | .02 [‡] | 0.63 | 0.53 [‡] |
| Cumulative Achilles force, BW/km [§] | 315.8 ± 44.4 | 270.8 ± 41.8 | <.001 [†] | 1.04 | 0.39 |
| Eccentric ankle power, W/kg·m | -3.15 ± 0.82 | -3.32 ± 0.67 | .25 | 0.23 | 0.69 [‡] |
| Concentric ankle power, W/kg·m | 6.19 ± 1.54 | 4.84 ± 0.75 | .001 [†] | 1.18 | 0.69 [‡] |

Abbreviation: BW, body weight.

*Values are mean ± SD unless otherwise indicated.

[†]Significant ($P < .005$).

[‡]Significant ($P < .05$).

[§]Cumulative Achilles load, in body weight, to run 1 km continuously.

treadmill running. Treadmill running was also associated with greater concentric ankle joint power ($P = .001$, $d = 1.18$; $r = 0.69$, $P < .001$), but there was no significant difference in eccentric joint power ($P = .25$, $d = 0.23$; $r = 0.69$, $P < .001$) between the 2 modes of running.

DISCUSSION

WE SOUGHT TO DETERMINE IF there were differences between running overground and running on a treadmill in regard to patellofemoral joint loading and Achilles tendon forces. We found no differences in peak patellofemoral joint reaction force or any measure of patellofemoral joint stress between overground and treadmill running. Due to moderate to strong correlations, this study suggests that findings from studies that utilize instrumented treadmills to assess loading of the patellofemoral joint may be largely applied to overground running, and vice versa. In contrast, ankle concentric power and all measures of Achilles tendon force were greater during treadmill running. While the Achilles tendon loads were moderately proportional between treadmill and overground running, caution should be used when extrapolating absolute values of Achilles tendon loads obtained via

instrumented treadmill running to overground running.

The cohort of runners in the present investigation was a sample of convenience representative of a typical university setting. However, the enrolled runners reported a relatively long length of continuous participation in endurance running of greater than 7 years. While the study was open to runners who ran as few as 10 km/wk, the range for running volume was 13.0 to 96.6 km/wk. Overall, we felt that the length of continuous participation in endurance running, coupled with a high level of comfort with treadmill running (9.6/10), was the best representation of running skill level. In contrast, running volume likely fluctuates throughout the year.

Counter to our hypothesis, we found no differences between overground and treadmill running in respect to sagittal knee joint mechanics, which exert major influences on patellofemoral joint reaction force and stress. Based on the previous literature, we expected reduced knee flexion kinematics and reduced knee extensor moments during treadmill running.^{19,40} There are several potential reasons for the discrepancy with the previous literature. First, the kinematic differences reported by Fellin et al¹⁹ were small (approximately 1.3° less knee flexion dur-

ing treadmill running) and may simply be due to small differences in running speed between overground and treadmill modes. Second, the only previous comparison of knee joint kinetics utilized different signal filtering parameters when processing treadmill and overground trials.⁴⁰ The present investigation utilized identical filtering parameters when processing overground and treadmill trials. The lower low-pass filter cutoff utilized by Riley et al⁴⁰ during treadmill running, when compared to their overground-running data, might have attenuated the knee extensor moment signal, resulting in the slightly lower peak knee extensor moment during treadmill running reported in their study. Finally, the present study examined runners during their normal endurance training pace (2.9 m/s), whereas previous investigations used the estimated 10-km race pace (approximately 3.8 m/s)⁴⁰ or a standardized pace (3.35 m/s).¹⁹ Therefore, differences in sagittal plane knee and patellofemoral joint kinetics between overground and treadmill running may occur at higher running speeds than what were sampled in the present investigation.

There were no differences in peak load, loading rate, and estimated cumulative patellofemoral joint stress per kilometer of continuous running. The 23 additional steps estimated to be required to run 1 km continuously on a treadmill were insufficient to increase the estimated cumulative patellofemoral joint stress per kilometer of continuous running. It has been suggested that the measures of peak load, loading rate, and cumulative joint stress play independent roles in the degradation of articular structures.⁸ Therefore, future study should be undertaken to determine whether return-to-running programs for the treatment of patellofemoral pain result in similar outcomes when conducted on a treadmill or overground. Further, strong relationships ($r \geq 0.81$) were found between overground and treadmill running for peak patellofemoral joint reaction force, peak and impulse patellofemoral joint stress, as well as the estimat-

ed cumulative patellofemoral joint stress to run 1 km continuously. Thus, treadmill and overground running appear to yield similar estimates of patellofemoral joint reaction force and stress measures.

In contrast to the patellofemoral joint, measures of Achilles tendon loading and concentric ankle joint power were considerably greater during treadmill running. Interestingly, peak ankle dorsiflexion was not different during treadmill running. Rather, the peak plantar flexion moment was greater during treadmill running, and this difference was associated with a large effect size. Thus, measures of peak load and loading rate of Achilles tendon force, as well as estimated cumulative Achilles tendon force to run 1 km continuously, were correspondingly greater ($d = 0.61-1.04$) during treadmill running. As stance duration was not different between overground and treadmill running, the greater peak Achilles tendon force was most likely responsible for the higher loading rate of the Achilles tendon. The sagittal ankle power data revealed that concentric ankle joint power was also greater during treadmill running, whereas eccentric ankle joint power was not. This finding contrasts with the previous investigation of ankle joint power during treadmill and overground running that found greater eccentric ankle joint power during treadmill running but similar concentric ankle joint power with overground running.⁴⁰ Potential reasons for this difference between investigations include differences in tested gait velocity (approximately 2.9 m/s in the present study versus 3.8 m/s in Riley et al⁴⁰) and differences in overground runway length (25 m in the present study versus 15 m in Riley et al⁴⁰). Nevertheless, we found moderate correlations for most of the Achilles, ankle joint power, and ankle kinematic measures between the 2 running modes. However, the moderate to large absolute differences that we found at the ankle suggest that caution should be exercised when interpreting Achilles data collected during treadmill running and extrapolating them to overground

running, and vice versa.

The greater estimated cumulative Achilles tendon force to run 1 km continuously during treadmill running may have implications for future study and potential clinical applications.^{11,27} We estimated that 1 km of continuous treadmill running would expose the Achilles tendon to an additional 45 body weights of cumulative force compared with overground running. The well-documented response of tendon to acute bouts of loading suggests that further investigation may be necessary to determine whether an acute bout of treadmill running would result in greater collagen turnover in the Achilles tendon compared to an equal volume of overground running. Further study is necessary to determine whether there are differences in Achilles tendon qualities or greater prevalence of Achilles tendinopathy in individuals who run solely on a treadmill versus solely overground.

Limitations

There are several limitations to the present investigation that should be kept in mind when interpreting these results. First, all participants were tested first on the treadmill, followed by overground. This testing order was chosen to establish a realistic self-selected running speed that could be maintained both overground and during treadmill running. Regardless, this might have introduced an order effect. Second, the musculoskeletal model used in this investigation was not entirely subject specific, utilized muscle architectural parameters from the literature, and represents estimates of in vivo tissue loads. However, any added benefit of subject-specific inputs to the model would have been negligible, due to the within-subject design. As implanted strain gauges are not presently feasible to measure in vivo joint and tendon loads, musculoskeletal models are generally accepted as estimates of these loads. Patellofemoral joint reaction force and Achilles tendon loads found in the present investigation are within the ranges reported in recently published investigations using different

musculoskeletal models.^{20,29,42} Third, the overground runway utilized in this investigation was 25 m in length, with the force plates embedded at approximately the halfway point. Due to the relatively short runway distance, it is possible that participants did not run at a constant speed when traversing the capture volume. This laboratory design is fairly standard and ubiquitous across gait laboratories that study running mechanics. The key papers of comparison for this investigation used runways of 15 m (Riley et al⁴⁰) and 25 m (Fellin et al¹⁹). As a longer track-based laboratory is neither common nor practical for most settings, the use of emerging wearable technologies during continuous outdoor running may provide the most valid comparison with continuous treadmill running. Additionally, the horizontal velocity of the sacral marker was used to provide feedback on running velocity during overground running trials, whereas the treadmill controller was used to control gait speed during treadmill trials. As a result, undetected variations in treadmill gait velocity may have occurred if subjects' positions drifted in the anterior-to-posterior direction on the treadmill during data collection. However, we only collected data when subjects' positions were stationary on the treadmill in an effort to minimize this potential influence. Finally, our participants were injury free and young, and there was a relatively wide range in habitual weekly running volume among the cohort. Therefore, care should be exercised when applying the results of this study to injured or older populations.

CONCLUSION

IN CONCLUSION, TREADMILL AND OVERGROUND running yielded similar estimates of patellofemoral joint reaction force and stress. In contrast, treadmill running resulted in greater Achilles tendon loads when compared to overground running. Further study is necessary to determine the clinical implications of these findings in return-to-running programs

or in assessing the risk of Achilles tendon injury in runners who undergo acute bouts of treadmill running. These findings also suggest that measures of patellofemoral joint reaction force and stress during instrumented treadmill running are a reasonable representation of those same loads during overground running. In contrast, Achilles tendon force estimates obtained during instrumented treadmill running appear to be moderately proportional to, yet greater than, those obtained during overground running. ●

KEY POINTS

FINDINGS: Estimates of patellofemoral joint loading did not differ between treadmill and overground running.

However, Achilles tendon loads and concentric ankle power were significantly greater during treadmill running compared with overground running.

IMPLICATIONS: Patellofemoral joint loading during treadmill running appears to be consistent with overground running. Therefore, the findings of studies examining patellofemoral joint loading during treadmill running can be applied to overground running. Conversely, measures of Achilles tendon loading during treadmill running were moderately correlated with, yet greater than, those during overground running. Future study should determine whether acute bouts of treadmill running place the Achilles tendon at risk for mechanical overload in runners who customarily perform their training overground.

CAUTION: Caution should be exercised when extrapolating these results to individuals with patellofemoral pain or Achilles tendinopathy.

REFERENCES

- Andriacchi TP, Andersson GB, Örtengren R, Mikosz RP. A study of factors influencing muscle activity about the knee joint. *J Orthop Res*. 1984;1:266-275. <http://dx.doi.org/10.1002/jor.1100010306>
- Arampatzis A, Brüggemann GP, Metzler V. The effect of speed on leg stiffness and

joint kinetics in human running. *J Biomech*. 1999;32:1349-1353. [http://dx.doi.org/10.1016/S0021-9290\(99\)00133-5](http://dx.doi.org/10.1016/S0021-9290(99)00133-5)

- Baggaley M, Willy RW, Meardon SA. Primary and secondary effects of real-time feedback to reduce vertical loading rate during running. *Scand J Med Sci Sports*. In press. <http://dx.doi.org/10.1111/sms.12670>
- Besier TF, Draper CE, Gold GE, Beaupré GS, Delp SL. Patellofemoral joint contact area increases with knee flexion and weight-bearing. *J Orthop Res*. 2005;23:345-350. <http://dx.doi.org/10.1016/j.orthres.2004.08.003>
- Bezodis NE, Salo AJ, Trewartha G. Excessive fluctuations in knee joint moments during early stance in sprinting are caused by digital filtering procedures. *Gait Posture*. 2013;38:653-657. <http://dx.doi.org/10.1016/j.gaitpost.2013.02.015>
- Challis JH. The variability in running gait caused by force plate targeting. *J Appl Biomech*. 2001;17:77-83.
- Chambon N, Delattre N, Guéguen N, Berton E, Rao G. Shoe drop has opposite influence on running pattern when running overground or on a treadmill. *Eur J Appl Physiol*. 2015;115:911-918. <http://dx.doi.org/10.1007/s00421-014-3072-x>
- Chen CT, Burton-Wurster N, Lust G, Bank RA, Tekoppele JM. Compositional and metabolic changes in damaged cartilage are peak-stress, stress-rate, and loading-duration dependent. *J Orthop Res*. 1999;17:870-879. <http://dx.doi.org/10.1002/jor.1100170612>
- Chen YJ, Powers CM. Comparison of three-dimensional patellofemoral joint reaction forces in persons with and without patellofemoral pain. *J Appl Biomech*. 2014;30:493-500. <http://dx.doi.org/10.1123/jab.2011-0250>
- Cohen J. A power primer. *Psychol Bull*. 1992;112:155-159.
- Cook JL, Purdam CR. Is tendon pathology a continuum? A pathology model to explain the clinical presentation of load-induced tendinopathy. *Br J Sports Med*. 2009;43:409-416. <http://dx.doi.org/10.1136/bjism.2008.051193>
- Davis IS, Futrell E. Gait retraining: altering the fingerprint of gait. *Phys Med Rehabil Clin N Am*. 2016;27:339-355. <http://dx.doi.org/10.1016/j.pmr.2015.09.002>
- Dempster WT, Gabel WC, Felts WJ. The anthropometry of the manual work space for the seated subject. *Am J Phys Anthropol*. 1959;17:289-317.
- Devereaux MD, Lachmann SM. Patello-femoral arthralgia in athletes attending a Sports Injury Clinic. *Br J Sports Med*. 1984;18:18-21. <http://dx.doi.org/10.1136/bjism.18.1.18>
- Devita P, Fellin RE, Seay JF, Ip E, Stavro N, Messier SP. The relationships between age and running biomechanics. *Med Sci Sports Exerc*. 2016;48:98-106. <http://dx.doi.org/10.1249/MSS.0000000000000744>
- DeVita P, Hortobagyi T. Functional knee brace alters predicted knee muscle and joint forces in people with ACL reconstruction during walking. *J Appl Biomech*. 2001;17:297-311.
- Elliott BC, Blanksby BA. A cinematographic analysis of overground and treadmill running by males and females. *Med Sci Sports*. 1976;8:84-87.
- Farrokhhi S, Keyak JH, Powers CM. Individuals with patellofemoral pain exhibit greater patellofemoral joint stress: a finite element analysis study. *Osteoarthritis Cartilage*. 2011;19:287-294. <http://dx.doi.org/10.1016/j.joca.2010.12.001>
- Fellin RE, Manal K, Davis IS. Comparison of lower extremity kinematic curves during overground and treadmill running. *J Appl Biomech*. 2010;26:407-414.
- Greenhalgh A, Sinclair J. Comparison of Achilles tendon loading between male and female recreational runners. *J Hum Kinet*. 2014;44:155-159. <http://dx.doi.org/10.2478/hukin-2014-0121>
- Gullstrand L, Halvorsen K, Tinmark F, Eriksson M, Nilsson J. Measurements of vertical displacement in running, a methodological comparison. *Gait Posture*. 2009;30:71-75. <http://dx.doi.org/10.1016/j.gaitpost.2009.03.001>
- Halvorsen K, Eriksson M, Gullstrand L, Tinmark F, Nilsson J. Minimal marker set for center of mass estimation in running. *Gait Posture*. 2009;30:552-555. <http://dx.doi.org/10.1016/j.gaitpost.2009.06.014>
- Heiderscheit BC, Chumanov ES, Michalski MP, Wille CM, Ryan MB. Effects of step rate manipulation on joint mechanics during running. *Med Sci Sports Exerc*. 2011;43:296-302. <http://dx.doi.org/10.1249/MSS.0b013e3181ebedf4>
- Herzog W, Read LJ. Lines of action and moment arms of the major force-carrying structures crossing the human knee joint. *J Anat*. 1993;182 pt 2:213-230.
- Klein P, Mattys S, Rooze M. Moment arm length variations of selected muscles acting on talocrural and subtalar joints during movement: an in vitro study. *J Biomech*. 1996;29:21-30. [http://dx.doi.org/10.1016/0021-9290\(95\)00025-9](http://dx.doi.org/10.1016/0021-9290(95)00025-9)
- Kristianslund E, Krosshaug T, van den Bogert AJ. Effect of low pass filtering on joint moments from inverse dynamics: implications for injury prevention. *J Biomech*. 2012;45:666-671. <http://dx.doi.org/10.1016/j.jbiomech.2011.12.011>
- Langberg H, Skovgaard D, Asp S, Kjaer M. Time pattern of exercise-induced changes in type I collagen turnover after prolonged endurance exercise in humans. *Calcif Tissue Int*. 2000;67:41-44. <http://dx.doi.org/10.1007/s00223001094>
- Leardini A, Cappozzo A, Catani F, et al. Validation of a functional method for the estimation of hip joint centre location. *J Biomech*. 1999;32:99-103. [http://dx.doi.org/10.1016/S0021-9290\(98\)00148-1](http://dx.doi.org/10.1016/S0021-9290(98)00148-1)
- Lenhart RL, Smith CR, Vignos MF, Kaiser J, Heiderscheit BC, Thelen DG. Influence of step rate and quadriceps load distribution on patellofemoral cartilage contact pressures during running. *J Biomech*. 2015;48:2871-2878. <http://dx.doi.org/10.1016/j.jbiomech.2015.04.036>
- Lenhart RL, Thelen DG, Wille CM, Chumanov ES,

Heiderscheid BC. Increasing running step rate reduces patellofemoral joint forces. *Med Sci Sports Exerc.* 2014;46:557-564. <http://dx.doi.org/10.1249/MSS.0b013e3182a78c3a>

31. Lopes AD, Hespanhol Junior LC, Yeung SS, Costa LO. What are the main running-related musculoskeletal injuries? A systematic review. *Sports Med.* 2012;42:891-905. <http://dx.doi.org/10.1007/BF03262301>
32. Lorimer AV, Hume PA. Achilles tendon injury risk factors associated with running. *Sports Med.* 2014;44:1459-1472. <http://dx.doi.org/10.1007/s40279-014-0209-3>
33. Mahieu NN, Witvrouw E, Stevens V, Van Tiggelen D, Roget P. Intrinsic risk factors for the development of Achilles tendon overuse injury: a prospective study. *Am J Sports Med.* 2006;34:226-235. <http://dx.doi.org/10.1177/0363546505279918>
34. Matsas A, Taylor N, McBurney H. Knee joint kinematics from familiarised treadmill walking can be generalised to overground walking in young unimpaired subjects. *Gait Posture.* 2000;11:46-53. [http://dx.doi.org/10.1016/S0966-6362\(99\)00048-X](http://dx.doi.org/10.1016/S0966-6362(99)00048-X)
35. Murray IR, Murray SA, MacKenzie K, Coleman S. How evidence based is the management of two common sports injuries in a sports injury clinic? *Br J Sports Med.* 2005;39:912-916; discussion 916. <http://dx.doi.org/10.1136/bjism.2004.017624>
36. Németh G, Ohlsén H. In vivo moment arm lengths for hip extensor muscles at different angles of hip flexion. *J Biomech.* 1985;18:129-140. [http://dx.doi.org/10.1016/0021-9290\(85\)90005-3](http://dx.doi.org/10.1016/0021-9290(85)90005-3)
37. Nielsen RO, Buist I, Sorensen H, Lind M, Rasmussen S. Training errors and running related injuries: a systematic review. *Int J Sports Phys Ther.* 2012;7:58-75.
38. Nigg BM, De Boer RW, Fisher V. A kinematic comparison of overground and treadmill running. *Med Sci Sports Exerc.* 1995;27:98-105.

39. Rasmussen CH, Nielsen RO, Juul MS, Rasmussen S. Weekly running volume and risk of running-related injuries among marathon runners. *Int J Sports Phys Ther.* 2013;8:111-120.
40. Riley PO, Dicharry J, Franz J, Della Croce U, Wilder RP, Kerrigan DC. A kinematics and kinetic comparison of overground and treadmill running. *Med Sci Sports Exerc.* 2008;40:1093-1100. <http://dx.doi.org/10.1249/MSS.0b013e3181677530>
41. Ruan Z, Zhao B, Qi H, et al. Elasticity of healthy Achilles tendon decreases with the increase of age as determined by acoustic radiation force impulse imaging. *Int J Clin Exp Med.* 2015;8:1043-1050.
42. Sinclair J, Selve J. Sex differences in knee loading in recreational runners. *J Biomech.* 2015;48:2171-2175. <http://dx.doi.org/10.1016/j.jbiomech.2015.05.016>
43. Souza RB. An evidence-based videotaped running biomechanics analysis. *Phys Med Rehabil Clin N Am.* 2016;27:217-236. <http://dx.doi.org/10.1016/j.pmr.2015.08.006>
44. Spoor CW, van Leeuwen JL. Knee muscle moment arms from MRI and from tendon travel. *J Biomech.* 1992;25:201-206. [http://dx.doi.org/10.1016/0021-9290\(92\)90276-7](http://dx.doi.org/10.1016/0021-9290(92)90276-7)
45. Spoor CW, van Leeuwen JL, Meskers CG, Titulaer AF, Huson A. Estimation of instantaneous moment arms of lower-leg muscles. *J Biomech.* 1990;23:1247-1259. [http://dx.doi.org/10.1016/0021-9290\(90\)90382-D](http://dx.doi.org/10.1016/0021-9290(90)90382-D)
46. Taunton JE, Ryan MB, Clement DB, McKenzie DC, Lloyd-Smith DR, Zumbo BD. A retrospective case-control analysis of 2002 running injuries. *Br J Sports Med.* 2002;36:95-101. <http://dx.doi.org/10.1136/bjism.36.2.95>
47. van Eijden TM, Kouwenhoven E, Verburg J, Weijs WA. A mathematical model of the patellofemoral joint. *J Biomech.* 1986;19:219-223, 225-229. [http://dx.doi.org/10.1016/0021-9290\(86\)90154-5](http://dx.doi.org/10.1016/0021-9290(86)90154-5)
48. van Eijden TM, Kouwenhoven E, Weijs WA.

Mechanics of the patellar articulation. Effects of patellar ligament length studied with a mathematical model. *Acta Orthop Scand.* 1987;58:560-566.

49. Visser JJ, Hoogkamer JE, Bobbert MF, Huijing PA. Length and moment arm of human leg muscles as a function of knee and hip-joint angles. *Eur J Appl Physiol Occup Physiol.* 1990;61:453-460. <http://dx.doi.org/10.1007/BF00236067>
50. Ward SR, Eng CM, Smallwood LH, Lieber RL. Are current measurements of lower extremity muscle architecture accurate? *Clin Orthop Relat Res.* 2009;467:1074-1082. <http://dx.doi.org/10.1007/s11999-008-0594-8>
51. Willson JD, Loss JR, Willy RW, Meardon SA. Sex differences in running mechanics and patellofemoral joint kinetics following an exhaustive run. *J Biomech.* 2015;48:4155-4159. <http://dx.doi.org/10.1016/j.jbiomech.2015.10.021>
52. Willson JD, Ratcliff OM, Meardon SA, Willy RW. Influence of step length and landing pattern on patellofemoral joint kinetics during running. *Scand J Med Sci Sports.* 2015;25:736-743. <http://dx.doi.org/10.1111/sms.12383>
53. Willy RW, Meardon SA, Schmidt A, Blaylock NR, Hadding SA, Willson JD. Changes in tibiofemoral contact forces during running in response to in-field gait retraining. *J Sports Sci.* 2016;34:1602-1611. <http://dx.doi.org/10.1080/02640414.2015.1125517>
54. Witvrouw E, Callaghan MJ, Stefanik JJ, et al. Patellofemoral pain: consensus statement from the 3rd International Patellofemoral Pain Research Retreat held in Vancouver, September 2013. *Br J Sports Med.* 2014;48:411-414. <http://dx.doi.org/10.1136/bjsports-2014-093450>



MORE INFORMATION
WWW.JOSPT.ORG

FIND Author Instructions & Tools on the Journal's Website

JOSPT's instructions to authors are available at www.jospt.org by clicking **Complete Author Instructions** in the right-hand Author Center widget on the home page, or by visiting the **Info Center for Authors**, located in the site's top navigation bar. The Journal's editors have assembled a list of **useful tools and links** for authors as well as reviewers.