

Annals of Thoracic Medicine

CASE REPORT

Year : 2019 | Volume : 14 | Issue : 4 | Page : 285--287

A novel diagnostic approach for *Pneumocystis jirovecii* pneumonia using fine-needle aspiration, electromagnetic navigational bronchoscopy and rapid on-site evaluation

Farnaz Houshmand¹, Fatima Zahra Aly², Mark Rollin Bowling³,

¹ Berkeley Medical Center, West Virginia University, Martinsburg, WV, USA

² Department of Pulmonary and Critical Care Medicine, University of Florida, Gainesville, FL, USA

³ Division of Pulmonary, Critical Care, and Sleep Medicine, Brody School of Medicine, East Carolina University, Greenville, NC, USA

Correspondence Address:

Dr. Mark Rollin Bowling

Division of Pulmonary, Critical Care, and Sleep Medicine, Brody School of Medicine, East Carolina University, Greenville, NC 28754 USA

Abstract

Cavitary lung lesions are common in patients with human immunodeficiency virus infections. Both atypical infections and thoracic malignancies can manifest as a cavitary pulmonary lesion. Standard bronchoscopy is commonly used to evaluate these abnormalities but is limited in its ability to fully assess for cancer and infection. Bronchoalveolar lavage samples are likely to aid in the diagnosis of infection but are less useful in the evaluation of malignancy. In addition, many of these pulmonary lesions are located in the periphery of the lung and are not accessible for tissue sampling by standard bronchoscopy. We present a unique presentation of *Pneumocystis jirovecii* pneumonia and discuss the utility of electromagnetic navigational bronchoscopy in the evaluation of immunocompromised patients with peripheral cavitary lung lesion.

How to cite this article:

Houshmand F, Aly FZ, Bowling MR. A novel diagnostic approach for *Pneumocystis jirovecii* pneumonia using fine-needle aspiration, electromagnetic navigational bronchoscopy and rapid on-site evaluation. Ann Thorac Med 2019;14:285-287

How to cite this URL:

Houshmand F, Aly FZ, Bowling MR. A novel diagnostic approach for *Pneumocystis jirovecii* pneumonia using fine-needle aspiration, electromagnetic navigational bronchoscopy and rapid on-site evaluation. Ann Thorac Med [serial online] 2019 [cited 2022 Mar 28];14:285-287

Available from: <https://www.thoracicmedicine.org/text.asp?2019/14/4/285/267949>

Full Text

Cavitary pulmonary lesions can be seen in both infectious and malignant conditions in human immunodeficiency virus (HIV)-infected individuals. Most are the result of fungal infections. *Pneumocystis jirovecii* (PJP) makes up to 2% of these occurrences.[1]

In addition, lung cancer is more prevalent and portends a worse prognosis in this population compared to nonHIV-infected individuals.[2] Approximately 67% of these cancers will be squamous cell carcinoma, of which 80% will present as a

cavitating lung lesion.[3]

Standard bronchoscopy is the initial standard approach in the evaluation of these lesions. However, it is limited in its ability to diagnose malignancy with bronchoalveolar lavage (BAL) (diagnostic yield of 33%)[4] but is excellent in the evaluation of infections (diagnostic yield of 98%).[5] Standard bronchoscopy is frequently restricted in accessing these lesions for tissue sampling if they are located in the periphery of lung. These drawbacks often lead to additional invasive procedures and therefore increased risk of a complication. This ultimately may cause a delay in diagnosis and therapy.

Electromagnetic navigational bronchoscopy (ENB) is an image-guided approach that uses a three-dimensional-reconstructed computed tomography (CT) scan and an electromagnetic sensor locator to access peripheral lung lesions beyond the reach of standard bronchoscopy. Once the lesion is located, tissue can be collected.

We present a unique and unexpected finding in the evaluation of a peripheral cavitary lung lesion in an immunocompromised patient utilizing ENB-guided fine-needle aspiration (FNA).

Case Report

A 40-year-old male with a history of smoking and untreated HIV infection presented to the hospital with concerns for a 10 lb weight loss, night sweats, and a nonproductive cough for 2 months. His initial laboratory studies demonstrated a white blood cell count of 2.6 K/ μ l, lymphocytes of 0.24 K/ μ L, and CD4 count of 0. A chest X-ray demonstrated a thick-walled cavitary lesion in the right upper lung, which was confirmed on a CT scan of the chest [Figure 1]. The right upper-lobe cavitary lesion was accessed by ENB, and several (seven FNA and seven forceps biopsy) samples were collected. Rapid on-site evaluation (ROSE) was available for immediate microscopic assessment of the tissue, and a preliminary diagnosis was rendered as concerning for the presence of *P. jirovecii*. This diagnosis was confirmed with the use of Grocott–Gomori's methenamine silver stain [Figure 2]. A bronchoalveolar sample was also collected from the right upper lobe, and *P. jirovecii* organisms were detected on cytologic evaluation of the BAL fluid. There was no malignancy or other infectious organism detected from the tissue of BAL samples.{Figure 1}{Figure 2}

Discussion

To our knowledge, this is the first reported case of an adult with the use of ENB-guided FNA of a lung lesion and immediate diagnosis of PJP with the use of ROSE. This case is noteworthy for several reasons; first, although cystic lung changes are seen with PJP, cavitary lung lesions are a rare finding in these infections. This adds to the uniqueness of the presented case in that other diagnoses such as malignancy and fungal or mycobacterial infections would have been considered a more likely etiology in immunocompromised patients with a solitary cavitary lung lesion.

Second, the diagnosis of PJP by FNA is rare. There are two previous reports of a diagnostic FNA of thyroid masses, which proved to be an extrapulmonary *P. jirovecii* infection.[6],[7] In addition, mediastinal lymph node sampling by the use of endobronchial ultrasound FNA has been reported to find PJP in both a HIV/AIDS patient [8] and a nonAIDS immunocompromised patient with a renal transplant.[9] Interestingly, Moualla and Saeed reported a simultaneous BAL to be nondiagnostic.[9] The choice of FNA with fiber-optic bronchoscopy in an immunocompromised pediatric population with diffuse pulmonary lesions and high clinical suspicion for PJP was reported by Choi et al., which showed a high yield and conveyed to be a safe procedure.[10]

Finally, ENB-guided FNA with ROSE is commonly used in the evaluation of thoracic malignancies, but there is no established use for it in the setting of the infectious disease workup. While bronchoscopy with BAL is a reasonable method for evaluating a cavitary lung lesion, a lavage sample would be unlikely to render a diagnosis of malignancy. In addition, many of these cavitary lesions may be located in the periphery of the lung and are difficult to access with conventional bronchoscopy and sampling via a transthoracic needle approach with radiologic guidance can lead to a pneumothorax (complication rates from 17% to 26%).[11],[12],[13] These methods may result in delays in treatment, the need for additional invasive procedures, and procedural complications. In our case, due to the presentation of a single cavitary lung lesion, with a history of heavy tobacco abuse, this patient could just as likely have had a malignancy as an infection; therefore, we felt that a simple BAL or sputum sample may not assess the possibility of malignancy as thoroughly as a

biopsy and BAL.

The use of ENB-guided biopsies in combination with ROSE may be an efficient approach to the management of these patients because both infectious and malignant etiologies can be evaluated safely. The complication rate of ENB-guided biopsies has been reported to be approximately 4%. [14] ENB both with and without ROSE has a diagnostic yield for malignancy that ranges from 38% to 97%. [14] In this case, we utilized the Super D™ (Medtronic, Minneapolis, MN, USA) ENB system because this is the navigational system we use at our institution. However, other image-guided modalities such as virtual bronchoscopy and radial ultrasound could also be employed.

Conclusion

The use of ENB-guided biopsies with ROSE in the assessment of cavitory lung lesions in immunocompromised patients is efficient and safe. Collaboration between an astute pathology team and chest physician is essential to maximize the utility of this procedural technique. Immediate feedback and direction from the pathologists concerning the need for additional tissue for culture is vital in maintaining the utility of the procedure and potentially avoiding unnecessary additional sampling or repeated invasive diagnostic studies.

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

References

- 1 Lin CY, Sun HY, Chen MY, Hsieh SM, Sheng WH, Lo YC, *et al.* Aetiology of cavitory lung lesions in patients with HIV infection. *HIV Med* 2009;10:191-8.
- 2 Pakkala S, Ramalingam SS. Lung cancer in HIV-positive patients. *J Thorac Oncol* 2010;5:1864-71.
- 3 Vourtsi A, Gouliamos A, Mouloupoulos L, Papacharalampous X, Chatjiioannou A, Kehagias D, *et al.* CT appearance of solitary and multiple cystic and cavitory lung lesions. *Eur Radiol* 2001;11:612-22.
- 4 Semenzato G, Spatafora M, Feruglio C, Pace E, Dipietro V. Bronchoalveolar lavage and the immunology of lung cancer. *Lung* 1990;168 Suppl: 1041-9.
- 5 Levine SJ, Kennedy D, Shelhamer JH, Kovacs A, Feuerstein IM, Gill VJ, *et al.* Diagnosis of *Pneumocystis carinii* pneumonia by multiple lobe, site-directed bronchoalveolar lavage with immunofluorescent monoclonal antibody staining in human immunodeficiency virus-infected patients receiving aerosolized pentamidine chemoprophylaxis. *Am Rev Respir Dis* 1992;146:838-43.
- 6 Battan R, Mariuz P, Raviglione MC, Sabatini MT, Mullen MP, Poretsky L. *Pneumocystis carinii* infection of the thyroid in a hypothyroid patient with AIDS: Diagnosis by fine needle aspiration biopsy. *J Clin Endocrinol Metab* 1991;72:724-6.
- 7 Keyhani-Rofagha S, Piquero C. *Pneumocystis carinii* thyroiditis diagnosis by fine needle aspiration cytology: A case report. *Acta Cytol* 1996;40:307-10.
- 8 Ellison E, Yuen SY, Lawson L, Chan NH. Fine-needle aspiration diagnosis of extrapulmonary *Pneumocystis carinii* lymphadenitis in a human immunodeficiency virus positive patient. *Diagn Cytopathol* 1995;12:251-3.
- 9 Moualla M, Saeed A. Endobronchial ultrasound with fine needle aspiration biopsy: A novel approach in diagnosing *Pneumocystis jirovecii* in mediastinal lymph node. *Chest Conf Case Rep* 2014;146:169A.
- 10 Choi YW, Kim SK, Jeon SC, Hahm CK, Choi CS. Fine needle aspiration biopsy of the lung in children with diffuse pulmonary lesions suggesting *Pneumocystis carinii* pneumonia *J Korean Radiol Soc* 1994;30:1147-50.
- 11 Yeow KM, Tsay PK, Cheung YC, Lui KW, Pan KT, Chou AS. Factors affecting diagnostic accuracy of CT-guided coaxial cutting needle lung biopsy: Retrospective analysis of 631 procedures. *J Vasc Interv Radiol* 2003;14:581-8.
- 12 Khan MF, Straub R, Moghaddam SR, Maataoui A, Gurung J, Wagner TO, *et al.* Variables affecting the risk of pneumothorax and intrapulmonary hemorrhage in CT-guided transthoracic biopsy. *Eur Radiol* 2008;18:1356-63.
- 13 Covey AM, Gandhi R, Brody LA, Getrajdman G, Thaler HT, Brown KT. Factors associated with pneumothorax and

pneumothorax requiring treatment after percutaneous lung biopsy in 443 consecutive patients. *J Vasc Interv Radiol* 2004;15:479-83.

- 14 Folch EE, Bowling MR, Gildea TR, Hood KL, Murgu SD, Toloza EM, *et al.* Design of a prospective, multicenter, global, cohort study of electromagnetic navigation bronchoscopy. *BMC Pulm Med* 2016;16:60.

Monday, March 28, 2022

[Site Map](#) | [Home](#) | [Contact Us](#) | [Feedback](#) | [Copyright and Disclaimer](#)