



Current Report

EXTENSION

Oklahoma Cooperative Extension Fact Sheets are also available on our website at: extension.okstate.edu

Management of Insect and Mite Pests in Sorghum

Tom A. Royer
Extension Entomologist

Sorghum pests, if not controlled when thresholds are exceeded, will reduce yield and quality of grain and forage. Pesticides should not be used as a substitute for good agronomic practices or as “preventative insurance” because it is rarely economically or environmentally justifiable. Many sorghum pest problems can be avoided by implementing an Integrated Pest Management (IPM) plan that includes preventive pest management practices, such as planting high-quality, vigorous, Oklahoma-proven hybrid seed; planting it at the proper time for optimal health and yield, providing proper fertilization and weed control; and, when possible, keeping sorghum fields as far away as possible from wheat.

The information herein is for educational purposes only. Reference to commercial products or trade names is made with the understanding that no discrimination is intended and no endorsement by the Cooperative Extension Service is implied.

Pesticide recommendations in this publication were correct as of the “Modified Date” but always check the label that came with the purchased insecticide for the most current rates and restrictions

The first name listed is the trade name of a product registered for use in corn for the listed pest. The name in

(parentheses) listed below the trade name is the name of the active ingredient. The active ingredient name is provided because in many cases, there are other registered products containing the same active ingredient that may cost less, so producers should compare prices.

The number [in brackets] following a product is its Mode of Action number [MOA]. The more frequently insecticides with the same MOA are used, the more likely resistance will occur. This number provides an easy way to select different modes of action to avoid selecting for pests that are resistant to a certain mode of action.

Refer to the following OSU publications for additional information.

- CR-2162 Grain Sorghum Performance Trials in Oklahoma, 2015
- EPP-7157 Field Key to Larvae in Sorghums
- EPP-7196 Grasshopper Management in Rangeland, Pastures, and Crops
- PSS-2113 Grain Sorghum Production Calendar
- PSS-2166 Use of Glyphosate as a Harvest Aid in Early Planted Grain Sorghum
- PT-2005-2010 Grain Sorghum Performance Trials in Oklahoma.

Management of Insect and Mite Pests in Sorghum

<i>Pest, Damage and Treatment Threshold</i>	<i>Insecticide, Formulation, [MOA Group] & (Active Ingredient)</i>	<i>Rate of Product (or AI) per Acre</i>	<i>Comments</i>
Chinch bug Adults are 1/8 inch long, black with white wings that are folded over the back into an “hour glass” shape. Nymphs are reddish to brown, with a white stripe across their “shoulders.” Damage: Feed at base of plants, in between leaf sheath and stem. Chinch bugs often migrate from small grains to sorghum. Feeding may kill small seedlings.	Planting Time		Seed treatments will generally provide 3 weeks of suppression. Use seed treatment if sorghum has suffered regular losses from chinch bug infestations. Do not feed leftover treated seed to livestock. Check table on last page for grazing and harvest restrictions for seed treatments. Best control is obtained when insecticide is applied by ground, with nozzles directed at the base of the plants using a minimum of 20 to 30 gallons of water.
	Gaucho 600 [4A] (imidacloprid)	6.4 fl oz/cwt seed (0.25 lb ai/cwt seed)	
	Cruiser 5FS [4A] (thiamethoxam)	5.1 to 7.6 fl oz/cwt seed (0.2 to 0.3 lb ai/A)	
	Poncho 600 [4A] (clothianidin)	5.1 to 6.4 fl oz/cwt seed (0.20 to 0.25 lb ai/A)	
	Post-Plant		
	Asana XL [3] (esfenvalerate)	5.8 to 9.6 fl oz (0.03 to 0.05 lb ai/A)	Do not apply more than 0.15 lb ai/season. 21 day wait for grazing or harvest.

<i>Pest, Damage and Treatment Threshold</i>	<i>Insecticide, Formulation, [MOA Group] & (Active Ingredient)</i>	<i>Rate of Product (or AI) per Acre</i>	<i>Comments</i>
Chinch bug (cont'd)			
<p>Threshold: Two to three bugs per plant on seedlings. Treat if large numbers are moving in to sorghum from grain. A border spray 30 to 60 feet wide on the margins of the field may be of value if chinch bug numbers are high in an adjacent wheat field.</p>	Baythroid XL [3] (beta-cyfluthrin)	2.0 to 2.8 fl oz (0.019 to 0.022 lb ai/A)	14-day wait for grazing or harvest.
	Cobalt [1B,3] (chlorpyrifos + gamma-cyhalothrin)	13 to 38 fl oz	30-day wait for applications of 26 fl oz/A or less, 60 day wait for applications over 26 fl oz/Acre.
	Delta Gold [3] (deltamethrin)	1.3 to 1.9 fl oz (0.015 to 0.022 lb ai/A)	14-day wait for grazing or harvest.
	Fastac EC [3] (alpha-cyhalothrin)	3.2 to 3.9 fl oz (0.020 to 0.025 lb ai/A)	14-day PHI for harvest, 45 day wait for forage.
	Lorsban 4E [1B] (chlorpyrifos)	1 to 2 pt (0.5 to 1 lb ai/A)	30- to 60-day wait for grazing or harvest.
	Mustang MAXX EC [3] (zeta-cypermethrin)	3.2 to 4.0 fl oz (0.02 to 0.025 lb ai/A)	14-day wait for harvest, 45 days for grazing.
	Proaxis 0.5 CS [3] (gamma-cyhalothrin)	3.84 fl oz (0.015 lb ai/A)	30-day wait for grazing or harvest.
	Sevin XLR [1A] (carbaryl)	1 to 2 qt (1 to 2 lb ai/A)	Sevin may cause spidermite buildup. 21-day wait for forage, 14 days for harvest or grazing.
	Stallion [1B, 3] (chlorpyrifos + zeta-cypermethrin)	9.25 to 11.75 oz	30-day wait for harvest, 45 days for forage.
	Warrior II with Zeon [3] (lambda-cyhalothrin)	1.92 fl oz (0.03 lb ai/A)	30-day wait for harvest or grazing.
Corn earworm (Headworm)			
<p>tive on very small (1st Up to 1 inch. Color varies from green, to brown to yellow and pink.</p> <p>Damage: Feed in whorl and ripening seed in head. Yield loss from whorl feeding is negligible. Are capable of causing damage to seed in head until grain reaches soft dough stage.</p> <p>Threshold: Two or more larvae per head before hard dough. A dynamic threshold that is based on plant population and crop value and control costs can be determined by accessing the sorghum headworm calculator http://entopl.okstate.edu/shwwweb/index.htm</p>			<p>Check labels, some state that product is only effective on very small (1st and 2nd instars) caterpillars.</p> <p>Do not apply more than 0.15 lb ai/season. 21-day wait for grazing or harvest.</p> <p>1st and 2nd instar only; 14-day wait for grazing or harvest.</p> <p>30-day wait for harvest.</p> <p>14-day wait for grazing, 7 days for harvest.</p> <p>30-day wait for applications of 26 fl oz/A or less, 60-day wait for applications over 26 fl oz/Acre.</p>
	Asana XL [3] (esfenvalerate)	5.8 to 9.6 fl oz (0.03 to 0.05 lb ai/A)	
	Baythroid XL [3] (beta-cyfluthrin)	1.3 to 2.8 fl oz (0.010 to 0.022 lb ai/A)	
	Besiege [28,3] (chlorantraniliprole + lambda-cyhalothrin)	6.0 to 10.0 fl oz	
	Blackhawk [5] (spinosad)	1.5 to 3.3 fl oz (0.034 to 0.094 lb ai/A)	
	Cobalt [1B,3] (chlorpyrifos + gamma-cyhalothrin)	19 to 38 fl oz	
	Concero [5,3] (spinosad + gamma-cyhalothrin)	2 to 2.85 fl oz/Acre (64 to 45 acres per gallon)	30-day wait for harvest or grazing.
	Coragen [28] (chlorantraniliprole)	3.5 to 7.5 fl oz 0.045 to 0.098 lb ai/a	1-day wait for harvest or grazing.
	Delta Gold [3] (deltamethrin)	1.0 to 1.5 fl oz (0.012 to 0.018 lb ai/A)	14-day wait for grazing or harvest.
	Diamond 0.8 EC [15] (novaluron)	9 to 12 fl oz	7-day wait for grazing, 14 days for grain; reapplication may be needed.
	Fastac EC [3] (alpha-cypermethrin)	1.8 to 3.8 fl oz (0.012 to 0.025 lb ai/A)	14-day PHI for harvest, 45-day wait for forage.

<i>Pest, Damage and Treatment Threshold</i>	<i>Insecticide, Formulation, [MOA Group] & (Active Ingredient)</i>	<i>Rate of Product (or AI) per Acre</i>	<i>Comments</i>	
Corn earworm (Headworm) (cont'd)	Lorsban 4E [1B] (chlorpyrifos)	2 pt (1 lb ai/A)	30- to 60-day wait for grazing or harvest.	
	Lannate LV [1A] (methomyl)	0.75 to 1.5 pt (0.225 to 0.45 lb ai/A)	14-day wait for grazing or harvest.	
	Mustang MAXX EC [3] (zeta-cypermethrin)	1.76 to 4.0 fl oz (0.011 to 0.025 lb ai/A)	14-day wait for harvest, 45 days for grazing.	
	Proaxis 0.5 CS [3] (gamma-cyhalothrin)	2.56 to 3.84 fl oz (0.01 to 0.015 lb ai/A)	30-day wait for grazing or harvest.	
	Sevin XLR [1A] (carbaryl)	1 to 2 qt (1 to 2 lb ai/A)	No wait for grazing, 21 days for harvest.	
	Stallion [1B,3] (chlorpyrifos + zeta-cypermethrin)	9.25 to 11.75 oz	30-day wait for harvest, 45 days for forage.	
	Warrior II with Zeon [3] (lambda-cyhalothrin)	1.28 to 1.92 fl oz (0.02 to 0.03 lb ai/A)	30-day wait for harvest or grazing.	
Corn leaf aphid Bluish-green, soft bodied aphid with black legs, antennae and cornicles. Typically found in whorl. Damage: Feed in whorl and may cause some delay of whorl emergence if numbers are high. Can mechanically transmit Maize Dwarf Mosaic virus disease. Threshold: Corn leaf aphids rarely cause significant yield loss, so no thresholds have been established.	Planting Time		Do not feed leftover treated seed to livestock. Check table on last page for grazing and harvest restrictions for seed treatments. Research indicates that yield losses occur only where corn leaf aphids cause stand loss on seedling plants. Chemical treatments, including seed treatments, are not likely to reduce potential for infection by Maize Dwarf Mosaic Virus because it can be transmitted within 30 seconds after an aphid begins feeding. Texas research suggests that corn leaf aphids serve as a food source for lady beetles which can help prevent greenbug outbreaks.	
	Gaucha 600 [4A] (imidacloprid)	6.4 fl oz/cwt seed (0.25 lb ai/cwt seed)		
	Cruiser 5FS [4A] (thiamethoxam)	5.1 to 7.6 fl oz/cwt seed (0.2 to 0.3 lb ai/A)		
	Poncho 600 [4A] (clothianidin)	5.1 to 6.4 fl oz/cwt seed (0.20 to 0.25 lb ai/A)		
	Post-Plant			
	Cobalt [1B,3] (chlorpyrifos + gamma-cyhalothrin)	7 to 13 fl oz		30-day wait for applications of 26 fl oz/A or less, 60-day wait for applications over 26 fl oz/Acre.
	Dimethoate 4E [1B] (dimethoate)	0.5 to 1 pt (0.25 to 0.5 lb ai/A)		28-day PHI.
	Fastac EC [3] (alpha-cypermethrin)	3.2 to 3.8 fl oz (0.02 to 0.025 lb ai/A)		14-day PHI for harvest, 45 PHI for grazing or forage.
	Lorsban 4E [1B] (chlorpyrifos)	0.5 to 1 pt (0.25 to 0.5 lb ai/acre)		30-day wait for grazing or harvest.
	Sivanto 200 SL [4D] (flupyradifurone)	7.0 to 10.5 fl oz (0.09 to 0.137 lb ai/A)		7-day wait for grazing, 21 days for harvest.
Stallion [1B,3] (chlorpyrifos + zeta-cypermethrin)	9.25 to 11.75 oz	30-day wait for harvest, 45 days for forage.		
Cutworms Robust caterpillars that "roll" up when disturbed, and prefer to live under ground. Damage: Cutworms generally feed at night, and live under the soil during the day. Plants will be cut at or slightly above the soil level. Threshold: Scout fields at seedling emergence. Treat when worms are less than ½ inch long, and skips are noticed.	Asana XL [3] (esfenvalerate)	5.8 to 9.6 fl oz (0.03 to 0.05 lb ai/A)	Do not apply more than 0.15 lb ai/season. 21-day wait for grazing or harvest.	
	Baythroid XL [3] (beta-cyfluthrin)	1.0 to 1.3 fl oz (0.008 to 0.010 lb ai/A)	14-day wait for grazing or harvest.	
	Besiege [28,3] (chlorantraniliprole + lambda-cyhalothrin)	5.0 to 6.0 fl oz	30-day wait for harvest.	
	Cobalt [1B,3] (chlorpyrifos + gamma-cyhalothrin)	13 to 38 fl oz	30-day wait for applications of 26 fl oz/A or less, 60-day wait for applications over 26 fl oz/Acre.	
	Delta Gold [3] (deltamethrin)	1.0 to 1.5 fl oz (0.012 to 0.018 lb ai/A)	14-day wait for grazing or harvest.	

<i>Pest, Damage and Treatment Threshold</i>	<i>Insecticide, Formulation, [MOA Group] & (Active Ingredient)</i>	<i>Rate of Product (or AI) per Acre</i>	<i>Comments</i>
Cutworms (cont'd)	Diamond 0.8 EC [15] (novaluron)	9 to 12 fl oz	7-day wait for forage, 14 days for grain, reapplication may be needed.
	Fastac EC [3] (alpha-cypermethrin)	1.3 to 3.8 fl oz (0.008 to 0.025 lb ai/A)	14-day PHI for harvest, 45 PHI for grazing or forage.
	Lorsban 4E [1B] (chlorpyrifos)	1 to 2 pt (0.5 to 1 lb ai/A)	30- to 60-day wait for grazing or harvest.
	Mustang MAXX EC [3] (zeta-cypermethrin)	1.28 to 4.0 fl oz (0.008 to 0.025 lb ai/A)	14-day wait for harvest, 45 days for grazing.
	Proaxis 0.5 CS [3] (gamma-cyhalothrin)	1.92 to 2.56 fl oz (0.0075 to 0.01 lb ai/A)	30-day wait for grazing or harvest.
	Stallion [1B, 3] (chlorpyrifos + zeta-cypermethrin)	3.75 to 11.75 oz	30-day wait for harvest, 45 days for forage.
	Warrior II with Zeon [3] (lambda-cyhalothrin)	0.96 to 1.28 fl oz (0.015 to 0.02 lb ai/A)	30-day wait for harvest or grazing.
Fall armyworm (Headworm) Large, striped, non-bristled caterpillar up to 1.5 inches. Has a light-colored inverted "Y" on head. Damage: Feed in whorl, and ripening seed in head. Yield loss from whorl feeding is negligible. Can damage seed in head until grain reaches soft dough stage. Threshold: Two or more larvae per head before hard dough. Open-headed varieties are less susceptible to attack than tight-headed varieties. A dynamic threshold that is based on plant population and crop value and control costs can be determined by accessing the sorghum headworm calculator entopl.okstate.edu/shwwweb/index.htm	Baythroid XL [3] (beta-cyfluthrin)	1.3 to 2.8 fl oz (0.010 to 0.022 lb ai/A)	Check labels, some state that product is only effective on very small (1st and 2nd instars) caterpillars. 1st and 2nd instar only; 14 day wait for grazing or harvest.
Blackhawk [5] (spinosad)	1.5 to 3.3 fl oz (0.034 to 0.094 lb ai/A)	14-day wait for grazing, 7 days for harvest.	
Besiege [28,3] (chlorantraniliprole + lambda-cyhalothrin)	6.0 to 10.0 fl oz	30-day wait for harvest.	
Cobalt [1B,3] (chlorpyrifos + gamma-cyhalothrin)	13 to 38 fl oz	30-day wait for applications of 26 fl oz/A or less, 60 day wait for applications over 26 fl oz/Acre.	
Concero [5,3] (spinosad + gamma-cyhalothrin)	2.85 fl oz/Acre (45 acres per gallon)	30-day wait for harvest or grazing.	
Coragen [28] (chlorantraniliprole)	3.5 to 7.5 fl oz (0.045-0.098 lb ai/A)	1-day wait for harvest or grazing.	
Delta Gold [3] (deltamethrin)	1.3 to 1.9 fl oz (0.015 to 0.022 lb ai/A)	14-day wait for grazing or harvest.	
Diamond 0.8 EC [15] (novaluron)	9 to 12 fl oz	7-day wait for grazing, 14 days for grain reapplication may be needed.	
Fastac EC [3] (alpha-cypermethrin)	1.8 to 3.8 fl oz 0.012 to 0.025 lb ai/A	14-day PHI for harvest, 45 days for grazing or forage.	
Intrepid 2F [18] (methoxyfenozide)	8 to 10 fl oz (0.12 to 0.16 lb ai/A)	21-day PIH for grain or stover harvest, 3 days for forage.	
Lorsban 4E [1B] (chlorpyrifos)	1 to 2 pt (0.5 to 1 lb ai/A)	30- to 60-day wait for grazing or harvest.	
Lannate LV[1A] (methomyl)	0.75 to 1.5 pt (0.225 to 0.45 lb ai/A)	14-day wait for grazing or harvest.	
Mustang MAXX EC [3] (zeta-cypermethrin)	1.76 to 4.0 fl oz (0.011 to 0.025 lb ai/A)	14-day wait for harvest, 45 days for grazing.	
Proaxisr 0.5 CS [3] (gamma-cyhalothrin)	2.56 to 3.84 fl oz (0.01 to 0.015 lb ai/A)	30-day wait for grazing or harvest.	

<i>Pest, Damage and Treatment Threshold</i>	<i>Insecticide, Formulation, [MOA Group] & (Active Ingredient)</i>	<i>Rate of Product (or AI) per Acre</i>	<i>Comments</i>
Fall armyworm (Headworm) (cont'd)	Sevin XLR [1A] (carbaryl)	1 to 2 qt (1 to 2 lb ai/A)	No wait for grazing, 21 days for harvest.
	Stallion [1B,3] (chlorpyrifos + zeta-cypermethrin)	9.25 to 11.75 oz	30-day wait for harvest, 45 days for forage.
	Warrior II with Zeon [3] (lambda-cyhalothrin)	1.28 to 1.92 fl oz (0.02 to 0.03 lb ai/A)	30-day wait for harvest or grazing.
False chinch bug Adults 1/8 inch long, dirty gray, with brown or black markings and piercing may be needed. mouthparts. Damage: Feed in groups. age. Large numbers may cause wilting of heads or small plants. Threshold: 140 or more per head.	Baythroid XL [3] (cyfluthrin)	1.3 to 2.8 fl oz (0.010 to 0.022 lb ai/A)	14-day wait for grazing or harvest.
	Diamond 0.8 EC [15] (novaluron)	9 to 12 fl oz	7-day wait for grazing, 14 days for grain reapplication
	Fastac EC [3] (alpha-cyhalothrin)	3.2 to 3.9 fl oz (0.02 to 0.025 lb ai/A)	14-day PHI for harvest, 45 days for grazing or forage.
	Mustang MAXX EC [3] (zeta-cypermethrin)	3.2 to 4.0 fl oz (0.02 to 0.025 lb ai/A)	14-day wait for harvest, 45 days for grazing.
	Stallion [1B,3] (chlorpyrifos + zeta-cypermethrin)	9.25 to 11.75 fl oz	30-day wait for harvest, 45 days for forage.
Grasshopper 1-2 inches, outer wings leathery, inner wings clear or colored. Enlarged hind legs designed for jumping. Damage: Chew leaves, leaving ragged edges or completely chew leaf blade. Damage emerging seed heads causing yield loss. Threshold: 15 to 20 per square yard. If nymph populations exceed threshold field borders age. (25-40 per square yard), treat before they move into sorghum. These products are for application in sorghum; See EPP-7196: Grasshopper Management in Rangeland, Pastures and Crops for treating non-crop areas.	Baythroid XL [3] (beta-cyfluthrin)	2 to 2.8 fl oz (0.019 to 0.022 lb ai/A)	14-day wait for grazing or harvest.
	Cobalt [1B,3] (chlorpyrifos + gamma-cyhalothrin)	7 to 13 fl oz	30-day wait for applications of 26 fl oz/A or less, 60 day wait for applications over 26 fl oz/Acre.
	Coragen [28] (chlorantraniliprole)	2.0 to 5.0 fl oz (0.026 to 0.065 lb ai/A)	1-day wait for harvest or grazing.
	Delta Gold [3] (deltamethrin)	1.0 to 1.5 fl oz (0.012 to 0.018 lb ai/A)	14-day wait for grazing or harvest.
	Dimethoate 4E [1B] (dimethoate)	1 pt (0.5 lb ai/A)	Only one post-plant application per season.
	Fastac EC [3] (alpha-cyhalothrin)	3.2 to 3.9 fl oz (0.02 to 0.025 lb ai/A)	14-day PHI for harvest, 45 days for grazing or forage.
	Lorsban 4E [1B] (chlorpyrifos)	0.5 to 1 pt (0.25 to 0.5 lb ai/A)	30-day wait for grazing or harvest.
	Mustang MAXX EC [3] (zeta-cypermethrin)	3.2 to 4.0 fl oz (0.02 to 0.025 lb ai/A)	14-day wait for harvest, 45 days for grazing.
	Proaxis 0.5 CS [3] (gamma-cyhalothrin)	2.56 to 3.84 fl oz (0.01 to 0.015 lb ai/A)	30-day wait for grazing or harvest.
	Stallion [1B,3] (chlorpyrifos + zeta-cypermethrin)	9.25 to 11.75 oz	30-day wait for harvest, 45 days for forage.
Warrior II with Zeon [3] (lambda-cyhalothrin)	1.28 to 1.92 fl oz (0.02 to 0.03 lb ai/A)	30 day wait for harvest or grazing.	

<i>Pest, Damage and Treatment Threshold</i>	<i>Insecticide, Formulation, [MOA Group] & (Active Ingredient)</i>	<i>Rate of Product (or AI) per Acre</i>	<i>Comments</i>	
<p>Greenbug Lime-green, soft bodied aphid with darker green stripe down back. Tips of legs, cornicles, and most of antennae are black.</p> <p>Damage: Injury can occur anytime from seedling emergence through soft dough stage. Greenbug feeding causes reddening of leaves which die as populations increase.</p> <p>Threshold: See Thresholds listed at end of publication. Need to treat is dependent upon greenbug numbers, plant size, variety, growing conditions, and the presence of predators and parasites. It is better to base treatment decision on presence of plant damage than on greenbug numbers alone.</p>	Seed Treatment		Do not feed leftover treated seed to livestock. Check table on last page for grazing and harvest restrictions for seed treatments.	
	Attendant 600 [4A] (imidacloprid)	6.4 fl oz/cwt seed (0.25 lb ai/cwt seed)		
	Cruiser 5FS [4A] (thiamethoxam)	5.1 to 7.6 fl oz/cwt seed (0.2 to 0.3 lb ai/A)		
		Poncho 600 [4A] (clothianidin)	5.1 to 6.4 fl oz/cwt seed (0.20 to 0.25 lb ai/A)	
	Planting Time			
		Counter 15G [1B]	"Lock 'n Load" or "Smartbox" applicator needed	Do not place granules in contact with seed. 50-day wait for grazing, 100 days for harvest.
	Post-Plant			
		Dimethoate 4E [1B] (dimethoate)	0.5 to 1 pt (0.25 to 0.5 lb ai/A)	28-day wait for harvest or grazing.
		Cobalt [1B,3] (chlorpyrifos + gamma-cyhalothrin)	13 to 38 fl oz	30-day wait for applications of 26 fl oz/A or less, 60-day wait for applications over 26 fl oz/Acre. See additional instructions on label.
		Fastac EC [3] (alpha-cyhalothrin)	3.2 to 3.9 fl oz (0.02 to 0.025 lb ai/A)	14-day PHI for harvest, 45 days for grazing or forage.
		Lorsban 4E [1B] (chlorpyrifos)	0.5 to 2 pt (0.25 to 1 lb ai/A)	30- to 60-day wait for grazing or harvest.
		Malathion 5E [1B] (malathion)	1.5 pt (0.93 lb ai/A)	7-day PHI for grain. Do not feed or graze forage, hay or straw to livestock.
		Sivanto 200 SL [4D] (flupyradifurone)	7.0 to 10.5 fl oz (0.09 to 0.137 lb ai/A)	7-day wait for forage, 21 days for harvest
		Stallion [1B,3] (chlorpyrifos + zeta-cypermethrin)	9.25 to 11.75 oz	30-day wait for harvest, 45 days for forage.
<p>Lesser cornstalk borer Caterpillar ¾ inch long when mature. Slender, blue-green with brown bands around each body segment. Make silken tunnels at feeding site.</p> <p>Damage Tunnels in roots and stems. Occurs in May through June.</p> <p>Threshold Treat before larva bore into stalk.</p>	Concero [5,3] (spinosad + gamma-cyhalothrin)	2 to 2.85 fl oz/Acre (64 to 45 acres per gallon)	30-day wait for harvest or grazing.	
		Delta Gold [3] (deltamethrin)	1.3 to 1.9 fl oz (0.015 to 0.022 lb ai/A)	14-day wait for grazing or harvest.
		Fastac EC [3] (alpha-cyhalothrin)	3.2 to 3.9 fl oz (0.02 to 0.025 lb ai/A)	14-day PHI for harvest, 45 days for grazing or forage.
		Lorsban 4E [1B] (chlorpyrifos)	1 to 2 pt (0.5 to 1 lb ai/A)	30- to 60-day wait for grazing or harvest.
		Mustang MAXX EC [3] (zeta-cypermethrin)	3.2 to 4.0 fl oz (0.02 to 0.025 lb ai/A)	14-day wait for harvest, 45 days for grazing.
		Proaxis 0.5 CS [3] (gamma-cyhalothrin)	2.56 to 3.84 fl oz (0.01 to 0.015 lb ai/A)	30-day wait for grazing or harvest.
		Stallion [1B, 3] (chlorpyrifos + zeta-cypermethrin)	9.25 to 11.75 oz	30-day wait for harvest, 45 days for forage.
	Warrior II with Zeon [3] (lambda-cyhalothrin)	1.28 to 1.92 fl oz (0.02 to 0.03 lb ai/A)	30-day wait for harvest or grazing.	

<i>Pest, Damage and Treatment Threshold</i>	<i>Insecticide, Formulation, [MOA Group] & (Active Ingredient)</i>	<i>Rate of Product (or AI) per Acre</i>	<i>Comments</i>	
Panicle feeding bugs Include stink bugs and leaf-footed bugs. Stink bugs: shield shaped bugs ranging from ½ to ¾ inch long. Leaf-footed bug: Brown, oblong about ¾ inch long with each age. hindleg leaf-like. Damage: Feed on seed, causing blasted heads, shrunken damaged seed. Most damage occurs before seed reaches hard dough stage. Thresholds: Milk stage: 5 bugs /head. Soft Dough: 9 bugs/head.	Concero [5,3] (spinosad + gamma-cyhalothrin)	2 to 2.85 fl oz/Acre (64 to 45 acres per gallon)	30-day wait for harvest or grazing.	
	Delta Gold [3] (deltamethrin)	1.3 to 1.9 fl oz (0.015 to 0.022 lb ai/A)	14-day wait for grazing or harvest.	
	Fastac EC [3] (alpha-cyhalothrin)	3.2 to 3.9 fl oz (0.02 to 0.025 lb ai/A)	14-day PHI for harvest, 45 days for grazing or forage.	
	Lorsban 4E [1B] (chlorpyrifos)	1 to 2 pt (0.5 to 1 lb ai/A)	30- to 60-day wait for grazing or harvest.	
	Mustang MAXX EC [3] (zeta-cypermethrin)	3.2 to 4.0 fl oz (0.02 to 0.025 lb ai/A)	14-day wait for harvest, 45 days for grazing.	
	Proaxis 0.5 CS [3] (gamma-cyhalothrin)	2.56 to 3.84 fl oz (0.01 to 0.015 lb ai/A)	30-day wait for grazing or harvest.	
	Stallion [1B, 3] (chlorpyrifos +zeta-cypermethrin)	9.25 to 11.75 oz	30-day wait for harvest, 45 days for forage.	
	Warrior II with Zeon [3] (lambda-cyhalothrin)	1.28 to 1.92 fl oz (0.02 to 0.03 lb ai/A)	30-day wait for harvest or grazing.	
	Sorghum midge Tiny, fragile orange-bodied fly that is active in early to mid-morning. Damage: Damaged heads appear to be "blasted" or "blighted" from high temperatures, infertility, or drought. Damage from sorghum midge generally restricted to sorghum that blooms after August 15. Threshold: Check fields before 11 am, when flies are most active Treat when 25-30% have begun bloom and adults average one or more per head.	Asana XL [3] (esfenvalerate)	2.9 to 5.8 fl oz (0.015 to 0.03 lb ai/A)	Check labels. May need to apply a second treatment 3-5 days after first. Uniform planting date is an option for management. Do not apply more than 0.15 lb ai/season. 21-day wait for grazing or harvest.
		Baythroid XL [3] (beta-cyfluthrin)	1.0 to 1.3 fl oz (0.008 to 0.010 lb ai/A)	14-day wait for grazing or harvest.
Blackhawk [5] (spinosad)		1.5 to 3.3 fl oz (0.034 to 0.094 lb ai/A)	14-day wait for grazing, 7 days for harvest.	
Cobalt [1B,3] (chlorpyrifos + gamma-cyhalothrin)		7 to 13 fl oz	30-day wait for applications of 26 fl oz/A or less, 60-day wait for applications over 26 fl oz/Acre.	
Delta Gold [3] (deltamethrin)		1.3 to 1.9 fl oz (0.015 to 0.022 lb ai/A)	14-day wait for grazing or harvest.	
Diamond 0.8 EC [15] (novaluron)		9 to 12 fl oz	7-day wait for grazing, 14 days for grain of heads reapplication may be needed.	
Fastac EC [3] (alpha-cypermethrin)		1.3 to 3.8 fl oz (0.008 to 0.025 lb ai/A)	14-day wait for harvest, 45 days for grazing or forage.	
Lorsban 4E [1B] (chlorpyrifos)		0.5 pt (0.25 lb ai/A)	30-day wait for grazing or harvest.	
Lannate LV [1A] (methomyl)		0.75 to 1.5 pt (0.225 to 0.45 lb ai/A)	14-day wait for grazing or harvest.	
Mustang MAXX [3] (zeta-cypermethrin)		1.28 to 4.0 fl oz (0.008 to 0.025 lb ai/A)	14-day wait for harvest, 45 days for grazing.	
Proaxis 0.5 CS [3] (gamma-cyhalothrin)		1.92 to 2.56 fl oz (0.0075 to 0.01 lb ai/A)	30-day wait for grazing or harvest.	
Stallion [1B, 3] (chlorpyrifos +zeta-cypermethrin)		3.75 to 11.75 oz	30-day wait for harvest, 45 days for forage.	
Warrior II with Zeon [3] (lambda-cyhalothrin)	0.96 to 1.28 fl oz (0.015 to 0.02 lb ai/A)	30-day wait for harvest or grazing.		

<i>Pest, Damage and Treatment Threshold</i>	<i>Insecticide, Formulation, [MOA Group] & (Active Ingredient)</i>	<i>Rate of Product (or AI) per Acre</i>	<i>Comments</i>
<p>Sorghum webworm Fuzzy, reddish to brown worms in head.</p> <p>Damage: Caterpillars feed on the seed, and hollow it out. Open-headed varieties are less susceptible than tight-headed varieties to attack.</p> <p>Threshold: 5 or more larvae per head before hard dough stage.</p>	Baythroid XL [3] (beta-cyfluthrin)	1.3 to 2.8 fl oz (0.010 to 0.022 lb ai/A)	14-day wait for grazing or harvest.
	Besiege [28,3] (chlorantraniliprole + lambda cyhalothrin)	6.0 to 10.0 fl oz	30-day wait for harvest.
	Blackhawk [5] (spinosad)	1.5 to 3.3 fl oz (0.034 to 0.094 lb ai/A)	14-day wait for grazing, 7 days for harvest.
	Cobalt [1B,3] (chlorpyrifos + gamma-cyhalothrin)	19 to 38 fl oz	30-day wait for applications of 26 fl oz/A or less, 60 day wait for applications over 26 fl oz/Acre.
	Coragen [28] (chlorantraniliprole)	3.5 to 7.5 fl oz (0.045 to 0.098 lb ai/A)	1-day wait for harvest or grazing.
	Delta Gold [3] (deltamethrin)	1 to 1.5 fl oz (0.012 to 0.018 lb ai/A)	14-day wait for grazing or harvest.
	Diamond 0.8 EC [15] (novaluron)	9 to 12 fl oz	7-day wait for grazing, 14 days for grain reapplication may be needed.
	Fastac EC [3] (alpha-cypermethrin)	1.8 to 3.8 fl oz (0.012 to 0.025 lb ai/A)	14-day PHI for harvest, 45 day for grazing or forage.
	Lorsban 4E [1B] (chlorpyrifos)	1 pt (0.5 lb ai/A)	30-day wait for grazing or harvest.
	Mustang MAXX EC [3] (zeta-cypermethrin)	1.76 to 4.0 fl oz (0.011 to 0.025 lb ai/A)	14-day wait for harvest, 45 days for grazing.
	Proaxis 0.5 CS [3] (gamma-cyhalothrin)	2.56 to 3.84 fl oz (0.01 to 0.015 lb ai/A)	30-day wait for grazing or harvest.
Stallion [1B, 3] (chlorpyrifos + zeta-cypermethrin)	5.0 to 11.75 oz	30-day wait for harvest, 45 days for forage.	
Warrior II with Zeon [3] (lambda-cyhalothrin)	1.28 to 1.92 fl oz (0.02 to 0.03 lb ai/a)	30-day wait for harvest or grazing.	
<p>Southwestern corn borer Full grown caterpillars are white with prominent dark spots on body.</p> <p>Damage: Tunnels throughout stalk. May girdle mature stalks.</p> <p>Threshold: Chemical control usually not warranted.</p>	Baythroid XL [3] (cyfluthrin)	1.3 to 2.8 fl oz (0.010 to 0.022 lb ai/A)	14-day wait for grazing or harvest.
	Besiege [28,3] (chlorantraniliprole + lambda cyhalothrin)	6.0 to 10.0 fl oz	30-day wait for harvest.
	Blackhawk [5] (spinosad)	1.5 to 3.3 fl oz (0.034 to 0.094 lb ai/A)	14-day wait for grazing, 7 days for harvest.
	Cobalt [1B,3] (chlorpyrifos + gamma-cyhalothrin)	19 to 38 fl oz	30-day wait for applications of 26 fl oz/A or less, 60 day wait for applications over 26 fl oz/Acre.
	Concero [5,3] (spinosad + gamma-cyhalothrin)	2 to 2.85 fl oz	30-day wait for harvest or grazing.
	Fastac EC[3] (alpha-cypermethrin)	1.8 to 3.8 fl oz (0.012 to 0.025 lb ai/A)	14-day PHI for harvest, 45 days for grazing or forage.
	Intrepid 2F [18] (methoxyfenozide)	8 to 10 fl oz (0.12 to 0.16 lb ai/A)	21-day PIH for grain or stover harvest, 3 days for forage.
	Lorsban 4E [1B] (chlorpyrifos)	1.5 to 2 pt (0.75 to 1 lb ai/A)	60-day wait for grazing or harvest.
	Mustang MAXX EC [3] (zeta-cypermethrin)	1.76 to 4.0 fl oz (0.011 to 0.025 lb ai/A)	14-day wait for harvest, 45 days for grazing.

<i>Pest, Damage and Treatment Threshold</i>	<i>Insecticide, Formulation, [MOA Group] & (Active Ingredient)</i>	<i>Rate of Product (or AI) per Acre</i>	<i>Comments</i>
Southwestern corn borer (cont'd)	Proaxis 0.5 CS [3] (gamma-cyhalothrin)	2.56 to 3.84 fl oz (0.01 to 0.015 lb ai/A)	30-day wait for grazing or harvest.
	Sevin XLR [1A] (carbaryl)	1.5 quarts (1.5 lb ai/A)	No wait for grazing, 21 days for harvest.
	Stallion [1B, 3] (chlorpyrifos + zeta-cypermethrin)	5.0 to 11.75 oz	30-day wait for harvest, 45 days for forage.
	Warrior II with Zeon [3] (lambda-cyhalothrin)	1.28 to 1.92 fl oz (0.02 to 0.03 lb ai/a)	30 day wait for harvest or grazing.
Spidermites			
	Post-Plant		
Small, less than 1/100 inch long. Causes brown stippling of leaves.	Comite II [14] (propargite)	24 to 36 fl oz (1.125 to 1.6875 lb ai/A)	30-day wait for grazing, 60 days for harvest.
Damage: Causes stippling of leaves; severe infestations can kill leaves.	Dimethoate 4E [1B] (dimethoate)	1 pt (0.5 lb ai/A)	Only one post-plant application per season.
	Onager [10A] (hexythiazox)	10 to 24 fl oz (0.078 to 0.1875 lb ai/A)	30-day waiting period for harvest, do not graze.
Threshold: No threshold established. Treat if majority of plants are infested with large, increasing mite infestations. Control is not be justified after head reaches hard dough stage	Supracide 2E [1B] (methidathion)	2 pt (0.5 lb ai/A)	30-day wait for grazing or harvest (24c label, OK050003).
Sugarcane aphid			
	Planting Time		
Whitish to light yellow, soft bodied aphid. Tips of legs, cornicles, and most of antennae are black. Colonies occur on underside of leaves, starting from the lower leaves.	Cruiser 5FS [4A] (thiamethoxam)	5.1 to 7.6 fl oz/cwt seed (0.2 to 0.3 lb ai/A)	Check table on last page for grazing and harvest restrictions for seed treatments.
	Post-Plant		
Damage: Injury can occur anytime from seedling emergence through harvest, but is more likely to occur from boot through soft dough. Heavy feeding causes early leaf senescence and reduces seed fill. Aphids produce large amounts of honeydew, which can affect harvest operations.	*Dimethoate 4E [1B] (dimethoate)	0.5 to 1 pint (0.25 to 0.5 lb ai/A)	*moderately effective, 28 day waiting period.
	*Lorsban 4E [1B] (chlorpyrifos)	0.5 to 2 pt (0.25 to 1 lb ai/A)	*moderately effective, 30 to 60-day wait for grazing or harvest.
	Sivanto Prime [4D] (flupyradifurone)	4.0 to 7.0 fl oz (0.05 to 0.09 lb ai/A)	7-day wait for grazing, 14 days for grain harvest or hay.
	Transform WD [4C] (sulfoxaflor)	0.75 to 1.5 oz (0.023 to 0.047 lb ai/A)	7-day waiting for grazing, 14 days for harvest. Do not spray less than 3 days before bloom, or until seed set. (Section 18 emergency use registration, expires 11/30/2017).
Threshold: Consult with the smartphone Glance-N-Go app for Sugarcane aphid at dasnr.okstate.edu/apps . It allows for calculation of a treatment threshold and can assist with scouting. If it is not available, thresholds are: 20% of plants are infested with 50 to 125 aphids per leaf before head emergence, and 30% of plants are infested with 50 to 125 aphids after head emergence.			

<i>Pest, Damage and Treatment Threshold</i>	<i>Insecticide, Formulation, [MOA Group] & (Active Ingredient)</i>	<i>Rate of Product (or AI) per Acre</i>	<i>Comments</i>
<p>White grub Large, "C" shaped grub with a white body and a brown head.</p> <p>Damage: Grubs feed on roots of seedling plants. Damage potential is dependent on planting date and speed of growth of the plant.</p> <p>Threshold: No treatment is available. An average of one grub per square foot may cause stand loss.</p>	NA	NA	No insecticide is currently registered for white grub control. Re-planting may be the best option.
<p>Wireworm Hard-shelled, smooth, cylindrical, yellowish to brown worms. Two- to six-year life cycle. More common in sorghum planted into a sod or grass pasture.</p> <p>Damage: Feed on seed, seedling. Cause stunting and stand loss.</p> <p>Threshold: Seed treatments are available. Treat if field history indicates a problem.</p>	<p>Seed Treatment</p> <p>Gaucho 600 [4A] (imidacloprid)</p> <p>Cruiser 5FS [4A] (thiamethoxam)</p> <p>Poncho 600 [4A] (clothianidin)</p> <p>Planting Time</p> <p>*Counter 15G [1B]</p>	<p>6.4 fl oz/cwt seed (0.25 lb ai/cwt seed)</p> <p>5.1 to 7.6 fl oz/cwt seed (0.2 to 0.3 lb ai/A)</p> <p>5.1 to 6.4 fl oz/cwt seed (0.20 to 0.25 lb ai/A)</p> <p>Apply per label.</p>	<p>Do not feed leftover treated seed to livestock. Check table on last page for grazing and harvest restrictions for seed treatments.</p> <p>* Counter 15 G can be used as a planting time treatment except in the Panhandle, but it requires a "Smartbox" or "Lock 'n Load" applicator, and has the potential to damage plants, and interact with several ALS-inhibiting herbicides. Check label for restrictions.</p>

Pre-harvest intervals and grazing restrictions

Asana XL	21-day PHI
Batallion/Delta Gold	14-day wait for grazing or harvest
Baythroidr XL	14-day PHI, 14 days grazing
Besiege	30-day PHI for harvest
Blackhawk	7-day PHI for harvest, 14 days for grazing
Cobalt	30-day wait for applications of 26 fl oz/A or less, 60-day wait for applications over 26 fl oz/Acre
Comite II	30-day PHI for silage, 60 days for grain harvest.
Concero	30-day PHI
Coragen	1-day PHI for harvest or grazing
Counterr 15G	100-day PHI for grain, 50 days for grazing
Cruiserr 5FS	no grazing restriction
Diamond 0.8 EC	7-day wait for grazing, 14 days for grain
Dimethoate	28-day PHI for grain or grazing, do not apply after heading.
Fastac	14-day PHI for harvest, 45-day PHI for forage/grazing
Lannate	14-day PHI for harvest or grazing
Lorsban 4E	30- to 60-day PHI for harvest or grazing, depending on rate applied.
Malathion	7-day PHI for grain. Do not feed or graze forage, hay or straw to livestock.
Mustang MAXX EC	14-day PHI for harvest, 45 days for grazing
Onager	30-day PHI for harvest, do not graze.
Poncho	no grazing restriction
Proaxis	30-Day PHI for harvest or grazing
Sevin XLR	21-day PHI for harvest, 0 days for forage.
Sivanto	14-day PHI for harvest, 7 days for forage
Stallion	30-day wait for harvest, 45 days for forage
Transform WD	14-day PHI for harvest, 7 days for forage
Warrior II with Zeon	30-day PHI for harvest or grazing

* MOA group numbers in brackets [#] following the insecticide name are used to designate the mode of action of the insecticide according to the classification system developed by the Insecticide Resistance Action Committee, (IRAC) in 2011. It is intended to help in the selection of insecticides for preventative resistance management. If you make multiple applications for a specific pest during a growing season, simply select a registered insecticide with a different number for each application. To further delay resistance from developing, integrate other control methods into your pest management programs.

Treatment Thresholds * For Greenbugs On Sorghum

<i>Plant Size When to Treat</i>	<i>Texas thresholds</i>	<i>Kansas thresholds</i>
0- to 1-leaf stage	20% of plants visibly damaged	25 to 50 greenbugs per plant
3-leaf stage	20% of plants visibly damaged	50 to 100 greenbugs per plant
Five-leaf stage	Visible damage on leaves, (red spots, yellow leaves) but before any entire leaves are killed on 20% of plants	
Mid-whorl stage	Visible damage on leaves (red spots, yellow leaves), but before any entire leaves are killed on 20% of plants	300 to 600 greenbugs per plant.
Boot to heading	Death of one functional leaf	700 to 1,000 greenbugs per plant
Heading through soft dough	Death of two functional leaves	700 to 1,000 greenbugs per plant

The pesticide information presented in this publication was current with federal and state regulations at the time of revision.
READ and FOLLOW all LABEL directions.

The Oklahoma Cooperative Extension Service

Bringing the University to You!

The Cooperative Extension Service is the largest, most successful informal educational organization in the world. It is a nationwide system funded and guided by a partnership of federal, state, and local governments that delivers information to help people help themselves through the land-grant university system.

Extension carries out programs in the broad categories of agriculture, natural resources and environment; family and consumer sciences; 4-H and other youth; and community resource development. Extension staff members live and work among the people they serve to help stimulate and educate Americans to plan ahead and cope with their problems.

Some characteristics of the Cooperative Extension system are:

- The federal, state, and local governments cooperatively share in its financial support and program direction.
- It is administered by the land-grant university as designated by the state legislature through an Extension director.
- Extension programs are nonpolitical, objective, and research-based information.
- It provides practical, problem-oriented education for people of all ages. It is designated to take the knowledge of the university to those persons who do not or cannot participate in the formal classroom instruction of the university.
- It utilizes research from university, government, and other sources to help people make their own decisions.
- More than a million volunteers help multiply the impact of the Extension professional staff.
- It dispenses no funds to the public.
- It is not a regulatory agency, but it does inform people of regulations and of their options in meeting them.
- Local programs are developed and carried out in full recognition of national problems and goals.
- The Extension staff educates people through personal contacts, meetings, demonstrations, and the mass media.
- Extension has the built-in flexibility to adjust its programs and subject matter to meet new needs. Activities shift from year to year as citizen groups and Extension workers close to the problems advise changes.

Oklahoma State University, as an equal opportunity employer, complies with all applicable federal and state laws regarding non-discrimination and affirmative action. Oklahoma State University is committed to a policy of equal opportunity for all individuals and does not discriminate based on race, religion, age, sex, color, national origin, marital status, sexual orientation, gender identity/ expression, disability, or veteran status with regard to employment, educational programs and activities, and/or admissions. For more information, visit <https://eeo.okstate.edu>.

Issued in furtherance of Cooperative Extension work, acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture, Director of Oklahoma Cooperative Extension Service, Oklahoma State University, Stillwater, Oklahoma. This publication is printed and issued by Oklahoma State University as authorized by the Vice President for Agricultural Programs and has been prepared and distributed at a cost of 40 cents per copy. Revised 12/2020 GH.