



UvA-DARE (Digital Academic Repository)

Decatungstate-Mediated C(sp³)-H Heteroarylation via Radical-Polar Crossover in Batch and Flow

Wan, T.; Capaldo, L.; Laudadio, G.; Nyuchev, A.V.; Rincón, J.A.; García-Losada, P.; Mateos, C.; Frederick, M.O.; Nuño, M.; Noël, T.

DOI

[10.1002/ange.202104682](https://doi.org/10.1002/ange.202104682)

[10.1002/anie.202104682](https://doi.org/10.1002/anie.202104682)

Publication date

2021

Document Version

Final published version

Published in

Angewandte Chemie

License

CC BY

[Link to publication](#)

Citation for published version (APA):

Wan, T., Capaldo, L., Laudadio, G., Nyuchev, A. V., Rincón, J. A., García-Losada, P., Mateos, C., Frederick, M. O., Nuño, M., & Noël, T. (2021). Decatungstate-Mediated C(sp³)-H Heteroarylation via Radical-Polar Crossover in Batch and Flow. *Angewandte Chemie*, 133(33), 18037-18041. <https://doi.org/10.1002/ange.202104682>, <https://doi.org/10.1002/anie.202104682>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

UvA-DARE is a service provided by the library of the University of Amsterdam (<https://dare.uva.nl>)

Flow Chemistry

Decatungstate-Mediated C(sp³)-H Heteroarylation via Radical-Polar Crossover in Batch and Flow

Ting Wan, Luca Capaldo, Gabriele Laudadio, Alexander V. Nyuchev, Juan A. Rincón, Pablo García-Losada, Carlos Mateos, Michael O. Frederick, Manuel Nuño, and Timothy Noël*

Abstract: Photocatalytic hydrogen atom transfer is a very powerful strategy for the regioselective C(sp³)-H functionalization of organic molecules. Herein, we report on the unprecedented combination of decatungstate hydrogen atom transfer photocatalysis with the oxidative radical-polar crossover concept to access the direct net-oxidative C(sp³)-H heteroarylation. The present methodology demonstrates a high functional group tolerance (40 examples) and is scalable when using continuous-flow reactor technology. The developed protocol is also amenable to the late-stage functionalization of biologically relevant molecules such as stanozolol, (-)-ambroxide, podophyllotoxin, and dideoxyribose.

Photocatalytic hydrogen atom transfer (HAT) is witnessing an ever-growing interest from the synthetic community as a versatile strategy for the late-stage functionalization of C(sp³)-H bonds.^[1-3] In this activation mode, the excited state of a photocatalyst can be conveniently exploited to cleave C(sp³)-H bonds to obtain carbon-centered radicals. By exploiting inherent electronic and steric properties of the parent molecule and by tuning the reaction conditions, these nucleophilic radicals can be obtained with high regioselectiv-

ity, thus obviating the need to use any directing or activating groups (Scheme 1 a).

Amongst the different HAT photocatalysts, the decatungstate anion (W; [W₁₀O₃₂]⁴⁻) has proven to be an ideal candidate owing to its unique selectivity, robustness and ease of preparation.^[4,5] The excited state of W (W*) can be readily obtained upon exposure to UV-A light ($\lambda > 365$ nm) and has been used for the activation of C(sp³)-H bonds within a wide variety of hydrogen donors such as ethers, aldehydes, amides and even alkanes. In most cases, the fleeting radical intermediates were used to forge C-C,^[6] C-F,^[7] and C-O^[8] bonds. In contrast, only a handful of examples demonstrate the formation of C-N bonds.^[9] These examples mainly rely on the trapping of the radical with a suitable Michael acceptor, e.g., diisopropyl azodicarboxylate (DIAD), delivering the corresponding hydrazides. Despite its synthetic utility to

[*] T. Wan, Dr. L. Capaldo, Dr. G. Laudadio, Prof. Dr. T. Noël
Flow Chemistry Group, Van't Hoff Institute for Molecular Sciences (HIMS), University of Amsterdam
Science Park 904, 1098 XH, Amsterdam (The Netherlands)
E-mail: t.noel@uva.nl
Homepage: <http://www.NoelResearchGroup.com>

Dr. A. V. Nyuchev
Department of Organic Chemistry
Lobachevsky State University of Nizhny Novgorod
Gagarina Avenue 23, 603950 Nizhny Novgorod (Russia)


Dr. J. A. Rincón, Dr. P. García-Losada, Dr. C. Mateos
Centro de Investigación Lilly S.A.
Avda. de la Industria 30, 28108 Alcobendas-Madrid (Spain)

Dr. M. O. Frederick
Small Molecule Design and Development, Eli Lilly and Company
Indianapolis, IN 46285 (USA)

Dr. M. Nuño
Vapourtec Ltd., Park Farm Business Centre
Fornham St Genevieve, Bury St Edmunds, Suffolk IP28 6TS (UK)

Supporting information and the ORCID identification number(s) for the author(s) of this article can be found under:
<https://doi.org/10.1002/anie.202104682>.

© 2021 The Authors. Angewandte Chemie published by Wiley-VCH GmbH. This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.



Scheme 1. a) Photocatalytic hydrogen atom transfer (HAT) enables the conversion of C-H bonds in complex biologically active molecules. b) Established mechanism for the formation of C-N bonds via TBADT-mediated HAT. c) Proposed approach to realize the regioselective C-H bond heteroarylation through combination of decatungstate-enabled HAT and Radical-Polar Crossover (RPC).

access hydroxamic acids and amides,^[10] this approach remains fairly specific, atom-inefficient and limited to the strongly electrophilic N=N double bond present in DIAD (Scheme 1b).

To expand the scope of C(sp³)-N bond forming reactions using HAT photocatalysis, we envisioned that an unprecedented combination of a decatungstate-induced HAT event with an oxidative radical-polar crossover (RPC) process might overcome this challenge.^[11-13] More specifically, we surmised that subsequent oxidation of carbon-centered radicals generated via HAT would lead to carbocations, which can be conveniently trapped with *N*-heteroaryl-based nucleophiles, thus establishing the targeted carbon-nitrogen bond. A crucial aspect in the development of such a methodology is the nature of the carbocation, which should not only be readily generated but also be sufficiently stable to enable interception with the selected nucleophile. We speculated that the generation of an oxocarbenium ion would provide the required stabilization of the carbocation (Scheme 1c).

Herein, we report the development of an efficient TBADT-mediated (TBADT: tetrabutylammonium decatungstate, (Bu₄N)₄W₁₀O₃₂) heteroarylation of C(sp³)-H bonds through the merger of HAT and RPC, thus demonstrating a new reactivity mode for this widely used HAT photocatalyst. Notably, as shown in this communication, the methodology is amenable to the late-stage functionalization of complex organic molecules and is scalable when using continuous-flow reactor technology.


Our initial investigations commenced with the coupling of tetrahydrofuran (**1a**) and pyrazole (**2a**) (Table 1). For this purpose, an acetonitrile solution of **1a** and **2a** in the presence

of TBADT (2 mol%) was irradiated with UV-A light ($\lambda = 365$ nm, 36 W) for 16 hours. In order to trigger the RPC event, we also added TBHP (*tert*-butylhydroperoxide) as a terminal oxidant. After extensive screening of potential reaction conditions (see the Supporting Information), we found that the target product **3** could be obtained in excellent yield using an excess of **1a** and 3 equivalents of TBHP (Table 1, Entry 1). Reducing the amount of TBHP from 3 to 1 equivalents resulted in decreased yields (Table 1, Entries 2,3). Replacing TBHP with any other oxidant did not lead to improved results (Table 1, Entries 4-7). As expected, when no oxidant was added, the RPC event failed and the nucleophilic addition was precluded; under these circumstances, the desired product **3** was formed only in traces (Table 1, Entry 8). No product was observed in the absence of light and photocatalyst (see the Supporting Information). Finally, in order to reduce the required reaction times and to enable scalability of the reaction protocol,^[14,15] we optimized a flow process (Table 1, Entries 9-11) by using the commercially available photochemical reactor Vapourtec UV-150 (PFA, inner diameter = 1.3 mm, see Supporting Information). Thus, we were able to increase efficiency up to 81% isolated yield (Table 1, Entry 11).

With optimal reaction conditions in both batch and flow established, we examined the generality of our photocatalytic transformation (Scheme 2). We commenced by combining model substrate **1a** with a set of structurally diverse nitrogen-containing heteroarenes. Our benchmark reaction between tetrahydrofuran (**1a**) and pyrazole (**2a**) could be readily scaled up in flow using the standard procedure for prolonged operation times (10 mmol, 80% isolated yield). Alkyl- and aryl-bearing pyrazoles afforded the desired cross-coupled products in fair to excellent yields (**4-8**, 44-94% yield). Furthermore, pyrazoles decorated with electron-withdrawing moieties (e.g. chloro **9**, bromo **10** and ethyl-ester **11**) proved to be competent substrates as well (68-83%). A pyrazole containing the boronic acid pinacol (Bpin) ester functionality was tolerated under the reaction conditions (**12**, 81% yield); notably, this boron-functionality can serve as a branching point for further diversification using Suzuki-Miyaura or Chan-Evans-Lam cross-coupling chemistry.^[16,17] Next, we successfully extended this C(sp³)-H heteroarylation protocol to other *N*-containing five-membered rings such as imidazoles (**13-14**, 42-76%), triazoles (**15**, 96%) and tetrazoles (**16**, 46%), as well as various benzo-fused heteroaromatic structures (**17-19**, 42-55%). Furthermore, several in nature-occurring heterocycles, such as xanthines (**20** and **21**, 18-60%) and purines (**22-25**, 24-63%), served as adequate coupling partners. Most of the latter reactions were run in batch owing to the limited solubility of the heteroaromatic nucleophiles. Remarkably, we also managed to functionalize anabolic steroid stanozolol to get product **26** with an excellent mass balance (34% yield, 87% brsm).


Next, we investigated the scope of suitable H-donors. Efficient α -to-O C-H functionalization was achieved for both tetrahydropyran and oxepane, providing the targeted compounds **27** and **28** in good yields (76% and 56%, respectively). Also α -to-S and α -to-N C-H bond functionalization was observed (**29-32**, 58-96%) with great selectivity over

Table 1: Optimization of reaction conditions.^[a]



Entry	Variation from conditions	Yield ^[c]
1	none	86
2	TBHP (2 equiv)	80
3	TBHP (1 equiv)	70
4 ^[d]	O ₂ atmosphere	4
5	H ₂ O ₂ (3 equiv)	32
6	DTP (3 equiv)	20
7	BPO (3 equiv)	52
8	No sacrificial oxidant	traces
9 ^[e]	flow, $t_R = 1$ h; TBADT (5 mol%)	64
10 ^[f]	flow, $t_R = 1$ h; TBADT (5 mol%), 60 W LED	74
11	flow, $t_R = 1$ h; 60 W LED, TBADT (5 mol%); 1a (18 equiv)	86 (81)

[a] **1a** (6 equiv), **2a** (0.2 mmol), TBADT (2 mol%), TBHP (3 equiv) in CH₃CN (1 mL); solution sparged with Ar prior to irradiation ($\lambda = 365$ nm, 16 h). [b] TBHP (5.5 M in decane or nonane). [c] Yields determined by ¹H NMR spectroscopy using pyrazine as external standard. [d] reaction mixture sparged with O₂ prior to irradiation and kept under aerobic atmosphere (balloon filled with O₂). [e] flow setup 1 (see SI): $V_R = 6$ mL, flow rate = 0.1 mL min⁻¹, $t_R = 1$ h, 36 W LED ($\lambda = 365$ nm). [f] flow setup 2 (see SI): $V_R = 10$ mL, flow rate = 0.167 mL min⁻¹, $t_R = 1$ h, 60 W LED ($\lambda = 365$ nm). In parenthesis is isolated yield.




Scheme 2. Substrate scope of the decatungstate-mediated C(sp³)-H heteroarylation via radical-polar crossover in batch and flow. **1a** (18 equiv), **2a** (1.0 mmol), TBADT (5 mol%), TBHP (3 equiv) in CH₃CN (5 mL); solution sparged with Ar prior to irradiation (λ = 365 nm 60 W, reactor volume: 10 mL, flow rate: 0.167 mL min⁻¹, t_R: 1 h). Isolated yields are given. [a] TBHP was used 5.5 M in decane or nonane. [b] Reaction carried out in batch (16 h); TBADT (5 mol%). [c] CH₃CN (12 mL). [d] CH₃CN/CHCl₃ 1:1 (10 mL); **1a** (18 equiv). [e] Solvent mixture: CH₃CN/**1a** 1:1 (10 mL). [f] Solvent mixture: CH₃CN/**1a** 2:5 (14 mL). [g] 6 equiv of the hydrogen donor were used. [h] CH₃CN/CH₂Cl₂ 5:1 (6 mL). [i] 2 equiv of hydrogen donor were used, CH₃CN (15 mL). [j] Reaction carried out as in Table 1, entry 1; 2 equiv of hydrogen donor were used. Brsm: based on remaining starting material.

competitive α-to-O C-H activation. In the case of very activated methylenes, such as in benzo[*b*]furan and 1,3-benzodioxole, excellent yields of the corresponding products **33** (95%) and **34** (86%) were observed. Acyclic ethers such as diethyl ether and methyl *tert*-butyl ether could be used as substrates as well, yielding products **35** and **36** in 68% and 33% yield, respectively. The ability of this method to enable the late-stage functionalization of complex organic molecules or natural scaffolds was demonstrated in the case of (-)-ambroxide and acetyl-protected podophyllotoxin; these substrates were functionalized in 88% (**37**) and 40% (**38**),

respectively. Finally, silyl-protected dideoxyribose was subjected to our reaction conditions and we found that it could be functionalized in satisfactory yields (**39–40**, 20–28% yield, 50–56% brsm).

A plausible mechanistic rationale for the C(sp³)-H heteroarylation is shown in Scheme 3a. Upon absorption of UV-A light, the excited state of TBADT (W*) is able to cleave the C-H bond in α-position to the heteroatom (i.e., O, N or S) yielding a carbon-centered nucleophilic α-oxyalkyl radical (**I**). This radical intermediate (**I**) is readily oxidized by TBHP resulting in the formation of a stabilized electrophilic



Scheme 3. Mechanistic investigation: a) Proposed mechanism with KIE and quenching rates. b) Radical trapping experiments. c) Oxocarbenium trapping experiments. [a] **1h**, **i** (18 equiv), **2a** (1.0 mmol), TBADT (5 mol%), TBHP (3 equiv) in CH_3CN (5 mL); solution sparged with Ar prior to irradiation ($\lambda = 365 \text{ nm}$ 60 W, reactor volume: 10 mL, flow rate: $0.167 \text{ mL min}^{-1}$, t_{R} : 1 h). [b] Yield determined by $^1\text{H NMR}$ spectroscopy using pyrazine as external standard. [c] Isolated yield.

oxocarbenium ion (I^+), which is trapped by the heteroarene establishing the targeted C–N bond.

Using laser flash photolysis, the quenching of the excited state of TBADT (W^*) was studied in more detail (see Supporting Information, Section 3.4). From the derived Stern–Volmer plots, the quenching rates with both substrates **1a** and **2a** were calculated. Interestingly, both tetrahydrofuran and pyrazole are able to quench W^* with comparable bimolecular rates ($k = 2.3$ and $1.6 \times 10^8 \text{ M}^{-1} \text{ s}^{-1}$, respectively). However, given the excess of **1a**, it is reasonable to state that tetrahydrofuran (**1a**) is the actual quencher in the reported experiments, thus delivering **I**. This assumption was further confirmed in radical trapping experiments by the detection of the corresponding adducts of the α -oxyalkyl radical with radical scavengers (**41** and **42**, as evidenced by HRMS) (Scheme 3b). Interestingly, a significant Kinetic Isotope Effect (KIE) of 2.3 was measured, which is in accordance with HAT being the rate-determining step (see Supporting Information, Section 3.3). Quantum yield measurements ($\Phi = 0.18$) further ruled out a radical chain mechanism (See Supporting Information, Section 3.5). Next, we set out to trap

the oxocarbenium ionic intermediate by adding an excess of CH_3OH or $t\text{-BuOH}$ (18 equiv) (Scheme 3c).^[18] In both cases, the corresponding orthoesters **43** and **44** could be detected via HRMS and $^1\text{H NMR}$ spectroscopy. Despite significant efforts, this $\text{C}(\text{sp}^3)\text{--H}$ heteroarylation method could not be extended to unactivated aliphatic $\text{C}(\text{sp}^3)\text{--H}$ bonds (see the Supporting Information, Section 5 for limitations of the scope). We surmise that the oxidation of **I** to I^+ is only favored when the latter species is strongly stabilized, e.g., as an oxocarbenium species.

It should be noted that our methodology compares favorably to thermal approaches relying on TBAI/TBHP^[19] (TBAI = tetrabutylammonium iodide) and $\text{Fe}^{\text{III}}/\text{TBHP}$ ^[20] systems as it requires more controlled and milder conditions to generate the oxocarbenium ion, resulting in a broader functional group tolerance. Furthermore, in comparison to other photocatalytic^[21] and (photo)electrochemical methods,^[22,23] the developed approach is based on a direct and mild $\text{C}(\text{sp}^3)\text{--H}$ cleavage, presents a broader scope and does not require pre-functionalization of the starting materials.

In conclusion, a convenient methodology to forge $\text{C}(\text{sp}^3)\text{--N}$ bonds by combining the radical–polar crossover concept with decatungstate HAT photocatalysis has been realized. Due to the mild reaction conditions, this protocol is amenable both to early and late-stage functionalization of organic molecules. As such, this synthetic method is valuable for the synthesis of medicinal and agrochemical intermediates. While this is the first report to combine decatungstate-induced HAT with oxidative RPC, we believe that the insights gained herein will inspire further advances in the use of this strategy to enable other challenging synthetic transformations.

Acknowledgements

The authors are appreciative of the funding received from the Lilly Research Award Program. T.W. has received support from the China Scholarship Council (CSC) for her PhD studies. L.C. acknowledges European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 101023615 (project name: HAT-TRICK).

Conflict of Interest

The authors declare no conflict of interest.

Keywords: flow chemistry · heteroarylation · hydrogen atom transfer · radical–polar crossover · TBADT

- [1] L. Capaldo, D. Ravelli, *Eur. J. Org. Chem.* **2017**, 2056–2071.
- [2] L. Capaldo, L. L. Quadri, D. Ravelli, *Green Chem.* **2020**, *22*, 3376–3396.
- [3] S. Govaerts, A. Nyuchev, T. Noël, *J. Flow Chem.* **2020**, *10*, 13–71.
- [4] D. Ravelli, S. Protti, M. Fagnoni, *Acc. Chem. Res.* **2016**, *49*, 2232–2242.
- [5] M. D. Tzirakis, I. N. Lykakis, M. Orfanopoulos, *Chem. Soc. Rev.* **2009**, *38*, 2609–2621.

- [6] For representative works on TBADT-catalyzed C–C bond formation, see: a) J. J. Murphy, D. Bastida, S. Paria, M. Fagnoni, P. Melchiorre, *Nature* **2016**, *532*, 218–222; b) I. B. Perry, T. F. Brewer, P. J. Sarver, D. M. Schultz, D. A. DiRocco, D. W. C. MacMillan, *Nature* **2018**, *560*, 70–75; c) G. Laudadio, Y. Deng, K. van der Wal, D. Ravelli, M. Nuño, M. Fagnoni, D. Guthrie, Y. Sun, T. Noël, *Science* **2020**, *369*, 92–96; d) H. Cao, Y. Kuang, X. Shi, K. L. Wong, B. B. Tan, J. M. C. Kwan, X. Liu, J. Wu, *Nat. Commun.* **2020**, *11*, 1956; e) Z. Wen, A. Maheshwari, C. Sambiagio, Y. Deng, G. Laudadio, K. Van Aken, Y. Sun, H. P. L. Gemoets, T. Noël, *Org. Process Res. Dev.* **2020**, *24*, 2356–2361; f) L. Capaldo, L. L. Quadri, D. Merli, D. Ravelli, *Chem. Commun.* **2021**, *57*, 4424–4427.
- [7] For representative works on TBADT-catalyzed C–F bond formation, see: a) S. D. Halperin, H. Fan, S. Chang, R. E. Martin, R. Britton, *Angew. Chem. Int. Ed.* **2014**, *53*, 4690–4693; *Angew. Chem.* **2014**, *126*, 4778–4781; b) M. B. Nodwell, H. Yang, M. Čolović, Z. Yuan, H. Merkens, R. E. Martin, F. Bénard, P. Schaffer, R. Britton, *J. Am. Chem. Soc.* **2017**, *139*, 3595–3598; c) Z. Yuan, M. B. Nodwell, H. Yang, N. Malik, H. Merkens, F. Bénard, R. E. Martin, P. Schaffer, R. Britton, *Angew. Chem. Int. Ed.* **2018**, *57*, 12733–12736; *Angew. Chem.* **2018**, *130*, 12915–12918; d) M. Meanwell, J. Lehmann, M. Eichenberger, R. E. Martin, R. Britton, *Chem. Commun.* **2018**, *54*, 9985–9988.
- [8] For representative works on TBADT-catalyzed C–O bond formation, see: a) D. M. Schultz, F. Lévesque, D. A. DiRocco, M. Reibarkh, Y. Ji, L. A. Joyce, J. F. Dropinski, H. Sheng, B. D. Sherry, I. W. Davies, *Angew. Chem. Int. Ed.* **2017**, *56*, 15274–15278; *Angew. Chem.* **2017**, *129*, 15476–15480; b) G. Laudadio, S. Govaerts, Y. Wang, D. Ravelli, H. F. Koolman, M. Fagnoni, S. W. Djuric, T. Noël, *Angew. Chem. Int. Ed.* **2018**, *57*, 4078–4082; *Angew. Chem.* **2018**, *130*, 4142–4146.
- [9] a) I. Ryu, A. Tani, T. Fukuyama, D. Ravelli, S. Montanaro, M. Fagnoni, *Org. Lett.* **2013**, *15*, 2554–2557; b) F. Bonassi, D. Ravelli, S. Protti, M. Fagnoni, *Adv. Synth. Catal.* **2015**, *357*, 3687–3695.
- [10] a) G. N. Papadopoulos, C. G. Kokotos, *J. Org. Chem.* **2016**, *81*, 7023–7028; b) G. N. Papadopoulos, C. G. Kokotos, *Chem. Eur. J.* **2016**, *22*, 6964–6967; c) G. N. Papadopoulos, M. G. Kokotou, N. Spiliopoulou, N. F. Nikitas, E. Voutyritsa, D. I. Tzaras, N. Kaplaneris, C. G. Kokotos, *ChemSusChem* **2020**, *13*, 5934–5944.
- [11] For reviews on radical–polar crossover (RPC) in photoredox catalysis, see: a) L. Pitzer, J. L. Schwarz, F. Glorius, *Chem. Sci.* **2019**, *10*, 8285–8291; b) R. J. Wiles, G. A. Molander, *Isr. J. Chem.* **2020**, *60*, 281–293.
- [12] K. Donabauer, M. Maity, A. L. Berger, G. S. Huff, S. Crespi, B. König, *Chem. Sci.* **2019**, *10*, 5162–5166.
- [13] K. Donabauer, K. Murugesan, U. Rozman, S. Crespi, B. König, *Chem. Eur. J.* **2020**, *26*, 12945–12950.
- [14] C. Sambiagio, T. Noël, *Trends Chem.* **2020**, *2*, 92–106.
- [15] D. Cambié, C. Bottecchia, N. J. W. Straathof, V. Hessel, T. Noël, *Chem. Rev.* **2016**, *116*, 10276–10341.
- [16] A. J. J. Lennox, G. C. Lloyd-Jones, *Chem. Soc. Rev.* **2014**, *43*, 412–443.
- [17] J. C. Vantourout, R. P. Law, A. Isidro-Llobet, S. J. Atkinson, A. J. B. Watson, *J. Org. Chem.* **2016**, *81*, 3942–3950.
- [18] a) X. Si, L. Zhang, Z. Wu, M. Rudolph, A. M. Asiri, A. S. K. Hashmi, *Org. Lett.* **2020**, *22*, 5844–5849; b) B. Ye, J. Zhao, K. Zhao, J. M. McKenna, F. D. Toste, *J. Am. Chem. Soc.* **2018**, *140*, 8350–8356.
- [19] a) H. Aruri, U. Singh, S. Sharma, S. Gudup, M. Bhogal, S. Kumar, D. Singh, V. K. Gupta, R. Kant, R. A. Vishwakarma, P. P. Singh, *J. Org. Chem.* **2015**, *80*, 1929–1936; b) H. Abebe, S. Vidavalur, V. R. Battula, *RSC Adv.* **2016**, *6*, 82289–82293; c) H. Abebe, S. Vidavalur, V. R. Battula, *ChemistrySelect* **2017**, *2*, 4165–4168.
- [20] S. Pan, J. Liu, H. Li, Z. Wang, X. Guo, Z. Li, *Org. Lett.* **2010**, *12*, 1932–1935.
- [21] L. Zhang, H. Yi, J. Wang, A. Lei, *J. Org. Chem.* **2017**, *82*, 10704–10709.
- [22] a) J. Wu, Y. Zhou, Y. Zhou, C.-W. Chiang, A. Lei, *ACS Catal.* **2017**, *7*, 8320–8323; b) T. Sheng, H.-J. Zhang, M. Shang, C. He, J. C. Vantourout, P. S. Baran, *Org. Lett.* **2020**, *22*, 7594–7598.
- [23] H. Huang, Z. M. Strater, T. H. Lambert, *J. Am. Chem. Soc.* **2020**, *142*, 1698–1703.

Manuscript received: April 5, 2021

Revised manuscript received: May 29, 2021

Accepted manuscript online: May 31, 2021

Version of record online: July 9, 2021