



UvA-DARE (Digital Academic Repository)

Operator preconditioning

the simplest case

Stevenson, R.; van Venetië, R.

DOI

[10.1016/j.apnum.2021.09.016](https://doi.org/10.1016/j.apnum.2021.09.016)

Publication date

2022

Document Version

Final published version

Published in

Applied Numerical Mathematics

License

CC BY

[Link to publication](#)

Citation for published version (APA):


Stevenson, R., & van Venetië, R. (2022). Operator preconditioning: the simplest case. *Applied Numerical Mathematics*, 172, 292-299. <https://doi.org/10.1016/j.apnum.2021.09.016>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.



Operator preconditioning: the simplest case

Rob Stevenson, Raymond van Venetië^{*,1}

Korteweg-de Vries Institute for Mathematics, University of Amsterdam, P.O. Box 94248, 1090 GE Amsterdam, the Netherlands



ARTICLE INFO

Article history:

Received 23 February 2021

Received in revised form 3 September 2021

Accepted 24 September 2021

Available online 30 September 2021

Keywords:

Operator preconditioning

Uniform preconditioners

Finite- and boundary elements

ABSTRACT

Using the framework of operator or Calderón preconditioning, uniform preconditioners are constructed for elliptic operators discretized with continuous finite (or boundary) elements. The preconditioners are constructed as the composition of an opposite order operator, discretized on the same ansatz space, and two identical diagonal scaling operators, whose matrix representation is the lumped mass matrix.

© 2021 The Authors. Published by Elsevier B.V. on behalf of IMACS. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

1. Introduction

This paper deals with the construction of uniform preconditioners for negative and positive order operators, discretized by continuous piecewise polynomial trial spaces, using the framework of ‘operator preconditioning’ [8], see also [12,11,3,9].

For some d -dimensional closed domain (or manifold) Ω and an $s \in [0, 1]$, we consider the (fractional) Sobolev space $H^s(\Omega)$ and its dual that we denote by $H^{-s}(\Omega)$. Let $(\mathcal{S}_{\mathcal{T}})_{\mathcal{T} \in \mathbb{T}}$ be a family of *continuous piecewise* polynomials of some fixed degree ℓ w.r.t. uniformly shape regular, possibly locally refined, partitions.

Given some families of uniformly boundedly invertible operators

$$A_{\mathcal{T}}: (\mathcal{S}_{\mathcal{T}}, \|\cdot\|_{H^{-s}(\Omega)}) \rightarrow (\mathcal{S}_{\mathcal{T}}, \|\cdot\|_{H^{-s}(\Omega)}),$$

$$B_{\mathcal{T}}: (\mathcal{S}_{\mathcal{T}}, \|\cdot\|_{H^s(\Omega)}) \rightarrow (\mathcal{S}_{\mathcal{T}}, \|\cdot\|_{H^s(\Omega)}),$$

we are interested in constructing a *preconditioner* for $A_{\mathcal{T}}$ using operator preconditioning with $B_{\mathcal{T}}$, and vice versa. To this end, we introduce a uniformly boundedly invertible operator $D_{\mathcal{T}}: (\mathcal{S}_{\mathcal{T}}, \|\cdot\|_{H^{-s}(\Omega)}) \rightarrow (\mathcal{S}_{\mathcal{T}}, \|\cdot\|_{H^s(\Omega)})$, yielding preconditioned systems $D_{\mathcal{T}}^{-1}B_{\mathcal{T}}(D'_{\mathcal{T}})^{-1}A_{\mathcal{T}}$ and $(D'_{\mathcal{T}})^{-1}A_{\mathcal{T}}D_{\mathcal{T}}^{-1}B_{\mathcal{T}}$ that are uniformly boundedly invertible.

In earlier research, [13,14], we already constructed such preconditioners in a more general setting where *different* ansatz spaces were used to define $A_{\mathcal{T}}$ and $B_{\mathcal{T}}$. The setting studied in the current work, however, allows for preconditioners with a remarkably simple implementation.

A typical setting is that for some $A: H^{-s}(\Omega) \rightarrow H^s(\Omega)$ and $B: H^s(\Omega) \rightarrow H^{-s}(\Omega)$, both boundedly invertible and coercive, it holds that $(A_{\mathcal{T}}u)(v) := (Au)(v)$ and $(B_{\mathcal{T}}u)(v) := (Bu)(v)$ with $u, v \in \mathcal{S}_{\mathcal{T}}$. An example for $s = \frac{1}{2}$ is that A is the Single Layer Integral operator and B is the Hypersingular Integral operator. For this case, continuity of piecewise polynomial trial functions is required for discretizing B , but not for A , for which often discontinuous piecewise polynomials are employed.

* Corresponding author.

E-mail addresses: r.p.stevenson@uva.nl (R. Stevenson), r.vanvenetie@uva.nl (R. van Venetië).

¹ The second author has been supported by the Netherlands Organisation for Scientific Research (NWO) under contract, no. 613.001.652.

Nevertheless, when the solution of the Single Layer Integral equation is expected to be smooth, e.g., when Ω is a smooth manifold, then it is advantageous to take an ansatz space of continuous (or even smoother) functions also for A .

An obvious choice for $D_{\mathcal{T}}$ would be to consider $(D_{\mathcal{T}}u)(v) := \langle u, v \rangle_{L_2(\Omega)}$. However, a problem becomes apparent when one considers the matrix representation $\mathbf{D}_{\mathcal{T}}$ of $D_{\mathcal{T}}$ in the standard basis being the mass matrix: the inverse matrix $\mathbf{D}_{\mathcal{T}}^{-1}$, that appears in the preconditioned system, is densely populated. In view of application cost, this inverse matrix has to be approximated, where it generally can be expected that, in order to obtain a uniform preconditioner, approximation errors have to decrease with a decreasing (minimal) mesh size, which will be confirmed in a numerical experiment. To circumvent this issue, we will introduce a $D_{\mathcal{T}}$ that has a *diagonal* matrix representation, so that its inverse can be exactly evaluated.

1.1. Notation

In this work, by $\lambda \lesssim \mu$ we mean that λ can be bounded by a multiple of μ , independently of parameters which λ and μ may depend on, with the sole exception of the space dimension d , or in the manifold case, on the parametrization of the manifold that is used to define the finite element spaces on it. Obviously, $\lambda \gtrsim \mu$ is defined as $\mu \lesssim \lambda$, and $\lambda \approx \mu$ as $\lambda \lesssim \mu$ and $\lambda \gtrsim \mu$.

For normed linear spaces \mathcal{Y} and \mathcal{Z} , in this paper for convenience over \mathbb{R} , $\mathcal{L}(\mathcal{Y}, \mathcal{Z})$ will denote the space of bounded linear mappings $\mathcal{Y} \rightarrow \mathcal{Z}$ endowed with the operator norm $\|\cdot\|_{\mathcal{L}(\mathcal{Y}, \mathcal{Z})}$. The subset of invertible operators in $\mathcal{L}(\mathcal{Y}, \mathcal{Z})$ with inverses in $\mathcal{L}(\mathcal{Z}, \mathcal{Y})$ will be denoted as $\text{Lis}(\mathcal{Y}, \mathcal{Z})$.

For \mathcal{Y} a reflexive Banach space and $C \in \mathcal{L}(\mathcal{Y}, \mathcal{Y}')$ being *coercive*, i.e.,

$$\inf_{0 \neq y \in \mathcal{Y}} \frac{(Cy)(y)}{\|y\|_{\mathcal{Y}}^2} > 0,$$

both C and $\Re(C) := \frac{1}{2}(C + C')$ are in $\text{Lis}(\mathcal{Y}, \mathcal{Y}')$ with

$$\begin{aligned} \|\Re(C)\|_{\mathcal{L}(\mathcal{Y}, \mathcal{Y}')} &\leq \|C\|_{\mathcal{L}(\mathcal{Y}, \mathcal{Y}')} \\ \|C^{-1}\|_{\mathcal{L}(\mathcal{Y}', \mathcal{Y})} &\leq \|\Re(C)^{-1}\|_{\mathcal{L}(\mathcal{Y}', \mathcal{Y})} = \left(\inf_{0 \neq y \in \mathcal{Y}} \frac{(Cy)(y)}{\|y\|_{\mathcal{Y}}^2} \right)^{-1}. \end{aligned}$$

The set of coercive $C \in \text{Lis}(\mathcal{Y}, \mathcal{Y}')$ is denoted as $\text{Lis}_c(\mathcal{Y}, \mathcal{Y}')$. If $C \in \text{Lis}_c(\mathcal{Y}, \mathcal{Y}')$, then $C^{-1} \in \text{Lis}_c(\mathcal{Y}', \mathcal{Y})$ and $\|\Re(C^{-1})^{-1}\|_{\mathcal{L}(\mathcal{Y}, \mathcal{Y}')} \leq \|C\|_{\mathcal{L}(\mathcal{Y}, \mathcal{Y}')}^2 \|\Re(C)^{-1}\|_{\mathcal{L}(\mathcal{Y}', \mathcal{Y})}$.

Given a family of operators $C_i \in \text{Lis}(\mathcal{Y}_i, \mathcal{Z}_i)$ ($\text{Lis}_c(\mathcal{Y}_i, \mathcal{Z}_i)$), we will write $C_i \in \text{Lis}(\mathcal{Y}_i, \mathcal{Z}_i)$ ($\text{Lis}_c(\mathcal{Y}_i, \mathcal{Z}_i)$) uniformly in i , or simply ‘uniform’, when

$$\sup_i \max(\|C_i\|_{\mathcal{L}(\mathcal{Y}_i, \mathcal{Z}_i)}, \|C_i^{-1}\|_{\mathcal{L}(\mathcal{Z}_i, \mathcal{Y}_i)}) < \infty,$$

or

$$\sup_i \max(\|C_i\|_{\mathcal{L}(\mathcal{Y}_i, \mathcal{Z}_i)}, \|\Re(C_i)^{-1}\|_{\mathcal{L}(\mathcal{Z}_i, \mathcal{Y}_i)}) < \infty.$$

2. Construction of $D_{\mathcal{T}}$ in the domain case

For some d -dimensional domain Ω and an $s \in [0, 1]$, we consider the Sobolev spaces

$$H^s(\Omega) := [L_2(\Omega), H^1(\Omega)]_{s,2}, \quad H^{-s}(\Omega) := H^s(\Omega)',$$

which form the Gelfand triple $H^s(\Omega) \hookrightarrow L_2(\Omega) \simeq L_2(\Omega)' \hookrightarrow H^{-s}(\Omega)$.

Remark 2.1. In this work, for convenience we restrict ourselves to Sobolev spaces with positive smoothness index which do not incorporate homogeneous Dirichlet boundary conditions and their duals. The proofs given below can however be extended to the setting with boundary conditions, see the arguments found in [13,14].

Let $(\mathcal{T})_{\mathcal{T} \in \mathbb{T}}$ be a family of *conforming* partitions of Ω into (open) *uniformly shape regular* d -simplices. Thanks to the conformity and the uniform shape regularity, for $d > 1$ we know that neighbouring $T, T' \in \mathcal{T}$, i.e. $\bar{T} \cap \bar{T}' \neq \emptyset$, have uniformly comparable sizes. For $d = 1$, we impose this uniform ‘ K -mesh property’ explicitly.

Fix $\ell > 0$. For $\mathcal{T} \in \mathbb{T}$, let $\mathcal{S}_{\mathcal{T}}$ denote the space of *continuous piecewise polynomials* of degree ℓ w.r.t. \mathcal{T} , i.e.,

$$\mathcal{S}_{\mathcal{T}} := \{u \in H^1(\Omega) : u|_T \in \mathcal{P}_{\ell} (T \in \mathcal{T})\}.$$

Additionally, for $r \in [-1, 1]$, we will write $\mathcal{S}_{\mathcal{T},r}$ as shorthand notation for the normed linear space $(\mathcal{S}_{\mathcal{T}}, \|\cdot\|_{H^r(\Omega)})$.

Denote $N_{\mathcal{T}}$ for the set of the usual Lagrange evaluation points of $\mathcal{S}_{\mathcal{T}}$, and equip the latter space with $\Phi_{\mathcal{T}} = \{\phi_{\mathcal{T},v} : v \in N_{\mathcal{T}}\}$, being the *canonical nodal basis* defined by $\phi_{\mathcal{T},v}(v') := \delta_{vv'}$ ($v, v' \in N_{\mathcal{T}}$). For $T \in \mathcal{T}$, set $h_T := |T|^{1/d}$ and let $N_T := \bar{T} \cap N_{\mathcal{T}}$ be the set of evaluation points in \bar{T} . We will omit notational dependence on \mathcal{T} if it is clear from the context, e.g., we will simply write ϕ_v .

2.1. Operator preconditioning

Given some family of opposite order operators $A_{\mathcal{T}} \in \text{Lis}_c(\mathcal{S}_{\mathcal{T},-s}, (\mathcal{S}_{\mathcal{T},-s})')$ and $B_{\mathcal{T}} \in \text{Lis}_c(\mathcal{S}_{\mathcal{T},s}, (\mathcal{S}_{\mathcal{T},s})')$, both uniformly in $\mathcal{T} \in \mathbb{T}$, we are interested in constructing *optimal* preconditioners for both $A_{\mathcal{T}}$ and $B_{\mathcal{T}}$, using the idea of opposite order preconditioning ([8]).

That is, if one has an additional family of operators $D_{\mathcal{T}} \in \text{Lis}(\mathcal{S}_{\mathcal{T},-s}, (\mathcal{S}_{\mathcal{T},s})')$ uniformly in $\mathcal{T} \in \mathbb{T}$, then uniformly preconditioned systems for $A_{\mathcal{T}}$ and $B_{\mathcal{T}}$ are given by

$$\begin{aligned} D_{\mathcal{T}}^{-1} B_{\mathcal{T}} (D'_{\mathcal{T}})^{-1} A_{\mathcal{T}} &\in \text{Lis}(\mathcal{S}_{\mathcal{T},-s}, \mathcal{S}_{\mathcal{T},-s}), \\ (D'_{\mathcal{T}})^{-1} A_{\mathcal{T}} D_{\mathcal{T}}^{-1} B_{\mathcal{T}} &\in \text{Lis}(\mathcal{S}_{\mathcal{T},s}, \mathcal{S}_{\mathcal{T},s}), \end{aligned} \tag{2.1}$$

see the following diagram:

$$\begin{array}{ccc} \mathcal{S}_{\mathcal{T},-s} & \xrightarrow{A_{\mathcal{T}}} & (\mathcal{S}_{\mathcal{T},-s})' \\ D_{\mathcal{T}}^{-1} \uparrow & & \downarrow (D'_{\mathcal{T}})^{-1} \cdot \\ (\mathcal{S}_{\mathcal{T},s})' & \xleftarrow{B_{\mathcal{T}}} & \mathcal{S}_{\mathcal{T},s} \end{array}$$

In the following we shall be concerned with constructing a suitable family $D_{\mathcal{T}}$.

2.1.1. An obvious but unsatisfactory choice for $D_{\mathcal{T}}$

An option would be to consider $(D_{\mathcal{T}}u)(v) := \langle u, v \rangle_{L_2(\Omega)}$ ($u, v \in \mathcal{S}_{\mathcal{T}}$), being uniformly in $\mathcal{L}(\mathcal{S}_{\mathcal{T},-s}, (\mathcal{S}_{\mathcal{T},s})')$. For showing boundedness of its inverse, let $Q_{\mathcal{T}}$ be the $L_2(\Omega)$ -orthogonal projector onto $\mathcal{S}_{\mathcal{T}}$ then

$$\begin{aligned} \|D_{\mathcal{T}}^{-1}\|_{\mathcal{L}((\mathcal{S}_{\mathcal{T},s})', \mathcal{S}_{\mathcal{T},-s})}^{-1} &= \inf_{0 \neq u \in \mathcal{S}_{\mathcal{T},-s}} \sup_{0 \neq v \in H^s(\Omega)} \frac{\langle u, v \rangle_{L_2(\Omega)}}{\|u\|_{H^{-s}(\Omega)} \|Q_{\mathcal{T}}v\|_{H^s(\Omega)}} \\ &\geq \|Q_{\mathcal{T}}\|_{\mathcal{L}(H^s(\Omega), H^s(\Omega))}^{-1}. \end{aligned}$$

As follows from [13, Prop. 2.3], the converse is also true, i.e., uniform boundedness of $\|D_{\mathcal{T}}^{-1}\|_{\mathcal{L}((\mathcal{S}_{\mathcal{T},s})', \mathcal{S}_{\mathcal{T},-s})}$ is actually *equivalent* to uniform boundedness of $\|Q_{\mathcal{T}}\|_{\mathcal{L}(H^s(\Omega), H^s(\Omega))}$.

This uniform boundedness of $\|Q_{\mathcal{T}}\|_{\mathcal{L}(H^s(\Omega), H^s(\Omega))}$ is well-known for families of *quasi-uniform*, uniformly shape regular conforming partitions of Ω into say d -simplices. It has also been demonstrated for families of locally refined partitions, for $d = 2$ including those that are generated by the newest vertex bisection (NVB) algorithm, see [4,6,5]. On the other hand, in [1] a one-dimensional counterexample was presented in which the $L_2(\Omega)$ -orthogonal projector on a family of sufficiently strongly graded, although uniform K meshes, is not $H^1(\Omega)$ -stable. Thus, in any case uniform $H^1(\Omega)$ -stability cannot hold without assuming some sufficiently mild grading of the meshes.

Aside from this latter theoretical shortcoming, more importantly, there is a computational problem with the current choice of $D_{\mathcal{T}}$. The matrix representation of $D_{\mathcal{T}}$ w.r.t. $\Phi_{\mathcal{T}}$ is the ‘mass matrix’ $\mathbf{D}_{\mathcal{T}} := \langle \Phi_{\mathcal{T}}, \Phi_{\mathcal{T}} \rangle_{L_2(\Omega)}$. Its inverse $\mathbf{D}_{\mathcal{T}}^{-1}$, appearing in the preconditioner, is densely populated, and therefore has to be approximated, where generally the error in such approximations has to decrease with a decreasing (minimal) mesh-size in order to arrive at a uniform preconditioner.

2.2. Constructing a practical $D_{\mathcal{T}}$

To avoid the aforementioned problems, we shall construct $D_{\mathcal{T}} \in \text{Lis}(\mathcal{S}_{\mathcal{T},-s}, (\mathcal{S}_{\mathcal{T},s})')$ with a diagonal matrix representation. To this end, we require some auxiliary space $\tilde{\mathcal{S}}_{\mathcal{T}} \subset H^1(\Omega)$ equipped with a local basis $\tilde{\Phi}_{\mathcal{T}}$ that is $L_2(\Omega)$ -biorthogonal to $\Phi_{\mathcal{T}}$ and that has ‘approximation properties’. To be precise, let $\tilde{\Phi}_{\mathcal{T}} := \{\tilde{\phi}_v \in H^1(\Omega) : v \in N_{\mathcal{T}}\}$ be some collection that satisfies²:

$$\begin{aligned} \langle \tilde{\phi}_v, \phi_{v'} \rangle_{L_2(\Omega)} &= \delta_{vv'} \langle \mathbb{1}, \phi_v \rangle_{L_2(\Omega)}, \quad \sum_{v \in N_{\mathcal{T}}} \tilde{\phi}_v = \mathbb{1}_{\Omega}, \\ \|\tilde{\phi}_v\|_{H^k(\Omega)} &\lesssim \|\phi_v\|_{H^k(\Omega)} \quad (k \in \{0, 1\}), \quad \text{supp } \tilde{\phi}_v \subseteq \text{supp } \phi_v. \end{aligned} \tag{2.2}$$

We will take $D_{\mathcal{T}} := I'_{\mathcal{T}} \tilde{D}_{\mathcal{T}}$ with $\tilde{D}_{\mathcal{T}}$ and $I_{\mathcal{T}}$ being defined and analyzed in the next two theorems.

Theorem 2.2. *The operator $\tilde{D}_{\mathcal{T}}: \mathcal{S}_{\mathcal{T},-s} \rightarrow (\tilde{\mathcal{S}}_{\mathcal{T},s})'$, defined by $(\tilde{D}_{\mathcal{T}}u)(v) := \langle u, v \rangle_{L_2(\Omega)}$, satisfies $\tilde{D}_{\mathcal{T}} \in \text{Lis}(\mathcal{S}_{\mathcal{T},-s}, (\tilde{\mathcal{S}}_{\mathcal{T},s})')$ uniformly in $\mathcal{T} \in \mathbb{T}$.*

² This last condition can be replaced by $\tilde{\phi}_v$ having (uniformly) local support.

Proof. This proof largely follows [13, Sect. 3.1], but because here we consider a Sobolev space $H^s(\Omega)$ that does not incorporate homogeneous boundary conditions, it allows for an easier proof.

From the assumptions (2.2), it follows that the biorthogonal ‘Fortin’ projector $P_{\mathcal{T}}: L_2(\Omega) \rightarrow H^1(\Omega)$ onto $\tilde{\mathcal{F}}_{\mathcal{T}}$ with $\text{ran}(\text{Id} - P_{\mathcal{T}}) = \mathcal{S}_{\mathcal{T}}^{\perp L_2(\Omega)}$ exists, and is given by

$$P_{\mathcal{T}}u = \sum_{v \in N_{\mathcal{T}}} \frac{\langle u, \phi_v \rangle_{L_2(\Omega)}}{\langle \tilde{\phi}_v, \phi_v \rangle_{L_2(\Omega)}} \tilde{\phi}_v.$$

Let $T \in \mathcal{T}$, by (2.2) and the fact that $\langle \mathbb{1}, \phi_v \rangle_{L_2(\Omega)} \approx \|\phi_v\|_{L_2(\Omega)}^2$, we find for $k \in \{0, 1\}$

$$\|P_{\mathcal{T}}u\|_{H^k(T)} \lesssim \sum_{v \in N_{\mathcal{T}}} \frac{\|\tilde{\phi}_v\|_{H^k(T)}}{\|\phi_v\|_{L_2(\Omega)}} \|u\|_{L_2(\text{supp } \phi_v)} \lesssim h_T^{-k} \|u\|_{L_2(\omega_{\mathcal{T}}(T))}, \tag{2.3}$$

with $\omega_{\mathcal{T}}(T) := \bigcup_{v \in N_{\mathcal{T}}} \text{supp } \phi_v$. This shows $\sup_{T \in \mathbb{T}} \|P_{\mathcal{T}}\|_{\mathcal{L}(L_2(\Omega), L_2(\Omega))} < \infty$.

From the above inequality, and $\sum_{v \in N_{\mathcal{T}}} \tilde{\phi}_v = \mathbb{1}$, we deduce that

$$\begin{aligned} \|(\text{Id} - P_{\mathcal{T}})u\|_{H^1(T)} &= \inf_{p \in \mathcal{P}_0} \|(\text{Id} - P_{\mathcal{T}})(u - p)\|_{H^1(T)} \\ &\lesssim \inf_{p \in \mathcal{P}_0} \|u - p\|_{H^1(T)} + h_T^{-1} \|u - p\|_{L_2(\omega_{\mathcal{T}}(T))} \\ &\lesssim \inf_{p \in \mathcal{P}_0} h_T^{-1} \|u - p\|_{L_2(\omega_{\mathcal{T}}(T))} + |u|_{H^1(T)} \\ &\lesssim |u|_{H^1(\omega_{\mathcal{T}}(T))}, \end{aligned}$$

with the last step following from the Bramble-Hilbert lemma. We conclude that $\sup_{T \in \mathbb{T}} \|P_{\mathcal{T}}\|_{\mathcal{L}(H^1(\Omega), H^1(\Omega))} < \infty$, and consequently by the Riesz-Thorin interpolation theorem, that

$$\sup_{T \in \mathbb{T}} \|P_{\mathcal{T}}\|_{\mathcal{L}(H^s(\Omega), H^s(\Omega))} < \infty.$$

This latter property guarantees that $\tilde{D}_{\mathcal{T}}$ is uniformly boundedly invertible:

$$\begin{aligned} \|\tilde{D}_{\mathcal{T}}\|_{\mathcal{L}(\mathcal{S}_{\mathcal{T},-s}, (\tilde{\mathcal{F}}_{\mathcal{T},s})')} &= \sup_{0 \neq u \in \mathcal{S}_{\mathcal{T},-s}} \sup_{0 \neq v \in \tilde{\mathcal{F}}_{\mathcal{T},s}} \frac{\langle u, v \rangle_{L_2(\Omega)}}{\|u\|_{H^{-s}(\Omega)} \|v\|_{H^s(\Omega)}} \leq 1, \\ \|\tilde{D}_{\mathcal{T}}^{-1}\|_{\mathcal{L}((\tilde{\mathcal{F}}_{\mathcal{T},s})', \mathcal{S}_{\mathcal{T},-s})}^{-1} &= \inf_{0 \neq u \in \mathcal{S}_{\mathcal{T},-s}} \sup_{0 \neq v \in \tilde{\mathcal{F}}_{\mathcal{T},s}} \frac{\langle u, v \rangle_{L_2(\Omega)}}{\|u\|_{H^{-s}(\Omega)} \|v\|_{H^s(\Omega)}} \\ &= \inf_{0 \neq u \in \mathcal{S}_{\mathcal{T},-s}} \sup_{0 \neq v \in H^s(\Omega)} \frac{\langle u, v \rangle_{L_2(\Omega)}}{\|u\|_{H^{-s}(\Omega)} \|P_{\mathcal{T}}v\|_{H^s(\Omega)}} \\ &\geq \|P_{\mathcal{T}}\|_{\mathcal{L}(H^s(\Omega), H^s(\Omega))}. \quad \square \end{aligned}$$

Theorem 2.3. For $I_{\mathcal{T}}: \mathcal{S}_{\mathcal{T},s} \rightarrow \tilde{\mathcal{F}}_{\mathcal{T},s}$ being the bijection given by $I_{\mathcal{T}}\phi_v = \tilde{\phi}_v$ ($v \in N_{\mathcal{T}}$), it holds that $I_{\mathcal{T}} \in \text{Lis}(\mathcal{S}_{\mathcal{T},s}, \tilde{\mathcal{F}}_{\mathcal{T},s})$ uniformly in $\mathcal{T} \in \mathbb{T}$.

Proof. Note that we may write

$$I_{\mathcal{T}}u = \sum_{v \in N_{\mathcal{T}}} \frac{\langle u, \tilde{\phi}_v \rangle_{L_2(\Omega)}}{\langle \phi_v, \tilde{\phi}_v \rangle_{L_2(\Omega)}} \tilde{\phi}_v \quad \text{and} \quad I_{\mathcal{T}}^{-1}u = \sum_{v \in N_{\mathcal{T}}} \frac{\langle u, \phi_v \rangle_{L_2(\Omega)}}{\langle \tilde{\phi}_v, \phi_v \rangle_{L_2(\Omega)}} \phi_v.$$

Equivalently to (2.3), we see for $k \in \{0, 1\}$ that

$$\|I_{\mathcal{T}}u\|_{H^k(T)} \lesssim \sum_{v \in N_{\mathcal{T}}} \frac{\|\tilde{\phi}_v\|_{H^k(T)} \|\tilde{\phi}_v\|_{L_2(\Omega)}}{\|\phi_v\|_{L_2(\Omega)}^2} \|u\|_{L_2(\text{supp } \phi_v)} \lesssim h_T^{-k} \|u\|_{L_2(\omega_{\mathcal{T}}(T))}.$$

Following the same arguments as in the proof of Theorem 2.2, using that $I_{\mathcal{T}}\mathbb{1} = \mathbb{1}$, then reveals that $I_{\mathcal{T}}$ is uniformly bounded. Uniformly boundedness of $I_{\mathcal{T}}^{-1}$ follows similarly. \square

As announced earlier, we define $D_{\mathcal{T}} \in \mathcal{L}(\mathcal{S}_{\mathcal{T},-s}, (\mathcal{S}_{\mathcal{T},s})')$ by $D_{\mathcal{T}} := I_{\mathcal{T}}^{-1} \tilde{D}_{\mathcal{T}}$, so $(D_{\mathcal{T}}u)(v) := \langle u, I_{\mathcal{T}}v \rangle_{L_2(\Omega)}$ ($u, v \in \mathcal{S}_{\mathcal{T}}$). Combining the previous theorems gives the following corollary.

Corollary 2.4. *The operator $D_{\mathcal{T}}$ is in $\text{Lis}(\mathcal{S}_{\mathcal{T},-s}, (\mathcal{S}_{\mathcal{T},s})')$ uniformly in $\mathcal{T} \in \mathbb{T}$.*

Remark 2.5. The matrix representation of $D_{\mathcal{T}}$ w.r.t. $\Phi_{\mathcal{T}}$ given by

$$\mathbf{D}_{\mathcal{T}} = \langle \Phi_{\mathcal{T}}, I_{\mathcal{T}} \Phi_{\mathcal{T}} \rangle_{L_2(\Omega)} = \text{diag}\{ \langle \mathbb{1}, \phi_{\nu} \rangle_{L_2(\Omega)} : \nu \in N_{\mathcal{T}} \},$$

which is *diagonal* and therefore easily invertible. The matrix $\mathbf{D}_{\mathcal{T}}$ is known as the *lumped mass matrix*.

Remark 2.6. The operator $D_{\mathcal{T}}$ depends merely on the *existence* of a biorthogonal basis $\tilde{\Phi}_{\mathcal{T}}$ that satisfies (2.2). Indeed, this basis does not appear in the implementation of $\mathbf{D}_{\mathcal{T}}$.

A possible construction of $\tilde{\Phi}_{\mathcal{T}}$ can be given using techniques from [13]. Consider some collection of local ‘bubble’ functions $\Theta_{\mathcal{T}} = \{ \theta_{\nu} \in H^1(\Omega) : \nu \in N_{\mathcal{T}} \}$ that satisfy: $|\langle \theta_{\nu}, \phi_{\nu'} \rangle_{L_2(\Omega)}| \approx \delta_{\nu\nu'} \|\phi_{\nu}\|_{L_2(\Omega)}^2$, $\|\theta_{\nu}\|_{H^k(\Omega)} \lesssim \|\phi_{\nu}\|_{H^k(\Omega)}$ ($k \in \{0, 1\}$), and $\text{supp } \theta_{\nu} \subseteq \text{supp } \phi_{\nu}$. Existence of such a collection can be shown by a construction on a reference d -simplex, and then using an affine bijection to transfer it to general elements, see [13, Sect. 4.1]. A suitable $\tilde{\Phi}_{\mathcal{T}}$ that satisfies (2.2) is then given by

$$\tilde{\phi}_{\nu} := \phi_{\nu} + \frac{\langle \mathbb{1}, \phi_{\nu} \rangle_{L_2(\Omega)}}{\langle \theta_{\nu}, \phi_{\nu} \rangle_{L_2(\Omega)}} \theta_{\nu} - \sum_{\nu' \in N_{\mathcal{T}}} \frac{\langle \phi_{\nu}, \phi_{\nu'} \rangle_{L_2(\Omega)}}{\langle \theta_{\nu'}, \phi_{\nu'} \rangle_{L_2(\Omega)}} \theta_{\nu'}.$$

We emphasize that the construction of a uniform preconditioner outlined in this subsection does not assume some sufficiently mild grading of the meshes.

2.2.1. Implementation

Taking $\Phi_{\mathcal{T}}$ as basis for both $\mathcal{S}_{\mathcal{T},-s}$ and $\mathcal{S}_{\mathcal{T},s}$, the *matrix representations* of the preconditioned systems from (2.1) read as

$$\mathbf{D}_{\mathcal{T}}^{-1} \mathbf{B}_{\mathcal{T}} \mathbf{D}_{\mathcal{T}}^{-\top} \mathbf{A}_{\mathcal{T}} \quad \text{and} \quad \mathbf{D}_{\mathcal{T}}^{-\top} \mathbf{A}_{\mathcal{T}} \mathbf{D}_{\mathcal{T}}^{-1} \mathbf{B}_{\mathcal{T}},$$

where

$$\begin{aligned} \mathbf{A}_{\mathcal{T}} &:= (A_{\mathcal{T}} \Phi_{\mathcal{T}})(\Phi_{\mathcal{T}}), & \mathbf{B}_{\mathcal{T}} &:= (B_{\mathcal{T}} \Phi_{\mathcal{T}})(\Phi_{\mathcal{T}}), \\ \mathbf{D}_{\mathcal{T}} &= \mathbf{D}_{\mathcal{T}}^{\top} := (D_{\mathcal{T}} \Phi_{\mathcal{T}})(\Phi_{\mathcal{T}}) = \text{diag}\{ \langle \mathbb{1}, \phi_{\nu} \rangle_{L_2(\Omega)} : \nu \in N_{\mathcal{T}} \}. \end{aligned}$$

Alternatively, we could equip the spaces with the *scaled* nodal basis $\check{\Phi}_{\mathcal{T}} := \mathbf{D}_{\mathcal{T}}^{-\frac{1}{2}} \Phi_{\mathcal{T}}$, so that the $L_2(\Omega)$ -norm of any basis function is proportional to 1, yielding

$$\begin{aligned} \check{\mathbf{A}}_{\mathcal{T}} &:= (A_{\mathcal{T}} \check{\Phi}_{\mathcal{T}})(\check{\Phi}_{\mathcal{T}}) = (\mathbf{D}_{\mathcal{T}}^{-\frac{1}{2}})^{\top} \mathbf{A}_{\mathcal{T}} \mathbf{D}_{\mathcal{T}}^{-\frac{1}{2}}, \\ \check{\mathbf{B}}_{\mathcal{T}} &:= (B_{\mathcal{T}} \check{\Phi}_{\mathcal{T}})(\check{\Phi}_{\mathcal{T}}) = (\mathbf{D}_{\mathcal{T}}^{-\frac{1}{2}})^{\top} \mathbf{B}_{\mathcal{T}} \mathbf{D}_{\mathcal{T}}^{-\frac{1}{2}}, \\ \check{\mathbf{D}}_{\mathcal{T}} &:= (D_{\mathcal{T}} \check{\Phi}_{\mathcal{T}})(\check{\Phi}_{\mathcal{T}}) = (\mathbf{D}_{\mathcal{T}}^{-\frac{1}{2}})^{\top} \mathbf{D}_{\mathcal{T}} \mathbf{D}_{\mathcal{T}}^{-\frac{1}{2}} = \mathbf{Id}, \end{aligned}$$

showing that $\check{\mathbf{B}}_{\mathcal{T}}$ is a uniform preconditioner for $\check{\mathbf{A}}_{\mathcal{T}}$ (and vice versa). To the best of our knowledge, so far this most easy form of operator preconditioning, where the stiffness matrix of some operator w.r.t. some basis is preconditioned by stiffness matrix of an opposite order operator w.r.t. the same basis, has not been shown to be optimal.

3. Manifold case

Let Γ be a compact d -dimensional Lipschitz, piecewise smooth manifold in $\mathbb{R}^{d'}$ for some $d' \geq d$ without boundary $\partial\Gamma$. For $s \in [0, 1]$, we consider the Sobolev spaces

$$H^s(\Gamma) := [L_2(\Gamma), H^1(\Gamma)]_{s,2}, \quad H^{-s}(\Gamma) := H^s(\Gamma)'$$

We assume that Γ is given as the closure of the disjoint union of $\cup_{i=1}^p \chi_i(\Omega_i)$, with, for $1 \leq i \leq p$, $\chi_i: \mathbb{R}^d \rightarrow \mathbb{R}^{d'}$ being some smooth regular parametrization, and $\Omega_i \subset \mathbb{R}^d$ an open polytope. W.l.o.g. assuming that for $i \neq j$, $\overline{\Omega}_i \cap \overline{\Omega}_j = \emptyset$, we define

$$\chi: \Omega := \cup_{i=1}^p \Omega_i \rightarrow \cup_{i=1}^p \chi_i(\Omega_i) \text{ by } \chi|_{\Omega_i} = \chi_i.$$

Let \mathbb{T} be a family of conforming partitions \mathcal{T} of Γ into ‘panels’ such that, for $1 \leq i \leq p$, $\chi^{-1}(\mathcal{T}) \cap \Omega_i$ is a uniformly shape regular conforming partition of Ω_i into d -simplices (that for $d = 1$ satisfies a uniform K -mesh property).

Fix $\ell > 0$, we set

$$\mathcal{S}_{\mathcal{T}} := \{ u \in H^1(\Gamma) : u \circ \chi|_{\chi^{-1}(T)} \in \mathcal{P}_{\ell} \text{ (} T \in \mathcal{T} \text{)}, \}$$

equipped with the canonical nodal basis $\Phi_{\mathcal{T}} = \{\phi_v : v \in N_{\mathcal{T}}\}$.

For construction of an operator $D_{\mathcal{T}} \in \text{Lis}(\mathcal{S}_{\mathcal{T},-s}, (\mathcal{S}_{\mathcal{T},s})')$ one can proceed as in the domain case. A suitable collection $\tilde{\Phi}_{\mathcal{T}}$ that is $L_2(\Gamma)$ -biorthogonal to $\Phi_{\mathcal{T}}$ exists. Moreover, the analysis from the domain case applies verbatim by only changing $\langle \cdot, \cdot \rangle_{L_2(\Omega)}$ into $\langle \cdot, \cdot \rangle_{L_2(\Gamma)}$. A *hidden problem*, however, is that the computation of $D_{\mathcal{T}} = \text{diag}\{\langle \mathbb{1}, \phi_v \rangle_{L_2(\Gamma)} : v \in N_{\mathcal{T}}\}$ involves integrals over Γ that generally have to be approximated using numerical quadrature.

In [13] we solved this issue by defining an additional ‘mesh-dependent’ scalar product

$$\langle u, v \rangle_{\mathcal{T}} := \sum_{T \in \mathcal{T}} \frac{|T|}{|\chi^{-1}(T)|} \int_{\chi^{-1}(T)} u(\chi(x))v(\chi(x))dx.$$

This is constructed by replacing on each $\chi^{-1}(T)$, the Jacobian $|\partial\chi|$ by its average $\frac{|T|}{|\chi^{-1}(T)|}$ over $\chi^{-1}(T)$.

By considering $\tilde{\Phi}_{\mathcal{T}}$ that is biorthogonal to $\Phi_{\mathcal{T}}$ with respect to $\langle \cdot, \cdot \rangle_{\mathcal{T}}$, and the linear bijection $I_{\mathcal{T}}$ given by $I_{\mathcal{T}}\phi_v = \tilde{\phi}_v$, one is able to show that the operator $D_{\mathcal{T}}$ defined as $(D_{\mathcal{T}}u)(v) := \langle u, I_{\mathcal{T}}v \rangle_{\mathcal{T}}$ satisfies the necessary requirements. For details we refer to [13]. The resulting matrix representation of $D_{\mathcal{T}}$ w.r.t. $\Phi_{\mathcal{T}}$ is then given by $D_{\mathcal{T}} = \text{diag}\{\langle \mathbb{1}, \phi_v \rangle_{\mathcal{T}} : v \in N_{\mathcal{T}}\}$.

4. Numerical results

Let $\Gamma = \partial[0, 1]^3 \subset \mathbb{R}^3$ be the two-dimensional manifold without boundary given as the boundary of the unit cube, $s = \frac{1}{2}$, and $\mathcal{S}_{\mathcal{T}}$ the space of continuous piecewise polynomials of degree ℓ w.r.t. a partition \mathcal{T} . We will evaluate preconditioning of the discretized Single Layer Integral operator $A_{\mathcal{T}} \in \text{Lis}_c(\mathcal{S}_{\mathcal{T},-s}, (\mathcal{S}_{\mathcal{T},s})')$ and an (essentially) discretized Hypersingular Integral operator $B_{\mathcal{T}} \in \text{Lis}_c(\mathcal{S}_{\mathcal{T},s}, (\mathcal{S}_{\mathcal{T},s})')$.

The Hypersingular Integral operator $\tilde{B} \in \mathcal{L}(H^{\frac{1}{2}}(\Gamma), H^{-\frac{1}{2}}(\Gamma))$, is only-semi coercive, but solving $\tilde{B}u = f$ for f with $f(\mathbb{1}) = 0$ is equivalent to solving $Bu = f$ with B given by $(Bu)(v) = (\tilde{B}u)(v) + \alpha \langle u, \mathbb{1} \rangle_{L_2(\Gamma)} \langle v, \mathbb{1} \rangle_{L_2(\Gamma)}$, for some fixed $\alpha > 0$. This operator B is in $\text{Lis}_c(H^{\frac{1}{2}}(\Gamma), H^{-\frac{1}{2}}(\Gamma))$, and we shall consider discretizations $B_{\mathcal{T}} \in \text{Lis}_c(\mathcal{S}_{\mathcal{T},s}, (\mathcal{S}_{\mathcal{T},s})')$ of B . We found $\alpha = 0.05$ to give good results in our examples.

Equipping both $\mathcal{S}_{\mathcal{T},s}$ and $\mathcal{S}_{\mathcal{T},-s}$ with the standard nodal basis $\Phi_{\mathcal{T}} = \{\phi_v : v \in N_{\mathcal{T}}\}$, the matrix representations of the preconditioned systems from Sect. 2.2 read as

$$D_{\mathcal{T}}^{-1}B_{\mathcal{T}}D_{\mathcal{T}}^{-\top}A_{\mathcal{T}} \quad \text{and} \quad D_{\mathcal{T}}^{-\top}A_{\mathcal{T}}D_{\mathcal{T}}^{-1}B_{\mathcal{T}},$$

for $D_{\mathcal{T}} = \text{diag}\{\langle \mathbb{1}, \phi_v \rangle_{L_2(\Gamma)} : v \in N_{\mathcal{T}}\}$, $A_{\mathcal{T}} = (A_{\mathcal{T}}\Phi_{\mathcal{T}})(\Phi_{\mathcal{T}})$ and $B_{\mathcal{T}} := (B_{\mathcal{T}}\Phi_{\mathcal{T}})(\Phi_{\mathcal{T}})$.

We calculated (spectral) condition numbers of these preconditioned systems, where this condition number is given by $\kappa_S(\mathbf{X}) := \rho(\mathbf{X})\rho(\mathbf{X}^{-1})$ with $\rho(\cdot)$ denoting the spectral radius. Note that the condition numbers of the preconditioned systems coincide, i.e.,

$$\kappa_S(D_{\mathcal{T}}^{-1}B_{\mathcal{T}}D_{\mathcal{T}}^{-\top}A_{\mathcal{T}}) = \kappa_S(D_{\mathcal{T}}^{-\top}A_{\mathcal{T}}D_{\mathcal{T}}^{-1}B_{\mathcal{T}}),$$

so we may restrict ourselves to results for preconditioning of $A_{\mathcal{T}}$.

We used the BEM++ software package [10] to approximate the application of the matrices $A_{\mathcal{T}}$ and $B_{\mathcal{T}}$ by hierarchical matrices based on adaptive cross approximation [7,2]. In particular, we assemble neither matrix $A_{\mathcal{T}}$ nor $B_{\mathcal{T}}$.

As initial partition $\mathcal{T}_{\perp} = \mathcal{T}_1$ of Γ we take a conforming partition consisting of 2 triangles per side, so 12 triangles in total, with an assignment of the newest vertices that satisfies the so-called matching condition. We let \mathbb{T} be the sequence $\{\mathcal{T}_k\}_{k \geq 1}$ where the (conforming) partition \mathcal{T}_k is found by applying both uniform and local refinements. To be precise, \mathcal{T}_k is constructed by first applying k uniform bisections to \mathcal{T}_{\perp} , and then $4k$ local refinements by repeatedly applying NVB to all triangles that touch a corner of the cube. These partitions share both the difficulties of locally refined partitions (the presence of triangles with strongly different sizes) and that of uniform partitions (the diagonally scaled stiffness matrix has a condition number $\gtrsim 2^{k|s|}$).

4.1. Comparison preconditioners

Write $G_{\mathcal{T}}^D := D_{\mathcal{T}}^{-1}B_{\mathcal{T}}D_{\mathcal{T}}^{-\top}$ for the preconditioner constructed in Sect. 2.2. We will compare this with the preconditioner described in Sect. 2.1.1, for which the matrix representation is given by $G_{\mathcal{T}}^M := M_{\mathcal{T}}^{-1}B_{\mathcal{T}}M_{\mathcal{T}}^{-\top}$ with mass matrix $M_{\mathcal{T}} = M_{\mathcal{T}}^{\top} = \langle \Phi_{\mathcal{T}}, \Phi_{\mathcal{T}} \rangle_{L_2(\Gamma)}$. Because our partitions of the two-dimensional surface are created with NVB, we know that also the latter preconditioner provides uniformly bounded condition numbers. In contrast to $D_{\mathcal{T}}^{-1}$, the inverse $M_{\mathcal{T}}^{-1}$ cannot be evaluated in linear complexity. We implemented the application of $M_{\mathcal{T}}^{-1}$ by computing an LU-factorization of $M_{\mathcal{T}}$.

Table 1 compares the spectral condition numbers for the preconditioned Single Layer systems with trial spaces given by continuous piecewise linears and those by continuous piecewise cubics. The condition numbers $\kappa_S(G_{\mathcal{T}}^D A_{\mathcal{T}})$ are uniformly bounded, but quantitatively the condition numbers $\kappa_S(G_{\mathcal{T}}^M A_{\mathcal{T}})$ are better.

For completeness, despite the small increase of $\kappa_S(G_{\mathcal{T}}^M A_{\mathcal{T}})$ when going from linear to cubic finite elements, there is no indication that with either $G_{\mathcal{T}}^D$ or $G_{\mathcal{T}}^M$ the condition numbers of the preconditioned system are not only uniformly bounded in the partition but also in the polynomial degree.

Table 1

Spectral condition numbers, $\kappa_S(\mathbf{G}_{\mathcal{T}}^{\circ} \mathbf{A}_{\mathcal{T}})$ for $\circ \in \{D, M\}$, of the preconditioned Single Layer system discretized on $\{\mathcal{T}_k\}_{k \geq 1}$, by continuous piecewise linears ($\ell = 1$) in the middle columns and discretized by continuous piecewise cubics ($\ell = 3$) in the right columns. Here $\mathbf{G}_{\mathcal{T}}^D$ is the preconditioner introduced in Sect. 2.2, whereas $\mathbf{G}_{\mathcal{T}}^M$ is the preconditioner described in Sect. 2.1.1 whose application requires an application of $\mathbf{M}_{\mathcal{T}}^{-1}$, which we implemented using an LU-factorization.

Partition \mathcal{T}		Linears ($\ell = 1$)			Cubics ($\ell = 3$)		
h_{\min}	h_{\max}	dofs	$\mathbf{G}_{\mathcal{T}}^D \mathbf{A}_{\mathcal{T}}$	$\mathbf{G}_{\mathcal{T}}^M \mathbf{A}_{\mathcal{T}}$	dofs	$\mathbf{G}_{\mathcal{T}}^D \mathbf{A}_{\mathcal{T}}$	$\mathbf{G}_{\mathcal{T}}^M \mathbf{A}_{\mathcal{T}}$
$1.4 \cdot 10^0$	$1.4 \cdot 10^0$	8	16.2	1.20	56	90.5	1.68
$4.4 \cdot 10^{-2}$	$5.0 \cdot 10^{-01}$	218	14.9	1.91	1946	87.9	2.08
$1.3 \cdot 10^{-3}$	$3.5 \cdot 10^{-01}$	482	14.7	2.04	4322	86.1	2.17
$4.3 \cdot 10^{-5}$	$1.7 \cdot 10^{-01}$	962	14.7	2.10	8642	85.0	2.21
$1.3 \cdot 10^{-6}$	$8.8 \cdot 10^{-02}$	2306	15.4	2.14	20738	84.9	2.23
$4.2 \cdot 10^{-8}$	$4.4 \cdot 10^{-02}$	7106	15.6	2.16	63938	84.9	2.24
$1.3 \cdot 10^{-9}$	$2.2 \cdot 10^{-02}$	25730	15.8	2.17	231554	84.8	2.25
$4.1 \cdot 10^{-11}$	$1.1 \cdot 10^{-02}$	99650	15.8	2.17	896834	84.7	2.25

Table 2

Spectral condition numbers $\kappa_S(\mathbf{G}_{\mathcal{T}}^{(k)} \mathbf{A}_{\mathcal{T}})$ with $\mathbf{G}_{\mathcal{T}}^{(k)}$ the preconditioner from (4.1) that incorporates k Richardson iterations. The systems are discretized by continuous piecewise linears in the left columns and discretized by continuous piecewise cubics in the right columns.

Linears ($\ell = 1$)				Cubics ($\ell = 3$)			
dofs	$k = 2$	$k = 4$	$k = 6$	dofs	$k = 2$	$k = 4$	$k = 6$
8	2.26	1.29	1.22	56	10.1	3.99	2.65
218	3.05	2.07	1.94	1946	8.96	3.57	2.52
482	3.53	2.28	2.08	4322	8.80	3.59	2.52
962	3.79	2.44	2.19	8642	8.63	3.59	2.52
2306	3.98	2.52	2.24	20738	8.54	3.59	2.52
7106	4.18	2.57	2.27	63938	8.54	3.59	2.52
25730	4.35	2.61	2.28	231554	8.54	3.59	2.52
99650	4.47	2.65	2.29	896834	8.54	3.59	2.52

4.2. Improving the preconditioner quality

As observed in Table 1, the preconditioner $\mathbf{G}_{\mathcal{T}}^M$ appears to be of superior quality, but it has unfavourable computational complexity. It does suggest a way for improving $\mathbf{G}_{\mathcal{T}}^D$: by replacing $\mathbf{D}_{\mathcal{T}}^{-1}$ with a better approximation of $\mathbf{M}_{\mathcal{T}}^{-1}$, one may hope to improve the quality. To this end, we introduce damped (preconditioned) Richardson. Let $0 < \lambda_- \leq \lambda_{\min}(\mathbf{D}_{\mathcal{T}}^{-1} \mathbf{M}_{\mathcal{T}})$, $\lambda_{\max}(\mathbf{D}_{\mathcal{T}}^{-1} \mathbf{M}_{\mathcal{T}}) \leq \lambda_+$, $\mathbf{R}_{\mathcal{T}}^{(0)} := \mathbf{0}$ and for $k \geq 0$ define

$$\mathbf{R}_{\mathcal{T}}^{(k+1)} := \mathbf{R}_{\mathcal{T}}^{(k)} + \omega \mathbf{D}_{\mathcal{T}}^{-1} (\text{Id} - \mathbf{M}_{\mathcal{T}} \mathbf{R}_{\mathcal{T}}^{(k)}), \quad \omega = \frac{2}{\lambda_- + \lambda_+},$$

being the result of k Richardson iterations. Correspondingly define

$$\mathbf{G}_{\mathcal{T}}^{(k)} := \mathbf{R}_{\mathcal{T}}^{(k)} \mathbf{B}_{\mathcal{T}} \mathbf{R}_{\mathcal{T}}^{(k)}. \tag{4.1}$$

It follows that $\mathbf{G}_{\mathcal{T}}^{(1)} = \mathbf{G}_{\mathcal{T}}^D$ and $\lim_{k \rightarrow \infty} \mathbf{G}_{\mathcal{T}}^{(k)} = \mathbf{G}_{\mathcal{T}}^M$. Although we have no proof, we suspect that $\mathbf{G}_{\mathcal{T}}^{(k)}$ provides a uniform preconditioner for $\mathbf{A}_{\mathcal{T}}$ due to the fact that $\mathbf{R}_{\mathcal{T}}^{(k)}$ approximates $\mathbf{M}_{\mathcal{T}}^{-1}$, while preserving constant functions, being a key ingredient in the proofs of Theorems 2.2 and 2.3.

Values for λ_- and λ_+ can be found by calculating the extremal eigenvalues of the corresponding preconditioned mass matrix on a reference simplex, see e.g. [15]. For $\ell = 1$ this gives $\omega = \frac{2(d+2)}{d+3}$, whereas for $\ell = 3$ and $d = 2$ we computed $\omega = 0.836$.

Table 2 compares the condition numbers $\kappa_S(\mathbf{G}_{\mathcal{T}}^{(k)} \mathbf{A}_{\mathcal{T}})$ for $k \in \{2, 4, 6\}$. We see that a few Richardson iterations drastically improve our preconditioner, making its quality on par with that of $\mathbf{G}_{\mathcal{T}}^M$ while having a favourable linear application cost.

Finally, to show that one cannot simply use any (iterative) method for approximating $\mathbf{M}_{\mathcal{T}}^{-1}$, we consider the case where one approximates this inverse using a Jacobi preconditioner. The resulting preconditioner is then given by

$$\mathbf{G}_{\mathcal{T}}^{\text{lac}} := (\text{diag } \mathbf{M}_{\mathcal{T}})^{-1} \mathbf{B}_{\mathcal{T}} (\text{diag } \mathbf{M}_{\mathcal{T}})^{-\top}. \tag{4.2}$$

Table 3 clearly displays that this is not a uniformly bounded preconditioner, which we assume is due to the fact that $(\text{diag } \mathbf{M}_{\mathcal{T}})^{-1}$ does not preserve constant functions for $\ell > 1$.

Table 3
Spectral condition numbers $\kappa_S(\mathbf{G}_{\mathcal{T}}^{\text{jac}} \mathbf{A}_{\mathcal{T}})$ with $\mathbf{G}_{\mathcal{T}}^{\text{jac}}$ from (4.2), and systems discretized by continuous piecewise cubics ($\ell = 3$).

dofs	$\mathbf{G}_{\mathcal{T}}^{\text{jac}} \mathbf{A}_{\mathcal{T}}$
56	62.6
1946	377.1
4322	495.6
8642	1016.9
20738	3067.8
63938	10928.3

5. Conclusion

Considering discretized opposite order operators $\mathbf{A}_{\mathcal{T}}$ and $\mathbf{B}_{\mathcal{T}}$ using the same ansatz space of continuous piecewise polynomial w.r.t. a possibly locally refined partition \mathcal{T} , we consider matrices $\mathbf{D}_{\mathcal{T}}$ such that $\mathbf{D}_{\mathcal{T}}^{-1} \mathbf{B}_{\mathcal{T}} \mathbf{D}_{\mathcal{T}}^{-\top}$ is a uniform preconditioner for $\mathbf{A}_{\mathcal{T}}$, and $\mathbf{D}_{\mathcal{T}}^{-\top} \mathbf{A}_{\mathcal{T}} \mathbf{D}_{\mathcal{T}}^{-1}$ for $\mathbf{B}_{\mathcal{T}}$. The obvious choice for $\mathbf{D}_{\mathcal{T}}$ would be the mass matrix, however, it yields uniformly bounded condition numbers only under a mildly grading assumption on the mesh, and more importantly, it has the disadvantage that its inverse is dense. We proved that when taking $\mathbf{D}_{\mathcal{T}}$ as the lumped mass matrix the condition numbers are uniformly bounded, remarkably without a sufficiently mild grading assumption on the mesh, while obviously its inverse can be applied in linear cost.

In our experiments with locally refined meshes generated by Newest Vertex Bisection, the condition numbers with $\mathbf{D}_{\mathcal{T}}$ being the mass matrix are quantitatively better than those found with $\mathbf{D}_{\mathcal{T}}$ being the lumped mass matrix though. Constructing $\mathbf{D}_{\mathcal{T}}^{-1}$ as an approximation for the inverse mass matrix by a few preconditioned damped Richardson steps with the lumped mass matrix as a preconditioner, both the resulting matrix can be applied at linear cost and the observed condition numbers are essentially as good as with the inverse mass matrix.

References

- [1] R.E. Bank, H. Yserentant, On the H^1 -stability of the L_2 -projection onto finite element spaces, *Numer. Math.* 126 (2) (2014) 361–381.
- [2] M. Bebendorf, Approximation of boundary element matrices, *Numer. Math.* 86 (4) (2000) 565–589.
- [3] A. Buffa, S.H. Christiansen, A dual finite element complex on the barycentric refinement, *Math. Comput.* 76 (260) (2007) 1743–1769.
- [4] C. Carstensen, Merging the Bramble-Pasciak-Steinbach and the Crouzeix-Thomée criterion for H^1 -stability of the L^2 -projection onto finite element spaces, *Math. Comput.* 71 (237) (2002) 157–163.
- [5] L. Diening, J. Storn, T. Tscherpel, On the Sobolev and L^p -stability of the L^2 -projection, 2020.
- [6] F.D. Gaspoz, C.-J. Heine, K.G. Siebert, Optimal grading of the newest vertex bisection and H^1 -stability of the L_2 -projection, *IMA J. Numer. Anal.* 36 (3) (2016) 1217–1241.
- [7] W. Hackbusch, A sparse matrix arithmetic based on \mathcal{H} -matrices. Part I: introduction to \mathcal{H} -matrices, *Computing* 62 (2) (1999) 89–108.
- [8] R. Hiptmair, Operator preconditioning, *Comput. Math. Appl.* 52 (5) (2006) 699–706.
- [9] R. Hiptmair, C. Jerez-Hanckes, C. Urzúa-Torres, Optimal operator preconditioning for Galerkin boundary element methods on 3-dimensional screens, *SIAM J. Numer. Anal.* 58 (1) (2020) 834–857.
- [10] W. Śmigaj, T. Betcke, S. Arridge, J. Phillips, M. Schweiger, Solving boundary integral problems with BEM++, *ACM Trans. Math. Softw. (TOMS)* 41 (2) (2015) 1–40.
- [11] O. Steinbach, On a generalized L_2 projection and some related stability estimates in Sobolev spaces, *Numer. Math.* 90 (4) (2002) 775–786.
- [12] O. Steinbach, W.L. Wendland, The construction of some efficient preconditioners in the boundary element method, *Adv. Comput. Math.* 9 (1–2) (1998) 191–216, Numerical treatment of boundary integral equations.
- [13] R.P. Stevenson, R. van Venetië, Uniform preconditioners for problems of negative order, *Math. Comput.* (2019).
- [14] R.P. Stevenson, R. van Venetië, Uniform preconditioners for problems of positive order, *Comput. Math. Appl.* 79 (12) (2020) 3516–3530.
- [15] A.J. Wathen, Realistic eigenvalue bounds for the Galerkin mass matrix, *IMA J. Numer. Anal.* 7 (4) (1987) 449–457.