

Draft Genome Sequences of *Dickeya* sp. Isolates B16 (NIB Z 2098) and S1 (NIB Z 2099) Causing Soft Rot of *Phalaenopsis* Orchids

Špela Alič,^{a,b} Tina Naglič,^{a,c*} Pablo Llop,^d Nataša Toplak,^e Simon Koren,^e Maja Ravnikar,^{a,c} Tanja Dreo^{a,c}

National Institute of Biology, Ljubljana, Slovenia^a; Jožef Stefan International Postgraduate School, Ljubljana, Slovenia^b; Centre of Excellence for Biosensors, Instrumentation and Process Control (COBIK), Laboratory for Bioanalytics (LBA), Ajdovščina, Slovenia^c; Department of Evolutionary Genetics, Institute Cavanilles, University of Valencia, Paterna, Valencia, Spain^d; OMEGA d.o.o., Ljubljana, Slovenia^e

* Present address: Tina Naglič, Institute of Microbiology and Immunology, Faculty of Medicine, University of Ljubljana, Ljubljana, Slovenia.

The genus *Dickeya* contains bacteria causing soft rot of economically important crops and ornamental plants. Here, we report the draft genome sequences of two *Dickeya* sp. isolates from rotted leaves of *Phalaenopsis* orchids.

Received 22 July 2015 Accepted 31 July 2015 Published 10 September 2015

Citation Alič Š, Naglič T, Llop P, Toplak N, Koren S, Ravnikar M, Dreo T. 2015. Draft genome sequences of *Dickeya* sp. isolates B16 (NIB Z 2098) and S1 (NIB Z 2099) causing soft rot of *Phalaenopsis* orchids. *Genome Announc* 3(5):e00973-15 doi:10.1128/genomeA.00973-15.

Copyright © 2015 Alič et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](https://creativecommons.org/licenses/by/3.0/).

Address correspondence to Tanja Dreo, tanja.dreo@nib.si.

Bacteria of the *Dickeya* genus of the family *Enterobacteriaceae* cause diseases on many plant species (1) and are currently divided into seven species: *D. dianthicola*, *D. dadantii*, *D. zea*, *D. chrysanthemi*, *D. paradisiaca* (2, 3), *D. solani* (4), and *D. aquatica* (5). Several *Dickeya* sp. isolates form distinct clades and may represent novel species (1, 6, 7). Here, we report the draft sequences of two strains, *Dickeya* sp. B16 (NIB Z 2098) and S1 (NIB Z 2099), belonging to the undefined lineages UDL-3 and UDL-4 (1), which were identified as the primary cause of soft rot of *Phalaenopsis* orchids.

Both strains were purified by dilution plating. Genomic DNA was extracted using the UltraClean microbial DNA isolation kit (Mo Bio Laboratories, Inc., Carlsbad, CA) and concentrated to an approximate concentration of 500 ng/μl (GeneVac EZ-2plus; Genevac, Ipswich, United Kingdom), as determined with the Qubit double-stranded DNA (dsDNA) HR assay (Thermo Fisher Scientific, Life Technologies, Carlsbad, CA). Two single-read libraries were constructed with the Ion Xpress Plus fragment library kit (Life Technologies), according to the manufacturer's protocol (8), with modifications: (i) 15 min DNA fragmentation time, (ii) six cycles in the library amplification, and (iii) library purification with 340 μl of Agencourt AMPure XP reagent (Beckman Coulter, Inc., Brea, CA). Fragments of approximately 150 to 200 bp were selected (E-Gel SizeSelect 2%; Life Technologies). Emulsion PCR was done with the Ion PGM template OT2 200 kit using the Ion OneTouch 2 system (both Life Technologies), according to the manufacturer's protocol (9), with a final library concentration of 13 pM. The quality of the libraries was checked with the Caliper LabChip GX using the DNA high-sensitivity assay (PerkinElmer, Waltham, MA). Ion 316 Chip version 2 was used to sequence each library on the Ion Torrent Personal Genome Machine (PGM) sequencer with the Ion PGM sequencing 200 kit version 2 (all Life Technologies).

Sequences in FASTQ format were quality checked, and adapters were trimmed using CLC Genomics Workbench version 7.5 (Aarhus, Denmark), with the following modifications: (i) se-

quence quality score was set at 0.1, (ii) eight nucleotides were trimmed off every read at the 5' terminal end, and (iii) reads of <15 nucleotides were discarded. Trimmed reads were independently assembled with CLC Genomics Workbench version 7.5 and the GS *de novo* assembler (454 Life Sciences, Roche Company, Branford, CT). Both assemblies were processed using standard program settings with minor modifications in the GS *de novo* assembler: (i) the all-contig threshold was set to 200 bp, and (ii) the long-contig threshold was changed to 1,000 bp. Assembled contigs were integrated using CISA from the Orione (Galaxy/CRS4) portal (<http://orione.crs4.it/>). Contigs with average coverage of <10×, contigs <1,000 bp, and all contaminants identified with BLASTn were discarded. The draft genomes comprise 61 and 59 contigs, with a total size of 4,885,657 bp and 4,892,969 bp and an N_{50} of 167,592 bp and 160,162 bp for strains B16 and S1, respectively.

Nucleotide sequence accession numbers. The draft sequences of these two *Dickeya* strains are available in GenBank under the accession numbers [JXBN000000000](https://ncbi.nlm.nih.gov/assembly/GCA015133.1) (BioProject PRJNA268269) and [JXBO000000000](https://ncbi.nlm.nih.gov/assembly/GCA015133.1) (BioProject PRJNA268270) for B16 and S1, respectively. The versions described in this paper are the first versions, [JXBN010000000](https://ncbi.nlm.nih.gov/assembly/GCA015133.1) and [JXBO010000000](https://ncbi.nlm.nih.gov/assembly/GCA015133.1).

ACKNOWLEDGMENTS

This work was supported by the Slovenian Research Agency (contracts P4-0165, L7-5534, and 1000-15-0105) and by the Euphresco Phytosanitary ERA-NET project on *Dickeya* and *Pectobacterium* species. The Centre of Excellence for Biosensors, Instrumentation and Process Control is an operation financed by the European Union, European Regional Development Fund and Republic of Slovenia, Ministry of Higher Education, Science and Technology.

We thank Tomaž Jevšnik, Ocean Orchids, for plant material and Denis Kutnjak, NIB, for help with genome analysis.

REFERENCES

1. Van Vaerenbergh J, Baeyen S, De Vos P, Maes M. 2012. Sequence diversity in the *Dickeya* *fliC* gene: phylogeny of the *Dickeya* genus and Taq-

- Man PCR for “*D. solani*”, new biovar 3 variant on potato in Europe. PLoS One 7:e35738. <http://dx.doi.org/10.1371/journal.pone.0035738>.
2. Samson R, Legendre JB, Christen R, Fischer-Le Saux M, Achouak W, Gardan L. 2005. Transfer of *Pectobacterium chrysanthemi* (Burkholder et al. 1953) Brenner et al. 1973 and *Brenneria paradisiaca* to the genus *Dickeya* gen. nov. as *Dickeya chrysanthemi* comb. nov. and *Dickeya paradisiaca* comb. nov. and delineation of four novel species, *Dickeya dadantii* sp. nov., *Dickeya dianthicola* sp. nov., *Dickeya dieffenbachiae* sp. nov. and *Dickeya zeae* sp. nov. Int J Syst Evol Microbiol 55:1415–1427. <http://dx.doi.org/10.1099/ijs.0.02791-0>.
 3. Brady CL, Cleenwerck I, Denman S, Venter SN, Rodríguez-Palenzuela P, Coutinho TA, De Vos P. 2012. Proposal to reclassify *Brenneria quercina* (Hildebrand and Schroth 1967) Hauben et al. 1999 into a new genus, *Lonsdalea* gen. nov., as *Lonsdalea quercina* comb. nov., descriptions of *Lonsdalea quercina* subsp. *quercina* comb. nov., *Lonsdalea quercina* subsp. *iberica* subsp. nov. and *Lonsdalea quercina* subsp. *britannica* subsp. nov., emendation of the description of the genus *Brenneria*, reclassification of *Dickeya dieffenbachiae* as *Dickeya dadantii* subsp. *dieffenbachiae* comb. nov., and emendation of the description of *Dickeya dadantii*. Int J Syst Evol Microbiol 62:1592–1602. <http://dx.doi.org/10.1099/ijs.0.035055-0>.
 4. van der Wolf JM, Nijhuis EH, Kowalewska MJ, Saddler GS, Parkinson N, Elphinstone JG, Pritchard L, Toth IK, Lojkowska E, Potrykus M, Waleron M, de Vos P, Cleenwerck I, Pirhonen M, Garlant L, Hélias V, Pothier JF, Pflüger V, Duffy B, Tsrör L, Manulis S. 2014. *Dickeya solani* sp. nov., a pectinolytic plant-pathogenic bacterium isolated from potato (*Solanum tuberosum*). Int J Syst Evol Microbiol 64:768–774. <http://dx.doi.org/10.1099/ijs.0.052944-0>.
 5. Parkinson N, DeVos P, Pirhonen M, Elphinstone J. 2014. *Dickeya aquatica* sp. nov., isolated from waterways. Int J Syst Evol Microbiol 64: 2264–2266. <http://dx.doi.org/10.1099/ijs.0.058693-0>.
 6. Sławiak M, van Beckhoven JRCM, Speksnijder AGCL, Czajkowski R, Grabe G, van der Wolf JM. 2009. Biochemical and genetical analysis reveal a new clade of biovar 3 *Dickeya* spp. strains isolated from potato in Europe. Eur J Plant Pathol 125:245–261. <http://dx.doi.org/10.1007/s10658-009-9479-2>.
 7. Parkinson N, Stead D, Bew J, Heeney J, Tsrör Lahkim L, Elphinstone J. 2009. *Dickeya* species relatedness and clade structure determined by comparison of *recA* sequences. Int J Syst Evol Microbiol 59:2388–2393.
 8. Life Technologies. 2014. Ion Xpress Plus gDNA Fragment Library Preparation. MA N0009847 Rev. C.0. Life Technologies, Carlsbad, CA.
 9. Life Technologies. 2013. Ion PGM Template OT2 200 Kit. MA N0007220 Rev. 5.0. Life Technologies, Carlsbad, CA.