

European perspectives on regional estimates of standing water bodies and the relevance of man-made ponds

Jaanus Terasmaa, Pascal Bartout, Agata Marzecova, Laurent Touchart, Tiiu Koff, Quentin Choffel, Galina Kapanen, Véronique Maleval, Camille Millot, Zoubida Qsair, et al.

► To cite this version:

Jaanus Terasmaa, Pascal Bartout, Agata Marzecova, Laurent Touchart, Tiiu Koff, et al.. European perspectives on regional estimates of standing water bodies and the relevance of manmade ponds. European Geosciences Union, General Assembly, Apr 2015, Vienne, Austria. <hr/> <hal-01147760>

HAL Id: hal-01147760 https://hal.archives-ouvertes.fr/hal-01147760

Submitted on 1 May 2015


HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Introduction

- The majority of terrestrial standing water bodies (SWB) are small in size, however, their abundance and distribution is not fully known and they are under-represented in legislation. For example, the EU Water Framework Directive 2000/60/EC proposes a threshold surface area for reporting as 50 ha.
- Small SWB can be well represented in the state-level inventories and studies, e.g. <10 ha in Meybeck, 1995, <1 ha in Rjanžin 2005, or <0.05 ha in Kuusisto & Raatikainen 1988, but these studies are region-specific and not suitable for estimating global inventories. Also, there is a bias towards studies of natural lakes while abundance of man-made water bodies is understudied.
- The models for global inventories of SWB are so far not sufficiently designed for estimating the relative abundance of small SWB (below 0.2 ha) and provide differing estimates (Downing et al. 2006; Verpoorter et al. 2014).
- In this pilot study, we : (1) suggest a bottom-up approach for estimating the number of SWB at EU-level that combines the ground-validated data on water bodies from state inventories and data from peer-generated map databases (Bartout & Touchart, 2013). (2) assess the inventories and relative importance of small terrestrial water bodies of two different countries, Estonia and France.

Geographical and historical settings




Estonia (45 215 km²) belongs to the Boreal region of Europe. The surface topography is flat (60% of the territory is < 50 m

The majority of lake basins formed during the ice retreat after the Last Glacial Maximum. Besides forests (50%), wetlands are the most common

landscape feature covering 20% of the

territory. Peat-rich mires with characteristic bog pools are abundant. Creation of man-made lakes and ponds has been relatively common throughout the centuries. The man-made water bodies serve range of ecological, cultural and agricultural purposes.



EU territory of France (547 030 km²) possesses a wide variety of coastal, lowland and mountain landscapes.

Except few mountain margins, France was not glaciated during the Würm period.

Forests account for 28% of land area. The territory has extensive river systems (Seine, Loire, Garonne and Rhone).

Most of the water bodies are artificial (~98%), smaller than Nordic lakes and built on fluvial continuum. The tradition of creating the fishponds was developed in Middle Ages and today they provide variety of agricultural functions. Historically, France has also a tradition of forming artificial wetlands.


References

Verpoorter et al. (2014) Geophysical Research Letters, 41. uchart (2013) Annales de Géographie, 691. epted) Norois. Bartout (ac Downing et al. (2006) Limnology and Oceanography, 51(5). Kuusisto & Raatikainen (1988) Terra, 102. Meybeck (1995) in Lerman et al., Physics and chemistry of lakes. Rjanžin (2005) Priroda, 4 Estonian Topographic Database 2014




Download poster as pdf.

Abundance and size distribution



Shoreline lenght VS area

The comparison of size classes of SWB in Estonian Topographic Database (ETD) and French national inventory. The green color indicates the calculated total shoreline lenght (perimeter) and blue color indicates the total area.



Natural VS man-made

In France, majority (98-99%) of SWB are man-made. In Estonia, 56% of total inventory of SWB are man-made and 44% natural SWB.




Table shows the total inventory of SWB in Estonia divided in size ranges.


	Perimeter, km		Area, ha		Сог
	Nat.	Mm.	Nat.	Mm.	Nat.
<0.01 ha	291	241	60	51	8 712
0.01-0.1 ha	2 443	3 076	973	1 386	31 791
0.1-1 ha	1 053	2 607	866	2 421	3 450
1-10 ha	976	1 213	3 085	2 857	963
10-100 ha	766	500	6 268	2 759	223
>100 ha*	626	305	38 032	12 017	45
Sum	6 156	7 942	49 284	21 490	45 184
Min.	0.02	0.01	0.00	0.00	-
Max.	130	211	26 901	10 553	-
Median	0.06	0.07	0.02	0.03	-
Mean	0.14	0.14	1.09	0.37	-

Acknowledgments

This project has been financially supported by the French-Estonian research grant program Parrot nr 29502WC. We thank also © OpenStreetMap contributors.

European perspectives on regional estimates of standing water bodies and the relevance of man-made ponds Jaanus Terasmaa⁽¹⁾, Pascal Bartout⁽²⁾, Agata Marzecova⁽¹⁾, Laurent Touchart⁽²⁾, Tiiu Koff⁽¹⁾, Quentin Choffel⁽²⁾, Galina Kapanen⁽¹⁾ Véronique Maleval⁽²⁾, Camille Millot⁽²⁾, Zoubida Qsair⁽²⁾, Egert Vandel⁽¹⁾

The distribution of the SWB in Estonia and France. Maps are showing SWB abundance in different size ranges. Every dot on the map represents single SWB. The number in corners indicates total counts. Pie-plots are summarising the findings.



Only ~ 1500 of natural water bodies are lakes (mean surface area – 30 ha excluding L. Peipsi), remaining SWB are small bog pools (mean surface area – 0.03 ha).

SWB at EU scale

50.6%

According to the analysis of the OpenStreetmap (OSM) database (Partout*, accepted*) the number os SWB in EU is 1 041 739. This number covers in average over 80% of SWB in the size class >10 ha, but it is strongly underestimated in smaller size

OpenStreetMap (OSM) is a map database built by a community of mappers that contribute and maintain data.

⁽¹⁾ Institute of Ecology at Tallinn University, Estonia ⁽²⁾ Laboratoire de géographie CEDETE de l'Université d'Orléans, France Contact: jaanus.terasmaa@tlu.ee; pascal.bartout@univ-orleans.fr

Conclusions

- In Estonia the density of the waterbodies is 2.3 per km2 and in France 1.1 per km2. The median size of standing water bodies (SWB) in Estonia and France is smaller (0.026 ha and 0.071 ha, respectively) than the detection limit (0.2 ha) of the most recent global inventory based on GLOWABO data from satellite imagery (Verpoorter & al., 2014). Our analysis of the abundance and distribution EU water bodies suggests that these global counts underrepresent the hydrologically complex terrain of European territory.
- Our present estimates of all EU inventory of water bodies in all size ranges based on OSM database is ~1 million. The comparison of ground-validated data from national inventories and OSM database shows that reliability of OSM data varies between countries. The number of small standing water bodies remains underestimated, especially in size class below 0.1 ha (the average representation of SWB for EU level in size range 0.01 - 0.1 ha is 18%, below 0.01 ha is inaccurate).
- The calculated perimeter in Estonian and French SWB is consistent with a hypothesis by Verpoorter et al. (2014) that the most of the global land-lake interface is in small SWB.
- The abundance and size distribution of SWB in Estonia and France shows the relative importance of very small and man-made water bodies in EU context. The abundance of man-made SWB is globally less studied, especially in the smallest size ranges, but they have important role in sustaining aquatic life while at the same time supporting human activities. Because of their small size and catchments, small SWB are far easier and cheaper to protect and restore than larger SWB.
- All SWB, including small and man-made ponds, play an important role in ecosystem services and require careful management to avoid hydrological and environmental deterioration
- Combining the national inventories of SWB proves to be the best available solution for estimating the regional inventories of SWB. For each geographical region specific function must be used in order to calculate the total number of SWB.

Quality of data



Estonian Topographic Database OpenStreetMap Missing from both

Repeated random check of OSM and Estonian national inventory datasets against aerial imagery reveals that both dataset have some limitations. In both cases, some SWB remain unrepresented (majority of bog pools is missing from OSM).





