



On the NP-Completeness of the Perfect Perfect Matching Free Subgraph Problem

Mathieu Lacroix, Ridha Mahjoub, Sébastien Martin, Christophe Picouleau

► **To cite this version:**

Mathieu Lacroix, Ridha Mahjoub, Sébastien Martin, Christophe Picouleau. On the NP-Completeness of the Perfect Perfect Matching Free Subgraph Problem. 2011. <hal-00875532>


HAL Id: hal-00875532

<https://hal.archives-ouvertes.fr/hal-00875532>

Submitted on 22 Oct 2013

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



CAHIER DU LAMSADE

312

Octobre 2011

Titre

**On the NP-Completeness of the Perfect
Perfect Matching Free Subgraph Problem**



AUTEURS

Mathieu Lacroix,
A. Ridha Mahjoub,
Sébastien Martin,
Christophe Picouleau.

On the NP-Completeness of the Perfect Matching Free Subgraph Problem

Mathieu Lacroix^a, A. Ridha Mahjoub^a, Sébastien Martin^a, Christophe Picouleau^{b,*}

^a *Université Paris-Dauphine LAMSADE, CNRS, Place du Maréchal de Lattre de Tassigny, 75116 Paris Cedex 16, France.*

^b *Conservatoire National des Arts et Métiers, CEDRIC laboratory, 292 Rue St Martin, 75141 Paris, France.*

Abstract

Given a bipartite graph $G = (U \cup V, E)$ such that $|U| = |V|$ and every edge is labelled true or false or both, the perfect matching free subgraph problem is to determine whether or not there exists a subgraph of G containing, for each node u of U , either all the edges labelled true or all the edges labelled false incident to u , and which does not contain a perfect matching. This problem arises in the structural analysis of differential-algebraic systems. The purpose of this paper is to show that this problem is NP-complete. We show that the problem is equivalent to the stable set problem in a restricted case of tripartite graphs. Then we show that the latter remains NP-complete in that case. We also prove the NP-completeness of the related minimum blocker problem in bipartite graphs with perfect matching.

Keywords: Bipartite graph, matching, blocker, stable set, tripartite graph, NP-complete, structural analysis problem.

1. Introduction

Given a graph $G = (V, E)$, a *matching* of G is a subset of edges such that no two edges share a common node. Matchings have shown to be useful for modeling various discrete structures [2, 8]. A graph is called *bipartite* (*tripartite*) if its nodes can be partitioned into two (three) disjoint sets such that every edge connects one node in a set to a node in a different set. A bipartite graph is called *complete* if there exists an edge between every pair of nodes of different sets. A complete bipartite graph is also called a *biclique*. A matching M in graph G is called *perfect* if every node of G is incident to some edge of M . Given a

*Corresponding author

Email addresses: lacroix@lamsade.dauphine.fr (Mathieu Lacroix),
mahjoub@lamsade.dauphine.fr (A. Ridha Mahjoub), martin@lamsade.dauphine.fr
(Sébastien Martin), christophe.picouleau@cnam.fr (Christophe Picouleau)

bipartite graph $G = (U \cup V, E)$ such that $|U| = |V| = n$, a matching M of G is then perfect if and only if $|M| = n$.

Let $G = (U \cup V, E)$ be a bipartite graph such that $|U| = |V| = n$. Let $U = \{u_1, \dots, u_n\}$ and $V = \{v_1, \dots, v_n\}$. Suppose that every edge of E is labelled *true* or *false*, where an edge may have both true and false labels. For a node $u_i \in U$, let E_i^t and E_i^f denote the sets of edges incident to u_i labelled true and false, respectively. The *perfect matching free subgraph problem* (PMFSP) in G is to determine whether or not there exists a subgraph containing for each node $u_i \in U$ either E_i^t or E_i^f (but not both), and which is perfect matching free. As it will be shown in the next section, this problem arises in the structural analysis of differential-algebraic systems. The purpose of this paper is to show that PMFSP is NP-complete. For this we first show that PMFSP is equivalent to the stable set problem in a restricted case of tripartite graphs. Then we show that the latter remains NP-complete in that case.

Given a graph $G = (V, E)$, a matching in G of maximum cardinality is called a *maximum matching*. Its size corresponds to the *matching number* of G which is denoted by $\nu(G)$. One of the most attractive and studied problems in combinatorial optimization is the *maximum matching problem*, which consists, given a graph G , in finding a maximum matching in G [3, 8]. This problem can be solved in polynomial time using the algorithm developed by Edmonds [3]. If the graph is bipartite, the problem is much simpler. It reduces to a maximum flow problem. A *vertex cover* of a graph G is a set T of nodes such that every edge of G has at least one end in T . A well known min-max relation in graph theory and combinatorics is the following. A *stable set* of a graph is a subset of nodes S such that no two nodes in S are adjacent. Given a graph $G = (V, E)$, the stable set problem in G consists in finding a stable set of maximum cardinality.

Theorem 1. (*König [8]*) *For a bipartite graph, the maximum cardinality of a matching is equal to the minimum cardinality of a vertex cover.* \square

As the complementary of a vertex cover in a graph is a stable set, a consequence of Theorem 1 is the following.

Corollary 2. *Given a bipartite graph, if M is a maximum cardinality matching and S is a maximum stable set, then $|M| + |S| = |V|$.* \square

For more details on matching theory, the reader is referred to [8].

The paper is organized as follows. In the next section we discuss the relation between PMFSP and the structural analysis problem in differential-algebraic systems. In Section 3 we show the equivalence between the PMFSP and the stable set problem in a special case of tripartite graphs. In Section 4 we show the NP-completeness of PMFSP. In Section 5 we consider the related minimum blocker problem in bipartite graphs with perfect matching, and in Section 6 we give some concluding remarks.

2. Differential-algebraic systems and the PMFSP

Differential-algebraic systems (DAS) are used for modeling complex physical systems as electrical networks and dynamic movements. Such a system can be given as $f(x, \dot{x}, u, p, t) = 0$ where x is the variable vector, \dot{x} denotes the derivative vector of x with respect to time, u is the input vector, p is the parameter vector and t is time. Establishing that a DAS definitely is not solvable can be helpful. A necessary (but not sufficient) condition for solving a DAS is that the number of variables and equations must agree. Simulation is the main tool for solving DASs. Object-oriented modeling languages like Modelica [4] enforce this as simulation is not possible if this is not case. Thus before solving a differential-algebraic system, one has to verify if there are as many equations as variables, and if there exists a mapping between the equations and the variables in such a way that each equation is related to only one variable and each variable is related to only one equation. If this is satisfied, then we say that the system is *well-constrained*. The *structural analysis problem* for a DAS consists in verifying if the system is well-constrained.

In many practical situations, physical systems yield differential-algebraic systems with conditional equations. A conditional equation is an equation whose form depends on the value (true or false) of a condition. A conditional equation can generate several equations. A conditional differential-algebraic system may then have different forms depending on the set of conditions that hold. Here we consider conditional DAS's such that any conditional equation may take two possible values, depending on whether the associated condition is true or false and may generate only one equation. Consider for example the following DAS :

$$\begin{aligned}
 eq_1 : & \quad \text{if } a > 0 \\
 & \quad \text{then } 0 = 4x^2 + 2\dot{x} + 4y + 2 \\
 & \quad \text{else } 0 = \dot{y} + 2z + 4 \\
 eq_2 : & \quad \text{if } b > 0 \\
 & \quad \text{then } 0 = 6\dot{y} + 2\dot{z} + 2 \\
 & \quad \text{else } 0 = x + \dot{y} + 1 \\
 eq_3 : & \quad \text{if } c > 0 \\
 & \quad \text{then } 0 = 6\dot{x} + y + 2 \\
 & \quad \text{else } 0 = 3\dot{y} + z + 3
 \end{aligned} \tag{4}$$

If $a > 0$, $b > 0$, $c > 0$, then system (4) is nothing but the following.

$$\begin{aligned}
 eq_1 : & \quad 0 = 4x^2 + 2\dot{x} + 4y + 2, \\
 eq_2 : & \quad 0 = 6\dot{y} + 2\dot{z} + 2, \\
 eq_3 : & \quad 0 = 6\dot{x} + y + 2.
 \end{aligned} \tag{5}$$

And if $a > 0$, $b > 0$, $c \leq 0$ then system (4) is nothing but the system.

$$\begin{aligned}
 eq_1 : & \quad 0 = 4x^2 + 2\dot{x} + 4y + 2, \\
 eq_2 : & \quad 0 = 6\dot{y} + 2\dot{z} + 2,
 \end{aligned} \tag{6}$$

$$eq_3 : \quad 0 = 3 \dot{y} + z + 3.$$

The structural analysis problem has been considered in the literature for non-conditional DAS's. In [9, 10], Murota introduces a formulation of the problem in terms of bipartite graphs and shows that a system of equations is well constrained if and only if there exists a perfect matching in the corresponding bipartite graph. Given a DAS, one can associate a bipartite graph $G = (U \cup V, E)$, called *incidence graph*, where U corresponds to the equations, V to the variables and there is an edge $u_i v_j \in E$ between a node $u_i \in U$ and a node $v_j \in V$ if and only if the variable corresponding to v_j appears in the equation corresponding to u_i . Checking if the system is well constrained then reduces to calculating a perfect matching in the associated incidence graph, which can then be done in polynomial time.

Given a conditional DAS, the associated structural analysis problem consists in verifying whether or not the system is well constrained for all the possible values. The SAP for a conditional DAS thus reduces to verifying whether or not the incidence bipartite graph, related to any configuration of the system, contains a perfect matching. More precisely, consider a conditional DAS with n equations (eq_1, \dots, eq_n) and n variables (x_1, \dots, x_n). Let $G = (U \cup V, E)$ be a bipartite graph where $U = \{u_1, \dots, u_n\}$ (resp. $V = \{v_1, \dots, v_n\}$) is associated with the equations (resp. variables). Between a node $u_i \in U$ and a node $v_j \in V$ we consider an edge, called *true edge* (resp. *false edge*) if the variable x_j appears in equation eq_i when the condition of eq_i is true (resp. false). Note that an edge may be at the same time true and false. Let E_i^t (resp. E_i^f) be the set of true (resp. false) edges incident to u_i , for $i = 1, \dots, n$. Then

$$E = \bigcup_{i=1, \dots, n} (E_i^t \cup E_i^f).$$

Hence, the SAP reduces to finding whether or not there exists a subgraph of G , containing, for each node u_i either E_i^t or E_i^f (but not both) and which does not contain a perfect matching. Therefore, the SAP reduces to the PMFSP [6]. In [7], an integer programming formulation is proposed for the problem and some algorithmic and polyhedral issues are discussed.

3. PMFSP and stable sets

The aim of this section is to show that PMFSP is equivalent to the stable set problem in a special case of tripartite graph. Let $H = (V^1 \cup V^2 \cup V^3, F)$ be a tripartite graph where $|V^1| = |V^2| = |V^3| = n$, $V^j = \{v_1^j, \dots, v_n^j\}$ for $j = 1, 2, 3$ and V^1 and V^2 are connected by the perfect matching $M = \{v_1^1 v_1^2, v_2^1 v_2^2, \dots, v_n^1 v_n^2\}$. We will consider the following problem : does there exist a stable set in H of size $n + 1$? We will call this problem the *tripartite stable set with perfect matching problem* (TSSPPM). In what follows we shall show that both problems TSSPPM and PMFSP are equivalent.

Theorem 3. *TSSPPM and PMFSP are polynomially equivalent.*

PROOF. Let $G = (U \cup V, E)$ and $H = (V^1 \cup V^2 \cup V^3, F)$ be the graphs on which the problems PMFSP and TSSPMS are considered, respectively. We will first show that an instance of TSSPMP can be transformed into an instance of PMFSP. For an edge $v_i^1 v_i^2$ of the perfect matching where $v_i^1 \in V^1$ and $v_i^2 \in V^2$, we consider a node u_i in U . And for a node v_i^3 of V^3 we consider a node v_i in V . Moreover, if $v_i^1 v_k^3$ (resp. $v_i^2 v_k^3$) is in F for some $i, k \in \{1, \dots, n\}$, then we add an edge $u_i v_k$ in E with label true (resp. false). Figure 1 illustrates this transformation.

Observe that graph $H = (V^1 \cup V^2 \cup V^3, F)$ can be obtained from graph $G = (U \cup V, E)$ by doing the reverse operations.

Let E_i^t (resp. E_i^f) be the set of edges incident to u_i labelled true (resp. false), for $i = 1, \dots, n$.




Figure 1: Two equivalent FPMSP and TSSPPM instances.

In what follows, we will show that there exists a stable set in H of size $n + 1$ if and only if there exists a subgraph $G' = (U \cup V, E')$ of G such that for each node $u_i \in U$, either $E_i^t \subseteq E'$ or $E_i^f \subseteq E'$, and G' is perfect matching free. In fact, suppose first that there exists a subgraph G' of G that satisfies the required properties. Since G' is perfect matching free, this implies that a maximum cardinality matching in G' contains less than n edges. As $|U \cup V| = 2n$ by Corollary 2 there exists a stable set in G' , say S' , of size $|S'| \geq n + 1$. Now from S' , we are going to construct a stable set in H with the same cardinality. Let S be the node subset of H obtained as follows. For every node $v_j \in V \cap S'$, add node v_j^3 in S . And for every node $u_i \in U \cap S'$, add node v_i^1 in S if $E_i^t \subseteq E'$ and node v_i^2 if $E_i^f \subseteq E'$. As $|S'| \geq n + 1$, we have $|S| \geq n + 1$. We now prove that S is indeed a stable set. Since the edges between V^1 and V^2 are only those of the perfect matching M , and since from each edge of M , we have taken exactly one node in S , clearly, the restriction of S on $V^1 \cup V^2$ is a stable set. Suppose now that S contains vertices, say $v_i^1 \in V^1$ and $v_j^3 \in V^3$ which are adjacent. This implies that u_i and v_j belong to S' , $u_i v_j \in E_i^t$ and $E_i^t \subseteq E'$. However, this contradicts the fact that S' is a stable set. Using the same argument we deduce that S does not contain adjacent nodes $v_i^2 \in V^2$ and $v_i^3 \in V^3$. Thus S is a solution of TSSPMS.

Conversely, Suppose that we have a stable set S in H of size greater or equal to $n + 1$. Let E' be the edge subset of E obtained as follows. For every node




Figure 2: A TSSPPM instance resulting from 1-in-3 3SAT instance $L = \{l_1, l_2, l_3\}$ and $C = \{(\bar{l}_1, l_2, \bar{l}_3), (l_1, l_2, l_3)\}$. Only the satisfiability edges are displayed.

$v_i^1 \in V^1$, add in E' edge set E_i^t if $v_i^1 \in S$ and E_i^f if not. We will show that $G' = (U \cup V, E')$ contains a stable set of size greater than or equal to $n + 1$ which by Corollary 2 implies that G' is perfect matching free. Let $S' \subseteq U \cup V$ be the node set obtained from S as follows. For every node v_i^3 of $V^3 \cap S$ add node v_i of V in S' . And for every node v_i^1 (resp. v_i^2) of $V^1 \cap S$ (resp. $V^2 \cap S$), add node u_i of U in S' . Since S does not contain both nodes v_i^1 and v_i^2 for some i , and $|S| \geq n + 1$, we have that $|S'| \geq n + 1$. Moreover S' is a stable set. Indeed, suppose that S' contains two nodes, say $u_i \in U$ and $v_j \in V$ such that $u_i v_j \in E'$. Without loss of generality, suppose that u_i comes from node v_i^1 in S (the proof is similar if $v_i^2 \in S$). By construction of E' , this implies that $E_i^t \subseteq E'$, and hence $u_i v_j \in E_i^t$. From the construction of H , it follows that $v_i^1 v_j^3 \in F$. As $v_i^1, v_j^3 \in S$, this is a contradiction, and the proof is complete. \square

4. The NP-completeness of PMFSP

In this section we show the NP-completeness of PMFSP. For this we shall show that TSSPMP is NP-complete. By Theorem 3, the result follows. In [11] it is shown that the stable set problem is NP-complete in tripartite graphs. (Recall that the problem is known to be polynomially solvable in bipartite graphs). What we are going to show in the following is that the more restricted variant TSSPMP is also NP-complete. In other words, the stable set problem in tripartite graphs remains NP-complete even when the sets of the partition of the graph have the same size and that the set of edges between two of the

three sets of the partition consists of a perfect matching. In order to show the NP-completeness of TSSPMP, we shall use the one-in-three 3SAT problem. An instance of *one-in-three 3SAT* (1-in-3 3SAT) consists of n variables l_1, \dots, l_n and m clauses C_1, \dots, C_m with three literals per clause. Each clause is the disjunction of three literals, where a literal is either a variable or its negation. If x_i is a variable which represents either l_i or \bar{l}_i , then $\bar{x}_i = \bar{l}_i$ (resp. $\bar{x}_i = l_i$) if $x_i = l_i$ (resp. $x_i = \bar{l}_i$), where a bar stands for negation. The question is whether or not there exists an assignment of truth values ("true" or "false") to the variables such that each clause has exactly one true literal.

Theorem 4. *TSSPPM is NP-complete.*

PROOF. It is clear that TSSPMP is in NP. To prove the theorem, we shall use a reduction from 1-in-3 3SAT. The proof uses ideas from [11]. So suppose we are given an instance of 1-in-3 3SAT with a set of n variables $L = \{l_1, \dots, l_n\}$ and a set of m clauses $C = \{C_1, \dots, C_m\}$. We shall construct an instance of TSSPMP on a graph $H = (V^1 \cup V^2 \cup V^3, F)$ where $|V^1| = |V^2| = |V^3| = p = 3n + m - 1$ and the set of edges between V^1 and V^2 consists of a perfect matching. We will show that H has a stable set of size $p + 1$ if and only if 1-in-3 3SAT admits a truth assignment. With each variable $l_i \in L$, we associate the nodes $v_i^1, \bar{v}_i^1 \in V^1$, $v_i^2, \bar{v}_i^2 \in V^2$ and $v_i^3, \bar{v}_i^3 \in V^3$. These will be called *variable nodes*. With each clause $C_j = (x_r, x_s, x_t)$, we associate the nodes $w_{j_r}^1 \in V^1$, $w_{j_s}^2 \in V^2$, $w_{j_t}^3 \in V^3$. These will be called *clause nodes*. Finally we add the nodes $z_q^1 \in V^1$, $z_q^2 \in V^2$, $z_q^3 \in V^3$ for $q = 1, \dots, n - 1$. These will be called *fictitious nodes*. Note that $|V^1| = |V^2| = |V^3| = p$. Now we construct the edge set F . For each variable $l_i \in L$, consider the edges $v_i^1 \bar{v}_i^2, \bar{v}_i^2 v_i^3, v_i^3 \bar{v}_i^1, \bar{v}_i^1 v_i^2, v_i^2 \bar{v}_i^3, \bar{v}_i^3 v_i^1$ in F . These will be called *variable edges*. Note that these edges form a cycle of length 6, which will be denoted by Γ_i for $i = 1, \dots, n$. For each clause $C_j = (x_r, x_s, x_t)$ add in F the edges $w_{j_r}^1 w_{j_s}^2, w_{j_s}^2 w_{j_t}^3, w_{j_t}^3 w_{j_r}^1$. These are called *clause edges*. Note that these edges form a triangle, which will be denoted by T_j , for $j = 1, \dots, m$. Also add in F the edges $z_q^1 z_q^2$ for $q = 1, \dots, n - 1$. Remark that the edges between V^1 and V^2 form a perfect matching given by the edges $v_i^1 \bar{v}_i^2, \bar{v}_i^1 v_i^2$, $i = 1, \dots, n$, $w_{j_r}^1 w_{j_s}^2$, $j = 1, \dots, m$, and $z_q^1 z_q^2$, $q = 1, \dots, n - 1$. Now according to the values of the literals, we add edges in E as follows. For every clause (x_r, x_s, x_t)

- if $x_r = l_r$, add the edges $w_{j_r}^1 \bar{v}_r^3, w_{j_s}^2 v_r^3, w_{j_t}^3 v_r^1, w_{j_t}^3 v_r^2$,
- if $x_r = \bar{l}_r$, add the edges $w_{j_r}^1 v_r^3, w_{j_s}^2 \bar{v}_r^3, w_{j_t}^3 \bar{v}_r^1, w_{j_t}^3 \bar{v}_r^2$,
- if $x_s = l_s$, add the edges $w_{j_r}^1 v_s^3, w_{j_s}^2 \bar{v}_s^3, w_{j_t}^3 v_s^1, w_{j_t}^3 v_s^2$,
- if $x_s = \bar{l}_s$, add the edges $w_{j_r}^1 \bar{v}_s^3, w_{j_s}^2 v_s^3, w_{j_t}^3 \bar{v}_s^1, w_{j_t}^3 \bar{v}_s^2$,
- if $x_t = l_t$, add the edges $w_{j_r}^1 v_t^3, w_{j_s}^2 v_t^3, w_{j_t}^3 \bar{v}_t^1, w_{j_t}^3 \bar{v}_t^2$,
- if $x_t = \bar{l}_t$, add the edges $w_{j_r}^1 \bar{v}_t^3, w_{j_s}^2 \bar{v}_t^3, w_{j_t}^3 v_t^1, w_{j_t}^3 v_t^2$,

These are called *satisfiability edges*. For each fictitious node in $V^1 \cup V^2$, add edges to connect all nodes in V^3 , and for each fictitious node in V^3 , add edges to connect all non fictitious nodes in $V^1 \cup V^2$.

Thus, from an instance of the 1-in-3 3SAT with n variables and m clauses, we obtain a tripartite graph with $9n + 3m - 3$ nodes and $10n^2 + 4nm - 5n + 14m + 1$ edges.

Figure 2 show an example of graph H when $L = \{l_1, l_2, l_3\}$ and $C = \{(\bar{l}_1, l_2, \bar{l}_3), (l_1, l_2, l_3)\}$. For sake of clarity, only the satisfiability edges are displayed.

Claim 5. *Any stable set in H cannot contain more than $3n + m$ nodes. Moreover, if a stable set contains $3n + m$ nodes, then it does not contain any fictitious node.*

PROOF. Let S be a stable set in H . First we show that if S contains a fictitious node, then $|S| \leq 3n + m - 1$. Suppose S contains a fictitious node z . If $z \in V^3$, as z is adjacent to all nodes in $V^1 \cup V^2$, $|S| \leq |V^3| = 3n + m - 1$. Now suppose, without loss of generality, that $z \in V^1$. As z is adjacent to all the nodes of V^3 , we have $S \cap V^3 = \emptyset$. Consider a cycle Γ_i of H corresponding to a variable l_i . As Γ_i alternates between the sets V^1, V^2, V^3 and four of the six edges of Γ_i are incident to nodes in V^3 , at most two nodes of Γ_i may belong to S . Moreover, S may contain at most one node from each triangle T_j , $j = 1, \dots, m$. Consequently, $|S| \leq 2n + m < 3n + m$.

Now suppose that S does not contain any fictitious node. Then all the nodes of S come from the cycles Γ_i , $i = 1, \dots, n$ and the triangles T_j , $j = 1, \dots, m$. Since S may intersect each Γ_i in at most 3 nodes and each triangle in at most one node, it follows that $|S| \leq 3n + m$. \square

In what follows we show that there exists in H a stable set of size $3n + m$ if and only if 1-in-3 3SAT admits a solution such that each clause has exactly one true literal.

(\Rightarrow) Let S be a stable set in H of size $3n + m$. By Claim 5, S does not contain any fictitious node. Thus, as $|S| = 3n + m$, S intersects each cycle Γ_i in exactly three nodes and each triangle T_j in exactly one node. Moreover, we have that either $S \cap \Gamma_i = \{v_i^1, v_i^2, v_i^3\}$ or $S \cap \Gamma_i = \{\bar{v}_i^1, \bar{v}_i^2, \bar{v}_i^3\}$, for $i = 1, \dots, n$. Consider the solution I for 1-in-3 3SAT defined as follows. If $v_i^k \in S$ (resp. $\bar{v}_i^k \in S$), $k = 1, 2, 3$, then associate the true (resp. false) value to the variable l_i , for $i = 1, \dots, n$. In what follows we will show that for each clause $C_j = (x_r, x_s, x_t)$, we have exactly one literal with value true. For this it suffices to show that a clause node of T_j is in S if and only if the corresponding literal is of true value. Indeed, suppose that $w_{j_r}^1 \in S$. We may suppose that $x_r = l_r$, the case where $x_r = \bar{l}_r$ is similar. By construction of H , as the satisfiability edge $w_{j_r}^1 \bar{v}_r^3$ belongs to F , it follows that $\bar{v}_r^3 \notin S$. By the remark above, this implies that v_r^1, v_r^2, v_r^3 belong to S . Therefore literal l_r has value true in solution I . Thus x_r has value true.

Conversely, if $x_r = \text{true}$ ($= l_r$), then by definition of I , $v_r^1, v_r^2, v_r^3 \in S$. Moreover, the satisfiability edges $w_{js}^2 v_r^3, w_{jt}^3 v_r^2$ belong to F . As $|S \cap T_r| = 1$, it follows that $w_{jr}^1 \in S$.

In consequence, as S contains exactly one clause node from each T_i , it follows that each clause has exactly one true literal.

(\Leftarrow) Suppose that there exists a solution I of 1-in-3 3SAT such that each clause has exactly one literal with true value. We will show that the maximum stable set in H is of size $3n + m$. Let S be the node set obtained as follows :

- if (in I) $l_i = \text{true}$, then add v_i^1, v_i^2, v_i^3 to S . If $l_i = \text{false}$, then add $\bar{v}_i^1, \bar{v}_i^2, \bar{v}_i^3$ to S .
- if for a clause $C_j = (x_r, x_s, x_t)$,
 $x_r = \text{true}$, add w_{jr}^1 to S ,
 $x_s = \text{true}$, add w_{js}^2 to S ,
 $x_t = \text{true}$, add w_{jt}^3 to S .

As each clause has exactly one true literal with respect to the solution I , we have that $|S| = 3n + m$. Now, it suffices to show that S is a stable set. For this it suffices to show that none of the variable nodes of S is adjacent to a clause node of S . Suppose, without loss of generality, that for some $r \in \{1, \dots, n\}$, $l_r = \text{true}$. Hence $v_r^1, v_r^2, v_r^3 \in S$. In S these nodes may only be adjacent to nodes coming from clauses containing literal l_r or its negation \bar{l}_r . Actually, if $C_j = (x_r, x_s, x_t)$, by the definition of the satisfiability edges, nodes v_r^1, v_r^2, v_r^3 may be adjacent to nodes among $\{w_{js}^2, w_{jt}^3\}$ if $x_r = l_r$ and to node w_{jr}^1 , if $x_r = \bar{l}_r$. If $x_r = \text{true}$ (that is $x_r = l_r$), then $w_{jr}^1 \in S$. However, in this case none of the nodes v_r^1, v_r^2, v_r^3 is adjacent to w_{jr}^1 . Thus, none of the variable nodes is adjacent to a clause node in S . Therefore, S is a stable. Since $|S| = 3n + m$, by Claim 5, S is of maximum size, and the proof is complete. \square

From Theorems 3 and 4, we deduce the following corollary.

Corollary 6. *PMFSP is NP-complete.* \square

Corollary 6 shows that PMFSP is NP-complete in general. To conclude this section let us remark that the PMFSP can be solved in polynomial time if there exists a vertex in V which is incident to no edge with both labels true and false. Indeed, suppose that all the edges incident to a vertex $v_j \in V$ are either true or false (but not both). In this case, we can consider the subgraph $G' = (U \cup V, E')$ where E' contains, for $i = 1, 2, \dots, n$, the edge set E_i^t if E_i^f contains an edge of type $u_i v_j$, and E_i^f otherwise. Observe that, E' does not contain any edge incident to v_j . Which implies that G' does not contain a perfect matching. Consequently, G' is a solution of PMFSP. We also remark that, in this case, the instance H of the TSSPMP associated with G is such that vertex v_j^3 is adjacent to at most one of the nodes v_i^1 and v_i^2 , for $i = 1, 2, \dots, n$. A solution of the

TSSPMP is thus obtained by considering the stable set containing v_j^3 and, for $i = 1, 2, \dots, n$, either v_i^1 , if v_i^1 is not adjacent to v_j^3 and v_i^2 if not.

5. The minimum blocker perfect matching problem

In this section, we consider a variant of the PMFSP when there is no labels on the edges. This problem can be stated as follows. Given a graph $G = (U \cup V, E)$ with a perfect matching and $|U| = |V|$, find a perfect matching free subgraph with a maximum number of edges and covering the vertices of U . As it will turn out, this problem is nothing but a special case of the so-called minimum blocker problem [12] (see also [1]).

Let $G = (U \cup V, E)$ be a bipartite graph with matching number $\nu(G)$. In [12], Zenklusen et al. define a *blocker* as a subset of edges $B \subset E$ such that $G' = (U \cup V, E \setminus B)$ has a matching number smaller than $\nu(G)$. They define the *minimum blocker problem* (MBP) as follows. Given a bipartite graph $G = (U \cup V, E)$ and a positive integer k , does there exist an edge subset B of E with $|B| \leq k$ such that B is a blocker? They prove that MBP is NP-complete. Here, we are interested in a special case of the MBP, hereafter called the *minimum blocker perfect matching problem* (MBPMP), where G contains a perfect matching. In what follows, we show that MBPMP is NP-complete. We also prove that it remains NP-complete in case where $G' = (U \cup V, E \setminus B)$ must cover U . Which corresponds to the PMFSP with no edge labels.

Theorem 7. *MBPMP is NP-complete.*

PROOF. It is shown in [12] that MBP is NP-complete even in the case where $\nu(G) = |U|$ (see the proof of Theorem 3.3 in [12]). We consider this subproblem. Moreover, we suppose that $|U| < |V|$ (otherwise MBP would be MBPMP). Let $\deg(U) = \min_{u_i \in U} \{\deg(u_i)\}$. We suppose that $k < \deg(U)$ (in the case where $k \geq \deg(U)$, the $\deg(U)$ edges incident to the vertex with minimum degree in U clearly form a blocker).

Let $\tilde{G} = (\tilde{U} \cup \tilde{V}, \tilde{E})$ be the graph obtained from G where $\tilde{U} = U \cup \bar{U}$ and $\tilde{V} = V \cup \bar{V}$ where $\bar{U} = \max\{|V| - |U|, k + 1\}$, $\bar{V} = \max\{k + 1 - |V| + |U|, 0\}$ and $\tilde{E} = E \cup \{\bar{u}\bar{v} : \bar{u} \in \bar{U}, \bar{v} \in \bar{V}\}$. Note that \bar{U} contains at least $k + 1$ nodes, $|\tilde{U}| = |\tilde{V}|$ and $(\bar{U} \cup \tilde{V}, \tilde{E} \setminus E)$ is a complete bipartite graph. Also note that, as $\nu(G) = |U|$, \tilde{G} contains a perfect matching.

In what follows we will show that G contains a blocker of cardinality less or equal than k if and only if \tilde{G} so does. For this we first give the following claim.

Claim 8. . *Let $H = (W_1 \cup W_2, F)$ be a complete bipartite graph such that $|W_1| = |W_2| \geq k + 1$ for some $k \geq 0$. Then H does not contain a blocker of size $\leq k$.*

PROOF. Suppose that there is a blocker B of size $|B| \leq k$. Then the subgraph $H' = (W_1 \cup W_2, F \setminus B)$ has no perfect matching. From Hall's theorem (see [8]) there exists $i \in \{1, 2\}$ and $W \subset W_i$ such that $|W| > |\Gamma(W)|$ in H' where $\Gamma(W)$

stands for the neighbor set of W . Since H is a complete bipartite graph, we have $|B| \geq |W| \times (|W_i| - |\Gamma(W)|)$. Now, since for each triplet of nonnegative integers x, y, z with $x \geq y > z$ we have $y(x - z) \geq x$, by considering $x = |W_i|$, $y = |W|$ and $z = |\Gamma(W)|$, we conclude that $|B| \geq |W| \times (|W_i| - |\Gamma(W)|) \geq |W_i| \geq k + 1$, a contradiction. \square

Now consider a blocker B of G with $|B| \leq k$, and suppose that B is not a blocker of \tilde{G} . Thus there exists a perfect matching of \tilde{G} , say M , which does not intersect B . Since $|M \cap E| = |U| = \nu(G)$ and B is a blocker of G , we have a contradiction. Thus B is also a blocker of \tilde{G} .

Conversely, suppose that G has no a blocker B with $|B| \leq k$. If \tilde{G} contains a blocker, say \tilde{B} with $|\tilde{B}| \leq k$, then let $\tilde{B}_1 = \tilde{B} \cap E$. Obviously, $|\tilde{B}_1| \leq k$. We claim that \tilde{B}_1 is a blocker of G . In fact, if this is not the case, then there must exist a matching M' in the graph $(U \cup V, E \setminus \tilde{B}_1)$ with $|M'| = |U| = \nu(G)$. Let V' be the subset of nodes of V covered by M' . Let $H = (W_1 \cup W_2, F)$ be the biclique with $W_1 = \bar{U} = \tilde{U} \setminus U$ and $W_2 = \tilde{V}'$ where $\tilde{V}' = \tilde{V} \setminus V'$. Clearly, $|W_1| = |W_2| \geq k + 1$. Let $\tilde{B}_2 = \tilde{B} \cap F$. As $|\tilde{B}_2| \leq k$, by the claim above, the subgraph $(W_1 \cup W_2, F \setminus \tilde{B}_2)$ contains a perfect matching say M'' . As $M' \cup M''$ is a perfect matching of $(\tilde{U} \cup \tilde{V}, \tilde{E} \setminus \tilde{B})$, this contradicts the fact that \tilde{B} is a blocker of \tilde{G} , and the proof is complete. \square

In the proof of Theorem 7, graph \tilde{G} is constructed in such a way that $\deg(u) \geq k + 1$ for all $u \in \tilde{U}$. Therefore, any graph obtained from \tilde{G} by removing the edges of any blocker B with $|B| \leq k$ covers the vertices of \tilde{U} . This implies that the variant of the PMFSP without label on the edges, considered in this section, is also NP-complete.

6. Concluding remarks

In this paper we have shown that the perfect matching free subgraph problem is NP-complete. For this, we have first proved that the problem is equivalent to the stable set problem in tripartite graphs when the set of edges between two elements of the partition of the graph is reduced to a perfect matching. Then we have shown that the latter is NP-complete. We have also proved that the related minimum blocker perfect matching problem is NP-complete.

The PMFSP can be easily generalized to the case where the edges incident to each vertex $u \in U$ are gathered in more than two (non-disjoint) edge sets (e.g. the true and false edge sets). This problem is clearly NP-complete since it contains PMFSP as a special case. Moreover, this latter more general problem has applications to the structural analysis problem for embedded conditional differential-algebraic systems. These consist of systems which may contain equations whose value may depend on more than one condition [6]. As it has been shown in [6], the latter can be reduced to the former one.

Acknowledgments

We would like to thank Sébastien Furic, Bruno Lacabanne and El Djillali Talbi from LMS-Imagine for stimulating discussions. This work has been supported by the projects ANR-06-TLOG-26-01 PARADE and ANR-09-JCJC-0068 DOPAGE. The financial support is much appreciated.

- [1] M. Costa, D. de Werra, C. Picouleau, Minimum d -blockers and d -transversals in graphs, To appear in *Journal of Combinatorial Optimization*. (2010).
- [2] A. L. Dulmage, N. S. Mendelsohn, Coverings of bipartite graphs, *Canadian Journal of Mathematics*. (1963) 517-534.
- [3] J. Edmonds, Maximum matching and a polyhedron with 0,1-vertices, *J. Res. Nat. Bur. Standards*. (1965) 69B, 125-130.
- [4] P. Fritzson, *Principles of Object-Oriented Modeling and Simulation with Modelica 2.1*, Wiley-Interscience. (2003).
- [5] Z. Galil, S. Micali, H. N. Gabow, An $O(EV \log V)$ algorithm for finding a maximal weighted matching in general graphs, *SIAM J. Computing*. (1986) 15:120 - 130.
- [6] M. Lacroix, A. R. Mahjoub, S. Martin, Structural analysis in Differential-Algebraic Systems and Combinatorial Optimization, *Proceedings of 39th International Conference on Computers & Industrial Engineering (CIE39)*. (2009) 331-337.
- [7] M. Lacroix, A. R. Mahjoub, S. Martin, Structural Analysis for Differential-Algebraic Systems : Complexity, Formulation and Facets. *ISCO 2010*, Hamamet, Tunisia. (2010).
- [8] L. Lovasz, M. D. Plummer, *Matching Theory*, North-Holland, 1986.
- [9] K. Murota, *Systems Analysis by Graphs and Matroids*, Springer-Verlag, 1987.
- [10] K. Murota, *Matrices and Matroids for Systems Analysis*, Springer-Verlag, 2000.
- [11] C. Phillips, T. J. Warnow, The asymmetric median tree A new model for building consensus trees, *Combinatorial Pattern Matching*. (1996) 234-252.
- [12] R. Zenklusen, B. Ries, C. Picouleau, D. de Werra, M.-C. Costa, C. Bentz, Blockers and transversals, *Discrete Mathematics* 309. (2009) 4306-4314.