

Exclusive Breast Feeding: Infant Feeding Pattern and Practices among Mothers in Santa Fe, Romblon

Ken N Falculan¹, Ralph N Falculan²

¹Assistant Professor, Romblon State University, Odiongan, Romblon, Philippines

²Development Management Officer, Department of Health- MIMAROPA, Philippines

Received: December 8, 2021

Received in Revised: January 9, 2021

Accepted: January 16, 2021

Abstract

If the following characteristics of respondents are taken into consideration: age, number of living children, marital status, income level, employment level, educational background, current feeding practices, and level of knowledge about the importance of EBF, it is hypothesized that these characteristics will have an impact on feeding among infants aged 0 to 6 months. A descriptive research employing a normative survey was conducted with a study population of 102 mothers from the Municipality of Santa Fe in the province of Romblon, after informed permission had been obtained. The following is the outcome: moms between the ages of 21 and 25 are the most prevalent, accounting for 39 percent of all respondents who use EBF. With more children in their household, the percentage of respondents who practice EBF increases. Additionally, 55 percent of respondents said that they are living with a partner, either married through a church wedding, civil marriage or in the state of cohabitation. Lower-income mothers (45 percent) practice EBF more frequently than mothers from other income classes. Mothers without a job are more likely to practice EBF (51 percent), while elementary graduate mothers are more likely to practice exclusive breastfeeding (63 percent). Forty-four (44) mothers out of one hundred and two (102) respondents are practicing exclusive breastfeeding, and at least 63 percent have revealed little knowledge about the importance of exclusive breastfeeding.

Keywords: Breastfeeding, Neurophysiological, Nuclear Family, Premature Babies

Introduction

It is preferable for our kid to be fed breast milk, and the advantages of breastfeeding extend well beyond the provision of basic nutrients. Breast milk, in addition to providing our infant with all of the vitamins and minerals he or she need during the first six months of life, is also a rich source of disease-fighting compounds that shield our child from sickness. As a result, the American Academy of Pediatrics recommends that babies should be breastfed exclusively for the first six months of their lives (although any amount of breastfeeding is beneficial). Furthermore, scientific research have shown that breastfeeding is beneficial to our health. Our kid is protected against a vast variety of ailments, including allergies and obesity. Breastfeeding may also help to improve the child's cognitive abilities. The stress level of moms and the risk of postpartum depression can be reduced by nursing. Many women report feeling calm when breastfeeding, and it may also lower the risk of some cancers in women who are breastfeeding. According to the World Health Organization (2013), infants should be exclusively breastfed for the first six months of life in order to achieve optimal growth, development, and health. The appropriate age at which solids should be introduced is around six months, due to the immaturity of the gastrointestinal tract and renal system, as well as the infant's neurophysiological state at that time. Following that, in order to satisfy their changing

nutritional needs, babies should be provided with nutritionally appropriate and safe supplemental meals while continuing to nurse for up to two years or longer if possible. When it comes to new moms, hospitals and maternity units set a tremendous example. The Baby-Friendly Hospital Initiative (BFHI), which was founded in 1991 by UNICEF and the World Health Organization, is a collaborative effort to guarantee that all maternity facilities, whether free standing or located in a hospital, are designated as breastfeeding support centers. Baby-friendly designations may be bestowed upon maternity facilities that do not accept free or low-cost breast milk substitutes, feeding bottles, or teats and that have taken 10 particular actions to assist in the successful breastfeeding of their infants.

Methods

Specifically, the descriptive approach of research will be used by the investigator. This survey, also known as a normative survey, is a fact-finding study that is used to gather demographic data about people's practices, beliefs, intentions, opinions, attitudes, interests, perceptions, and the like. After the data has been collected, it is analyzed, organized, and interpreted to determine trends (Calderon, 2003).

Respondents of the Study

The respondents of this study included one hundred two mothers from the Municipality of Santa Fe, Romblon.

Research Instrument

A questionnaire will be the gathering device used to obtain the data needed to answer the specific problems particularly the profile of the target respondents who would possibly respond to the statement of the problem. It is composed of survey questionnaire distributed to randomly selected mothers who gave birth using stratified sampling coming from eleven barangays of the municipality.

Data Gathering Procedure

The researchers sought permission from 203 mothers. Likewise, a letter of request will be endorsed to the Municipal Health Officer to allow me to administer the questionnaire. Lastly, the setting of time frame in distributing and retrieving the instruments will be accomplished to set forth data needed in the study.

Results and Discussion

This section was the data gathered by the researcher. These were analyzed and interpreted below.

Table 1. Profile of the Respondents in terms of Age

AGE	FREQUENCY	TOTAL
15-20	25	24%
21-25	33	32%
26-30	30	31%
31-35	10	9%
36-40	3	4%
41-45	1	2%
TOTAL	102	100%

From the table, it is indicated that mothers, whose age ranges from 21 to 25 were the most prevailing, with 32 percent of the total respondents, 17 out of 33 in this age group or 39%, of the total mother practicing EBF followed by mother in age group 26 to 30 with 15 out of 30 or 34 %. Mixed feeding is also predominant in age group 21-25 yrs. old with 14 out of 40 mothers who is mixed feeding or 35%. Followed by mother in age group 15-20 yrs. old with 13 out of 40 or 33%, while age group from 26-35 yrs. old, bottle feeding is predominant or 78% of the total bottle-feeding mothers.

Table 2. Profile of the Respondents in terms the Number of Living Children

No. of children	FREQUENCY	TOTAL
1-3	68	67%
4-7	32	31%
8-11	2	2%
12-15	0	0%
TOTAL	102	100

Table 2 notifies us about the profile of the mothers in terms of the number of their living children. The table above shows majority of the respondents has children ranging from 1 to 3, (67%) followed by mothers with 4 to 7 children (31%), and only 2 respondent has children ranging from 8-11. The result shows that, the more the children the respondent had the higher the percentage of practicing EBF, respondent with children 8-11 are practicing 100% EBF, followed by mother with 4-7 children 69%, while mixed feeding is predominant in mother with 1-3 children and bottle feeding is predominant in mother with 4-7 children 8 out 32 children (25%).

Table 3. Profile of the Respondents according to Marital Status of the Respondents

Status of Mother	FREQUENCY	PERCENTAGE
Single	46	45%
Living in	56	55%
TOTAL	102	100%

**Single includes with children but no spouse; being widowed*

**Living in includes being married through church wedding, civil rights or in the state of cohabitation*

Table 3 displays the profile of the respondents in terms of marital status. Of the 102 respondents, fifty-six or 55% said that they are living in with a partner, either married through church wedding, civil rights or in the state of cohabitation. On the other hand, 45% or 46 mothers divulged that they were actually single, characterized in this study as mother/s with children but no spouse, the result shows that EBF practice is common in married mother with 38 out of 56 (67%) compared to 13% among single mother. mixed feeding and bottle feeding are common to single mother with result as follows 25 out of 46 single mother or 54% and 14 out of 46 or 32% respectively.

Table 4. Profile of the Respondents in terms of Economic Status

Income Class	FREQUENCY	PERCENTAGE
Low Income Bracket	96	95%
Middle Income Bracket	6	5%
Upper Income Bracket	0	0%
TOTAL	102	100%

**Low income includes P15, 780 and below*

**Middle income includes monthly family income of P15, 781—P78,900*

**Upper Income bracket includes monthly income of P78, 901 and above*

Table 4 highlights the profile of the mothers in terms of their economic status. From the table above, result shows that lower income mother 43 out of 96 (45%) practice EBF more compared to middle income mother 1 out of 6 (17%) are practicing EBF. While mixed feeding is predominant in middle income mother with 4 out of 6 (67%) compared to 38% among low income mother. Bottle feeding is also dominant in low income mother 17 out of 96 or 18% of low-income mother.

Table 5. Profile of the Mothers in terms Employment Status

Employment	FREQUENCY	PERCENTAGE
None	79	77%
Government-Permanent	4	3%
Government-Job Order	9	8%
Private—Fulltime	10	12%
Private-Part-time	0	0%
TOTAL	102	100%

Table 5 expresses the profile of the respondents as regards to their employment. From the table, it is clear that seventy-seven per cent of the participating respondents said they were having no job at all to attend to their family’s financial needs. A least 10 mothers are working full time to privately-owned companies. Four respondents, on the other hand, revealed that they work in a government agency, while none made mentioned that they have part-time jobs in private employers or offices.

In terms of feeding practices based on employment status the result shows that mother without job practice EBF 40 out of 79 mothers (51%) followed by mother working full time in private jobs 40% 4 out of 10 mothers, while bottle feeding is common in mothers working permanently in government 3 out of 4 or 75%. Mixed feeding is dominant in respondent working in government as job order 6 out of 9 respondent or 67% of the total working in government as job order.

Table 6. Profile of the Mothers in terms of Educational Attainment

Education	FREQUENCY	PERCENTAGE
None	0	0%
Elementary Under Grad	0	0%
Elementary Graduate	30	29%
High School	40	38%
College Level	18	17%
College Graduate	14	16%
Vocational Course	0	0%
Others	0	0%
TOTAL	102	100%

Table 6 exhibits the profile of the respondents in terms of educational attainment. It is clearly stated in the table that 40 mothers or 38% said they were graduate of high school; a staggering number of 30 mothers reported that they had graduated elementary while only sixteen (16)

revealed they were college graduates. No respondents admitted they had never attended any schooling, vocational courses or an undergraduate of elementary.

Result based on educational attainment revealed that exclusive breastfeeding is practiced more by elementary graduate mothers 19 out of 30 (63%) followed by high school graduate mothers 21 out of 40 mother or 53%. College graduate mothers are the least who practice EBF 1 out of 14 (7%) mainly because these mothers are working regularly in government or private offices.

Mixed feeding is common among college level mother 10 out of 18 (56%), followed by high school graduate 16 out of 40 mothers (40%) and elementary graduate 10 out of 30 mothers (33%). Bottle feeding is predominant in college graduate and college level with result as follows; 9 out of 14 (64%) and 5 out of 18 (28%) respectively.

Table 7. Profile of the Mothers in terms their CURRENT FEEDING PRACTICES

Practices	FREQUENCY	PERCENTAGE
Exclusive Breast Feeding	44	43%
Mixed Feeding	40	39%
Formula Feeding	18	18%
TOTAL	102	100%

Table 7 dowries the profile of the mothers in terms of their current feeding practices. From the table, it points out to the fact that forty-four (44) mothers out of one hundred two (102) respondents said that for their babies, they are doing exclusive breastfeeding. Forty (40) respondents said they are into mixed feeding; while only eighteen (18) are using formula feeding. Penultimate, these findings reveal that most mothers believe that exclusive breastfeeding—with no other food, drink, not even water, except breast milk or from a wet nurse for six months, is important for their babies’ growth and development.

Table 8. Profile of the Mothers regarding the Level of Knowledge on the Importance of Exclusive Feeding

Level of Knowledge	FREQUENCY	PERCENTAGE
None	0	0
Little Knowledge	64	63%
Highly Knowledgeable	38	37%
TOTAL	102	100%

Table 8 presents profile of the mothers pertaining to a level of knowledge on the importance of exclusive feeding. The findings vividly depict the respondents’ unimpressive awareness on informed choice. Sixty-four (64) respondents or at least sixty-three per cent have revealed little knowledge on the importance of exclusive feeding; in direct contrast to such findings, only thirty-eight have clearly said that they are highly knowledgeable on the importance of exclusive feeding.

The data effectively characterize the penchant of the respondents on the health and emotional benefits of exclusive breastfeeding from a mother to her child. This is a testament to the seemingly inadequate information drive and advocacy by the Rural Health Unit staff to keep the mothers and other family members aware on the importance of exclusive breastfeeding. Knowledge on the importance of EBF among mothers of infant ages 0-6 months has bases and impact on the success and failure of the breastfeeding program of the government.

Mothers, whose age ranges from 21 to 25 were the most prevailing, with 32 percent of the total respondents, 17 out of 33 in this age group or 39%, of the total mother practicing EBF followed by mother in age group 26 to 30 with 15 out of 30 or 34 %. Mixed feeding is also predominant in age group 21-25 yrs. old with 14 out of 40 mothers who is mixed feeding or 35%. Followed by mother in age group 15-20 yrs. old with 13 out of 40 or 33%, while age group from 26-35 yrs. old, bottle feeding is predominant or 78% of the total bottle-feeding mothers. In Mauritius, Breastfeeding Practices among participants of the study ranged from 18 to 45 years old whereby the majority of the participants belonged to the age group 25–31 years which is almost the same with this study.

The table above shows majority of the respondents has children ranging from 1 to 3, (67%) followed by mothers with 4 to 7 children (31%), and only 2 respondent has children ranging from 8-11. The result shows that, the more the children the respondent had the higher the percentage of practicing EBF, respondent with children 8-11 are practicing 100% EBF, followed by mother with 4-7 children 69%, while mixed feeding is predominant in mother with 1-3 children and bottle feeding is predominant in mother with 4-7 children 8 out 32 children (25%).

Of the 102 respondents, fifty-six or 55% said that they are living in with a partner, either married through church wedding, civil rights or in the state of cohabitation. On the other hand, 45% or 46 mothers divulged that they were actually single, characterized in this study as mother/s with children but no spouse, the result shows that EBF practice is common in married mother with 38 out of 56 (67%) compared to 13% among single mother. mixed feeding and bottle feeding is common to single mother with result as follows 25 out of 46 single mother or 54% and 14 out of 46 or 32% respectively.

Result shows that lower income mother 43 out of 96 (45%) practice EBF more compared to middle income mother 1 out of 6 (17%) are practicing EBF. While mixed feeding is predominant in middle income mother with 4 out 6 (67%) compared to 38% among low income mother. Bottle feeding is also dominant in low income mother 17 out of 96 or 18% of low-income mother. Study of Millman (1981), Popkin (1978), Mohiuddin (1986), Mott (1984) and Othman (1985) prove that socioeconomic status is negatively correlated with breastfeeding duration.

In terms of feeding practices based on employment status the result shows that mother without job practice EBF 40 out of 79 mothers (51%) followed by mother working full time in private jobs 40% 4 out of 10 mothers, while bottle feeding is common in mothers working permanently in government 3 out of 4 or 75%. Mixed feeding is dominant in respondent working in government as job order 6 out of 9 respondent or 67% of the total working in government as job order.

revealed that exclusive breastfeeding are practiced more by elementary graduate mothers 19 out of 30 (63%) followed by high school graduate mothers 21 out of 40 mother or 53%. College graduate mothers are the least who practice EBF 1 out of 14 (7%) mainly because this mother is working regularly in government or private offices.

In other study, mixed feeding is common among college level mother 10 out of 18 (56%), followed by high school graduate 16 out of 40 mothers (40%) and elementary graduate 10 out of 30 mothers (33%). Using a sample of 2,293 urban and 712 rural Cebu households, they have proven that better- educated women and women with higher socioeconomic status are likely to breastfed.

Bottle feeding is predominant in college graduate and college level with result as follows; 9

out of 14 (64%) and 5 out of 18 (28%) respectively.

From the table, it points out to the fact that forty-four (44) mothers out of one hundred two (102) respondents said that for their babies, they are doing exclusive breastfeeding. Forty (40) respondents said they are into mixed feeding; while only eighteen (18) are using formula feeding.

The findings vividly depict the respondents' limited awareness on informed choice. Sixty-four (64) respondents or at least sixty-three per cent have revealed little knowledge on the importance of exclusive feeding; in direct contrast to such findings, only thirty-eight have clearly said that they are highly knowledgeable on the importance of exclusive feeding. I find out based on others previous studies that factors like the planned duration of RBFF, socioeconomic status, infant's gender, maternal employment outside the household, birth weight, and maternity ward are the determinants of EBF common to more than one country

Conclusion

On the age of the respondents, mother belonging to 21-25 age groups has the higher rate of Exclusive Breastfeeding practice. Mothers with living children from 4 to 11 children are practicing exclusively breastfeeding. Exclusive breastfeeding practice is high on married mothers. Result shows that lower income mother 43 out of 96 (45%) practice EBF. In terms of feeding practices based on employment status the result shows that majority of mother without job practice Exclusive Breast Feeding 40 out of 79 mothers (51%). With regards to the educational attainment of mothers, respondents who graduated from elementary is the one who practice more exclusive breastfeeding. Majority of the respondents, currently practicing breastfeeding. Majority of the mothers has little knowledge on the importance of exclusive breastfeeding.

Recommendation

Proper breastfeeding practices are effective ways in reducing childhood morbidity and mortality. While many mothers understand the importance of breastfeeding, others are less knowledgeable on the benefits of breastfeeding and weaning. The aim here is to provide adequate information continuously thru bench conferences in the barangay level for all age group especially the very young and less educated mothers.

Adequate nutrition using breast milk during the first 6 months is essential to ensure the growth, health, and development of children to their full potential. It is necessary that first time mothers understand the health messages conveyed to them by using local dialect in health teaching.

Single mothers' perception about breastfeeding practice should be corrected through counseling early on during prenatal visit. Encourage family support by doing home visitation of the RHU or health center staff. Providing incentives to low income mothers who EBF her child like Phil-health enrollment, free medical check-up will encourage mother to fully breastfed.

Government should encourage employer to provide space in their offices as breastfeeding areas to provide privacy for nursing mother. Especial time allocation/schedule should also be provided to mothers to make sure that the child can be fed well. Nurseries are also encouraged to be put up in offices so that mothers and baby are nearer with each other during feeding time.

Importance, advantages of breastfeeding should be taught in elementary grade 5 and 6 to instill awareness. Governmental health services need to emphasize and support the importance of regular training programs to all Rural Health Unit and hospital staff, especially those involved

in antenatal clinics and maternity wards, as they can influence the early infant feeding practices among the new generation of mothers.

Conflict of interest

The authors declare no conflict of interest.

Contributions from the authors

All the authors contributed to the realization of this work. All authors also declare that they have read and approved the final version of the manuscript.

References

- Ali S, Ali SF, Imam A.M, Ayub S, Billoo A.G. (2011). Perception and practices of breastfeeding of infants 0-6 months in an urban and a semi-urban community in Pakistan: a cross-sectional study. *J Pak Med Assoc.* 61(1):99-104. PMID: 22368919.
- Calderon C. (2003). National demographic health survey. Manila: National Statistics Office; The direction of causality between financial development and economic growth. *Journal of Development Economics*, 72(1), 321-334.
- Donald Sherbourne', C and Stewart A. I. (1991). The Mos Social support survey. *Sot. Sci. Med.* Vol. 32, No. 6, pp. 705-714, 1991. 0277-9536/91, Printed in Great Britain, Pergamon Press plc.
- Jason Millman. (1981). Handbook of Teacher Evaluation. *Beverly Hills, Calif.: Sage Publications, 1981, 66 (451), page(s): 120-121*
<https://doi.org/10.1177/019263658206645122>
- Mohiuddin, S.M. Reddy, M.V. Reddy, M.M. (1986). Ramakrishna, K. "Studies on phagocytic activity and hematological changes in aflatoxicosis in poultry". *Indian Veterinary Journal (India) ISSN: 0019-6479, 63 (6) Pp. 442 – 445.*
- Motee A, Ramasawmy D, Pugo-Gunsam P, Jeewon R. (2013). An Assessment of the Breastfeeding Practices and Infant Feeding Pattern among Mothers in Mauritius. *J Nutr Metab.* 243852. doi: 10.1155/2013/243852.
- Oliver E. (2009). Williamson A Comparison of Alternative Approaches to Economic Organization, *Journal of Institutional and Theoretical Economics (JITE)* 146 (1) pp. 61-71, <https://www.jstor.org/stable/40751303>
- Onah, S., Osuorah, D.I.C., Ebenebe, J. (2014) Infant feeding practices and maternal socio-demographic factors that influence practice of exclusive breastfeeding among mothers in Nnewi South-East Nigeria: a cross-sectional and analytical study. *Int Breastfeed J* 9, 6., <https://doi.org/10.1186/1746-4358-9-6>.
- Popkin, R.H. Pre-adamism in 19th century American thought: "Speculative biology" and racism. *Philosophia* 8, 205–239 (1978). <https://doi.org/10.1007/BF02379241>
- Rafael Pérez-Escamilla, Chessa Lutter, Maria Segall, Ada Rivera, Sandra Treviño-Siller, Tina Sanghvi. Exclusive Breast-Feeding Duration Is Associated with Attitudinal, Socioeconomic and Biocultural Determinants in Three Latin American Countries, *The Journal of Nutrition*, 125 (12), 2972–2984, <https://doi.org/10.1093/jn/125.12.2972>
- Sonia Isoyama Venancio and Carlos Augusto Monteiro. Individual and contextual determinants of exclusive breast-feeding in São Paulo, Brazil: a multilevel analysis *Public Health Nutrition* 9(1), 40–46, 2006. DOI: 10.1079/PHN2005760

Tuladhar JM. Effect of family planning availability and accessibility on contraceptive use in Nepal. *Stud Fam Plann.* Jan-Feb;18 (1), 1987:49-53. PMID: 3824424 United Nations Children's Fund. *At a glance: Philippines.* In: United Nations Children's Fund [website]. New York: United Nations Children's Fund; 2013. (http://www.unicef.org/infobycountry/philippines_statistics.html, accessed May 2015.)

World Health Organization. *Infant and young child feeding counselling: an integrated course.* Geneva: UNICEF and WHO; 2006.