



## OPTIMIZATION OF SOLVENT AND CONCENTRATION OF TURMERIC (*Curcuma longa* Linn.) EXTRACT FOR STRIP-TEST AS BORAX DETECTION TOOL

By

Yulianita Pratiwi Indah Lestari<sup>1</sup>, Rizki Ramadani<sup>2</sup>, Rahmawati<sup>3</sup>  
<sup>1,2,3</sup>Faculty of Pharmacy, Universitas Muhammadiyah Banjarmasin  
Email : [rahmawati@umbjm.ac.id](mailto:rahmawati@umbjm.ac.id)

### Article Info

#### Article history:

Received Feb 11, 2022

Revised Feb 20, 2022

Accepted Mar 25, 2022

#### Keywords:

Turmeric, Borax, Borax  
Detector, Qualitative Analysis,  
Concentration Optimization

### ABSTRACT (10 PT)

Borax, a well-known toxic substance, had been banned as food additive in many countries, including Indonesia. Vomiting, fatigue, renal failure, cancer and even sudden death were reported as its toxic effect. There are many studies about Borax identification had been carried out, including instrumental procedures and alternative tests using natural indicator from the plants, such as turmeric. Alternative tests were low-cost, simple and easy to do. However, a certain solvent and concentration of turmeric for borax detection has not studied yet, which made a gap in borax analysis using a natural indicator. This research aims to determine the effect of turmeric solvent extraction and its concentration on borax detection results using the Turmeric Paper Test Kit. Turmeric powder was macerated with three different solvents (methanol, ethyl acetate and n-hexane) for 3 days. The highest yielded extract was methanol extract, followed by ethyl acetate and n-hexane extract. Solution of Turmeric extract in various solvents were prepared in many concentrations and dripped on paper of strip-test detection tool and leaved to dry. Borax detection was performed with dripped on strip-test detection tools, and discoloration of turmeric paper was observed. The most optimal solvent in this research is methanol extract with 10.000 ppm concentration, and discoloration to reddish-brown color.

*This is an open access article under the [CC BY-SA](https://creativecommons.org/licenses/by-sa/4.0/) license.*



### Corresponding Author:

Rahmawati  
Faculty of Pharmacy  
Universitas Muhammadiyah Banjarmasin  
Email: [rahmawati@umbjm.ac.id](mailto:rahmawati@umbjm.ac.id)

## 1. INTRODUCTION

Food additives are mixed materials added to food in the manufacture, processing, preparation, treatment, packing, packaging, storage, or transportation of food to produce or are expected to produce (directly or indirectly) with the aim of making the nature or form of food in accordance with what is intended, expected[1]. According to the Regulation of the Head of BPOM RI Number 36 of 2013 concerning the Maximum Limit for the Use of Food Additives, preservatives are food additives to prevent or inhibit fermentation, acidification, decomposition, and other damage to food caused by microorganisms [2].

Previously, borax commonly used as preservative[3], increase food elasticity[4], strengthen and brighten the food[5], also make the chewy texture[6]. Until the Joint FAO/WHO Expert Committee on Food Additives (JECFA) banned this compound, because it is inaplicable as a food additive[4]. Borax is a crystalline compound, white in color, odorless and stable at normal pressure temperatures but also very toxic. Borax accumulation in the body could give some adverse reaction such as vomiting, fatigue, and renal failure because it has a toxic effect that can harm the metabolic system of human health such as irritation of the respiratory tract, skin, eyes, and target organs such as blood, kidneys, heart, respiratory system, central nervous system, liver, lymph, digestive system, eyes, reproductive system, and skin[7]. The use of borax for a long time and in large quantities can cause cancer. Increasing the dose of borax can result in edema, cell inflammation, neovascularization, and very high doses which result in sudden death[8][9]. As a result, many countries have banned this substance as food additive[4], [5], including Indonesia[1]. According to the Regulation of the Minister of Health of the Republic of Indonesia No. 033 of 2012 concerning food additives, borax is one of the types of food additives that are prohibited from being used in food products, even consuming 5 – 10 g/kg body weight, or 5,000 – 10,000 ppm can cause weakness, kidney damage, even shock and death.

Borax is a sodium salt of  $\text{Na}_2\text{B}_4\text{O}_7 \cdot 10\text{H}_2\text{O}$  which is often used in various non-food industries including soldering materials, paper, cleaning agents, glass, antiseptics, wood preservatives, cockroach control, and

ceramics. Pyrex glass which is often used in laboratory glassware is made with a mixture of borax[10]. However, some sellers are still stubborn about adding borax with the excuse of making a profit.

Various studies on borax identification have been carried out, both instrumental procedures and simple evaluation. Borax could be identified using spectrophotometer UV-Vis (Colorimetry)[15], [16],[17] FTIR-ATR[18], optic fiber[19], liquid chromatography inductively coupled plasma mass spectrometry (LC-ICP-MS)[4], and high-throughput SELEX for screening borax-specific ssDNA aptamer[5]. Meanwhile, a simpler way to identify borax also growing rapidly, such as a drop test [20], [21], tooth picks[22] flame test and turmeric paper[23], [24]. Those simple evaluation, mostly using a natural indicator from turmeric[21], purple cabbage and sappanwood[25], and fluorescent probe from water hyacinth leaves[26]. Instrumental methods described before, were expensive and need special procedures, while the alternative tests are low cost, simple and easy to do.

In Indonesia, turmeric is easy to obtain, and the presence of Borax could be confirmed by discoloration, to a red orange to red color in food products containing borax[27]. Simply put, curcumin in turmeric can break the bonds of borax into boric acid and form curcumin boron cyano complex, a complex rosin compound[22]. This reaction is easy to observe, but still there are no exploration about turmeric concentration that could detect the presence of borax in food. Hence, it is become an uncertain condition for detecting Borax using a natural indicator from turmeric. In this study, various concentrations of turmeric extract and various solvents were used. The purpose of this study is as follows, namely to determine whether there is an effect of turmeric concentration on the detection results on the Turmeric Paper Strip-Test Kit, and to determine whether there is an effect of solvent on the detection results on the Turmeric Paper Strip-Test Kit.

## 2. RESEARCH METHOD

### TOOLS

Laboratory equipment made of glass, filter paper, glass bottles, oven, dropper,

### PLANT RAW MATERIAL

Turmeric (*Curcuma longa* Linn.).

### CHEMICAL MATERIAL

Aquadest, methanol, ethyl acetate, and n-hexane (PT. Smart Lab Indonesia, Indonesia), Borax (Merck).

### TURMERIC POWDER PREPARATION

The turmeric tuber flesh is dried in an oven at a temperature of around 60-70°C to remove the moisture content. After drying, turmeric is ground using a grinder until it becomes a fine powder. Then the sample was sieved through a 60 mesh sieve.

### TURMERIC EXTRACT PREPARATION

The turmeric powder that has been made is then added with a solvent in a ratio of 1:5 (until the simplicia powder is submerged in the solvent). Then do the maceration for 3 days. The collected macerate was then aerated at a cool temperature (air-conditioned room) until it became a thick extract.

### PREPARATION OF TURMERIC PAPER STRIP-TEST KIT

The process of making this borax detector is as follows:

1. Prepare a solution of Turmeric extract in various solvents (methanol, ethyl acetate, n-hexane) with various concentration series (1,000 ppm; 2,500 ppm; 5,000 ppm; 7,500 ppm; 10,000 ppm). Each made as much as 10 mL.
  2. Stir until smooth and transfer to a wider container.
    3. Take mica plastic size 7x1 cm
    4. Take filter paper, cut a 1x1 cm square and dip it in the turmeric solution, back and forth using tweezers until evenly distributed over the entire surface of the filter paper.
    5. Then place it on a baking sheet to dry by aerating.
    6. After drying, store in a closed container. Turmeric Paper Strip-Test Kit is ready to use.
- (Hartati, 2017 with modification).




Figure 1. Test strip sketch



### Preparation Of Borax Solution

A 1000 ppm borax solution was made by weighing as much as 100 mg of borax into a 100 mL volumetric flask. Dissolve with distilled water up to 100 mL (Adawiyah, 2014 with modification).

### Qualitative Test On The Turmeric Paper Test Kit

The test was carried out by dripping 2-3 drops of a 1000 ppm borax solution onto the turmeric paper test kit that had been made. Let stand a few moments until the paper dries and the resulting color is stable. Observe the color change that occurs.

### DATA PROCESSING

Data obtained from laboratory test results is then processed and entered into a table which is then described.

## 3. RESULTS AND ANALYSIS (10 PT)

### 3.1 Turmeric Powder Preparation



Figure 2. Turmeric powder

The yield of symplisia was 62.145% while the powder yield was 25.866%. Making simplisia powder can reduce the size of the turmeric rhizome so that it will be able to increase the surface area of the turmeric rhizome and will optimize the extraction process.

### 3.2 Turmeric Extract Preparation





Figure 3. Turmeric extract preparation

(a) Maceration process, (b) Metanol Extract, (c) Etil Acetate Extract, (d) n-Hexane Extract

*Curcuma longa* L. a family of Zingiberaceae, contain curcumin, as the main component of turmeric, demethoxycurcumin and bisdemethoxycurcumin, as secondary components, as part of the “curcuminoid” family [28][29]. Curcumin is a hydrophobic polyphenolic compound derived from turmeric rhizome, which consists about 2-5% of the total rhizome content and is a more valuable component of turmeric[30]. Due to the activity of natural indicator for detecting borax, turmeric powder macerated using three different solvent methanol, ethyl acetate and n-hexane. In this study, polar and non-polar organic solvents have been used to extract curcumin from *C. longa* L. The highest yield extract obtained from methanol solvent, yields 5,14 g extract (14,69%), followed by ethyl acetate (3,63 g; 10,37%) and n-hexane (1,71 g; 4,89%). This result showed that methanol is the best solvent to extract turmeric powder. This result relevant to Priyadarsini’s study which states that curcumin known readily soluble in polar solvents like methanol and ethyl acetate[31]. Curcumin is almost insoluble in water in acidic and neutral pH, so water extraction could be impossible to extract the curcumin. Creating a detection tool could be simple and easy

to do, but using an inappropriate solvent to extract the main natural indicator, could give a wrong analysis. Even in Kiamahalleh et al study, they reported that the maximum amount of curcumin extracted at the optimum condition with subcritical water extraction was only 3.8wt% [30].





### 3.3 Qualitative Test on The Turmeric Strip-Test




**Figure 4. Result of qualitative test on the turmeric strip-test**

Qualitative test of borax using turmeric strip-test showed discoloration of turmeric paper. Curcumin, which contained in turmeric paper reacted to borax, and showed color changes. Curcumin compounds found in turmeric could break down borax bonds into boric acid and bind them into a complex color of rosocyanine (brownish red) in an acidic atmosphere, thus causing a red orange to red color in food products containing borax [27]. Curcumin is very sensitive to Ph change, that can cause structure formation [20]. The discoloration of turmeric paper from yellow to reddish, was the reason of turmeric utilization as natural indicator material to detect the presence of borax in alternative test. Another content contained in turmeric is essential oil (volatile oil) 1-3% which has characteristics of sharp smell and taste. The smell and taste come from several substances contained in the oil so that turmeric can be used as a traditional medicine to prevent some diseases [32]. Curcumin which is known as the most biologically active form of curcuminoids family in turmeric, can form a molecular complexes with various charged species, including tetraedric boron complexes (rosocyanine) [27], which inclined the use of it as natural indicator.

Methanol extract has given the most intense color of turmeric extract, with yellow color on turmeric paper compared to ethyl acetate and n-hexane extract. It could be happened, prompt to curcumin levels as major compound in turmeric extract. The higher the concentration of turmeric extract used, the more intense the color changes on the paper (Table 1). Those concentrations of various turmeric extract, also influenced borax identification using turmeric strip-test detection tool. Turmeric extract that produced the most intense color in this study was EM 10.000 ppm with the color result in the form was yellow orange, meanwhile after dripping Borax solution, it become the most optimal discoloration with the result reddish-brown color.

Concentration	Warna yang dihasilkan	
	Before	After
EH 1.000 ppm		
EH 2.500 ppm		



\*EH = Extract of n-Hexane; EE = Extract of Etil Acetate; EM = Extract of Metanol

#### 4. CONCLUSION

Thus, based on the results of the tests that carried out, choosing the right solvent to extract the curcumin, as main natural indicator in detecting the presence of Borax must be done. In accordance to that, deciding a concentration

of turmeric extract used in a detection tool requisite a proper concentration to simplify the color changes observation. Increasing the concentration of turmeric has an effect in the form of increasing the intensity of the color results in the test kit. The magnitude of the increase in color was in line with the increase in the concentration of turmeric extract. The higher the concentration of turmeric extract used, the more intense the color produced. The concentration of turmeric extract that produced the most intense color in this study was EM 10.000 ppm with the color result in the form was yellow orange. The type of solvent also affects the intensity of the color results in the test kit. The most optimal solvent in this research is EM (Metanol Extract) 10.000 ppm with the result reddish-brown color.

#### ACKNOWLEDGEMENTS

Thanks to all those who contributed to this research

#### REFERENCES

- [1] Kementerian Kesehatan RI, "PerMenKes Republik Indonesia No. 33 tentang Bahan Tambahan Pangan," *PerMenKes Republik Indones. No. 33*, pp. 1–10, 2012.
- [2] BPOM, "Peraturan Kepala Badan Pengawas Obat Dan Makanan Republik Indonesia Nomor 36 Tahun 2013 Tentang Batas Maksimum Penggunaan Bahan Tambahan Pangan Pengawet," pp. 1–16, 2013.
- [3] Sugiatmi and I. A. Poetra, "Knowledge and Practice of the Use of Borax in Meatballs Seller At the Campus a University of Muhammadiyah Jakarta," in *Proceedings: The 2nd International Multidisciplinary Conference*, 2016, vol. 1, no. 1, pp. 744–750.
- [4] C. Dinh Viet *et al.*, "Simultaneous determination of borax and polyphosphates content in food by liquid chromatography inductively coupled plasma mass spectrometry (LC-ICP-MS)," *Heavy Met. Arsen. Conc. water, Agric. soil, rice Ngan Son Dist. Bac Kan Prov. Vietnam*, vol. 4, no. 2, pp. 115–126, 2021, doi: 10.47866/2615-9252/vjfc.3788.
- [5] L. Jing *et al.*, "A novel borax-specific ssDNA aptamer screened by high-throughput SELEX and its colorimetric assay with aggregation of AuNPs," *J. Food Compos. Anal.*, vol. 101, no. November 2020, p. 103947, 2021, doi: 10.1016/j.jfca.2021.103947.
- [6] I. P. Sari, F. A. Yanti, D. I. Saefullah, and B. T. Yunianto, "Identification of borax in meatballs at Ciroyom Market, Bandung City, Indonesia," *J. Sustain. Sci. Technol.*, vol. 1, no. 1, pp. 44–51, 2021, doi: 10.23960/josst.v1i1.8.
- [7] Aseptianova, D. Afriansyah, and M. Astriani, "Penyuluhan Bahan Makanan Yang Mengandung Boraks Di Kelurahan Kebun Bunga Kota Palembang," *J. Batoboh*, vol. 2, no. 1, pp. 5–24, 2017.
- [8] M. Kabu, M. Tosun, B. Elitok, and M. S. Akosman, "Evaluación histológica de los efectos del borax obtenido de diversas fuentes en diferentes Órganos de rata," *Int. J. Morphol.*, vol. 33, no. 1, pp. 255–261, 2015, doi: 10.4067/S0717-95022015000100040.
- [9] A. W. Prasetya and L. Dewi, "Deteksi Kandungan Rhodamin B Pada Saus Serta Cemaran Boraks Dan Bakteri Salmonella Sp. Padacilok Keliling Salatiga," *Agric*, vol. 28, no. 1, p. 69, 2017, doi: 10.24246/agric.2016.v28.i1.p69-78.
- [10] L. K. Muharrami, "Analisis kualitatif kandungan Boraks pada krupuk puli di Kecamatan Kamal," *J. Pena Sains*, vol. 2, no. 2, 2015.
- [11] Ernati, "Tingkat Pendidikan, Pengetahuan, Sikap Pedagang Bakso dan Penggunaan Boraks Pada Bakso di SDN Lemahputro III Sidoarjo," *J. Kesehat. Lingkungan*, vol. 9, no. 2, pp. 209–216, 2017.
- [12] D. Suseno, "Analisis Kualitatif dan Kuantitatif Kandungan Boraks Pada Bakso Menggunakan Kertas Turmeric, FT – IR Spektrometer dan Spektrofotometer Uv -Vis," *Indones. J. Halal*, vol. 2, no. 1, p. 1, 2019, doi: 10.14710/halal.v2i1.4968.
- [13] Nurkholidah, M. Ilza, and C. Jose, "Analisis Kandungan Boraks Pada Jajanan Bakso Tusuk Di Sekolah Dasar Di Kecamatan Bangkinang Kabupaten Kampar," *J. Ilmu Lingkungan*, vol. 6, no. 2, pp. 134–145, 2013, [Online]. Available: <http://ejournal.unri.ac.id/index.php/JIL/article/view/963>.
- [14] C. Rahma and Y. Hidjrawan, "Qualitative Identification of Borax Content in Meatball Snacks Using Turmeric Paper and Shallot Extract," *J. Pangan dan Agroindustri*, vol. 9, no. 1, pp. 56–63, 2021, doi: 10.21776/ub.jpa.2021.009.01.6.
- [15] N. Ratnarathorn and W. Dungchai, "Paper-based Analytical Device (PAD) for the Determination of Borax, Salicylic Acid, Nitrite, and Nitrate by Colorimetric Methods," *J. Anal. Chem.*, vol. 75, no. 4, pp. 487–494, 2020, doi: 10.1134/S1061934820040127.
- [16] R. E. Y. Adu, R. Roto, and A. Kuncaka, "Spectrophotometric Determination of Boron in Food Products By Ester Borate Distillation Into Curcumin," vol. 15, no. 1, pp. 67–73, 2021.
- [17] Q. A. Lathifah, L. Turista, Dora Dayu Rahma Azizah, and A. E. Khulaifi, "Identification of formalin and borax on tuna in Ngemplak market Tulungagung regency," *Med. Lab. Anal. Sci. J.*, vol. 1, no. 1, pp. 1–5, 2019.
- [18] T. Septiani and A. P. Roswien, "Analisis Kualitatif Kandungan Boraks Pada Bahan Pangan Daging Olahan



- dan Identifikasi Sumber Boron dengan FTIR – ATR,” *Indones. J. Halal*, vol. 1, no. 1, p. 48, 2018, doi: 10.14710/halal.v1i1.3403.
- [19] M. Yuniyanto, D. P. Andaresta, F. Anwar, and T. E. Saraswati, “Preliminary study on the use of optical fiber to detect borax content in meat,” *J. Phys. Conf. Ser.*, vol. 1825, no. 1, 2021, doi: 10.1088/1742-6596/1825/1/012078.
- [20] N. Rahman, “Formalin and Borax Qualitative Test Use Natural Indicator,” *J. Phys. Conf. Ser.*, vol. 1363, no. 1, 2019, doi: 10.1088/1742-6596/1363/1/012092.
- [21] E. D. Astuti and W. S. Nugroho, “Kemampuan Reagen Curcumax Mendeteksi Boraks dalam Bakso yang Direbus,” *J. Sain Vet.*, vol. 35, no. 1, p. 42, 2017, doi: 10.22146/jsv.29289.
- [22] R. Samson, “BoDex : A User-friendly Qualitative Test Kit,” pp. 1–18, 2015.
- [23] S. Harimurti, I. H. Bariroh, A. Setiyawan, R. I. Permatasari, F. D. Putri, and L. Y. Fajriana, “Identification of the Spread of Borax Use in Meatball Skewers in Bantul District, Special Region of Yogyakarta,” *J. Pharm. Sci. Community*, vol. 17, no. 2, pp. 120–130, 2020, doi: 10.24071/jpsc.002369.
- [24] H. Rahman, M. Prajuwita, D. Z. Yanni, P. M. Sari, and I. Lestari, “Analysis of Borax in Ground Red Chilies in the Traditional Markets of Jambi City,” *EduChemia (Jurnal Kim. dan Pendidikan)*, vol. 5, no. 1, p. 10, 2020, doi: 10.30870/educhemia.v5i1.6520.
- [25] T. E. Purbaningias, I. D. Lestari, B. Wiyantoko, P. Kurniawati, and D. Sriadryani, “Utilization of natural indicators for borax identification in the Indonesian tofu,” *AIP Conf. Proc.*, vol. 1823, no. March 2017, 2017, doi: 10.1063/1.4978130.
- [26] T. Prathumsuwan, P. Jaiyong, I. In, and P. Paoprasert, “Label-free carbon dots from water hyacinth leaves as a highly fluorescent probe for selective and sensitive detection of borax,” *Sensors Actuators, B Chem.*, vol. 299, no. July, p. 126936, 2019, doi: 10.1016/j.snb.2019.126936.
- [27] G. Grynkwicz and P. Ślifirski, “Curcumin and curcuminoids in quest for medicinal status,” *Acta Biochim. Pol.*, vol. 59, no. 2, pp. 201–212, 2012, doi: 10.18388/abp.2012\_2139.
- [28] K. Doldolova, M. Bener, M. Lalikoğlu, Y. S. Aşçı, R. Arat, and R. Apak, “Optimization and modeling of microwave-assisted extraction of curcumin and antioxidant compounds from turmeric by using natural deep eutectic solvents,” *Food Chem.*, vol. 353, no. January, 2021, doi: 10.1016/j.foodchem.2021.129337.
- [29] S. Tripathy *et al.*, “Curcumin Extraction, Isolation, Quantification and Its Application in Functional Foods: A Review With a Focus on Immune Enhancement Activities and COVID-19,” *Front. Nutr.*, vol. 8, no. September, 2021, doi: 10.3389/fnut.2021.747956.
- [30] M. Valizadeh Kiamahalleh, G. Najafpour-Darzi, M. Rahimnejad, A. A. Moghadamnia, and M. Valizadeh Kiamahalleh, “High performance curcumin subcritical water extraction from turmeric (*Curcuma longa* L.),” *J. Chromatogr. B Anal. Technol. Biomed. Life Sci.*, vol. 1022, pp. 191–198, 2016, doi: 10.1016/j.jchromb.2016.04.021.
- [31] K. I. Priyadarsini, “The Chemistry of Curcumin: From Extraction to Therapeutic Agent,” *Molecules*, vol. 19, pp. 20091–20112, 2014, doi: 10.3390/molecules191220091.
- [32] A. Rohman, “Analysis of curcuminoids in food and pharmaceutical products,” in *International Food Research Journal*, 2012, vol. 19, no. 1, pp. 19–27.

HIS PAGE HAS INTENTIONALLY BEEN LEFT BLANK