



How behave the typical L^q -dimensions of measures?

Frédéric Bayart

► **To cite this version:**

| Frédéric Bayart. How behave the typical L^q -dimensions of measures?. 2012. <hal-00678595>

HAL Id: hal-00678595

<https://hal.archives-ouvertes.fr/hal-00678595>

Submitted on 13 Mar 2012

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

HOW BEHAVE THE TYPICAL L^q -DIMENSIONS OF MEASURES?

FRÉDÉRIC BAYART

ABSTRACT. We compute, for a compact set $K \subset \mathbb{R}^d$, the value of the upper and of the lower L^q -dimension of a typical probability measure with support contained in K , for any $q \in \mathbb{R}$. Different definitions of the “dimension” of K are involved to compute these values, following $q \in \mathbb{R}$.

1. INTRODUCTION

Let K be a compact subset of \mathbb{R}^d , and let $\mathcal{P}(K)$ be the set of Borel probability measures on K ; we endow $\mathcal{P}(K)$ with the weak topology. In this paper, we are interested in properties shared by typical measures of $\mathcal{P}(K)$. By a property true for a typical measure of $\mathcal{P}(K)$, we mean a property which is satisfied by a dense G_δ set of elements of $\mathcal{P}(K)$. Specifically we deal with the upper and lower L^q -dimensions of measures. Let $\mu \in \mathcal{P}(K)$, $r > 0$ and $q \in \mathbb{R} \setminus \{1\}$. We write

$$I_\mu(r, q) = \int_K \mu(B(x, r))^{q-1} d\mu(x).$$

The lower and upper L^q -dimensions are now defined, for $q \neq 1$, by

$$\begin{aligned} \underline{D}_\mu(q) &= \liminf_{r \rightarrow 0} \frac{1}{q-1} \times \frac{\log I_\mu(r, q)}{\log r} \\ \overline{D}_\mu(q) &= \limsup_{r \rightarrow 0} \frac{1}{q-1} \times \frac{\log I_\mu(r, q)}{\log r}. \end{aligned}$$

When $q = 1$, the definitions involve a logarithmic factor:

$$\begin{aligned} I_\mu(r, 1) &= \int_K \log \mu(B(x, r)) d\mu(x) \\ \underline{D}_\mu(1) &= \liminf_{r \rightarrow 0} \frac{\int_K \log \mu(B(x, r)) d\mu(x)}{\log r} \\ \overline{D}_\mu(1) &= \limsup_{r \rightarrow 0} \frac{\int_K \log \mu(B(x, r)) d\mu(x)}{\log r}. \end{aligned}$$

These dimensions were introduced by Hentschel and Procaccia in [HP83] in order to generalize the information dimension of a measure. They are important for their relationship with the multifractal formalism. This formalism, which was conjectured in the 1980s in the physic literature, asserts that for “good” measures, $\underline{D}_\mu(q) = \overline{D}_\mu(q)$ for any $q \in \mathbb{R}$ and that the Hausdorff multifractal spectrum of μ coincides with the Legendre transform of $\tau_\mu : \mathbb{R} \rightarrow \mathbb{R}$ defined by $\tau_\mu(q) = (1-q)\underline{D}_\mu(q) = (1-q)\overline{D}_\mu(q)$. This multifractal formalism has been verified for various classes of measures on \mathbb{R}^d , see [Fal97] and the references therein.

Date: March 13, 2012.

In a series of papers ([Ols05], [Ols07] and [Ols08]), L. Olsen made a first approach to estimate the typical value of $\underline{D}_\mu(q)$ and of $\overline{D}_\mu(q)$. To state his results, we need to introduce some classical terminology which can be found e.g. in [Fal03]. For a subset $E \subset \mathbb{R}^d$, we denote the lower box dimension of E and the upper box dimension of E by $\underline{\dim}_B(E)$ and $\overline{\dim}_B(E)$, respectively. The Hausdorff and the packing dimension of E are denoted respectively by $\dim_{\mathcal{H}}(E)$ and $\dim_{\mathcal{P}}(E)$. Also, for a subset K of \mathbb{R}^d and $x \in K$, we define the lower local box dimension of K at x and the upper local box dimension of K at x by

$$\begin{aligned}\underline{\dim}_{B,\text{loc}}(x, K) &= \lim_{r \rightarrow 0} \underline{\dim}_B(K \cap B(x, r)) \\ \overline{\dim}_{B,\text{loc}}(x, K) &= \lim_{r \rightarrow 0} \overline{\dim}_B(K \cap B(x, r)).\end{aligned}$$

Theorem A (Olsen). *Let K be a compact subset of \mathbb{R}^d . Write*

$$\begin{aligned}s_- &= \inf_{x \in K} \underline{\dim}_{B,\text{loc}}(x, K) = \inf_{x \in K} \inf_{r > 0} \underline{\dim}_B(B(x, r)) \\ s^+ &= \inf_{x \in K} \overline{\dim}_{B,\text{loc}}(x, K) = \inf_{x \in K} \inf_{r > 0} \overline{\dim}_B(B(x, r)) \\ s &= \overline{\dim}_B(K).\end{aligned}$$

Then the following holds:

	All measures $\mu \in \mathcal{P}(K)$ satisfy	A typical measure $\mu \in \mathcal{P}(K)$ satisfies
$q > 1$	$0 \leq \underline{D}_\mu(q) \leq \overline{D}_\mu(q) \leq s$	$\underline{D}_\mu(q) = 0$ $s_- \leq \overline{D}_\mu(q) \leq s^+$
$q \in [0, 1]$	$0 \leq \underline{D}_\mu(q) \leq \overline{D}_\mu(q) \leq s$	$s_- \leq \overline{D}_\mu(q) \leq s.$

Remark 1.1. In this paper, we adopt the normalization of [Ols07] instead of that of [Ols05], because it avoids unpleasant factors $(1 - q)$ all over the proofs, and also because it seems clearer if a dimension is always nonnegative. We nevertheless mention that this inverts the definition of $\underline{D}_\mu(q)$ and $\overline{D}_\mu(q)$ of [Ols05] when $q > 1$.

Remark 1.2. In [Ols08], the inequality $s_- \leq \overline{D}_\mu(q) \leq s^+$ is announced for a typical measure when $q \in [0, 1)$. However, there is a mistake in the proof, precisely in Lemma 2.3.1. A correct statement of this lemma should be:

Let $E \subset K$ and let $\mu \in \mathcal{P}(E)$. Then

$$\overline{D}_\mu(q) \leq \overline{\dim}_B(E)$$

for all $q \in [0, 1)$.

This correct statement forces to replace s^+ by s in Theorem A, which is less good. See also the forthcoming Theorem 1.6, which shows that typically $\overline{D}_\mu(q) \geq s^+$ for $q \in [0, 1)$.

Remark 1.3. It should be observed that the precise statement of Theorem A is:

a typical measure satisfies for every $q > 1 \dots$

That is formally stronger than:

for every $q > 1$, a typical measure satisfies \dots

The work of Olsen leaves open several questions:

- What happens in the remaining cases? Olsen conjectured that, for any $q < 1$, $\underline{D}_\mu(q) = 0$ and that, for any $q < 0$, $\overline{D}_\mu(q) = +\infty$.
- Can we say more for the upper L^q -dimension? Precisely, does there exist for any $q \geq 0$ a real number $f(q)$ such that a typical measure $\mu \in \mathcal{P}(K)$ satisfies $\overline{D}_\mu(q) = f(q)$? In this case, Olsen conjectured that we cannot do better than $s_- \leq \overline{D}_\mu(q) \leq s^+$ for a typical $\mu \in \mathcal{P}(K)$.

Our aim, in this paper, is to answer these questions. To this intention, we need to introduce two new ways to measure the size of a compact set. The first one measures how behaves locally the upper box dimension, uniformly in K . For a set $E \subset \mathbb{R}^d$ and $r > 0$, we denote by $\mathbf{P}_r(E)$ the largest number of pairwise disjoint balls of radius r with centers in E and by $\mathbf{N}_r(E)$ the smallest number of balls of radius r which are needed to cover E .

Definition 1.4. Let K be a compact subset of \mathbb{R}^d . The *local uniform upper box dimension* of K is the real number $\overline{\dim}_{B,\text{loc},\text{unif}}(K)$ defined by

$$\begin{aligned} \overline{\dim}_{B,\text{loc},\text{unif}}(K) &= \inf_{N \geq 1} \inf_{x_1, \dots, x_N \in K} \inf_{r_0 > 0} \limsup_{r \rightarrow 0} \inf_{i=1, \dots, N} \frac{\log \mathbf{P}_r(K \cap B(x_i, r_0))}{-\log r} \\ &= \inf_{N \geq 1} \inf_{x_1, \dots, x_N \in K} \inf_{r_0 > 0} \limsup_{r \rightarrow 0} \inf_{i=1, \dots, N} \frac{\log \mathbf{N}_r(K \cap B(x_i, r_0))}{-\log r} \end{aligned}$$

(the equivalence between these two definitions is a standard property of box-like dimensions.) It is easy to check that

$$\inf_{x \in K} \underline{\dim}_{B,\text{loc}}(x, K) \leq \overline{\dim}_{B,\text{loc},\text{unif}}(K) \leq \inf_{x \in K} \overline{\dim}_{B,\text{loc}}(x, K).$$

We shall see later that these inequalities can be strict.

We will also need another variant of the lower box dimension, which measures both the size and the connectivity of a set. For K a compact subset of \mathbb{R}^d , let $\mathcal{C}_n(K)$ be the collection of the half-closed dyadic cubes of size 2^{-n} intersecting K , namely

$$\mathcal{C}_n(K) = \left\{ \prod_{j=1}^n \left[\frac{k_j}{2^n}, \frac{k_j+1}{2^n} \right); k_j \in \mathbb{Z} \text{ and } \prod_{j=1}^n \left[\frac{k_j}{2^n}, \frac{k_j+1}{2^n} \right) \cap K \neq \emptyset \right\}.$$

Let $K_n = \bigcup_{C \in \mathcal{C}_n(K)} C$ and let $C_n(K)$ be the number of connected components of K_n . We may observe that the sequence $(C_n(K))$ is nondecreasing.

Definition 1.5. Let K be a compact subset of \mathbb{R}^d . The *box separation index* of K is the real number $\text{bsi}(K)$ defined by

$$\text{bsi}(K) = \liminf_{n \rightarrow +\infty} \frac{\log(C_n(K))}{n \log 2}.$$

It is clear that $\text{bsi}(K) \leq \underline{\dim}_B(K)$. Heuristically speaking, $\text{bsi}(K)$ is large if, for each $n \geq 1$, you need many cubes of size 2^{-n} to cover it and if these cubes are far away from each other.

It should be pointed out that neither the local uniform upper box dimension nor the box separation index can be considered as a dimension. For instance, $E \subset F$ does not imply $\overline{\dim}_{B,\text{loc},\text{unif}}(E) \leq \overline{\dim}_{B,\text{loc},\text{unif}}(F)$ or $\text{bsi}(E) \leq \text{bsi}(F)$. As an example, if you set $E = [1, 2]$ and $F = \{0\} \cup [1, 2]$, then $\overline{\dim}_{B,\text{loc},\text{unif}}(E) = 1$ whereas $\overline{\dim}_{B,\text{loc},\text{unif}}(F) = 0$. Regarding the box separation index, $\text{bsi}([0, 1]) = 0$ (the 2^{-n} -mesh cubes intersecting $[0, 1]$ are connected),

whereas $\text{bsi}(K) = 1/2$ when $K = \{0\} \cup \{1/n; n \geq 1\}$. This last fact follows easily from the standard proof of $\underline{\dim}_B(K) = 1/2$.

Our first main result now reads:

Theorem 1.6. *Let K be an infinite compact subset of \mathbb{R}^d . Write*

$$\begin{aligned} s_{\text{sep}} &= \text{bsi}(K) \\ s_u &= \overline{\dim}_{B,\text{loc},\text{unif}}(K) \\ s &= \overline{\dim}_B(K) \\ s_{\mathcal{P}} &= \dim_{\mathcal{P}}(K). \end{aligned}$$

Then the following holds:

	All measures $\mu \in \mathcal{P}(K)$ satisfy	A typical measure $\mu \in \mathcal{P}(K)$ satisfies
$q > 1$	$0 \leq \underline{D}_\mu(q) \leq \overline{D}_\mu(q) \leq s_{\mathcal{P}}$	$\overline{D}_\mu(q) = s_u$ $\underline{D}_\mu(q) = 0$
$q \in (0, 1)$	$0 \leq \underline{D}_\mu(q) \leq \overline{D}_\mu(q) \leq s$	$\overline{D}_\mu(q) = s$ $\underline{D}_\mu(q) = 0$
$q = 0$	$0 \leq \underline{D}_\mu(q) \leq \overline{D}_\mu(q) \leq s$	$\overline{D}_\mu(q) = s$ $\underline{D}_\mu(q) = s_{\text{sep}}$
$q < 0$	$0 \leq \underline{D}_\mu(q) \leq \overline{D}_\mu(q)$	$\overline{D}_\mu(q) = +\infty$ $\underline{D}_\mu(q) = s_{\text{sep}}$.

The typical values of $\overline{D}_\mu(q)$, for $q > 1$, and of $\underline{D}_\mu(q)$, for $q \leq 0$, are very interesting. Indeed, they did not take the worst possible values, what is rather atypical in analysis! From a technical point of view, these results force us to prove two inequalities instead of just one, since we cannot use the results of the second column. They also point out that the typical L^q -dimensions of measures supported by K depend heavily both on the size and on the local structure of K , since they involve the local uniform upper box dimension and the box separation index of K .

As one can guess, the proof of Theorem 1.6 requires delicate constructions of measures as well as a careful examination of the local structure of a compact set. More surprisingly, it also involves elements of graph theory. It should be noted that the unusual notions of dimension used in Theorem 1.6 can be easily computed for natural compact sets arising in multifractal analysis: Cantor sets, self-similar compact sets, finite disjoint unions of these sets,...

We turn now to the L^1 -case. This is maybe the most important case. In the literature, the L^1 -dimensions are also known as the (upper and lower) information dimension or as the (upper and lower) entropy. From Theorem 1.6, we can get immediately the typical value of $\underline{D}_\mu(1)$. However, that of $\overline{D}_\mu(1)$ is unpredictable: it is not clear if it should be s_u ,

like for $q > 1$, or s , like for $q < 1$. It turns out that neither s_u nor s is convenient. In fact, the situation breaks down dramatically for the upper L^1 -dimension: in general, there is no typical value for $\overline{D}_\mu(1)$! We cannot say more that $\overline{D}_\mu(1)$ belongs typically to some interval, and we shall give soon the optimal interval. Nevertheless, we need to introduce yet another couple of definitions.

Definition 1.7. Let K be a compact subset of \mathbb{R}^d . The *convex upper box dimension* of K is the real number $\overline{\dim}_{B,\text{conv}}(K)$ defined by

$$\overline{\dim}_{B,\text{conv}}(K) = \inf_{N \geq 1} \inf_{x_1, \dots, x_N \in K} \inf_{\rho > 0} \inf_{p_i > 0, \sum_i p_i = 1} \limsup_{r \rightarrow 0} \frac{\sum_i p_i \log \mathbf{P}_r(K \cap B(x_i, \rho))}{-\log r}.$$

The *maximal convex upper box dimension* of K is the real number $\overline{\dim}_{B,\text{conv},\text{max}}(K)$ defined by

$$\overline{\dim}_{B,\text{conv},\text{max}}(K) = \sup_{y \in K, \rho > 0} \overline{\dim}_{B,\text{conv}}(K \cap B(y, \rho)).$$

It is clear from the definition that

$$\overline{\dim}_{B,\text{loc},\text{unif}}(K) \leq \overline{\dim}_{B,\text{conv}}(K) \leq \overline{\dim}_{B,\text{conv},\text{max}}(K).$$

We shall see later that these inequalities can be strict.

Our main theorem on L^1 now reads:

Theorem 1.8. *Let K be an infinite compact subset of \mathbb{R}^d . Then a typical measure $\mu \in \mathcal{P}(K)$ satisfies*

$$\underline{D}_\mu(1) = 0 \text{ and } \overline{D}_\mu(1) \in [\overline{\dim}_{B,\text{conv}}(K), \overline{\dim}_{B,\text{conv},\text{max}}(K)].$$

Moreover, if $[a, b]$ is any interval such that a typical measure $\mu \in \mathcal{P}(K)$ satisfies $\overline{D}_\mu(1) \in [a, b]$, then

$$a \leq \overline{\dim}_{B,\text{conv}}(K) \text{ and } b \geq \overline{\dim}_{B,\text{conv},\text{max}}(K).$$

As before, even though the definition of the (maximal) convex upper box dimension is not very appealing, it can be easily computed for many compact sets, like Cantor sets, self-similar compact sets, finite disjoint unions of these sets... Here is an example to see how Theorem 1.8 reads on a very easy compact set. It points out that we cannot expect to get a typical value for the upper L^1 -dimension. This happens only for this value of q .

Example 1.9. Let $K = \{0\} \cup [1, 2]$. Then a typical measure $\mu \in \mathcal{P}(K)$ satisfies $\overline{D}_\mu(1) \in [0, 1]$ and this interval is the best possible.

The paper is organized as follows. In Section 2, we introduce the tools which are needed throughout the paper. In Section 3,4,5, we prove Theorem 1.6, whereas Section 6 is devoted to the study of the upper L^1 -dimension. We conclude in Section 7 by remarks and open questions.

2. PRELIMINARIES

2.1. The topology on $\mathcal{P}(K)$. Throughout this paper, $\mathcal{P}(K)$ will be endowed with the weak topology. It is well known (see for instance [Par67]) that this topology is completely metrizable by the Fortet-Mourier distance defined as follows. Let $\text{Lip}(K)$ denote the family

of Lipschitz functions $f : K \rightarrow \mathbb{R}$, with $|f| \leq 1$ and $\text{Lip}(f) \leq 1$, where $\text{Lip}(f)$ denotes the Lipschitz constant of f . The metric L is defined by

$$L(\mu, \nu) = \sup_{f \in \text{Lip}(K)} \left| \int f d\mu - \int f d\nu \right|$$

for any $\mu, \nu \in \mathcal{P}(K)$. We endow $\mathcal{P}(K)$ with the metric L . In particular, for $\mu \in \mathcal{P}(K)$ and $\delta > 0$, $B_L(\mu, \delta) = \{\nu \in \mathcal{P}(K); L(\mu, \nu) < \delta\}$ will stand for the ball with center at μ and radius equal to δ .

We shall use several times the following lemma.

Lemma 2.1. *For any $\alpha \in (0, 1)$, for any $\beta > 0$, there exists $\delta > 0$ such that, for any E a Borel subset of K , for any $\mu, \nu \in \mathcal{P}(K)$,*

$$L(\mu, \nu) < \delta \implies \mu(E) \leq \nu(E(\alpha)) + \beta,$$

where $E(\alpha) = \{x \in K; \text{dist}(x, E) < \alpha\}$.

Proof. We set

$$f(t) = \begin{cases} \alpha & \text{provided } t \in \overline{E} \\ \alpha - \text{dist}(x, E) & \text{provided } 0 < \text{dist}(x, E) \leq \alpha \\ 0 & \text{otherwise.} \end{cases}$$

Then f is Lipschitz, with $|f| \leq 1$ and $\text{Lip}(f) \leq 1$. Thus,

$$\begin{aligned} \mu(E) &\leq \frac{1}{\alpha} \int f d\mu \\ &\leq \frac{1}{\alpha} \left[\int f d\nu + \delta \right] \\ &\leq \nu(E(\alpha)) + \frac{\delta}{\alpha}. \end{aligned}$$

Hence, it suffices to take $\delta = \alpha\beta$. □

Our first application of Lemma 2.1 is that a small perturbation of a finite measure μ does not change dramatically the value of $I_\mu(r, q)$, provided we allow to change slightly the radius. Here is the statement that we can get for $q > 1$. $\mathcal{F}(K)$ denotes the set of probability measures with finite support in K .

Corollary 2.2. *Let $q > 1$. There exists $C_q > 0$ such that, for any $\mu \in \mathcal{F}(K)$, for any $r > 0$, there exists $\delta > 0$ such that, for any $\nu \in \mathcal{P}(K)$ with $L(\mu, \nu) < \delta$,*

$$I_\nu(r, q) \leq C_q I_\mu(2r, q) \text{ and } I_\nu(2r, q) \geq C_q^{-1} I_\mu(r, q).$$

Proof. We begin by fixing a pair (t, η) with $t \in (r, 2r)$ and $\eta > 0$ such that

$$\begin{cases} [t - \eta, t + \eta] \cap \{\|x - y\|; x, y \in \text{supp}(\mu)\} = \emptyset; \\ \forall x, y \in \text{supp}(\mu), x \neq y \implies \|x - y\| > 2\eta. \end{cases}$$

Observe that this choice of t and η guarantees that, for any $x \in \text{supp}(\mu)$ and any $z \in B(x, \eta)$, then

$$\text{supp}(\mu) \cap B(x, t) = \text{supp}(\mu) \cap B(z, t)$$

so that $\mu(B(x, t)) = \mu(B(z, t))$. Now, suppose that $\delta > 0$ has been chosen so small that any $\nu \in \mathcal{P}(K)$ with $L(\mu, \nu) < \delta$ satisfies

- for any $x \in \text{supp}(\mu)$,

$$\frac{1}{2}\mu(\{x\}) = \frac{1}{2}\mu(B(x, \eta/2)) \leq \nu(B(x, \eta)) \leq 2\mu(B(x, 2\eta)) = 2\mu(\{x\})$$

(this is possible by applying Lemma 2.1 with $\alpha = \eta/2$ and $\beta = \inf_{x \in \text{supp}(\mu)} \mu(\{x\})$).

- for any $x \in \text{supp}(\mu)$,

$$\begin{aligned} \nu(B(x, \eta)) &\geq \mu(B(x, \eta/2)) - \frac{I_\mu(2r, q)}{N} \\ &\geq \mu(\{x\}) - \frac{I_\mu(2r, q)}{N} \end{aligned}$$

where N denotes the cardinal of the support of μ . What is important here is that, setting $Y = \bigcup_{x \in \text{supp}(\mu)} B(x, \eta)$, one gets

$$\nu(Y) \geq 1 - I_\mu(2r, q).$$

- for any $x \in \text{supp}(\mu)$ and any $z \in B(x, \eta)$,

$$\nu(B(z, r)) \leq 2\mu(B(z, t)) = 2\mu(B(x, t))$$

$$\nu(B(z, 2r)) \geq \frac{1}{2}\mu(B(z, t)) = \frac{1}{2}\mu(B(x, t)).$$

Again, this is possible thanks to Lemma 2.1 and because μ has finite support.

Hence, on the one hand, we get

$$\begin{aligned} I_\nu(r, q) &= \sum_{x \in \text{supp}(\mu)} \int_{B(x, \eta)} \nu(B(z, r))^{q-1} d\nu(z) + \int_{Y^c} \nu(B(z, r))^{q-1} d\nu(z) \\ &\leq 2^q \sum_{x \in \text{supp}(\mu)} \mu(B(x, t))^{q-1} \mu(\{x\}) + \nu(Y^c) \\ &\leq 2^q I_\mu(t, q) + I_\mu(2r, q) \\ &\leq (2^q + 1) I_\mu(2r, q). \end{aligned}$$

On the other hand, we also have

$$\begin{aligned} I_\nu(2r, q) &\geq \sum_{x \in \text{supp}(\mu)} \int_{B(x, \eta)} \nu(B(z, 2r))^{q-1} d\nu(z) \\ &\geq 2^{-q} \sum_{x \in \text{supp}(\mu)} \mu(B(x, t))^{q-1} \mu(\{x\}) \\ &\geq 2^{-q} I_\mu(t, q) \geq 2^{-q} I_\mu(r, q). \end{aligned}$$

□

When $q < 1$, the situation is more difficult. Indeed, for an arbitrary close to μ measure $\nu \in \mathcal{P}(K)$, $\nu(B(x, r))^{q-1}$ may be very large even if $x \notin \text{supp}(\mu)$. However, adding an assumption to avoid singularities, we are able to prove the following corollary.

Corollary 2.3. *Let $q < 1$. There exists $C_q > 0$ such that, for any $\mu \in \mathcal{F}(K)$, for any $r > 0$ with $\mu(B(x, r)) > 0$ for any $x \in K$, there exists $\delta > 0$ such that, for any $\nu \in \mathcal{P}(K)$ with $L(\mu, \nu) < \delta$,*

$$I_\nu(2r, q) \leq C_q I_\mu(r, q) \text{ and } I_\nu(r, q) \geq C_q^{-1} I_\mu(2r, q).$$

Proof. Let $a = \min_{x \in K} \mu(B(x, r)) > 0$. a is positive since μ has finite support so that $\mu(B(x, r))$ can only take a finite number of values. As before, let $t \in (r, 2r)$ and let $\eta > 0$ be such that

$$\begin{cases} [t - \eta, t + \eta] \cap \{\|x - y\|; x, y \in \text{supp}(\mu)\} = \emptyset; \\ \forall x, y \in \text{supp}(\mu), x \neq y \implies \|x - y\| > 2\eta. \end{cases}$$

Then, for any $x \in \text{supp}(\mu)$ and any $z \in B(x, \eta)$, one gets

$$\mu(B(x, t)) = \mu(B(z, t)).$$

Now, provided $\delta > 0$ is small enough, Lemma 2.1 tells us that, for any $\nu \in \mathcal{P}(K)$ with $L(\mu, \nu) < \delta$,

$$\forall z \in K, \nu(B(z, 2r)) \geq \frac{1}{2}\mu(B(z, t)) \geq \frac{1}{4}\nu(B(z, r))$$

(recall that $\mu(B(z, t)) \geq a > 0$). Let $F = K \setminus \bigcup_{x \in \text{supp}(\mu)} B(x, \eta)$. Then

$$\begin{aligned} I_\nu(2r, q) &= \sum_{x \in \text{supp}(\mu)} \int_{B(x, \eta)} \frac{d\nu(z)}{\nu(B(z, 2r))^{1-q}} + \int_F \frac{d\nu(z)}{\nu(B(z, 2r))^{1-q}} \\ &\leq 2^{1-q} \sum_{x \in \text{supp}(\mu)} \int_{B(x, \eta)} \frac{d\nu(z)}{\mu(B(z, t))^{1-q}} + 2^{1-q} \int_F \frac{d\nu(z)}{\mu(B(z, t))^{1-q}} \\ &\leq 2^{1-q} \sum_{x \in \text{supp}(\mu)} \frac{\nu(B(x, \eta))}{\mu(B(x, t))^{1-q}} + 2^{1-q} a^{q-1} \nu(F). \end{aligned}$$

We apply Lemma 2.1 again to observe that, provided δ is small enough, for any $x \in \text{supp}(\mu)$,

$$\nu(B(x, \eta)) \leq 2\mu(B(x, 3\eta/2)) = 2\mu(\{x\})$$

and that

$$\nu(F) \leq \mu(F(\eta/2)) + \varepsilon = \varepsilon,$$

where $\varepsilon > 0$ is arbitrary. Thus we conclude that

$$\begin{aligned} I_\nu(2r, q) &\leq 2^{2-q} I_\mu(t, q) + 2^{1-q} a^{q-1} \varepsilon \\ &\leq 2^{3-q} I_\mu(r, q) \end{aligned}$$

if $\varepsilon > 0$ is small enough. For the other inequality, we simply write

$$\begin{aligned} I_\nu(r, q) &\geq \sum_{x \in \text{supp} \mu} \int_{B(x, \eta)} \frac{d\nu(z)}{\nu(B(z, r))^{1-q}} \\ &\geq \sum_{x \in \text{supp} \mu} 2^{q-1} \int_{B(x, \eta)} \frac{d\nu(z)}{\mu(B(x, t))^{1-q}} \\ &\geq \sum_{x \in \text{supp} \mu} 2^{q-1} \frac{\nu(B(x, \eta))}{\mu(B(x, t))^{1-q}}. \end{aligned}$$

Now, we can choose $\delta > 0$ to ensure that $\nu(B(x, \eta)) \geq 2^{-1}\mu(B(x, \eta/2)) = 2^{-1}\mu(\{x\})$, so that

$$I_\nu(r, q) \geq 2^{q-2} I_\mu(t, q) \geq 2^{q-2} I_\mu(2r, q).$$

□

Remark 2.4. The assumption $\mu(B(x, r)) > 0$ for any $x \in K$ is needed only to prove $I_\nu(2r, q) \leq C_q I_\mu(r, q)$.

Of course, there is a similar statement for $q = 1$. Nevertheless, we will need later a more precise result, because of the logarithm. We restrict ourselves to one inequality.

Corollary 2.5. *Let $\varepsilon > 0$, $r > 0$ and $\mu \in \mathcal{F}(K)$. There exists $\delta > 0$ such that for any $\nu \in \mathcal{P}(K)$ with $L(\mu, \nu) < \delta$,*

$$I_\nu(r, 1) \leq \log 2 + (1 - \varepsilon)I_\mu(2r, 1).$$

Proof. Again, let $t \in (r, 2r)$ and let $\eta > 0$ be such that

$$\begin{cases} [t - \eta, t + \eta] \cap \{\|x - y\|; x, y \in \text{supp}(\mu)\} = \emptyset; \\ \forall x, y \in \text{supp}(\mu), x \neq y \implies \|x - y\| > 2\eta. \end{cases}$$

Then, for any $x \in \text{supp}(\mu)$ and any $z \in B(x, \eta)$, one gets

$$\mu(B(x, t)) = \mu(B(z, t)).$$

Let $\delta > 0$ be such that, for any $x \in \text{supp}(\mu)$ and any $z \in B(x, \eta)$, for any $\nu \in \mathcal{P}(K)$ with $L(\mu, \nu) < \delta$,

$$\begin{aligned} \nu(B(z, r)) &\leq 2\mu(B(z, t)) = 2\mu(B(x, t)) \leq 2\mu(B(x, 2r)), \\ \nu(B(x, \eta)) &\geq (1 - \varepsilon)\mu(B(x, \eta/2)) = (1 - \varepsilon)\mu(\{x\}). \end{aligned}$$

Then,

$$\begin{aligned} I_\nu(r, 1) &\leq \sum_{x \in \text{supp}(\mu)} \int_{B(x, \eta)} \log(\nu(B(z, r))) d\nu(z) \\ &\leq \sum_{x \in \text{supp}(\mu)} \log(2)\nu(B(x, \eta)) + \sum_{x \in \text{supp}(\mu)} \log \mu(B(x, 2r))\nu(B(x, \eta)) \\ &\leq \log 2 + (1 - \varepsilon) \sum_{x \in \text{supp}(\mu)} \log \mu(B(x, 2r))\mu(\{x\}) \\ &\leq \log 2 + (1 - \varepsilon)I_\mu(2r, 1). \end{aligned}$$

□

Another application of Lemma 2.1 is the following result on open subsets of $\mathcal{P}(K)$:

Lemma 2.6. (a) *Let $x \in K$, $a \in \mathbb{R}$ and $r > 0$. Then $\{\mu \in \mathcal{P}(K); \mu(B(x, r)) > a\}$ is open.*

(b) *Let $E \subset K$ be such that there exists $\alpha > 0$ with $E(\alpha) \cap K = E$. Then $\{\mu \in \mathcal{P}(K); \mu(E) > 0\}$ is open.*

Proof. (a) If a does not belong to $[0, 1)$, then the set is either empty or equal to $\mathcal{P}(K)$.

Otherwise, let $\mu \in \mathcal{P}(K)$ be such that $\mu(B(x, r)) > a$. One may find $\varepsilon > 0$ such that $\mu(B(x, (1 - \varepsilon)r)) > a$. Thus the result follows from Lemma 2.1 applied with $E = B(x, (1 - \varepsilon)r)$, $\alpha = \varepsilon r$ and $\beta = (\mu(B(x, (1 - \varepsilon)r)) - a)/2$.

(b) The proof is similar (and even easier).

□

Finally, we will need that some subsets of $\mathcal{P}(K)$ are dense in $\mathcal{P}(K)$. We first recall a result which can be found e.g. in [Ols05, Lemma 2.2.4.].

Lemma 2.7. *Let $(x_i)_{i \geq 1}$ be a dense sequence of K . Let $(\mu_{n,i})_{n \geq 1, 1 \leq i \leq n}$ be a sequence of $\mathcal{P}(K)$ such that, for any $n \geq 1$ and any $1 \leq i \leq n$, $\text{supp}(\mu_{n,i}) \subset K \cap B(x_i, 1/n)$. Then, for any $m \geq 1$, $\bigcup_{n \geq m} \{\sum_{i=1}^n p_i \mu_{n,i}; p_i \geq 0, \sum_i p_i = 1\}$ is dense in $\mathcal{P}(K)$.*

Since a compact set is separable, the previous lemma yields in particular that the set $\mathcal{F}(K)$ of probability measures on K with finite support is dense in $\mathcal{P}(K)$. Moreover, taking the p_i in \mathbb{Q} , we can always consider a sequence (μ_n) of $\mathcal{F}(K)$ which is dense in $\mathcal{P}(K)$.

We shall also need several times the following result.

Lemma 2.8. *Let $n \geq 1$ and let K_1, \dots, K_n be nonempty subsets of K . Then $\{\mu \in \mathcal{P}(K); \mu(K_j) > 0 \text{ for any } j = 1, \dots, n\}$ is dense in $\mathcal{P}(K)$.*

Proof. Let $\nu \in \mathcal{P}(K)$, $\varepsilon > 0$ and for each $j = 1, \dots, n$, let $x_j \in K_j$. We set

$$\mu = (1 - \varepsilon)\nu + \frac{\varepsilon}{n} \sum_{j=1}^n \delta_{x_j}.$$

Then $L(\mu, \nu) \leq 2\varepsilon$ and $\mu(K_j) > 0$ for any $j = 1, \dots, n$. □

2.2. Graph theory. Surprisingly enough, our constructions of measures with prescribed properties need some results from graph theory. Let us first recall some terminology, which can be found e.g. in [Die10]. Let $G = (V, E)$ be a graph. We recall that two vertices v, w of E are *adjacent* or *neighbour* if vw is an edge of G . A *path* between two vertices v and w is a sequence of vertices $v_0 = v, v_1, \dots, v_N = w$ such that $v_i v_{i+1}$ belongs to the set of edges E for any $i \in \{0, \dots, N-1\}$. We say that the graph is *connected* provided any two vertices of G can be linked by a path in G .

If U is a subset of V , the induced subgraph $G(U)$ is the graph whose set of vertices is U and whose set of edges is the subset of E containing all edges vw with $v, w \in U$.

Our first lemma is a well known result in graph theory. For convenience, we provide a proof.

Lemma 2.9. *Let $G = (V, E)$ be a connected graph. There exists a vertex $v \in V$ such that $G(V \setminus \{v\})$ remains connected.*

Proof. Let $[v_0, \dots, v_N]$ be a path such that its vertices are pairwise different, and whose length is maximal in the set of such paths. Then $G \setminus \{v_0\}$ is connected. Indeed, take any $v, w \in V \setminus \{v_0\}$ and let $[w_0, \dots, w_p]$ be the shortest path from $v = w_0$ to $w = w_p$ in G . In particular, $w_i = v_0$ for at most one $i \in \{1, \dots, p-1\}$.

If for any $i \in \{1, \dots, p-1\}$, w_i is not equal to v_0 , there is nothing to prove : the path stays already in $G(V \setminus \{v_0\})$. Thus, suppose that $w_i = v_0$ for some i . If w_{i-1} or w_{i+1} did not belong to $\{v_1, \dots, v_N\}$, then we could add the edge $w_{i-1}v_0$ or $w_{i+1}v_0$ to the path $[v_0, \dots, v_N]$, with a new vertex. This would contradict the maximality of this path. Thus, w_{i-1} and w_{i+1} both belong to $\{v_1, \dots, v_N\}$, and we can replace in $[w_0, \dots, w_p]$ the subpath $[w_{i-1}, w_i, w_{i+1}]$ by the corresponding subpath joining w_{i-1} to w_{i+1} using only vertices in $\{v_1, \dots, v_N\}$, and thus avoiding v_0 . □

This lemma allows us to give weights on the vertices of a connected graph with particular properties.

Lemma 2.10. *Let $G = (V, E)$ be a connected graph, let $\varepsilon > 0$ and let $\rho \geq 1$. There exist $v_0 \in V$ and positive real numbers $(\omega_v)_{v \in V}$ such that*

- for any $v \in V \setminus \{v_0\}$, one can find a neighbour u of v such that $\omega_v \leq \varepsilon \omega_u^\rho$;
- $\omega_{v_0} > 1/2$;
- $\sum_{v \in V} \omega_v = 1$.

Proof. We proceed by induction on the number of vertices of G , the case where this number equals 1 or 2 being trivial. So, suppose that V contains $n \geq 3$ elements and let v_1 be given by Lemma 2.9. The graph $G(V \setminus \{v_1\})$ remaining connected, we can apply the induction hypothesis with some $\varepsilon_0 \in (0, \varepsilon)$ to the graph $G(V \setminus \{v_1\})$. We get positive numbers $(\theta_v)_{v \in V \setminus \{v_1\}}$ and $v_0 \in V \setminus \{v_1\}$. Let $\alpha > 0$ be very small and let

$$\begin{cases} \omega_v &= (1 - \alpha)\theta_v & \text{if } v \neq v_1 \\ \omega_{v_1} &= \alpha. \end{cases}$$

Provided $\alpha > 0$ is small enough, ω_{v_1} is much smaller than ω_v , for v any neighbour of v_1 . Moreover, when $v \in V \setminus \{v_0, v_1\}$, one can find another vertex u which is a neighbour of v in $G(V \setminus \{v_1\})$, in particular in G , such that

$$\theta_v \leq \varepsilon_0 \theta_u^\rho.$$

This leads to

$$\omega_v \leq \frac{\varepsilon_0}{(1 - \alpha)^{\rho-1}} \omega_u^\rho \leq \varepsilon \omega_u^\rho,$$

provided $\alpha > 0$ is small enough again. The property $\omega_{v_0} > 1/2$ is also true, under the same restriction on α . \square

2.3. Measures with prescribed properties. Our results will depend on the construction of probability measures on K having prescribed properties depending on the dimension of K . Of course, the choice of the definition of the dimension will determine the properties of the measure on K that we can expect. In this direction, the most famous result is the Frostman lemma (see [Fal03]) which is based on the Hausdorff dimension.

Lemma 2.11. *Let K be a nonempty compact subset of \mathbb{R}^d . Then, for every $0 \leq t < \dim_{\mathcal{H}}(K)$, there exists $\mu \in \mathcal{P}(K)$ and constants $C > 0, r_0 > 0$ such that $\mu(B(x, r)) \leq Cr^t$ for all $x \in K$ and $0 < r \leq r_0$.*

This lemma has a counterpart for the upper-box dimension, due to Tricot ([Tri82]).

Lemma 2.12. *Let K be a nonempty compact subset of \mathbb{R}^d . Then, for every $t > \overline{\dim}_B(K)$, there exists $\mu \in \mathcal{P}(K)$ and constants $C > 0, r_0 > 0$ such that $\mu(B(x, r)) \geq Cr^t$ for all $x \in K$ and $0 < r \leq r_0$.*

One can also ask to weaken the assumption in the anti-Frostman lemma, by the use of the lower-box dimension instead of the upper-box dimension. This is possible, even if there is a price to pay: the conclusion is also weaker, since it is only obtained for one value of r , arbitrarily small.

Lemma 2.13. *Let K be a nonempty compact subset of \mathbb{R}^d . Then, for every $t > \underline{\dim}_B(K)$, for every $\alpha > 0$, there exists $\mu \in \mathcal{P}(K)$ and $r \in (0, \alpha)$ such that $\mu(B(x, r)) \geq 2^{-t} r^t$ for all $x \in K$.*

Proof. One can find $r \in (0, \alpha)$ such that $\mathbf{P}_{r/2}(K) \leq 2^t r^{-t}$. We set $P = \mathbf{P}_{r/2}(K)$ and let $x_1, \dots, x_P \in K$ be centers of disjoint balls of radius $r/2$. We define

$$\mu = \frac{1}{P} \sum_{j=1}^P \delta_{x_j}.$$

Then, for any $x \in K$, one can find $i \in \{1, \dots, P\}$ such that $x_i \in B(x, r)$ (otherwise $B(x, r/2)$ would not intersect any $B(x_j, r/2)$). Thus

$$\mu(B(x, r)) \geq \frac{1}{P} \geq 2^{-t} r^t.$$

□

We can, like in Frostman lemma, obtain measures with an upper estimate of the size of balls using the upper-box dimension only. However, here too, we have to work with a fixed radius.

Lemma 2.14. *Let K be a nonempty compact subset of \mathbb{R}^d and let $0 \leq t < \overline{\dim}_B(K)$. Then, for any $\alpha > 0$, one can find $r \in (0, \alpha)$ and $\mu \in \mathcal{P}(K)$ such that, for any $x \in K$, $\mu(B(x, r)) \leq r^t$.*

Proof. The proof is similar to that of Lemma 2.13. By definition of the upper-box dimension, one can find $r \in (0, \alpha)$ such that

$$r^{-t} \leq \mathbf{P}_r(K).$$

Let $P = \mathbf{P}_r(K)$ and let $x_1, \dots, x_P \in K$ be the centers of disjoint balls with radius r . Then the measure

$$\mu = \frac{1}{P} \sum_{j=1}^P \delta_{x_j}$$

is convenient. Indeed, for any $x \in K$, the ball $B(x, r)$ contains at most one of the x_j , so that $\mu(B(x, r)) \leq P^{-1} \leq r^t$. □

We finally use the box separation index. This is much more delicate, and we work directly with $I_\mu(r, q)$ instead of $\mu(B(x, r))$.

Lemma 2.15. *Let K be a nonempty compact subset of \mathbb{R}^d , let $n \geq 1$ and let $q \leq 0$. There exists $\nu_n \in \mathcal{P}(K)$ with finite support such that, for any $\mu = \sum_{i=1}^N p_i \delta_{x_i} \in \mathcal{P}(K)$ with finite support, for any $\theta \in (0, 1)$, setting $\nu = (1 - \theta)\mu + \theta\nu_n$, then*

$$I_\nu(3 \cdot 2^{-n}, q) \leq N(1 - \theta)^q a^q + \theta^q + \frac{\theta^q}{2^q} C_n(K)^{1-q},$$

where $a = \min_i(p_i)$. Moreover, $\nu(B(x, 3 \cdot 2^{-n})) > 0$ for any $x \in K$.

Proof. We use the notations of the introduction. We decompose K_n into its connected components $\mathcal{C}_{n,1}(K), \dots, \mathcal{C}_{n,\kappa_n}(K)$. For any $j = 1, \dots, \kappa_n(K)$, we denote by C_j^i , $i = 1, \dots, \kappa_j$, the half-closed dyadic cubes of size 2^{-n} appearing in \mathcal{C}_j . By definition, $C_j^i \cap K$ is never empty, and we can fix y_j^i a point in $C_j^i \cap K$.

We endow each $\mathcal{C}_{n,j}$ with a graph structure. Precisely, we say that two cubes C_j^i and C_j^l are adjacent provided their closure do intersect. The graph $\mathcal{C}_{n,j}$ is then a connected graph. Moreover, it is easy to check that if C_j^i and C_j^l are adjacent, then for any $x \in C_j^i$, C_j^l is contained in $B(x, 3 \cdot 2^{-n})$ (at this part of the proof, we find convenient to use the ℓ^∞ -norm on \mathbb{R}^d , and to suppose that $B(x, r)$ is the ball corresponding to this norm. Working with a different norm would only replace here 3 by a constant depending on d . This would not change at all the content of our main theorem.)

Let $\varepsilon > 0$ be such that $\varepsilon(\kappa_1 + \dots + \kappa_{\kappa_n(K)}) \leq C_n(K)^q$. We apply Lemma 2.10 to each connected graph $\mathcal{C}_{n,j}$ with $\rho = 1 - q$. This gives weights ω_j^i , $i = 1, \dots, \kappa_j$. Renumbering the cubes if necessary, we can always assume that, for any i in $1, \dots, \kappa_j - 1$, there exists $l \neq i$ such that the cubes C_j^i and C_j^l are adjacent, and $\omega_j^i \leq \varepsilon(\omega_j^l)^{1-q}$ (that is we suppose that the vertex v_0 in Lemma 2.10 is $C_j^{\kappa_j}$.)

We finally set

$$\begin{aligned}\lambda_j &= \sum_{i=1}^{\kappa_j} \omega_j^i \delta_{y_j^i} \\ \nu_n &= \sum_{j=1}^{C_n(K)} \frac{1}{C_n(K)} \lambda_j.\end{aligned}$$

Let now μ and ν as in the assumptions of the lemma. We first observe that for any $x \in K$, $\nu(B(x, 3.2^{-n})) > 0$. Indeed, any $x \in K$ belongs to some C_j^i and $\nu(B(x, 3.2^{-n})) \geq \nu(\{y_j^i\}) > 0$. Let us now control $I_\nu(3.2^{-n}, q)$:

$$I_\nu(3.2^{-n}, q) = \sum_{j=1}^{C_n(K)} \sum_{i=1}^{\kappa_j} \int_{C_j^i \cap K} \frac{d\nu(x)}{\nu(B(x, 3.2^{-n}))}.$$

To estimate the integral, we distinguish three different cases:

- (a) There exists (at least) one point in the support of μ which lies in $C_j^i \cap K$. Then we just use that, for any $x \in C_j^i \cap K$, $\nu(B(x, 3.2^{-n})) \geq \nu(C_j^i \cap K) \geq (1 - \theta)a$, so that

$$\int_{C_j^i \cap K} \frac{d\nu(x)}{\nu(B(x, 3.2^{-n}))^{1-q}} \leq \nu(C_j^i \cap K)^q \leq (1 - \theta)^q a^q.$$

Observe that this estimate concerns at most N cubes.

- (b) $\text{supp}(\mu) \cap C_j^i$ is empty, and $i \leq \kappa_j - 1$. Let C_j^l be a neighbour of C_j^i such that $\omega_j^i \leq \varepsilon(\omega_j^l)^{1-q}$. For any $x \in C_j^i \cap K$, we already observed that $B(x, 3.2^{-n})$ contains $C_j^l \cap K$, so that $\nu(B(x, 3.2^{-n})) \geq \nu(C_j^l \cap K) \geq \frac{\theta}{C_n(K)} \omega_j^l$. Thus,

$$\int_{C_j^i \cap K} \frac{d\nu(x)}{\nu(B(x, 3.2^{-n}))^{1-q}} \leq \frac{\nu(C_j^i \cap K)}{\theta^{1-q}(\omega_j^l)^{1-q} C_n(K)^{q-1}} \leq \frac{\theta^q \omega_j^i}{(\omega_j^l)^{1-q} C_n(K)^q} \leq \frac{\theta^q \varepsilon}{C_n(K)^q}.$$

- (c) For the last cubes, namely $\text{supp}(\mu) \cap C_j^i$ is empty and $i = \kappa_j$, we argue as in case (a), except that now the mass of the cube $C_j^i \cap K$ is given by ν_n :

$$\begin{aligned}\int_{C_j^i \cap K} \frac{d\nu(x)}{\nu(B(x, 3.2^{-n}))^{1-q}} &\leq \nu(C_j^i \cap K)^q \\ &\leq \frac{\theta^q}{C_n(K)^q} (\omega_j^{\kappa_j})^q \\ &\leq \left(\frac{\theta}{2}\right)^q C_n(K)^{-q}.\end{aligned}$$

Putting this together, and remembering the value of ε , we get that

$$\begin{aligned}I_\nu(3.2^{-n}, q) &\leq N(1 - \theta)^q a^q + \sum_{j=1}^{C_n(K)} \sum_{i=1}^{\kappa_j - 1} \frac{\theta^q \varepsilon}{C_n(K)^q} + \sum_{j=1}^{C_n(K)} \left(\frac{\theta}{2}\right)^q C_n(K)^{-q} \\ (1) \quad &\leq N(1 - \theta)^q a^q + \theta^q + \frac{\theta^q}{2^q} C_n(K)^{1-q}.\end{aligned}$$

□

2.4. Results true for all measures. The results true for all measures which are quoted in Theorem 1.6 are not new. All of them can be found in [Ols05] or in [Ols08], except $\overline{D}_\mu(q) \leq s_{\mathcal{P}}$ for $q \geq 1$. But this inequality can be found in [Cut95] for $q = 1$, and the remaining cases follow easily from the next lemma, which is an easy application of Jensen inequality.

Lemma 2.16. *Let $q_1 \leq q_2$ be two real numbers and let $\mu \in \mathcal{P}(K)$. Then*

$$\underline{D}_\mu(q_1) \geq \underline{D}_\mu(q_2) \text{ and } \overline{D}_\mu(q_1) \geq \overline{D}_\mu(q_2).$$

3. TYPICAL LOWER L^q -DIMENSIONS, $q \leq 0$

Let K be an infinite compact subset of \mathbb{R}^d . In this section, we prove that a typical measure $\mu \in \mathcal{P}(K)$ satisfies $\underline{D}_\mu(q) = s_{\text{sep}}$ for any $q \leq 0$. We first observe that we can work with a fixed value of q . Indeed, suppose that we are able to prove that, for any $q \leq 0$, the set

$$\mathcal{M}(q) = \{\mu \in \mathcal{P}(K); \underline{D}_\mu(q) = s_{\text{sep}}\}$$

contains a dense G_δ -set. Then consider (q_n) a dense sequence in $(-\infty, 0]$, with $q_0 = 0$ and let

$$\mathcal{M} = \bigcap_n \mathcal{M}(q_n)$$

which itself contains a dense G_δ -set. Let $\mu \in \mathcal{M}$ and let $q \in (-\infty, 0]$. One can find two subsequences $(q_{\phi(n)})$ and $(q_{\psi(n)})$ such that $q_{\phi(n)} \leq q \leq q_{\psi(n)}$. Since $\underline{D}_\mu(q_{\psi(n)}) \leq \underline{D}_\mu(q) \leq \underline{D}_\mu(q_{\phi(n)})$, taking the limit, we get $\underline{D}_\mu(q) = s_{\text{sep}}$.

This trick works in any case. In particular, in this section, from now on, we fix $q \leq 0$.

3.1. The lower estimate. We first prove that a typical measure $\mu \in \mathcal{P}(K)$ satisfies $\underline{D}_\mu(q) \geq s_{\text{sep}}$. Let $n \geq 1$, and recall that $\mathcal{C}_{n,1}, \dots, \mathcal{C}_{n,C_n(K)}$ denote the connected components of K_n , the union of the dyadic cubes of size 2^{-n} intersecting K . We also define

$$\mathcal{A}_{n,j} = \bigcup_{C \in \mathcal{C}_{n,j}} C \cap K.$$

It is important to notice that $\text{dist}(\mathcal{A}_{n,j}, \mathcal{A}_{n,k}) \geq 2^{-n}$ provided $j \neq k$, and that K is the (disjoint) union of the $\mathcal{A}_{n,j}$. In particular, $\mathcal{A}_{n,j}(2^{-n-1}) \cap K = \mathcal{A}_{n,j}$. We finally set

$$\begin{aligned} \mathcal{R}_n &= \{\mu \in \mathcal{P}(K); \forall j \in \{1, \dots, C_n(K)\}, \mu(\mathcal{A}_{n,j}) > 0\} \\ \mathcal{R} &= \bigcap_{n \geq 1} \mathcal{R}_n, \end{aligned}$$

and we claim that \mathcal{R} is the dense G_δ -set we are looking for. Indeed, each \mathcal{R}_n is dense by Lemma 2.8 and open by Lemma 2.6. Moreover, any μ in \mathcal{R} has its lower L^q -dimension greater than or equal to s_{sep} . Indeed, let $r > 0$ be small, and let $n \geq 1$ be such that $2^{-(n+1)} < r \leq 2^{-n}$. We know that

$$\begin{aligned} I_\mu(r, q) &= \sum_{j=1}^{C_n(K)} \int_{\mathcal{A}_{n,j}} \frac{d\mu(x)}{\mu(B(x, r))^{1-q}} \\ &\geq \sum_{j=1}^{C_n(K)} \int_{\mathcal{A}_{n,j}} \frac{d\mu(x)}{\mu(\mathcal{A}_{n,j})^{1-q}}, \end{aligned}$$

because the $\mathcal{A}_{n,j}$ are well separated, so that $\mathcal{A}_{n,j} \supset B(x, r) \cap K$ for any $x \in \mathcal{A}_{n,j}$ and any $r \leq 2^{-n}$. Thus we find

$$I_\mu(r, q) \geq \sum_{j=1}^{C_n(K)} \mu(\mathcal{A}_{n,j})^q.$$

Now, since $\sum_{j=1}^{C_n(K)} \mu(\mathcal{A}_{n,j}) = 1$, it is well known that this quantity is minimal when $\mu(\mathcal{A}_{n,j}) = \frac{1}{C_n(K)}$ for any $j = 1, \dots, C_n(K)$. This shows that

$$I_\mu(r, q) \geq C_n(K)^{1-q}.$$

Taking the logarithm, dividing by $(q-1)\log r$, which is positive, and taking the liminf, we get the aforementioned lower bound for $\underline{D}_\mu(q)$.

3.2. The upper estimate. We now prove that a typical measure $\mu \in \mathcal{P}(K)$ satisfies $\underline{D}_\mu(q) \leq s_{\text{sep}}$. It is sufficient to prove that, for any $t > s_{\text{sep}}$, a typical measure $\mu \in \mathcal{P}(K)$ satisfies $\underline{D}_\mu(q) \leq t$.

Let $\mu \in \mathcal{F}(K)$, $\mu = \sum_{i=1}^N p_i \delta_{x_i}$, $a = \min_i(p_i) > 0$ and let $l \geq 1$. By definition of the box separation index, one can find $n \geq l$ very large such that

$$N(1 - N^{-1})^q a^q + N^{-q} + 2^{-q} N^{-q} C_n(K)^{1-q} \leq 2^{nt(1-q)}.$$

Let ν_n be the measure given by Lemma 2.15 and let

$$\nu_{\mu,l} = \left(1 - \frac{1}{N}\right) \mu + \frac{1}{N} \nu_n.$$

We can apply Corollary 2.3 to the measure $\nu_{\mu,l}$ and to $r = 3 \cdot 2^{-n}$ to get some positive $\delta_{\mu,l}$ such that, for any ν with $L(\nu, \nu_{\mu,l}) < \delta_{\mu,l}$,

$$\begin{aligned} I_\nu(6 \cdot 2^{-n}, q) &\leq C_q I_{\nu_{\mu,l}}(3 \cdot 2^{-n}, q) \\ &\leq C_q (N(1 - N^{-1})^q a^q + N^{-q} + 2^{-q} N^{-q} C_n(K)^{1-q}) \\ &\leq C_q 2^{nt(1-q)}. \end{aligned}$$

We finally set

$$\mathcal{R} = \bigcap_{l \geq 1} \bigcup_{\mu \in \mathcal{F}(K)} B_L(\nu_{\mu,l}, \delta_{\mu,l}).$$

For any fixed $l \geq 1$, $\{\nu_{\mu,l}; \mu \in \mathcal{F}(K)\}$ is dense in $\mathcal{P}(K)$. Indeed, let $\lambda \in \mathcal{P}(K)$ and let $\varepsilon > 0$. Since K is infinite, there exists $\mu \in \mathcal{F}(K)$ such that $L(\mu, \lambda) < \varepsilon/2$ and the cardinal number of the support of μ , denoted by N , satisfies $N > 4/\varepsilon$. Now,

$$L(\nu_{\mu,l}, \lambda) < \frac{\varepsilon}{2} + \frac{2}{N} < \varepsilon.$$

Thus, \mathcal{R} is a dense G_δ -set. Moreover, for any $\nu \in \mathcal{R}$, there exist arbitrarily large integers n such that

$$I_\nu(6 \cdot 2^{-n}, q) \leq C_q 2^{nt(1-q)}.$$

Taking the logarithm and dividing by $(q-1)\log(6 \cdot 2^{-n})$, this shows that $\underline{D}_\nu(q) \leq t$.

3.3. The box separation index of self-similar compact sets. We end up this section by computing the box separation index of a self-similar compact set. Fix an integer $N \geq 2$ and let $S_i : \mathbb{R}^d \rightarrow \mathbb{R}^d$, $i = 1, \dots, N$ be contracting similarities with respective ratio r_i . Let K be the self-similar compact set associated to these similarities, namely K is the unique nonempty compact subset of \mathbb{R}^d satisfying

$$K = \bigcup_i S_i(K).$$

We say that (S_1, \dots, S_N) satisfies the *open set condition* if there exists an open nonempty and bounded subset \mathcal{U} of \mathbb{R}^d with $S_i\mathcal{U} \subset \mathcal{U}$ and $S_i\mathcal{U} \cap S_j\mathcal{U} = \emptyset$ for all i, j with $i \neq j$. We also say that (S_1, \dots, S_N) satisfies the *strong separation condition* if $S_i K \cap S_j K = \emptyset$ for all $i \neq j$.

When (S_1, \dots, S_N) satisfies the open set condition, the Hausdorff dimension and the box dimension of K are well known. Define β as the unique solution of $\sum_{i=1}^N r_i^\beta = 1$. Then $\dim_{\mathcal{H}}(K) = \underline{\dim}_B(K) = \overline{\dim}_B(K) = \beta$ (all details can be found in [Fal03]). Under a slightly stronger assumption, this is also the value of the box separation index.

Theorem 3.1. *Let (S_1, \dots, S_N) be contracting similarities of \mathbb{R}^d with ratios (r_1, \dots, r_N) and let K be the associated self-similar compact set. If (S_1, \dots, S_N) satisfies the strong separation condition, then*

$$\text{bsi}(K) = \dim_{\mathcal{H}}(K) = \underline{\dim}_B(K) = \overline{\dim}_B(K) = \beta$$

where β is the unique solution of $\sum_{i=1}^N r_i^\beta = 1$.

Proof. Since $\text{bsi}(K) \leq \underline{\dim}_B(K)$, we just need to prove that $\liminf_n \frac{\log C_n(K)}{n \log 2} \geq \beta$. For a word $\mathbf{i} = i_1 \dots i_p$ with entries in $\{1, \dots, N\}$, define

$$S_{\mathbf{i}} = S_{i_1} \circ \dots \circ S_{i_p}, \quad r_{\mathbf{i}} = r_{i_1} \dots r_{i_p}, \quad \rho_{\mathbf{i}} = r_{\mathbf{i}}^\beta.$$

Since the strong separation condition is satisfied, there exists $m \geq 0$ such that

$$\text{dist}(S_{\mathbf{i}}(K), S_{\mathbf{j}}(K)) \geq 2^{-m} \text{ for any } \mathbf{i} \neq \mathbf{j}.$$

For $r > 0$, denote by $I(r)$ the set of words \mathbf{i} such that

$$r_{i_1} \dots r_{i_p} \leq r \text{ and } r_{i_1} \dots r_{i_{p-1}} > r.$$

Such a word is called *minimal* with respect to r . The key points are the following facts which can be found e.g. in the proof of [Fal03, Theorem 9.3]:

- For any $r > 0$, $\sum_{\mathbf{i} \in I(r)} \rho_{\mathbf{i}} = 1$;
- For any $r > 0$, $K \subset \bigcup_{\mathbf{i} \in I(r)} S_{\mathbf{i}}(K)$.

Now, by definition of $I(r)$, $\rho_{\mathbf{i}} \leq r^\beta$ for any $\mathbf{i} \in I(r)$, so that

$$\text{card}(I(r)) \geq r^{-\beta}.$$

Pick now $\mathbf{i} \neq \mathbf{j}$ in $I(r)$. By minimality of \mathbf{i} and \mathbf{j} , \mathbf{i} is not the beginning of \mathbf{j} , and conversely. Let k be the smallest index with $i_k \neq j_k$ and let \mathbf{u} be the word $\mathbf{u} = i_1 \dots i_{k-1}$. Then

$$\begin{aligned} \text{dist}(S_{\mathbf{i}}(K), S_{\mathbf{j}}(K)) &\geq \text{dist}(S_{\mathbf{u}}S_{i_k}(K), S_{\mathbf{u}}S_{j_k}(K)) \\ &\geq r_{i_1} \dots r_{i_{k-1}} \text{dist}(S_{i_k}(K), S_{j_k}(K)) \\ &\geq 2^{-m} r. \end{aligned}$$

We specialize this inequality for $r = 2^{-n}$. In each $S_{\mathbf{i}}(K)$, $\mathbf{i} \in I(2^{-n})$, one can find a point belonging to K . Furthermore, the distance between $S_{\mathbf{i}}(K)$ and $S_{\mathbf{j}}(K)$ is greater than

$2^{-(n+m)}$ when $\mathbf{i} \neq \mathbf{j} \in I(2^{-n})$. Thus, the closure of the dyadic cubes of size $2^{-(n+m+2)}$ intersecting $S_{\mathbf{i}}(K)$ and the closure of the dyadic cubes of size $2^{-(n+m+2)}$ intersecting $S_{\mathbf{j}}(K)$, $\mathbf{i} \neq \mathbf{j} \in I(2^{-n})$, are disjoint. Hence,

$$C_{n+m+2}(K) \geq \text{card}(I(2^{-n})) \geq 2^{-n\beta}.$$

Taking the logarithm and the liminf, this yields $\text{bsi}(K) \geq \beta$. \square

Remark 3.2. The example of $K = [0, 1]$ shows that we cannot only assume the open set condition.

4. LOCAL UNIFORM UPPER BOX DIMENSION AND TYPICAL UPPER L^q -DIMENSION

In this section, we study the link between the local uniform upper box dimension of the compact set K and the typical upper L^q -dimension, $q > 1$, of the probability measures on K . We also show that the local uniform upper box dimension does not coincide with the local upper box or the local lower box dimensions.

From now on, we fix $q > 1$ and set $s_u = \overline{\dim}_{B, \text{loc}, \text{unif}}(K)$.

4.1. The lower estimate. Let $t < s_u$. One intends to show that $\overline{D}_\mu(q) \geq t$ for a typical measure $\mu \in \mathcal{P}(K)$. Let $(x_i)_{i \geq 1}$ be a dense sequence in K and let $n \geq 1$. By definition of s_u , one may find $r_n < 1/n$ such that, for any $i = 1, \dots, n$,

$$\mathbf{P}_{r_n}(K \cap B(x_i, 1/n)) \geq r_n^{-t}.$$

Arguing like in the proof of Lemma 2.14, we can construct for each $n \geq 1$ and each $i \leq n$ a measure $\mu_{n,i} \in \mathcal{P}(K)$ such that $\text{supp}(\mu_{n,i}) \subset B(x_i, 1/n)$ and, for any $x \in K$, $\mu_{n,i}(B(x, r_n)) \leq r_n^t$.

Let also $\delta_n > 0$ be given by Lemma 2.1 for $\alpha = r_n/2$ and $\beta = r_n^t$. We finally set

$$\begin{aligned} \Lambda_n &= \left\{ \sum_{i=1}^n p_i \mu_{n,i}; p_i \geq 0, \sum_{i=1}^n p_i = 1 \right\} \\ \mathcal{R}_n &= \bigcup_{\mu \in \Lambda_n} B_L(\mu, \delta_n) \\ \mathcal{R} &= \bigcap_{m \geq 1} \bigcup_{n \geq m} \mathcal{R}_n. \end{aligned}$$

By Lemma 2.7, for any $m \geq 1$, $\bigcup_{n \geq m} \mathcal{R}_n$ is dense, so that \mathcal{R} is a dense G_δ -subset of $\mathcal{P}(K)$. Pick now $\nu \in \mathcal{R}$. There exists an arbitrarily large integer n and $\mu \in \Lambda_n$ such that $L(\mu, \nu) < \delta_n$. Now, μ satisfies, for any $x \in K$,

$$\mu(B(x, r_n)) = \sum_{i=1}^n p_i \mu_{n,i}(B(x, r_n)) \leq r_n^t.$$

Taking into account the value of δ_n , this yields

$$\nu(B(x, r_n/2)) \leq 2r_n^t.$$

Integrating this inequality, we get

$$I_\nu(r_n/2, q) \leq 2^{q-1} r_n^{t(q-1)}.$$

Since r_n can be taken arbitrarily small, this implies $\overline{D}_\nu(q) \geq t$.

4.2. The upper estimate. Let $t > s_u$. Our aim is to show that, generically, $\overline{D}_\mu(q) \leq t$. By definition of the local uniform upper box dimension, one can find $x_1, \dots, x_n \in K$, all different, $r_0 > 0$, with $4r_0 < \min_{i \neq j} \|x_i - x_j\|$ and $\alpha > 0$, such that, for any $r \in (0, \alpha)$, one may find $i \in \{1, \dots, n\}$ with

$$\frac{\log \mathbf{N}_r(K \cap B(x_i, r_0))}{-\log r} \leq t.$$

The crucial point here is a result which appears in the proof of [Ols05, Lemma 2.3.1.]. It is based on Jensen's inequality.

Lemma 4.1. *Let $\mu \in \mathcal{P}(K)$, let E be a Borel subset of K with $\mu(E) > 0$. Then, for any $r > 0$,*

$$I_\mu(2r, q) \geq \frac{\mu(E)^q}{\mathbf{N}_r(E)^{q-1}}.$$

We then set

$$\mathcal{R} = \{\mu \in \mathcal{P}(K); \forall i \in \{1, \dots, n\}, \mu(B(x_i, r_0)) > 0\}.$$

The set \mathcal{R} is dense (Lemma 2.8) and open (Lemma 2.6). Moreover, for any $\mu \in \mathcal{R}$ and any $r > 0$,

$$I_\mu(2r, q) \geq \sup_{i=1, \dots, n} \frac{\mu(B(x_i, r_0))^q}{\mathbf{N}_r(K \cap B(x_i, r_0))^{q-1}} \geq \frac{C}{\inf_{i=1, \dots, n} (\mathbf{N}_r(K \cap B(x_i, r_0))^{q-1})},$$

where $C = \inf_{i=1, \dots, n} \mu(B(x_i, r_0))^q$. Taking the logarithm, we find

$$\log I_\mu(2r, q) \geq \log C - (q-1) \inf_{i=1, \dots, n} \log \mathbf{N}_r(K \cap B(x_i, r_0)).$$

Now, for any r in $(0, \alpha)$, this becomes

$$\log I_\mu(2r, q) \geq \log C + t(q-1) \log r.$$

We now divide this inequality by $(q-1) \log r$ (which is negative) and we take the limsup to obtain that $\overline{D}_\mu(q) \leq t$.

4.3. Comparison of three notions of dimensions. We now show that the local uniform upper box dimension is not always equal to the (local) lower box dimension or to the (local) upper box dimension. For simplicity, we set

$$\begin{aligned} s^+ &= \inf_{x \in K} \overline{\dim}_{B, \text{loc}}(x, K) \\ s_- &= \inf_{x \in K} \underline{\dim}_{B, \text{loc}}(x, K). \end{aligned}$$

That $s_u > s_-$ for some compact sets is easy: it suffices to take a Cantor set K for which $\underline{\dim}_B(K) < \overline{\dim}_B(K)$. The uniformity in the construction of the Cantor set ensures that

$$s_- = \underline{\dim}_B(K) \text{ and } s_u = s^+ = \overline{\dim}_B(K)$$

(see [Tri82] for details).

To prove that it is possible that $s_u < s^+$, we also use Cantor sets. Let (n_k) be a sequence of integers increasing to $+\infty$ with $n_0 = 1$ and $n_{k+1} > 5n_k$. We define two sequences (α_n) and (β_n) by $\alpha_0 = \beta_0 = 1$ and

$$\alpha_{n+1} = \begin{cases} 27 & \text{provided } n \in [n_k, 2n_k) \\ 3 & \text{provided } n \in [2n_k, 3n_k) \\ 9 & \text{otherwise} \end{cases}$$

$$\beta_{n+1} = \begin{cases} 27 & \text{provided } n \in [3n_k, 4n_k) \\ 3 & \text{provided } n \in [4n_k, 5n_k) \\ 9 & \text{otherwise.} \end{cases}$$

Define E (resp. F) a Cantor subset of $[0, 1]$ (resp. of $[2, 3]$) as follows : $E = \bigcap_n E_n$ (resp. $F = \bigcap_n F_n$) where at each step we divide each subinterval of E_n (resp. F_n) into α_{n+1} (resp. β_{n+1}) intervals of size $100^{-(n+1)}$. Thus E_n (resp. F_n) consists of $\alpha_1 \times \dots \times \alpha_n$ (resp. $\beta_1 \times \dots \times \beta_n$) intervals of size 100^{-n} . We finally set $K = E \cup F$. A key point in the construction is that, for each n ,

$$\min(\alpha_1 \dots \alpha_n, \beta_1 \dots \beta_n) = 9^n.$$

Thus we need at most 9^n intervals of size 100^{-n} to cover either E or F . On the contrary, when we look separately at E or at F , we need for certain values of n more intervals.

Let us proceed with the details. By uniformity of the construction of each Cantor set, it is not hard to show that

$$\inf_{x \in K} \overline{\dim}_{B, \text{loc}}(x, K) = \inf(\overline{\dim}_B(E), \overline{\dim}_B(F)).$$

Now,

$$\overline{\dim}_B(E) = \limsup_{n \rightarrow +\infty} \frac{\log \alpha_1 \dots \alpha_n}{n \log 100} = \lim_{k \rightarrow +\infty} \frac{\log 9^{n_k} 27^{n_k}}{2n_k \log 100} = \frac{5 \log 3}{2 \log 100},$$

and, in the same way,

$$\overline{\dim}_B(F) = \limsup_{n \rightarrow +\infty} \frac{\log \beta_1 \dots \beta_n}{n \log 100} = \lim_{k \rightarrow +\infty} \frac{\log 9^{3n_k} 27^{n_k}}{4n_k \log 100} = \frac{9 \log 3}{4 \log 100}.$$

On the contrary, we shall see that for this compact set K , $s_u \leq \frac{2 \log 3}{\log 100}$ (actually, s_u is equal to $\frac{2 \log 3}{\log 100}$). Indeed, it is enough to show that

$$\limsup_{r \rightarrow 0} \left(\frac{\inf(\log \mathbf{N}_r(E), \log \mathbf{N}_r(F))}{-\log r} \right) \leq \frac{2 \log 3}{\log 100}.$$

This follows easily from

$$\limsup_{n \rightarrow +\infty} \frac{\log 9^n}{n \log 100} = \frac{2 \log 3}{\log 100}.$$

Remark 4.2. The local uniform upper box dimension can also be easily computed for a regular compact set. We recall that a compact set $K \subset \mathbb{R}^d$ with $\dim_{\mathcal{H}}(K) = s$ is called *Ahlfors regular* if there are positive constants $c_1, c_2, r_0 > 0$ such that

$$c_1(2r)^s \leq \mathcal{H}^s(K \cap B(x, r)) \leq c_2(2r)^s$$

for all $x \in K$ and all $0 < r < r_0$. For a Ahlfors-regular compact set, it is easy (and classical) to check that

$$\inf_{x \in K} \underline{\dim}_{B, \text{loc}}(x, K) = \overline{\dim}_{B, \text{loc}, \text{unif}}(K) = \dim_{\mathcal{H}}(K) = s.$$

This happens in particular if K is a self-similar compact set satisfying the open set condition.

5. TYPICAL L^q -DIMENSIONS, THE REMAINING CASES, $q \neq 1$

In this section, we turn to the remaining cases, for $q \neq 1$. More precisely, we prove that a typical measure $\mu \in \mathcal{P}(K)$ satisfies

- $\overline{D}_\mu(q) = \overline{\dim}_B(K)$ provided $q \in [0, 1)$;
- $\overline{D}_\mu(q) = +\infty$ provided $q < 0$;
- $\underline{D}_\mu(q) = 0$ provided $q \in (0, 1)$.

As previously, we can work with a fixed $q \in \mathbb{R}$.

5.1. Typical upper L^q -dimension, $q \in [0, 1)$. In [Ols08], it is proved that any measure $\mu \in \mathcal{P}(K)$ satisfies $\overline{D}_\mu(q) \leq s$ where $s = \overline{\dim}_B(K)$. So, since the result is trivial if $\overline{\dim}_B(K) = 0$, we may assume that s is positive, and we have just to prove that, given any $t \in (0, s)$, a typical $\mu \in \mathcal{P}(K)$ satisfies $\overline{D}_\mu(q) \geq t$.

Let (μ_n) be a sequence of $\mathcal{F}(K)$ which is dense in $\mathcal{P}(K)$. Write $\mu_n = \sum_{i=1}^N p_i \delta_{x_i}$, where the x_i are all different and each p_i is nonzero. Let $\alpha_n > 0$ be such that

$$\alpha_n \leq \frac{1}{n}, \quad N\alpha_n^t \leq \frac{1}{2} \quad \text{and} \quad 4\alpha_n < \min_{i \neq j} \|x_i - x_j\|.$$

For these values of t and α_n , Lemma 2.14 gives us a real number $r_n \in (0, \alpha_n)$ and a measure $m_n \in \mathcal{P}(K)$ such that, for any $x \in K$, $m_n(B(x, r_n)) \leq r_n^t$. We then set

$$\varepsilon_n = \frac{1}{-\log r_n}, \quad \nu_n = (1 - \varepsilon_n)\mu_n + \varepsilon_n m_n \quad \text{and} \quad Y_n = \bigcup_{i=1}^N B(x_i, r_n).$$

We first observe that

$$\begin{aligned} \nu_n(Y_n) &\leq 1 - \varepsilon_n + \varepsilon_n \sum_{i=1}^N m_n(B(x_i, r_n)) \\ &\leq 1 - \varepsilon_n + \varepsilon_n N r_n^t \\ &\leq 1 - \frac{\varepsilon_n}{2}. \end{aligned}$$

Moreover, for any $x \notin Y_n$, $\nu_n(B(x, r_n)) \leq \varepsilon_n r_n^t$, so that,

$$\begin{aligned} I_{\nu_n}(r_n, q) &\geq \int_{Y_n^c} \frac{d\nu_n(x)}{\nu_n(B(x, r_n))^{1-q}} \\ &\geq \nu_n(Y_n^c) \varepsilon_n^{q-1} r_n^{t(q-1)} \\ &\geq \frac{1}{2} \varepsilon_n^q r_n^{t(q-1)} \end{aligned}$$

Now, Corollary 2.3 (see also the remark following it) gives us a real number $\delta_n > 0$ such that, for any $\nu \in \mathcal{P}(K)$ satisfying $L(\nu, \nu_n) < \delta_n$, one has

$$I_\nu(r_n/2, q) \geq \frac{1}{4} \varepsilon_n^q r_n^{t(q-1)}.$$

Taking the logarithm and dividing by $(q-1) \log r_n$, which is positive, we get

$$\frac{\log I_\nu(r_n/2, q)}{(q-1) \log(r_n)} \geq C_1 + C_2 \frac{\log(-\log(r_n))}{\log r_n} + t.$$

Finally, we observe that since (ε_n) is going to zero, (ν_n) keeps dense in $\mathcal{P}(K)$. We then consider the dense G_δ -set

$$\mathcal{R} = \bigcap_{m \geq 0} \bigcup_{n \geq m} B_L(\nu_n, \delta_n).$$

The work done before implies immediately that any $\nu \in \mathcal{R}$ satisfies $\overline{D}_\nu(q) \geq t$.

5.2. Typical upper L^q -dimension, $q < 0$. Let (μ_n) be a sequence in $\mathcal{F}(K)$ which is dense in $\mathcal{P}(K)$. For any $n \in \mathbb{N}$, let $m_n = \delta_{y_n}$ where y_n does not belong to $\text{supp}(\mu_n)$. Let (r_n) be a sequence decreasing to zero such that $2r_n < \text{dist}(y_n, \text{supp}(\mu_n))$. We set

$$\varepsilon_n = r_n^n \text{ and } \nu_n = (1 - \varepsilon_n)\mu_n + \varepsilon_n m_n.$$

Observe that (ν_n) remains dense in $\mathcal{P}(K)$. Moreover, the choice of r_n ensures that

$$I_{\nu_n}(r_n, q) \geq \nu_n(\{y_n\})\nu_n(B(y_n, r_n))^{q-1} = \nu_n(\{y_n\})^q = r_n^{-nq}.$$

Corollary 2.3 gives us a real number $\delta_n > 0$ such that any $\nu \in \mathcal{P}(K)$ satisfying $L(\nu, \nu_n) < \delta_n$ also verifies

$$(2) \quad I_\nu(r_n/2, q) \geq C_q r_n^{-nq}.$$

We then consider the dense G_δ -set

$$\mathcal{R} = \bigcap_{m \geq 0} \bigcup_{n \geq m} B_L(\nu_n, \delta_n).$$

Picking any $\nu \in \mathcal{R}$, we can find n as large as we want and the corresponding r_n as small as we want such that (2) holds true. Dividing by $(q-1)\log r_n$, which is positive, and taking the limsup, this yields $\overline{D}_\nu(q) = +\infty$, and this holds for a typical $\nu \in \mathcal{P}(K)$.

5.3. Typical lower L^q -dimensions, $q \in (0, 1)$. We begin by applying the anti-Frostman lemma 2.12 to get a measure $m \in \mathcal{P}(K)$ and some $C, r_0 > 0$ such that $m(B(x, r)) \geq Cr^t$ for all $x \in K$ and all $0 < r \leq r_0$, where $t > \overline{\dim}_B(K)$. Let (μ_n) be a sequence of $\mathcal{F}(K)$ which is dense in $\mathcal{P}(K)$. We write $\mu_n = \sum_{i=1}^N p_i \delta_{x_i}$ and we also set

$$r_n = \min\left(\frac{1}{n}, \min_i(p_i)^{n(1-q)}\right), \quad \varepsilon_n = r_n^{t(1-q)/q} \text{ and } \nu_n = (1 - \varepsilon_n)\mu_n + \varepsilon_n m.$$

Observe that (ε_n) goes to zero, so that (ν_n) remains dense in $\mathcal{P}(K)$. Let also $Y_n = \bigcup_{i=1}^N B(x_i, r_n)$. It is easy to check that

- $\forall x \in Y_n, \nu_n(B(x, r_n)) \geq \min_i(p_i)/2$;
- $\forall x \notin Y_n, \nu_n(B(x, r_n)) \geq C\varepsilon_n r_n^t$.

Thus, one obtains

$$\begin{aligned} I_{\nu_n}(r_n, q) &= \int_{Y_n} \frac{d\nu_n(x)}{\nu_n(B(x, r_n))^{1-q}} + \int_{Y_n^c} \frac{d\nu_n(x)}{\nu_n(B(x, r_n))^{1-q}} \\ &\leq C \left(\min(p_i)^{q-1} + \varepsilon_n^q r_n^{-t(1-q)} \right). \end{aligned}$$

(in this proof, the letter C denotes a positive real number which may change from line to line, remaining always independent of n). The values that we impose on r_n and ε_n ensure that

$$I_{\nu_n}(r_n, q) \leq C \left(\frac{1}{r_n^{1/n}} + 1 \right).$$

Now, Corollary 2.3 gives us some $\delta_n > 0$ such that any $\nu \in \mathcal{P}(K)$ with $L(\nu, \nu_n) < \delta_n$ satisfies

$$I_{\nu_n}(2r_n, q) \leq C \left(\frac{1}{r_n^{1/n}} + 1 \right).$$

It is now routine to prove that

$$\mathcal{R} = \bigcap_{m \geq 0} \bigcup_{n \geq m} B_L(\nu_n, \delta_n)$$

is the dense G_δ -set we are looking for to prove that, typically, $\underline{D}_\mu(q) = 0$ when $q \in (0, 1)$.

6. TYPICAL L^1 -DIMENSION

We now turn to the L^1 -dimensions. Lemma 2.16 yields that, for a typical measure $\mu \in \mathcal{P}(K)$,

$$0 \leq \underline{D}_\mu(1) \leq \underline{D}_\mu(1/2) = 0 \text{ and } s_u \leq \overline{D}_\mu(2) \leq \overline{D}_\mu(1) \leq \overline{D}_\mu(1/2) = s.$$

We now prove the nontrivial part of Theorem 1.8. We shall need several times the following analog of Lemma 4.1.

Lemma 6.1. *Let K be a compact subset of \mathbb{R}^d , let $A \subset K$, let $\mu \in \mathcal{P}(K)$ and let $r > 0$. Then*

$$\int_A \log \mu(B(x, 2r)) d\mu(x) \geq -\mu(A(r)) \log \mathbf{N}_r(A) - e^{-1}.$$

Proof. We follow [Ols07]. For brevity, write $N = \mathbf{N}_r(A)$. Let $x_1, \dots, x_N \in K$ be such that $A \subset \bigcup_{i=1}^N B(x_i, r)$. Put $E_1 = B(x_1, r)$ and $E_i = B(x_i, r) \setminus \bigcup_{j=1}^{i-1} B(x_j, r)$ for $i = 2, \dots, N$. Since for any $x \in E_i$, $E_i \subset B(x, 2r)$, we may write

$$\begin{aligned} \int_A \log \mu(B(x, 2r)) d\mu(x) &= \sum_i \int_{A \cap E_i} \log \mu(B(x, 2r)) d\mu(x) \\ &\geq \sum_i \int_{A \cap E_i} \log (\mu(E_i \cap K)) d\mu(x) \\ &\geq \sum_i \mu(E_i \cap K) \log (\mu(E_i \cap K)). \end{aligned}$$

Now, we apply Jensen's inequality to the convex function $\phi : (0, +\infty) \rightarrow \mathbb{R}$, $t \mapsto t \log t$. Observing that $\bigcup_{i=1}^N E_i \cap K \subset A(r)$ and that ϕ is bounded from below by e^{-1} , we get

$$\begin{aligned} \int_A \log \mu(B(x, 2r)) d\mu(x) &\geq N \sum_{i=1}^N \frac{1}{N} \phi(\mu(E_i \cap K)) \\ &\geq N \phi \left(\sum_i \frac{1}{N} \mu(E_i \cap K) \right) \\ &\geq \phi \left(\mu \left(\bigcup_i E_i \cap K \right) \right) - \mu \left(\bigcup_i E_i \cap K \right) \log N \\ &\geq -e^{-1} - \mu(A(r)) \log N. \end{aligned}$$

□

We divide the proof of Theorem 1.8 into four parts. For convenience, we set :

$$s_{\text{conv}} = \overline{\dim}_{B, \text{conv}}(K) \text{ and } s_{\text{conv}}^{\max} = \overline{\dim}_{B, \text{conv}, \max}(K).$$

6.1. The lower bound. Let $t < s_{\text{conv}}$. One intends to show that $\overline{D}_\mu(1) \geq t$ for a typical measure $\mu \in \mathcal{P}(K)$. Let $(x_i)_{i \geq 1}$ be a dense sequence of distinct points in K and let $n \geq 1$. Let $\rho_n > 0$ be such that $4\rho_n < \min(\|x_i - x_j\|; 1 \leq i < j \leq n)$ and $\rho_n < 1/n$. Let finally $p_1, \dots, p_n > 0$ be such that $\sum_i p_i = 1$. By definition of s_{conv} , one may find $r_n < \rho_n$ such that, setting $P_i = \mathbf{P}_{r_n}(K \cap B(x_i, \rho_n))$,

$$\sum_{i=1}^n p_i \log(P_i) \geq -t \log r_n.$$

We then consider, for a fixed i in $1, \dots, n$, points x_i^j , $j = 1, \dots, P_i$, which are the centers in $K \cap B(x_i, \rho_n)$ of disjoint balls of radius r_n . We observe that our choice of ρ_n ensures that all balls $B(x_i^j, r_n)$, $i = 1, \dots, n$, $j = 1, \dots, P_i$ are pairwise disjoint. We set

$$\mu_{n, (p_i)} = \sum_{i=1}^n \frac{p_i}{P_i} \sum_{j=1}^{P_i} \delta_{x_i^j}$$

and we majorize $I_{\mu_{n, (p_i)}}(r_n, 1)$:

$$\begin{aligned} I_{\mu_{n, (p_i)}}(r_n, 1) &= \sum_{i=1}^n \frac{p_i}{P_i} \sum_{j=1}^{P_i} \log \left(\mu_{n, (p_i)}(B(x_i^j, r_n)) \right) \\ &= \sum_{i=1}^n p_i \log(p_i) - \sum_{i=1}^n p_i \log(P_i) \\ &\leq t \log r_n. \end{aligned}$$

Now, Corollary 2.5 gives us a real number $\delta_{n, (p_i)} > 0$ such that $L(\nu, \mu_{n, (p_i)}) < \delta_{n, (p_i)}$ implies $I_\nu(r_n/2, 1) \leq \log 2 + (1 - \frac{1}{n}) t \log r_n$. We conclude by setting

$$\begin{aligned} \Lambda_n &= \left\{ \mu_{n, (p_i)}; p_i > 0, \sum_i p_i = 1 \right\} \\ \mathcal{R}_n &= \bigcup_{\mu_{n, (p_i)} \in \Lambda_n} B_L(\mu_{n, (p_i)}, \delta_{n, (p_i)}) \\ \mathcal{R} &= \bigcap_{m \geq 1} \bigcup_{n \geq m} \mathcal{R}_n. \end{aligned}$$

\mathcal{R} is a dense G_δ subset of $\mathcal{P}(K)$ and any $\nu \in \mathcal{R}$ satisfies

$$\overline{D}_\nu(1) \geq \limsup_{n \rightarrow +\infty} \left(1 - \frac{1}{n} \right) t = t.$$

6.2. The upper bound. Let $t > s_{\text{conv}}^{\max}$. One intends to show that there exists a dense and open set of measures μ of $\mathcal{P}(K)$ such that $\overline{D}_\mu(1) \leq t$. We first fix $t_0 \in (s_{\text{conv}}^{\max}, t)$ and $\eta > 0$ such that $(1 + 2\eta)t_0 < t$. Let (y_n) be a dense sequence of distinct points in K and let (ρ_n) be a sequence of $(0, 1)$ such that

$$\begin{cases} \rho_n \leq \frac{1}{n} \\ B(y_i, 2\rho_n) \cap B(y_j, 2\rho_n) = \emptyset \text{ for any } i \neq j \text{ in } \{1, \dots, n\}. \end{cases}$$

Let $N \geq 1$ and let $n \leq N$. By definition of s_{conv}^{\max} , we know that we may find an integer $M_{n,N} > 0$, elements $x_n^1, \dots, x_n^{M_{n,N}}$ contained in $B(y_n, \rho_n)$, a real number $\delta_{n,N} \in (0, \rho_N)$ and real numbers $p_n^1, \dots, p_n^{M_{n,N}}$ in $(0, 1)$ with $\sum_i p_n^i = 1$ satisfying

$$\sum_i \frac{p_n^i \log \mathbf{N}_r(K \cap B(x_n^i, \delta_{n,N}))}{-\log r} \leq t_0$$

provided r is small enough. Reducing $\delta_{n,N}$ if necessary, we can always assume that the balls $B(x_n^i, 2\delta_{n,N})$, for $i = 1, \dots, M_{n,N}$, are pairwise disjoint and that they are contained in $B(y_n, \rho_N)$. Let also \mathcal{Q}_N be defined by

$$\mathcal{Q}_N = \left\{ (q_1, \dots, q_N) \in (0, 1); \sum_i q_i = 1 \right\}.$$

For any $\mathbf{q} = (q_1, \dots, q_N)$ in \mathcal{Q}_N , we also fix $\varepsilon_{N,\mathbf{q}} \in (0, 1)$ satisfying

$$\begin{cases} (d+1)\varepsilon_{N,\mathbf{q}} \leq \eta t_0 \\ \varepsilon_{N,\mathbf{q}}(1 - p_n^i q_n) \leq \eta p_n^i q_n \text{ for any } n = 1, \dots, N \text{ and any } i = 1, \dots, M_{n,N}. \end{cases}$$

We finally set, for $N \geq 1$ and $\mathbf{q} \in \mathcal{Q}_N$,

$$\mathcal{U}_{N,\mathbf{q}} = \left\{ \mu \in \mathcal{P}(K) \ ; \ \forall n = 1, \dots, N, \forall i = 1, \dots, M_{n,N}, \right. \\ \left. \begin{aligned} \mu(B(x_n^i, \delta_{n,N}/2)) &> (1 - \varepsilon_{N,\mathbf{q}}) p_n^i q_n \text{ and} \\ \mu(B(y_n, \rho_N)) &> (1 - \varepsilon_{N,\mathbf{q}}) q_n \end{aligned} \right\}.$$

Each $\mathcal{U}_{N,\mathbf{q}}$ is open by Lemma 2.6 and $\bigcup_{N,\mathbf{q}} \mathcal{U}_{N,\mathbf{q}}$ is dense by Lemma 2.7. Let us show that any $\mu \in \mathcal{U}_{N,\mathbf{q}}$ satisfies $\overline{D}_\mu(1) \leq t$. We set

$$Y_N = K \setminus \bigcup_{n=1}^N \bigcup_{i=1}^{M_{n,N}} B(x_n^i, \delta_{n,N})$$

and we apply Lemma 6.1 to each $B(x_n^i, \delta_{n,N})$ and to Y_N . Taking into account that we have a partition of K , we thus obtain

$$\begin{aligned} I_\mu(2r, 1) &\geq - \sum_{n=1}^N \sum_{i=1}^{M_{n,N}} \mu(B(x_n^i, \delta_{n,N} + r)) \log \mathbf{N}_r(K \cap B(x_n^i, \delta_{n,N})) \\ &\quad - \mu(Y_N(r)) \log \mathbf{N}_r(Y_N) + C_\mu, \end{aligned}$$

where C_μ does not depend on r . We first look at the last term. Provided r is small enough,

$$Y_N(r) \subset K \setminus \bigcup_{n=1}^N \bigcup_{i=1}^{M_{n,N}} B(x_n^i, \delta_{n,N}/2)$$

so that

$$\begin{aligned} \mu(Y_N(r)) &\leq 1 - \sum_n \sum_i (1 - \varepsilon_{N,\mathbf{q}}) p_n^i q_n \\ &\leq \varepsilon_{N,\mathbf{q}}. \end{aligned}$$

On the other hand, since Y_N is contained in \mathbb{R}^d , which has dimension d , we know that

$$\log(\mathbf{N}_r(Y_N)) \leq -(d+1) \log r$$

provided r is small enough. Thus, taking into account the value of $\varepsilon_{N,\mathbf{q}}$,

$$-\mu(Y_N(r)) \log \mathbf{N}_r(Y_N) \geq \eta t_0 \log r.$$

The analysis of the first term is slightly more delicate. We first observe that, provided r is small enough,

$$\begin{aligned} \mu(B(x_n^i, \delta_{n,N} + r)) &\leq \mu(B(x_n^i, 2\delta_{n,N})) \\ &\leq \mu(B(y_n, \rho_N)) - \sum_{j \neq i} \mu(B(x_n^j, \delta_{n,N})) \\ &\leq 1 - \sum_{m \neq n} \mu(B(y_m, \rho_N)) - \sum_{j \neq i} (1 - \varepsilon_{N,\mathbf{q}}) p_n^j q_n \\ &\leq 1 - (1 - \varepsilon_{N,\mathbf{q}}) \sum_{m \neq n} q_m - (1 - \varepsilon_{N,\mathbf{q}})(1 - p_n^i) q_n \\ &\leq 1 - (1 - \varepsilon_{N,\mathbf{q}})(1 - q_n) - (1 - \varepsilon_{N,\mathbf{q}})(1 - p_n^i) q_n \\ &\leq p_n^i q_n + \varepsilon_{N,\mathbf{q}}(1 - p_n^i) q_n \\ &\leq (1 + \eta) p_n^i q_n. \end{aligned}$$

Thus,

$$\begin{aligned} I_\mu(2r, 1) &\geq -(1 + \eta) \sum_n q_n \sum_i p_n^i \log \mathbf{N}_r(K \cap B(x_n^i, \delta_{n,N})) + \eta t_0 \log r + C_\mu \\ &\geq (1 + 2\eta) t_0 \log r + C_\mu, \end{aligned}$$

provided again that r is small enough. Dividing by $\log r$ and taking the limsup, we find

$$\overline{D}_\mu(1) \leq (1 + 2\eta) t_0 \leq t.$$

6.3. Optimality of the lower bound. Suppose that there exists $t > s_{\text{conv}}$ such that $\mathcal{R} = \{\mu \in \mathcal{P}(K); \overline{D}_\mu(1) \geq t\}$ is residual. We shall arrive at a contradiction. The argument follows rather closely that of the previous subsection. We fix $t_0 \in (s_{\text{conv}}, t)$ and $\eta > 0$ such that $(1 + 2\eta)t_0 < t$. By definition of s_{conv} , one may find $x_1, \dots, x_N \in K$, $\rho > 0$ with $4\rho < \min(\|x_i - x_j\|; i \neq j)$ and $p_1, \dots, p_N > 0$ with $\sum_i p_i = 1$ such that any $r > 0$ sufficiently small verifies

$$\sum_{i=1}^N p_i \log(\mathbf{N}_r(K \cap B(x_i, \rho))) \leq (-\log r) t_0.$$

Now, let $\varepsilon > 0$ satisfying the following conditions:

$$\begin{cases} (d+1)\varepsilon &\leq \eta t_0 \\ \varepsilon &\leq \eta \frac{p_i}{1-p_i} \text{ for any } i = 1, \dots, N. \end{cases}$$

Let finally

$$\mathcal{U} = \bigcap_{i=1}^N \{\mu \in \mathcal{P}(K); \mu(B(x_i, \rho/2)) > (1 - \varepsilon) p_i\}.$$

\mathcal{U} is open and nonempty and one can pick $\mu \in \mathcal{U} \cap \mathcal{R}$. We set $Y = K \setminus \bigcup_{i=1}^N B(x_i, \rho)$ and we apply Lemma 6.1 to get

$$\begin{aligned} I_\mu(2r, 1) &\geq - \sum_{i=1}^N \mu(B(x_i, \rho + r)) \log \mathbf{N}_r(K \cap B(x_i, \rho)) - \mu(Y(r)) \log \mathbf{N}_r(Y) + C_\mu \\ &\geq - \sum_{i=1}^N \mu(B(x_i, 2\rho)) \log \mathbf{N}_r(K \cap B(x_i, \rho)) - \mu(Y(\rho/2)) \log \mathbf{N}_r(Y) + C_\mu \end{aligned}$$

provided $r < \rho/2$. Now we observe that

$$\begin{aligned} \mu(B(x_i, 2\rho)) &\leq 1 - \sum_{j \neq i} \mu(B(x_j, \rho)) \\ &\leq 1 - \sum_{j \neq i} p_j(1 - \varepsilon) \\ &\leq p_i + \varepsilon(1 - p_i) \leq (1 + \eta)p_i. \end{aligned}$$

In the same vein, we also get

$$\mu(Y(\rho/2)) \leq 1 - \sum_j \mu(B(x_j, \rho/2)) \leq \varepsilon,$$

whereas, provided r is small enough,

$$\log \mathbf{N}_r(Y) \leq -(d+1) \log r.$$

Coming back to $I_\mu(2r, 1)$ and summarizing the previous results, we conclude that

$$\begin{aligned} I_\mu(2r, 1) &\geq - \sum_{i=1}^N (1 + \eta)p_i \log \mathbf{N}_r(K \cap B(x_i, \rho)) + (d+1)\varepsilon \log r + C_\mu \\ &\geq (1 + 2\eta)t_0 \log r + C_\mu \end{aligned}$$

where C_μ does not depend on r . Dividing by $\log r$ and taking the limsup, we obtain that $\overline{D}_\mu(1) \leq (1 + 2\eta)t_0 < t$, a contradiction with $\mu \in \mathcal{R}$.

6.4. Optimality of the upper bound. Suppose that there exists $t < s_{\text{conv}}^{\max}$ such that $\mathcal{R} = \{\mu \in \mathcal{P}(K); \overline{D}_\mu(1) \leq t\}$ is residual. Let $t_0 \in (t, s_{\text{conv}}^{\max})$ and let $y \in K$, $\rho > 0$ be such that, setting $E = K \cap B(y, \rho)$, $\overline{\dim}_{B, \text{conv}}(E) > t_0$. Let also $\beta > 0$ be such that $\frac{(1-\beta)^3}{1+\beta} t_0 > t$. We start from a sequence $(\mu_n) \subset \mathcal{F}(K)$ which is dense in $\mathcal{P}(K)$ and such that $\mu_n(E) > 0$ for any $n \geq 1$. We fix $n \geq 1$ and we write

$$\mu_n = \sum_{i=1}^q q_i \delta_{y_i} + \sum_{i=1}^p p_i \delta_{x_i}$$

where $x_i \in E$, $y_i \notin E$, $p_i, q_i > 0$. Let $\alpha > 0$ be such that $4\alpha < \min(\|x_i - x_j\|; i \neq j)$, $4\alpha < \min(\|y_i - y_j\|; i \neq j)$, $4\alpha < \min(\|x_i - y_j\|)$ and $\alpha < 1/n$. By definition of the convex upper box dimension, we know that one can find $r_n < \min(\alpha, 1/n)$ as small as we want such that

$$\sum_{i=1}^p p_i \log \mathbf{P}_{r_n}(E \cap B(x_i, \alpha)) \geq (-\log r_n) t_0 \sum_{i=1}^p p_i = (-\log r_n) t_0 \mu_n(E).$$

For simplicity, we set $P_i = \mathbf{P}_{r_n}(E \cap B(x_i, \alpha))$ and we consider points $x_i^1, \dots, x_i^{P_i}$ all in $E \cap B(x_i, \alpha)$ such that the balls $B(x_i^j, r_n)$ are disjoint. We then set

$$\begin{aligned}\nu_{n,i} &= \frac{1}{P_i} \sum_{j=1}^{P_i} \delta_{x_i^j} \\ \nu_n &= \sum_{i=1}^q q_i \delta_{y_i} + \sum_{i=1}^p p_i \nu_{n,i}.\end{aligned}$$

ν_n is close to μ_n . Indeed, for any $f \in \text{Lip}(K)$,

$$\left| \int_K f d\nu_n - \int_K f d\mu_n \right| \leq \sum_{i=1}^p p_i \|x_i - x_i^j\| \leq \alpha \leq \frac{1}{n}.$$

In particular, the sequence (ν_n) keeps dense in $\mathcal{P}(K)$. Furthermore, $\nu_n(E) = \mu_n(E) > 0$ and

$$\begin{aligned}I_{\nu_n}(r_n, 1) &= \sum_{i=1}^q q_i \log q_i + \sum_{i=1}^p \frac{p_i}{P_i} \sum_{j=1}^{P_i} \log \left(\frac{p_i}{P_i} \right) \\ &= \sum_{i=1}^q q_i \log q_i + \sum_{i=1}^p p_i \log(p_i) - \sum_{i=1}^p p_i \log(P_i) \\ &\leq \sum_{i=1}^q q_i \log(q_i) + \sum_{i=1}^p p_i \log(p_i) + \log(r_n) t_0 \nu_n(E).\end{aligned}$$

Provided r_n is small enough, this implies

$$I_{\nu_n}(r_n, 1) \leq (1 - \beta) t_0 \log(r_n) \nu_n(E).$$

Now, Lemma 2.1 and Corollary 2.5 give us a real number $\delta_n > 0$ such that $L(\nu, \nu_n) < \delta_n$ implies

$$\begin{cases} I_{\nu}(r_n/2, 1) &\leq (1 - \beta) I_{\nu_n}(r_n, 1) + \log 2 \\ \nu(E) &\leq \nu_n(E(\alpha)) + \beta \nu_n(E) = (1 + \beta) \nu_n(E). \end{cases}$$

Thus,

$$I_{\nu}(r_n/2, 1) \leq \frac{(1 - \beta)^2}{1 + \beta} t_0 \log(r_n) \nu(E) + \log 2.$$

We define

$$\mathcal{R}' = \bigcap_{m \geq 1} \bigcup_{n \geq m} B_L(\nu_n, \delta_n)$$

which is a dense G_δ -subset of $\mathcal{P}(K)$ such that any $\nu \in \mathcal{R}'$ verifies $\overline{D}_\nu(1) \geq \frac{(1 - \beta)^2}{1 + \beta} t_0 \nu(E)$. Finally, we considered

$$\mathcal{U} = \{ \mu \in \mathcal{P}(K); \mu(E) > 1 - \beta \}$$

which is open and nonempty. Then any measure μ in the nonempty intersection $\mathcal{R} \cap \mathcal{R}' \cap \mathcal{U}$ satisfies the contradictory conditions

$$\overline{D}_\mu(1) \leq t \text{ and } \overline{D}_\mu(1) \geq \frac{(1 - \beta)^3}{1 + \beta} t_0 > t.$$

6.5. Examples. To apply Theorem 1.8, we need to be able to compute the dimensions which are involved. This is indeed possible for simple compact spaces.

Example 6.2. Let $K = \{0\} \cup [1, 2]$. Then a typical measure $\mu \in \mathcal{P}(K)$ satisfies $\overline{D}_\mu(1) \in [0, 1]$ and this interval is the best possible.

Proof. That $s_{\text{conv}} = 0$ is easy. It suffices to take in the definition of the convex upper box dimension $N = 1$ and $x_1 = 0$. To prove that $s_{\text{conv}}^{\text{max}} = 1$, one may observe that $\overline{\dim}_{B, \text{conv}}([1, 2]) = 1$. \square

Example 6.3. Let E and F be the two Cantor sets defined in Section 4.3 and let $K = E \cup F$. Then a typical measure $\mu \in \mathcal{P}(K)$ satisfies $\overline{D}_\mu(1) \in \left[\frac{13 \log 3}{6 \log 100}, \frac{5 \log 3}{2 \log 100} \right]$ and this interval is the best possible.

An interesting feature of the previous example is that $s_{\text{conv}} > s_u$.

Proof. By homogeneity of the two Cantor sets, to compute s_{conv} , we just need to compute

$$\inf_{\alpha+\beta=1} \limsup_{n \rightarrow +\infty} \frac{\alpha \mathbf{N}_{100^{-n}}(E) + \beta \mathbf{N}_{100^{-n}}(F)}{n \log 100}.$$

From the construction of the Cantor sets, it is easy to check that, for a fixed choice of α and β , the limsup will be attained along either the sequence $(2n_k)_k$ or the sequence $(4n_k)_k$. Moreover

$$\begin{aligned} \mathbf{N}_{100^{-2n_k}}(E) &= 9^{n_k} 27^{n_k} = 3^{5n_k} & \mathbf{N}_{100^{-2n_k}}(F) &= 3^{4n_k} \\ \mathbf{N}_{100^{-4n_k}}(E) &= 3^{8n_k} & \mathbf{N}_{100^{-4n_k}}(F) &= 3^{9n_k}. \end{aligned}$$

Thus,

$$\begin{aligned} s_{\text{conv}} &= \inf_{\alpha+\beta=1} \max \left((5\alpha + 4\beta) \frac{\log 3}{2 \log 100}, (8\alpha + 9\beta) \frac{\log 3}{4 \log 100} \right) \\ &= \frac{\log 3}{4 \log 100} \inf_{\alpha \in [0, 1]} (8 + 2\alpha, 9 - \alpha). \end{aligned}$$

The minimum is obtained for $\alpha = 1/3$ (when the two terms are equal) so that

$$s_{\text{conv}} = \frac{13 \log 3}{6 \log 100}.$$

That $s_{\text{conv}}^{\text{max}} = \dim_{\mathcal{P}}(K) = \frac{5 \log 3}{2 \log 100}$ is easier. Indeed, we can restrict ourselves to E where, by homogeneity, $\overline{\dim}_{B, \text{conv}}(E) = \overline{\dim}_B(E)$. \square

Example 6.4. Let K be a Ahlfors-regular compact set with Hausdorff dimension s . Then a typical measure $\mu \in \mathcal{P}(K)$ satisfies $\overline{D}_\mu(1) = s$.

Proof. For such a compact set,

$$s \leq \inf_{x \in K} \underline{\dim}_{B, \text{loc}}(x, K) \leq s_{\text{conv}} \leq s_{\text{conv}}^{\text{max}} \leq \overline{\dim}_B(K) = s.$$

\square

7. CONCLUDING REMARKS

7.1. **Case $q = \pm\infty$.** The L^q -dimensions of a probability measure μ have also a meaning for $q = -\infty$ and $q = +\infty$. Their definitions are

$$\begin{aligned}\overline{D}_\mu(-\infty) &= \limsup_{r \rightarrow 0} \frac{\log \inf_{x \in \text{supp}(\mu)} \mu(B(x, r))}{\log r} \\ \underline{D}_\mu(-\infty) &= \liminf_{r \rightarrow 0} \frac{\log \inf_{x \in \text{supp}(\mu)} \mu(B(x, r))}{\log r} \\ \overline{D}_\mu(+\infty) &= \limsup_{r \rightarrow 0} \frac{\log \sup_{x \in \text{supp}(\mu)} \mu(B(x, r))}{\log r} \\ \underline{D}_\mu(+\infty) &= \liminf_{r \rightarrow 0} \frac{\log \sup_{x \in \text{supp}(\mu)} \mu(B(x, r))}{\log r}.\end{aligned}$$

In [Ols07], Olsen shows that a typical $\mu \in \mathcal{P}(K)$ satisfies

$$\underline{D}_\mu(+\infty) = 0 \text{ and } \inf_{x \in K} \underline{\dim}_{B, \text{loc}}(x, K) \leq \overline{D}_\mu(+\infty) \leq \inf_{x \in K} \overline{\dim}_{B, \text{loc}}(x, K).$$

Our methods allow us to determine the exact typical value of the L^q -dimensions, for $q = -\infty$ and $q = +\infty$.

Theorem 7.1. *Let K be an infinite compact subset of \mathbb{R}^d . Then a typical $\mu \in \mathcal{P}(K)$ satisfies*

$$\begin{aligned}\underline{D}_\mu(+\infty) &= 0 & \overline{D}_\mu(+\infty) &= \overline{\dim}_{B, \text{loc}, \text{unif}}(K) \\ \underline{D}_\mu(-\infty) &= \underline{\dim}_B(K) & \overline{D}_\mu(-\infty) &= +\infty.\end{aligned}$$

Among these values, that of $\underline{D}_\mu(-\infty)$ is surprising, because it is different from the value of $\underline{D}_\mu(q)$ when $q < 0$.

Proof. For commodity reasons, we set $s_u = \overline{\dim}_{B, \text{loc}, \text{unif}}(K)$. It is easy to show that, for any $q \in \mathbb{R}$,

$$\underline{D}_\mu(+\infty) \leq \underline{D}_\mu(q) \leq \underline{D}_\mu(-\infty) \text{ and } \overline{D}_\mu(+\infty) \leq \overline{D}_\mu(q) \leq \overline{D}_\mu(-\infty).$$

Thus, Theorem 1.6 already implies that $\overline{D}_\mu(+\infty) \leq s_u$ and that $\overline{D}_\mu(-\infty) = +\infty$ for a typical $\mu \in \mathcal{P}(K)$. Thus, it remains to show the typical (in)equalities $\overline{D}_\mu(+\infty) \geq s_u$ and $\underline{D}_\mu(-\infty) = \underline{\dim}_B(K)$. We begin by showing that $\overline{D}_\mu(+\infty) \geq s_u$ for a typical $\mu \in \mathcal{P}(K)$. Let $t < s_u$ and let us apply the results of Section 4.1. They provide a dense G_δ -set \mathcal{R} and a sequence (r_n) going to zero such that, for any $\nu \in \mathcal{R}$, we can find n as large as we want such that

$$\nu(B(x, r_n)) \leq Cr_n^t.$$

This immediately yields $\overline{D}_\nu(+\infty) \geq t$.

We now prove that a typical $\mu \in \mathcal{P}(K)$ satisfies $\underline{D}_\mu(-\infty) \geq \underline{\dim}_B(K)$. For $n \geq 1$, let $P = \mathbf{P}_{2^{-n}}(K)$ be the maximal number of balls of radius 2^{-n} with center in K which do not intersect. Let $B(x_1, 2^{-n}), \dots, B(x_P, 2^{-n})$ be such a family of balls. We set $U_{j,n} = B(x_j, 2^{-n})$, $U'_{j,n} = B(x_j, 2^{-(n+1)})$ and

$$\begin{aligned}\mathcal{R}_n &= \{\mu \in \mathcal{P}(K); \forall j \in \{1, \dots, \mathbf{P}_{2^{-n}}(K)\}, \mu(U'_{j,n}) > 0\} \\ \mathcal{R} &= \bigcap_{n \geq 1} \mathcal{R}_n.\end{aligned}$$

Each \mathcal{R}_n is dense by Lemma 2.8 and open by Lemma 2.6, so that \mathcal{R} is a dense G_δ -subset of $\mathcal{P}(K)$. Pick now any $\mu \in \mathcal{R}$ and let $r \in (0, 1)$. There exists $n \geq 1$ such that $2^{-(n+1)} \leq r < 2^{-n}$. Moreover,

$$\sum_{j=1}^{\mathbf{P}_{2^{-n}}(K)} \mu(U_{j,n}) \leq 1$$

so that there exists $j \in \{1, \dots, \mathbf{P}_{2^{-n}}(K)\}$ with $\mu(U_{j,n}) \leq (\mathbf{P}_{2^{-n}}(K))^{-1}$. Now, since $\mu(U'_{j,n}) > 0$, one may find $y_j \in \text{supp}(\mu) \cap U'_{j,n}$. Noticing that $B(y_j, r/2)$ is contained in $U_{j,n}$, we get

$$\log \mu(B(y_j, r/2)) \leq \log \mu(U_{j,n}) \leq -\log \mathbf{P}_{2^{-n}}(K)$$

which itself yields

$$\frac{\log \inf_{x \in \text{supp}(\mu)} \mu(B(x, r/2))}{\log(r/2)} \geq \frac{-\log \mathbf{P}_{2^{-n}}(K)}{\log(r/2)} \geq \frac{\log \mathbf{P}_{2^{-n}}(K)}{(n+2) \log 2}.$$

Taking the liminf, this shows that $\underline{D}_\mu(-\infty) \geq \underline{\dim}_B(K)$.

We finally prove that $\underline{D}_\mu(-\infty) \leq \underline{\dim}_B(K)$. Let $t > t' > \underline{\dim}_B(K)$. Lemma 2.13 gives us, for each $n \geq 1$, a measure $m_n \in \mathcal{P}(K)$ and a real number $r_n \in (0, 1/n)$ such that $m_n(B(x, r_n/2)) \geq Cr_n^{t'}$ for each $x \in K$, where C just depends on t' . Let now $\mu \in \mathcal{F}(K)$ and let us set

$$\nu_{\mu,n} = (1 - r_n^{(t-t')})\mu + r_n^{t-t'} m_n.$$

For any $x \in K$, $\nu_{\mu,n}(B(x, r_n)) \geq Cr_n^t$. By Lemma 2.1, we may find $\delta_n > 0$ such that any $\nu \in \mathcal{P}(K)$ with $L(\nu, \nu_{\mu,n}) < \delta_n$ satisfies

$$\nu(B(x, 2r_n)) \geq 2Cr_n^t.$$

We then consider

$$\mathcal{R} = \bigcap_{n \geq 1} \bigcup_{\mu \in \mathcal{F}(K)} B(\nu_{\mu,n}, \delta_n)$$

which is a dense G_δ -set. For any $\nu \in \mathcal{R}$, there exists r_n as small as desired such that

$$\frac{\log \inf_{x \in \text{supp} \mu} \nu(B(x, 2r_n))}{\log r_n} \leq t + \frac{\log(2C)}{\log r_n},$$

so that $\underline{D}_\nu(-\infty) \leq t$. □

7.2. On the optimality. It is natural to ask whether the inequalities appearing in the second column of Theorem 1.6 are optimal or not. The third column gives the answer, except for $\overline{D}_\mu(q)$, $q \geq 1$ and $\underline{D}_\mu(q)$, $q \leq 0$. It is very easy to exhibit a measure satisfying $\underline{D}_\mu(q) = 0$, $q < 0$: any Dirac mass does the job. The situation is different for $\overline{D}_\mu(q)$, $q \geq 1$; we just know that $\overline{D}_\mu(1) \leq \dim_{\mathcal{P}}(K)$, but we do not know whether this is always the best bound.

Question 7.2. *What is the biggest possible value of $\overline{D}_\mu(q)$, $q \geq 1$?*

It is conceivable that, at least for $q = 1$, we need a kind of convex version of the packing dimension.

7.3. L^q -dimension along subsequences. When, for a measure $\mu \in \mathcal{P}(K)$, $\underline{D}_\mu(q) < \overline{D}_\mu(q)$, it is natural to ask the following question: for which $\tau \in (\underline{D}_\mu(q), \overline{D}_\mu(q))$ can we find a sequence (r_n) going to zero such that

$$\frac{\log I_\mu(r_n, q)}{(q-1) \log r_n} \rightarrow \tau?$$

The most interesting case is when all values of $(\underline{D}_\mu(q), \overline{D}_\mu(q))$ can be attained. For continuity reasons, this is always the case.

Proposition 7.3. *Let K be a compact subset of \mathbb{R}^d , let $\mu \in \mathcal{P}(K)$ and let $q \in \mathbb{R}$. Then, for any $\tau \in (\underline{D}_\mu(q), \overline{D}_\mu(q))$, there exists a sequence (r_n) going to zero such that*

$$\frac{\log I_\mu(r_n, q)}{(q-1) \log r_n} \rightarrow \tau \text{ when } q \neq 1, \quad \frac{\log I_\mu(r_n, 1)}{\log r_n} \rightarrow \tau \text{ otherwise.}$$

The proof of this proposition is based on an application of the intermediate value theorem. Unfortunately, the map $r \mapsto \frac{1}{q-1} \times \frac{\log I_\mu(r, q)}{\log r}$ does not need to be continuous. However, an enhancement of the intermediate value theorem to semicontinuous functions will be sufficient in our context. It can be found e.g. in [Gui95].

Lemma 7.4. *Let $f : [u, v] \rightarrow \mathbb{R}$ be upper semicontinuous on the right and lower semicontinuous on the left. Suppose moreover that $f(u) \geq f(v)$. Then for any $\lambda \in (f(v), f(u))$, there exists $x \in [u, v]$ such that $f(x) = \lambda$.*

The regularity of $I_\mu(\cdot, q)$ will depend on the position of q with respect to 1.

Lemma 7.5. *Let $\mu \in \mathcal{P}(K)$ and let $q \in \mathbb{R}$. Suppose moreover that, for any $r > 0$, $I_\mu(r, q) \neq +\infty$.*

- (a) *For $q > 1$, $r \mapsto \log I_\mu(r, q)$ is negative, nondecreasing and continuous on the left;*
- (b) *For $q = 1$, $r \mapsto I_\mu(r, 1)$ is negative, nondecreasing and continuous on the left;*
- (c) *For $q < 1$, $r \mapsto \log I_\mu(r, q)$ is positive, nonincreasing and continuous on the left.*

Proof. We just prove the continuity statement. We fix $r > 0$ and we pick a sequence (r_n) increasing to r . Then $(\mu(B(x, r_n)))$ increases to $\mu(B(x, r))$ for any $x \in \text{supp}(\mu)$. Thus (a) follows from the monotone convergence theorem, whereas (b) and (c) follow from Lebesgue's theorem. \square

In order to apply Lemma 7.4 to $r \mapsto \frac{1}{q-1} \times \frac{\log I_\mu(r, q)}{\log r}$, we need a last lemma.

Lemma 7.6. *Let I be an interval, let $g : I \rightarrow (-\infty, 0)$ be continuous and let $f : I \rightarrow (-\infty, 0)$ be nondecreasing. Then fg is upper semicontinuous on the right.*

Proof. Let $r_0 \in I$, let $\varepsilon > 0$ and let $\eta > 0$ such that any $r \in [r_0, r_0 + \eta) \cap I$ satisfies

$$g(r_0) \leq g(r) + \varepsilon.$$

We also know that for these values of r ,

$$f(r_0) \leq f(r) \leq 0,$$

so that

$$\begin{aligned} f(r_0)g(r_0) &\geq f(r)g(r_0) \\ &\geq f(r)g(r) + \varepsilon f(r) \\ &\geq f(r)g(r) + \varepsilon f(r_0). \end{aligned}$$

This shows that $\limsup_{r \rightarrow r_0^+} f(r)g(r) \leq f(r_0)g(r_0)$, which means that fg is upper semi-continuous on the right. \square

Proof of Proposition 7.3. Let $\tau \in (\underline{D}_\mu(q), \overline{D}_\mu(q))$ (if $\underline{D}_\mu(q) = \overline{D}_\mu(q)$), there is nothing to say. Define

$$\phi(r) = \begin{cases} \frac{1}{q-1} \times \frac{\log I_\mu(r, q)}{\log r} & \text{if } q \neq 1 \\ \frac{I_\mu(r, 1)}{\log r} & \text{if } q = 1. \end{cases}$$

By the above lemmas, ϕ is continuous on the left and upper semicontinuous on the right. Moreover, for any $\varepsilon > 0$, one may find $0 < r_1 < r_2 < \varepsilon$ such that

$$\phi(r_1) > \tau \text{ and } \phi(r_2) < \tau.$$

By Lemma 7.4, one can find $r_3 \in (r_1, r_2)$ such that

$$\phi(r_3) = \tau.$$

The conclusion of Proposition 7.3 follows easily. \square

Combining this proposition with Theorem 1.6 yields:

Corollary 7.7. *Let K be an infinite compact subset of \mathbb{R}^d . Write*

$$\begin{aligned} s_{\text{sep}} &= \text{bsi}(K) \\ s_u &= \overline{\dim}_{B, \text{loc}, \text{unif}}(K) \\ s &= \overline{\dim}_B(K). \end{aligned}$$

Define, for $q \in \mathbb{R} \setminus \{1\}$,

$$E_q = \begin{cases} [0, s_u] & \text{for } q > 1 \\ [0, s] & \text{for } q \in (0, 1) \\ [s_{\text{sep}}, s] & \text{for } q = 0 \\ [s_{\text{sep}}, +\infty) & \text{for } q < 0. \end{cases}$$

Then a typical measure $\mu \in \mathcal{P}(K)$ satisfies, for any $q \in \mathbb{R} \setminus \{1\}$, for any $\tau \in E_q$, there exists a sequence (r_n) going to zero such that

$$\frac{\log I_\mu(r_n, q)}{(q-1) \log r_n} \rightarrow \tau.$$

7.4. Prevalence. The notion of genericity used in this paper is in the sense of Baire theorem. One could also consider other notions, in particular that of prevalence. In [Ols10], Olsen studied the typical values of $\underline{D}_\mu(q)$ and $\overline{D}_\mu(q)$ for $q \geq 0$ for this notion of genericity. Surprisingly enough, the results are rather different from the results in the Baire point of view.

Problem. What can be said on $\underline{D}_\mu(q)$ and $\overline{D}_\mu(q)$ for a prevalent measure, when $q < 0$?

REFERENCES

- [Cut95] C.D. Cutler, *Strong and weak duality principles for fractal dimension in euclidean space*, Math. Proc. Camb. Phil. Soc. **118** (1995), 393–410.
- [Die10] R. Diestel, *Graph theory*, Graduate Text in Mathematics, vol. 173, Springer, 2010.
- [Fal97] K. Falconer, *Techniques in fractal geometry*, Wiley, 1997.
- [Fal03] ———, *Fractal geometry: Mathematical foundations and applications*, Wiley, 2003.
- [Gui95] J. Guillerme, *Intermediate value theorems and fixed point theorems for semi-continuous functions in product spaces*, Proc. Amer. Math. Soc. **123** (1995), 2119–2122.

- [HP83] H. Hentschel and I. Procaccia, *The infinite number of generalized dimensions of fractals and strange attractors*, Phys. D **8** (1983), 435–444.
- [Ols05] L. Olsen, *Typical L^q -dimensions of measures*, Monatsh. Math. **146** (2005), 143–157.
- [Ols07] ———, *Typical Rényi dimensions of measures. The cases: $q = 1$ and $q = \infty$* , J. Math. Anal. Appl. **331** (2007), 1425–1439.
- [Ols08] ———, *Typical upper L^q -dimensions of measures for $q \in [0, 1]$* , Bull. Sci. Math. **132** (2008), 551–561.
- [Ols10] ———, *Prevalent L^q -dimensions of measures*, Math. Proc. Camb. Phil. Soc. **149** (2010), 553–571.
- [Par67] K.R. Parthasarathy, *Probability measures on metric spaces*, Probability and Mathematical Statistics, Academic Press, 1967.
- [Tri82] C. Tricot, *Two definitions of fractional dimension*, Math. Proc. Camb. Phil. Soc. **91** (1982), 57–74.

CLERMONT UNIVERSITÉ, UNIVERSITÉ BLAISE PASCAL, LABORATOIRE DE MATHÉMATIQUES, BP 10448, F-63000 CLERMONT-FERRAND - CNRS, UMR 6620, LABORATOIRE DE MATHÉMATIQUES, F-63177 AUBIERE

E-mail address: `Frederic.Bayart@math.univ-bpclermont.fr`