



Universiteit  
Leiden  
The Netherlands

## Mitigation of *Pseudomonas syringae* virulence by signal inactivation

Sieber, S.; Mathew, A.J.C.; Kohler, T.; Bar, M.; Carrion Bravo, V.J.; Cazorla, F.M.; ... ; Gademann, K.

### Citation

Sieber, S., Mathew, A. J. C., Kohler, T., Bar, M., Carrion Bravo, V. J., Cazorla, F. M., ... Gademann, K. (2021). Mitigation of *Pseudomonas syringae* virulence by signal inactivation. *Science Advances*, 7(37). doi:10.1126/sciadv.abg2293

Version: Publisher's Version  
License: [Creative Commons CC BY-NC 4.0 license](https://creativecommons.org/licenses/by-nc/4.0/)  
Downloaded from: <https://hdl.handle.net/1887/3275219>

**Note:** To cite this publication please use the final published version (if applicable).

## PLANT SCIENCES

Mitigation of *Pseudomonas syringae* virulence by signal inactivation

Simon Sieber<sup>1†</sup>, Anugraha Mathew<sup>2†</sup>, Christian Jenul<sup>2</sup>, Tobias Kohler<sup>1</sup>, Max Bär<sup>1</sup>, Víctor J. Carrión<sup>3</sup>, Francisco M. Cazorla<sup>4</sup>, Urs Stalder<sup>1</sup>, Ya-Chu Hsieh<sup>1</sup>, Laurent Bigler<sup>1</sup>, Leo Eberl<sup>2\*</sup>, Karl Gademann<sup>1\*</sup>

*Pseudomonas syringae* is an important plant pathogen of many valuable crops worldwide, with more than 60 identified pathovars. The phytotoxins produced by these organisms were related to the severity of the damage caused to the plant. An emerging strategy to treat bacterial infections relies on interference with their signaling systems. In this study, we investigated *P. syringae* pv. *syringae*, which produces the virulence factor mangotoxin that causes bacterial apical necrosis on mango leaves. A previously unknown signaling molecule named leudiazin was identified, determined to be unstable and volatile, and responsible for mangotoxin production. A strategy using potassium permanganate, compatible with organic farming, was developed to degrade leudiazin and thus to attenuate the pathogenicity of *P. syringae* pv. *syringae*.

## INTRODUCTION

*Pseudomonas* represent an excellent example of both benign and pathogenic plant-bacteria interactions (1). While some strains are beneficial for plants (2), others, such as members of the genus *Pseudomonas syringae* (*P. syringae*), are important pathogens, which infect almost all economically important crop species (3). Decades of studies have revealed the broad repertoire of virulence strategies used by *P. syringae*, which include large numbers of functionally redundant type III secretion system (T3SS) effectors and phytotoxins (4). A toxin that has a crucial ecological impact is mangotoxin (5), which is produced by *P. syringae* pv. *syringae* (Pss) and identified as the causal agent of bacterial apical necrosis (BAN) of mango trees (6). While the structure of the toxin remains unknown, the *mbo* gene cluster encoding mangotoxin biosynthesis genes was first described in Pss (7) and was later also identified in many other *P. syringae* pathovars (8, 9).

Investigations of the regulation of mangotoxin production has led to the identification of the *mgo* operon (*mgoBCAD*), which was shown to be required for the production of an unknown signaling molecule (Fig. 1A) (10, 11). The *mgo* gene cluster comprises four genes including a nonribosomal peptide synthetase (NRPS) complex (MgoA), and homologous clusters have been identified in other strains, such as *Pseudomonas entomophila* (*pvf* cluster) (12), *Pseudomonas aurantiaca* PB-St2 (13), *Pseudomonas* sp. SH-C52 (14, 15), several *Pseudomonas fluorescens* (16, 17), and various *Burkholderia* species (*ham* cluster) (18, 19). Recently, we found that the *ham* cluster directs the synthesis of the diazeniumdiolate valdiazin (1), which was identified as a signal molecule that positively autoregulates its own expression (18). Furthermore, in 2018, Morgan *et al.* investigated the substrate specificity of the NRPS domain of MgoA and its homologs PvfC and HamD and concluded that the different strains produce at least two distinct signaling molecules (20). A comparison

of the *mgo*, *pvf*, and *ham* clusters (Fig. 1) reveals that the four genes required for the biosynthesis of the signals are highly conserved in *P. syringae*, *Pseudomonas entomophila*, and *Burkholderia cenocepacia*, respectively (tables S1 to S4). Here, we demonstrate that mangotoxin production is controlled by leudiazin (2, Fig. 1B), a new member of the diazeniumdiolate family of bacterial signal molecules. Furthermore, we speculated that the knowledge of the signaling molecule structure might pave the way to develop a strategy for plant protection.


## RESULTS

Given the homology of the NRPS complexes MgoA and HamD (fig. S1), we speculated that the *mgo* signal could be a related diazeniumdiolate compound. We used a transcriptional fusion of the promoter of the mangotoxin gene cluster (*mbo*) to the *lacZ* gene in a Pss  $\Delta$ *mgoA* mutant background ( $\Delta$ *mgoA*) to detect the signal-inducing mangotoxin production. The biosensor was activated by the wild-type extract but not by the  $\Delta$ *mgoA* extract (fig. S2). Valdiazin (1) only weakly induced the biosensor relative to the Pss extract, suggesting that the unknown signaling molecule may not be valdiazin (fig. S2). Bioassay-guided fractionation under various conditions did not result in a pure product, as bioactivity was lost during the purification procedure, suggesting that the signal is either unstable or volatile.


In a complementary approach, we synthesized various derivatives of 1 and tested them for activation of the *mbo* promoter with the underlying anticipation that the indigenous signal shows the highest bioactivity. The choice of the derivatives was guided by a bioinformatic analysis of the predicted specificity of the MgoA NRPS (fig. S1 and table S5) (20). On this basis, we selected the aliphatic amino acids glycine, alanine, leucine, and isoleucine as starting material. Leudiazin (2) and isoleudiazin (3) were synthesized using a four-step procedure, and glydiazin (4) and aladiazin (5) [also known as nitrosofungin (21)] were synthesized using an alternative route because of the instability of the intermediates (5, Fig. 2A). The five derivatives were tested, and leudiazin (2) was found to be the most potent inducer of the *mbo* promoter (Fig. 2B). No differences in the activities of the (R) and (S) enantiomer of leudiazin were observed (fig. S3). Using a targeted ultra-high-performance liquid

<sup>1</sup>Department of Chemistry, University of Zurich, 8057 Zurich, Switzerland. <sup>2</sup>Department of Plant and Microbial Biology, University of Zurich, 8008 Zurich, Switzerland. <sup>3</sup>Institute of Biology, Leiden University, 2333 BE Leiden, Netherlands. <sup>4</sup>IHSM-UMA-CSIC, Department of Microbiology, University of Málaga, 29071 Málaga, Spain. \*Corresponding author. Email: karl.gademann@uzh.ch (K.G.); leberl@botinst.uzh.ch (L.E.)

†These authors contributed equally to this work.




**Fig. 1. Mitigation of plant pathogenicity.** (A) *P. syringae* plant pathogens represent a global problem as for Pss, which is responsible for BAN caused by the virulence factor mangotoxin that is regulated by an unknown signal (6, 10, 11). (B) Discovery of leudiazan (2) as the unknown signal and developing a mitigation strategy compatible with organic farming procedures. Photo Credit: Anugraha Mathew, University of Zurich.



**Fig. 2. Leudiazan: A novel signal.** (A) Valdiazan derivatives that have been designed, synthesized, and tested. (B) Activity of the *P<sub>mboA</sub>-lacZ* fusion in the presence of different valdiazan derivatives. Error bars represent the SEM, and statistical analysis was performed using one-way analysis of variance (ANOVA) and Tukey's multiple comparisons as posttest; ns, not significant ( $P > 0.05$ ),  $n = 3$  (\*\* $P < 0.01$  and \*\*\* $P < 0.001$ ). (C) Leudiazan (2) detection by UHPLC-MS/MS in the SRM mode. From top to bottom: Pss extract,  $\Delta mgoA$  mutant extract, and synthetic leudiazan ((S)-2).

chromatography–tandem mass spectrometry (UHPLC-MS/MS) approach (163 to 83 Da), we found  $74 \pm 7$  nM leudiazan (2) in the Pss extract (fig. S4). No leudiazan (2) was detected in the extract of the  $\Delta mgoA$  mutant (Fig. 2C).

The biosensor was activated in a dose-dependent manner and  $10 \mu\text{M}$  leudiazan ((S)-2), where the  $\beta$ -galactosidase activity was higher than the level observed for Pss::*P<sub>mboA</sub>-lacZ* (Fig. 3A and fig. S5). Leudiazan's (2) ability to trigger mangotoxin production was evaluated



**Fig. 3. Positive regulation of mangotoxin by leudiazin.** (A) Activities of the *mboA* promoter in the presence of leudiazin (**2**). Error bars represent the SEM, and statistical analysis was performed using one-way ANOVA and Tukey's multiple comparisons as posttest. ns, not significant ( $P > 0.05$ ),  $n = 3$  ( $***P < 0.001$ ). (B) Mangotoxin production assay using *E. coli* JM 105 as an indicator. The halos, indicating mangotoxin production, are highlighted with a dashed line. (C) Tomato leaves were infected with Pss and the  $\Delta mgoA$  mutant in the absence or presence of 10  $\mu M$  leudiazin ((S)-**2**). Pictures were taken after 5 and 10 days postinfection (dpi). Photo Credit: Anugraha Mathew, University of Zurich.

using an *Escherichia coli* (*E. coli*) indicator strain. The antimicrobial activity, which indicates mangotoxin production, of the  $\Delta mgoA$  mutant was restored to the level of the wild-type strain by the addition of 10  $\mu M$  leudiazin ((S)-**2**) but not below this concentration (Fig. 3B and fig. S6). Likewise, 10  $\mu M$  leudiazin ((S)-**2**) was needed to rescue the virulence of the mutant strain to the level of the wild-type in a tomato leaflet infection model (Fig. 3C).

Preliminary results suggested that leudiazin (**2**) could easily evaporate and be present in the gas phase (figs. S7 to S9). To confirm the volatility of **2**, we performed a split plate assay, where one compartment contained the *E. coli* JM 105 (DSM 3949) reporter strain overlaid with the  $\Delta mgoA$  mutant strain while the other compartment contained 20  $\mu M$  leudiazin (**2**, Fig. 4A). After incubation for 24 hours, a halo was observed around the  $\Delta mgoA$  mutant strain, suggesting mangotoxin production by the *mbo* gene cluster. In a control experiment, in which charcoal was added to a third compartment of the petri dish, growth of the *E. coli* JM 105 indicator strain was not inhibited, suggesting that leudiazin (**2**) present in the gas phase was adsorbed by the charcoal (fig. S10). Furthermore, the degradation of leudiazin (**2**) was studied by nuclear magnetic resonance spectroscopy, and the results indicated that the stability of **2** is solvent dependent (figs. S11 and S12).


We hypothesized that the *mgo* signaling system's inhibition could be an effective way to control Pss pathogenicity. A chemical strategy was developed for leudiazin (**2**) degradation and potentially other diazeniumdiolates. In agreement with previous work reporting that  $KMnO_4$  efficiently decomposes the diazeniumdiolate functional group (22), we observed that leudiazin ((S)-**2**) was inactivated by treatment with 300 parts per million (ppm) of  $KMnO_4$  within 10 min

(Fig. 4B and fig. S13). Spraying of infected tomato leaves with an aqueous  $KMnO_4$  solution (1000 ppm) suppressed necrotic symptoms (Fig. 4C). We recovered Pss cells from the treated and the untreated leaves (Fig. 4D), indicating that the chemical treatment abolished toxin production and did not inhibit bacterial growth.

## DISCUSSION

Our study adds leudiazin (**2**) as only the second member of the diazeniumdiolate signal family. Homologs of the leudiazin gene cluster are widespread among *Pseudomonas* strains (tables S1 to S4), indicating that the signal may regulate the production of diverse secondary metabolites in these organisms. While valdiazin (**1**) controls the production of an antifungal agent in the opportunistic human pathogen H111 (18), we show that leudiazin (**2**) controls mangotoxin production in the plant pathogen Pss. The biosynthesis and chemical properties of diazeniumdiolate natural products have been recently investigated in several studies (18–20, 23–26). A remarkable aspect of leudiazin in contrast with other diazeniumdiolates is its volatility and the ability to be transmitted through the air (Fig. 4A). Only very few volatile signals in bacteria have been described to date (27–29), and only two of those were determined to control the virulence in important plant pathogens. Volatile signals may be advantageous for long-distance communication, a feature that may be important for bacteria that live on plant surfaces.

Interference with bacterial signaling has emerged as a valuable strategy for developing novel anti-infective drugs that do not aim at killing the pathogen but attenuate their virulence (30). Our data



**Fig. 4. Volatile signal and novel treatment of BAN disease.** (A) Split plate assay with leudiazin (2) into two compartments and  $\Delta mgoA$  stabbed on two spots over the lawn of *E. coli* JM 105 (DSM 3949) in the third compartment. Halos denote growth inhibition due to mangotoxin production. (B) Leudiazin [(S)-2], 10  $\mu\text{g/ml}$  solution treated with  $\text{KMnO}_4$  solution at concentrations varying from 0 to 500 ppm. UV, ultraviolet. (C) Tomato leaves infected with Pss and analyzed after 5 or 10 dpi and either treated with  $\text{KMnO}_4$  solution or left untreated. (D) Amount of *P. syringae* cells recovered from infected plants at 10 dpi. Error bars represent the SEM, and the statistical analysis was performed using a t test; ns, not significant ( $P > 0.05$ ). Photo Credit: Anugraha Mathew, University of Zurich.

demonstrate that leudiazin (2) can be inactivated by treatment with a dilute  $\text{KMnO}_4$  solution without affecting the bacteria's growth. A common strategy to interfere with bacterial communication is based on the enzymatic degradation of signaling molecules (31, 32). Here, we present the first example of a chemical approach to inactivate a bacterial signaling molecule to attenuate virulence. Notably, the concentration of the  $\text{KMnO}_4$  solution used in our experiments is regulatorily approved for use in organic farming within the European Union (33).

## MATERIALS AND METHODS

### Microbiological and analytical procedures

Strains and plasmids used in this study are listed in table S6. Pss wild-type UMAF0158, Pss  $\Delta mgoA$  mutant as well as strains carrying transcriptional *lacZ* fusions, Pss pMP/*P<sub>mboA</sub>-lacZ*, and  $\Delta mgoA$  pMP/*P<sub>mboA</sub>-lacZ* strains were provided by V.J.C. and F.M.C. The construction of both Pss  $\Delta mgoA$  mutant and the transcriptional fusions has been reported previously (7, 10). *E. coli* strains were routinely grown in LB medium (BD Difco, USA) at 37°C, while *P. syringae* strains were grown at 30°C. For the extraction of leudiazin (2), as well as for promoter activity measurements, *P. syringae* strains were grown in *Pseudomonas* minimal salts (PMS) medium [consists of 1 g of  $(\text{NH}_4)_2\text{H}_2\text{PO}_4$  per liter, 0.2 g of KCl per liter, and 0.2 g of anhydrous  $\text{MgSO}_4$  per liter] (34). The pH was adjusted to 7.0 before autoclaving, and 0.2% glucose was added as a carbon source. Yeast extract (0.005%) was routinely added to the PMS medium to boost growth. When required, media were supplemented with antibiotics

at the following concentrations: kanamycin (100  $\mu\text{g/ml}$ ), gentamicin (20  $\mu\text{g/ml}$ ), tetracycline (20  $\mu\text{g/ml}$ ), and chloramphenicol (20  $\mu\text{g/ml}$ ).

### Assessment of promoter activity by $\beta$ -galactosidase assay

Promoter activity of the transcriptional *lacZ* fusions in liquid cultures was assessed by  $\beta$ -galactosidase assays as described before with minor modifications (35). Briefly, bacterial cells were grown overnight in PMS medium. Where indicated, synthetic valdiazin (1) or leudiazin (2) or ((S)-2) or ((R)-2) and different valdiazin derivatives dissolved in methanol were added to the cultures. Fifty to 200  $\mu\text{l}$  of cells were harvested and resuspended in Z buffer (1 ml), and  $\text{OD}_{600}$  (optical density at 600 nm) values were recorded. To permeabilize the cell membrane, chloroform (25  $\mu\text{l}$ ) and sodium dodecyl sulfate (SDS, 0.1%) were added to the residual bacterial suspension (1 ml), vortexed for 10 s, and incubated at 28°C for 10 min. The reaction was initiated by adding *o*-nitrophenyl- $\beta$ -D-galactoside (200  $\mu\text{l}$ ) solution (4 mg/ml in Z buffer) to each sample, vortexed briefly, and incubated at room temperature (RT). The reaction was stopped by the addition of an aqueous  $\text{Na}_2\text{CO}_3$  solution (500  $\mu\text{l}$ , 1 M). The samples were centrifuged at 16,000 rpm for 10 min, and cell debris-free supernatant (1 ml) was used to measure the absorbance at 420 and 550 nm. The promoter activity (expressed as Miller units) was determined using the following formula:

1 Miller unit =  $1000 \times \frac{\text{OD}_{420} - 1.75 \times \text{OD}_{550}}{t \times v \times \text{OD}_{600}}$ . Where  $t$  = reaction time in minutes and  $v$  = volume of assayed sample in milliliters. Data were based on three independent biological replicates ( $n = 3$ ). For

every  $\beta$ -galactosidase assay, a sample containing only the growth medium was processed as described above and used as a blank.

### Extraction procedure

Bacterial strains were grown in PMS medium containing 0.005% yeast extract for 7 days with agitation (220 rpm) at 30°C. Bacterial cultures were centrifuged for 10 min at 8000 rpm in a Sorvall RC-5C plus centrifuge at 4°C. Culture supernatants were subsequently filtered with a Millipore Express Plus 0.22- $\mu$ m system to remove the cell debris. The cell-free supernatant was acidified to pH 5 with an aqueous HCl solution (10 M) and extracted thrice with 0.5 volumes of dichloromethane. The organic phases were combined, dried with MgSO<sub>4</sub>, filtered, and concentrated under reduced pressure to obtain the crude extract.

### Method for the detection and quantification of leudiazin (2)

Leudiazin (2) was quantified by UHPLC–ESI (electrospray ionization)–MS/MS (Ultimate 3000 LC, Thermo Fisher Scientific; coupled to a TSQ Quantum Ultra, Thermo Fisher Scientific) using a Kinetex EVO C18 column (50 mm by 2.1 mm, 1.7  $\mu$ m) at a flow rate of 0.4 ml/min. The solvent system was composed of A (H<sub>2</sub>O with 0.1% HCO<sub>2</sub>H) and B (MeCN with 0.1% HCO<sub>2</sub>H). After isocratic elution at 5% B for 1 min, the gradient varied from 5% to 60% of B in 2.5 min, 60 to 95% of B in 1 min, and 95 to 100% of B in 0.05 min, and the column was finally washed with 100% B for 1.24 min. The detection was achieved in the selected reaction monitoring (SRM) mode using the specific fragmentation of the protonated molecules [M + H]<sup>+</sup> at mass/charge ratio (*m/z*) 163 into the fragment ion at *m/z* 83 Da at a collision energy of 14 eV. The quantification was done using a calibration curve obtained from analytical standard solutions prepared in MeOH at the following concentrations: 1, 5, 10, 20, 50, and 100  $\mu$ g/ml. A sample volume of 3  $\mu$ l was injected. Cultures of Pss (2 liters) and the  $\Delta$ ngo mutant (2 liters) were prepared following the extraction procedure described above. The crude extract was resuspended in MeOH (1 ml). A concentration of  $24 \pm 3$   $\mu$ g/ml of leudiazin (2) was determined in the Pss extract using a calibration curve (fig. S4), which represents a leudiazin (2) concentration of  $74 \pm 7$  nM in the supernatant of Pss. Leudiazin (2) was not detected in the  $\Delta$ ngo crude extract.

### Volatility experiment

A glass vial (2 ml) containing leudiazin [((S)-2), ~5 mg] was placed into a vacuum chamber at 0.02 mbar, and another sample was placed under a gentle flow of nitrogen at 40°C. The samples were weighted each 2 hours for 8 hours (fig. S7). The headspace experiment was performed by heating leudiazin [((S)-2), 0.9 mg] in a 10-ml vial for 10 min at 100°C (figs. S8 and S9). Then, 10 ml of the gas phase was transferred into another vial containing 1-ml acetonitrile with a gas-tight syringe. This vial was shaken for 30 s, and an aliquot of the acetonitrile layer (3  $\mu$ l/min) was measured with ESI–high-resolution mass spectrometry (ESI–HRMS). ESI–HRMS: *m/z* 161.0930 (C<sub>6</sub>H<sub>13</sub>N<sub>2</sub>O<sub>3</sub><sup>−</sup> [M – H]<sup>−</sup>; calc. 161.0932, −1.2 ppm). A control experiment was performed by injecting a MeCN solution (fig. S9).

### Mangotoxin production assay

Mangotoxin production was determined using an *E. coli* strain as an indicator using a previously described procedure (34) with minor modifications. Briefly, a double layer of the indicator microorganism,

an *E. coli* K12 strain, was prepared, and after solidification, the Pss strains to be tested were stab-inoculated [Pss wild type,  $\Delta$ ngo mutant, and  $\Delta$ ngo mutant complemented with different amounts of leudiazin (2) or ((S)-2) or ((R)-2)] on to the agar seeded with *E. coli*. The plates were incubated at 30°C for 48 hours and observed for inhibition zones around the colony.

*E. coli* killing assay was also performed on split petri dishes to confirm the volatile-mediated effect of leudiazin. PMS medium mixed with *E. coli* [10<sup>7</sup> colony-forming unit (CFU) final concentration] was included in one compartment of the petri dish, and the  $\Delta$ ngoA mutant was stab-inoculated on to it while 10  $\mu$ l of 20  $\mu$ M leudiazin was spotted in the adjacent compartments. A control plate, containing charcoal in a third compartment to trap the volatile leudiazin, was also included in the assay (fig. S10).

### Virulence assay using tomato leaflets as model host

To assess whether the putative signal molecules restored the virulence in the mutant strain, detached tomato leaflets (Hellfrucht Frühstamm variety) were used as model hosts, and the virulence assay was performed as previously described (36) with some modifications. Exponentially growing cultures of Pss in PMS were adjusted to 10<sup>8</sup> CFU ml<sup>−1</sup>. In one set of virulence assay, 10- $\mu$ l drops of the bacterial suspension were injected on three different points on one side of a leaflet, while three 10- $\mu$ l drops of an aqueous MgSO<sub>4</sub> solution (10 mM) were injected on the other side of the same leaflet as a control. Alternatively, in another set of virulence assay, six 10- $\mu$ l drops of the bacterial suspension were injected on six different points on the same leaflet, and detached leaflets inoculated with an aqueous MgSO<sub>4</sub> solution (10 mM) were included in all experiments as a control. Six tomato leaflets were used per strain, and the infected leaflets were maintained at 22°C and a 16:8-hour light:dark photoperiod for a period of 10 days. These experiments were repeated three times. The development and severity of necrosis on the leaflets were determined every 2 days for a period of 10 days. Bacterial strains were retrieved from the infected leaves, and colony counts were calculated on several days after inoculation.

### KMnO<sub>4</sub> treatment of tomato leaves infected with Pss

To test whether KMnO<sub>4</sub> protects the tomato leaves from Pss infection, detached tomato leaves infected with Pss, as described above, were treated with an aqueous KMnO<sub>4</sub> solution. Leaflets (6) infected with Pss were dipped in an aqueous KMnO<sub>4</sub> solution (1000 ppm) with manual shaking at RT, allowing a total contact time of around 20 s. The solution was decanted, and the leaves were then dipped in fresh sterile distilled water to remove any traces of KMnO<sub>4</sub>. The tomato leaflets (6) infected with Pss without KMnO<sub>4</sub> treatment were included in the assay. Both sets of leaflets were maintained for a period of 10 days, and development of necrotic symptoms was recorded. Bacterial strains were recovered from both sets of leaflets after 10 days, and presence of Pss was confirmed by polymerase chain reaction using Pss-specific primers.

### Degradation procedures

Aqueous KMnO<sub>4</sub> solution (500  $\mu$ l at a concentration of 20, 200, 600, or 1000 ppm) were added to 500  $\mu$ l of an aqueous solution of leudiazin [((S)-2), 10  $\mu$ g/ml] to obtain solutions having a concentration of KMnO<sub>4</sub> of 10, 100, 300, and 500 ppm. The samples were stirred at RT for 10 min, filtered, and 3  $\mu$ l injected to the UHPLC–ultraviolet system (254 nm).

## SUPPLEMENTARY MATERIALS

Supplementary material for this article is available at <https://science.org/doi/10.1126/sciadv.abg2293>

## REFERENCES AND NOTES

1. A. Passera, S. Compant, P. Casati, M. G. Maturo, G. Battelli, F. Quaglino, L. Antonielli, D. Salerno, M. Brasca, S. L. Toffolatti, F. Mantegazza, M. Delledonne, B. Mitter, Not just a pathogen? Description of a plant-beneficial *Pseudomonas syringae* strain. *Front. Microbiol.* **10**, 1409 (2019).
2. C. M. J. Pieterse, C. Zamioudis, R. L. Berendsen, D. M. Weller, S. C. M. Van Wees, P. A. H. M. Bakker, Induced systemic resistance by beneficial microbes. *Annu. Rev. Phytopathol.* **52**, 347–375 (2014).
3. J. R. Lamichhane, A. Messéan, C. E. Morris, Insights into epidemiology and control of diseases of annual plants caused by the *Pseudomonas syringae* species complex. *J. Gen. Plant. Pathol.* **81**, 331–350 (2015).
4. X.-F. Xin, B. Kvitko, S. Y. He, *Pseudomonas syringae*: What it takes to be a pathogen. *Nat. Rev. Microbiol.* **16**, 316–328 (2018).
5. E. Arrebola, F. M. Cazorla, V. E. Durán, E. Rivera, F. Olea, J. C. Codina, A. Pérez-García, A. de Vicente, Mangotoxin: A novel antimetabolite toxin produced by *Pseudomonas syringae* inhibiting ornithine/arginine biosynthesis. *Physiol. Mol. Plant Pathol.* **63**, 117–127 (2003).
6. E. Arrebola, F. M. Cazorla, D. Romero, A. Pérez-García, A. de Vicente, A nonribosomal peptide synthetase gene (*mgoA*) of *Pseudomonas syringae* pv. *syringae* is involved in mangotoxin biosynthesis and is required for full virulence. *Mol. Plant Microbe Interact.* **20**, 500–509 (2007).
7. V. J. Carrión, E. Arrebola, F. M. Cazorla, J. Murillo, A. de Vicente, The *mbo* operon is specific and essential for biosynthesis of mangotoxin in *Pseudomonas syringae*. *PLOS ONE* **7**, e36709 (2012).
8. R. I. Sultanov, G. P. Arapidi, S. V. Vinogradova, V. M. Govorun, D. G. Luster, A. N. Ignatov, Comprehensive analysis of draft genomes of two closely related *pseudomonas syringae* phylogroup 2b strains infecting mono- and dicotyledon host plants. *BMC Genomics* **17**, 1010 (2016).
9. S. Marcelletti, M. Scortichini, Comparative genomic analyses of multiple *Pseudomonas* strains infecting *Corylus avellana* trees reveal the occurrence of two genetic clusters with both common and distinctive virulence and fitness traits. *PLOS ONE* **10**, e0131112 (2015).
10. E. Arrebola, V. J. Carrión, F. M. Cazorla, A. Pérez-García, J. Murillo, A. de Vicente, Characterisation of the *mgo* operon in *Pseudomonas syringae* pv. *syringae* UMAF0158 that is required for mangotoxin production. *BMC Microbiol.* **12**, 10 (2012).
11. V. J. Carrión, M. van der Voort, E. Arrebola, J. A. Gutiérrez-Barranquero, A. de Vicente, J. M. Raaijmakers, F. M. Cazorla, Mangotoxin production of *Pseudomonas syringae* pv. *syringae* is regulated by MgoA. *BMC Microbiol.* **14**, 46 (2014).
12. I. Vallet-Gely, O. Opoata, A. Boniface, A. Novikov, B. Lemaître, A secondary metabolite acting as a signalling molecule controls *Pseudomonas entomophila* virulence. *Cell. Microbiol.* **12**, 1666–1679 (2010).
13. S. Mehnaz, J. S. Bauer, H. Gross, Complete genome sequence of the sugar cane endophyte *Pseudomonas aurantiaca* PB-St2, a disease-suppressive bacterium with antifungal activity toward the plant pathogen *Colletotrichum falcatum*. *Genome Announc.* **2**, e01108-13 (2014).
14. M. Van Der Voort, H. J. G. Meijer, Y. Schmidt, J. Watrous, E. Dekkers, R. Mendes, P. C. Dorrestein, H. Gross, J. M. Raaijmakers, Genome mining and metabolic profiling of the rhizosphere bacterium *Pseudomonas* sp. SH-C52 for antimicrobial compounds. *Front. Microbiol.* **6**, 693 (2015).
15. N. Kirchner, C. Cano-Prieto, M. van der Voort, J. M. Raaijmakers, H. Gross, Draft genome sequence of lipopeptide-producing strain *Pseudomonas fluorescens* DSM 11579 and comparative genomics with *Pseudomonas* sp. strain SH-C52, a closely related lipopeptide-producing strain. *Microbiol. Resour. Announc.* **9**, e00304–20 (2020).
16. J. E. Loper, K. A. Hassan, D. V. Mavrodi, E. W. Davis II, C. K. Lim, B. T. Shaffer, L. D. H. Elbourne, V. O. Stockwell, S. L. Hartnett, K. Breakwell, M. D. Henkels, S. G. Tetu, L. I. Rangel, T. A. Kidarsa, N. L. Wilson, J. E. van de Mortel, C. Song, R. Blumhagen, D. Radune, J. B. Hostetler, L. M. Brinkac, A. S. Durkin, D. A. Kluepfel, W. P. Wechter, A. J. Anderson, Y. C. Kim, L. S. Pierson III, E. A. Pierson, S. E. Lindow, D. Y. Kobayashi, J. M. Raaijmakers, D. M. Weller, L. S. Thomashow, A. E. Allen, I. T. Paulsen, Comparative genomics of plant-associated *Pseudomonas* spp.: Insights into diversity and inheritance of traits involved in multitrophic interactions. *PLOS Genet.* **8**, e1002784 (2012).
17. M. De Vrieze, A. R. Varadarajan, K. Schneeberger, A. Bailly, R. P. Rohr, C. H. Ahrens, L. Weisskopf, Linking comparative genomics of nine potato-associated *Pseudomonas* isolates with their differing biocontrol potential against late blight. *Front. Microbiol.* **11**, 857 (2020).
18. C. Jenul, S. Sieber, C. Daepfen, A. Mathew, M. Lardi, G. Pessi, D. Hoepfner, M. Neuburger, A. Linden, K. Gademann, L. Eberl, Biosynthesis of fragin is controlled by a novel quorum sensing signal. *Nat. Commun.* **9**, 1297 (2018).
19. S. Sieber, C. Daepfen, C. Jenul, V. Mannancheril, L. Eberl, K. Gademann, Biosynthesis and structure-activity relationship investigations of the diazeniumdiolate antifungal agent fragin. *ChemBioChem* **21**, 1587–1592 (2020).
20. G. L. Morgan, A. M. Kretsch, K. C. Santa Maria, S. J. Weeks, B. Li, Specificity of nonribosomal peptide synthetases in the biosynthesis of the *Pseudomonas* virulence factor. *Biochemistry* **58**, 5249–5254 (2019).
21. L. A. Dolak, T. M. Castle, B. R. Hannon, A. D. Argoudelis, F. Reusser, Fermentation, isolation, characterization and structure of nitrosfungin. *J. Antibiot.* **36**, 1425–1430 (1983).
22. J. A. Hrabie, L. K. Keefer, Chemistry of the nitric oxide-releasing diazeniumdiolate (“nitrosohydroxylamine”) functional group and its oxygen-substituted derivatives. *Chem. Rev.* **102**, 1135–1154 (2002).
23. R. Hermenau, J. L. Mehl, K. Ishida, B. Dose, S. J. Pidot, T. P. Stinear, C. Hertweck, Genomics-driven discovery of NO-donating diazeniumdiolate siderophores in diverse plant-associated bacteria. *Angew. Chem. Int. Ed.* **58**, 13024–13029 (2019).
24. J. Jiao, J. Du, A. Frediansyah, G. Jahanshah, H. Gross, Structure elucidation and biosynthetic locus of trinickiabin from the plant pathogenic bacterium *Trinickia caryophylli*. *J. Antibiot.* **73**, 28–34 (2020).
25. T. L. Ng, M. E. McCallum, C. R. Zheng, J. X. Wang, K. J. Y. Wu, E. P. Balskus, The L-alanosine gene cluster encodes a pathway for diazeniumdiolate biosynthesis. *ChemBioChem* **21**, 1155–1160 (2020).
26. M. Wang, H. Niikura, H.-Y. He, P. Daniel-lvad, K. S. Ryan, Biosynthesis of the N-N-bond-containing compound L-alanosine. *Angew. Chem. Int. Ed.* **59**, 3881–3885 (2020).
27. K. Kai, H. Ohnishi, M. Shimatani, S. Ishikawa, Y. Mori, A. Kiba, K. Ohnishi, M. Tabuchi, Y. Hikichi, Methyl 3-hydroxymyristate, a diffusible signal mediating *phc* quorum sensing in *Ralstonia solanacearum*. *ChemBioChem* **16**, 2309–2318 (2015).
28. D. Wang, X. Ding, P. N. Rather, Indole can act as an extracellular signal in *Escherichia coli*. *J. Bacteriol.* **183**, 4210–4216 (2001).
29. A. B. Flavier, S. J. Clough, M. A. Schell, T. P. Denny, Identification of 3-hydroxypalmitic acid methyl ester as a novel autoregulator controlling virulence in *Ralstonia solanacearum*. *Mol. Microbiol.* **26**, 251–259 (1997).
30. S. W. Dickey, G. Y. C. Cheung, M. Otto, Different drugs for bad bugs: Antivirulence strategies in the age of antibiotic resistance. *Nat. Rev. Drug Discov.* **16**, 457–471 (2017).
31. B. LaSarre, M. J. Federle, Exploiting quorum sensing to confuse bacterial pathogens. *Microbiol. Mol. Biol. Rev.* **77**, 73–111 (2013).
32. Y.-H. Dong, L.-H. Wang, J.-L. Xu, H.-B. Zhang, X.-F. Zhang, L.-H. Zhang, Quenching quorum-sensing-dependent bacterial infection by an N-acyl homoserine lactonase. *Nature* **411**, 813–817 (2001).
33. Commission Regulation EC No 889/2008, Laying down detailed rules for the implementation of council regulation (EC) no 834/2007 on organic production and labelling of organic products with regard to organic production, labelling and control. *Off. J. Eur. Communities: Legis* **L250**, 1–84 (2008).
34. M. J. Gasson, Indicator technique for antimetabolic toxin production by phytopathogenic species of *Pseudomonas*. *Appl. Environ. Microbiol.* **39**, 25–29 (1980).
35. S. E. Stachel, G. An, C. Flores, E. W. Nester, A Tn3 *lacZ* transposon for the random generation of β-galactosidase gene fusions: Application to the analysis of gene expression in *Agrobacterium*. *EMBO J.* **4**, 891–898 (1985).
36. E. Arrebola, F. M. Cazorla, J. C. Codina, J. A. Gutiérrez-Barranquero, A. Pérez-García, A. de Vicente, Contribution of mangotoxin to the virulence and epiphytic fitness of *Pseudomonas syringae* pv. *syringae*. *Int. Microbiol.* **12**, 87–95 (2009).
37. Y. Takenaka, T. Kiyosu, J.-C. Choi, T. Sakakura, H. Yasuda, Selective synthesis of N-alkyl hydroxylamines by hydrogenation of nitroalkanes using supported palladium catalysts. *ChemSusChem* **3**, 1166–1168 (2010).
38. F. M. Cazorla, J. A. Torés, L. Olalla, A. Pérez-García, J. M. Farré, A. de Vicente, Bacterial apical necrosis of mango in southern Spain: A disease caused by *Pseudomonas syringae* pv. *syringae*. *Phytopathology* **88**, 614–620 (1998).

**Acknowledgments:** We thank K. Agnoli for help with the signal extraction procedures and C. Fabbri and M. Müller for the technical assistance. **Funding:** The research was funded, in part, by the Swiss National Science Foundation (CRSII5\_186410 and 200021\_182043). **Author contributions:** The study was designed by S.S., A.M., L.E., and K.G. Experiments were planned by S.S., A.M., L.E., and K.G. S.S., A.M., C.J., T.K., M.B., U.S., and Y.-C.H. performed experiments. S.S., A.M., C.J., T.K., M.B., V.J.C., F.M.C., U.S., Y.-C.H., L.B., L.E., and K.G. analyzed and discussed data. All authors contributed to the redaction of the manuscript. **Competing interests:** The authors declare that they have no competing interests. **Data and materials availability:** All data needed to evaluate the conclusions in the paper are present in the paper and/or the Supplementary Materials.

Submitted 18 December 2020

Accepted 19 July 2021

Published 8 September 2021

10.1126/sciadv.abg2293

**Citation:** S. Sieber, A. Mathew, C. Jenul, T. Kohler, M. Bär, V. J. Carrión, F. M. Cazorla, U. Stalder, Y.-C. Hsieh, L. Bigler, L. Eberl, K. Gademann, Mitigation of *Pseudomonas syringae* virulence by signal inactivation. *Sci. Adv.* **7**, eabg2293 (2021).

## Mitigation of *Pseudomonas syringae* virulence by signal inactivation

Simon SieberAnugraha MathewChristian JenulTobias KohlerMax BärVíctor J. CarriónFrancisco M. CazorlaUrs StalderYa-Chu HsiehLaurent BiglerLeo EberlKarl Gademann

*Sci. Adv.*, 7 (37), eabg2293. • DOI: 10.1126/sciadv.abg2293

### View the article online

<https://www.science.org/doi/10.1126/sciadv.abg2293>

### Permissions

<https://www.science.org/help/reprints-and-permissions>

Use of this article is subject to the [Terms of service](#)

---

*Science Advances* (ISSN ) is published by the American Association for the Advancement of Science, 1200 New York Avenue NW, Washington, DC 20005. The title *Science Advances* is a registered trademark of AAAS. Copyright © 2021 The Authors, some rights reserved; exclusive licensee American Association for the Advancement of Science. No claim to original U.S. Government Works. Distributed under a Creative Commons Attribution NonCommercial License 4.0 (CC BY-NC).