



Universiteit
Leiden
The Netherlands

Pim1 maintains telomere length in mouse cardiomyocytes by inhibiting TGF beta signalling

Ebeid, D.E.; Khalafalla, F.G.; Broughton, K.M.; Monsanto, M.M.; Esquer, C.Y.; Sacchi, V.; ...
; Sussman, M.A.

Citation

Ebeid, D. E., Khalafalla, F. G., Broughton, K. M., Monsanto, M. M., Esquer, C. Y., Sacchi, V., ... Sussman, M. A. (2021). Pim1 maintains telomere length in mouse cardiomyocytes by inhibiting TGF beta signalling. *Cardiovascular Research*, 117(1), 201-211.
doi:10.1093/cvr/cvaa066

Version: Publisher's Version

License: [Creative Commons CC BY 4.0 license](#)

Downloaded from: <https://hdl.handle.net/1887/3276207>

Note: To cite this publication please use the final published version (if applicable).

Pim1 maintains telomere length in mouse cardiomyocytes by inhibiting TGFβ signalling

David E. Ebeid [†], Farid G. Khalafalla ^{†‡}, Kathleen M. Broughton, Megan M. Monsanto, Carolina Y. Esquer , Veronica Sacchi [¶], Nirmala Hariharan [§], Kelli I. Korski, Maryam Moshref ^{||}, Jacqueline Emathing , Christopher T. Cottage , Pearl J. Quijada [#], Jonathan H. Nguyen, Roberto Alvarez Jr, Mirko Völkers^{**}, Mathias H. Konstandin^{**}, Bingyan J. Wang , Fareheh Firouzi, Julian M. Navarrete , Natalie A. Gude, Marie-Jose Goumans ^{††}, and Mark A. Sussman ^{*}

Department of Biology, San Diego State University, North Life Sciences, 426, 5500 Campanile Drive, San Diego, CA 92182, USA

Received 30 July 2019; accepted 13 March 2019; online publish-ahead-of-print 16 March 2020

Time for primary review: 36 days

Aims

Telomere attrition in cardiomyocytes is associated with decreased contractility, cellular senescence, and up-regulation of proapoptotic transcription factors. Pim1 is a cardioprotective kinase that antagonizes the aging phenotype of cardiomyocytes and delays cellular senescence by maintaining telomere length, but the mechanism remains unknown. Another pathway responsible for regulating telomere length is the transforming growth factor beta (TGFβ) signalling pathway where inhibiting TGFβ signalling maintains telomere length. The relationship between Pim1 and TGFβ has not been explored. This study delineates the mechanism of telomere length regulation by the interplay between Pim1 and components of TGFβ signalling pathways in proliferating A549 cells and post-mitotic cardiomyocytes.

Methods and results

Telomere length was maintained by lentiviral-mediated overexpression of PIM1 and inhibition of TGFβ signalling in A549 cells. Telomere length maintenance was further demonstrated in isolated cardiomyocytes from mice with cardiac-specific overexpression of PIM1 and by pharmacological inhibition of TGFβ signalling. Mechanistically, Pim1 inhibited phosphorylation of Smad2, preventing its translocation into the nucleus and repressing expression of TGFβ pathway genes.

Conclusion

Pim1 maintains telomere lengths in cardiomyocytes by inhibiting phosphorylation of the TGFβ pathway downstream effectors Smad2 and Smad3, which prevents repression of telomerase reverse transcriptase. Findings from this study demonstrate a novel mechanism of telomere length maintenance and provide a potential target for preserving cardiac function.

*Corresponding author. Tel: +1 619 594 2983, E-mail: heartman4ever@icloud.com

[†]The first two authors contributed equally to the study.

[‡]Present address. Department of Pharmaceutical and Biomedical Sciences, College of Pharmacy, California Health Sciences University, Clovis, CA, USA.

[¶]Present address. Regenerative Medicine Department, Genomic Institute of the Novartis Research Foundation, San Diego, CA, USA.

[§]Present address. Department of Pharmacology, University of California, Davis, Davis, CA, USA.


^{||}Present address. Department of Pathology, Microbiology, and Immunology, School of Veterinary Medicine, University of California, Davis, Davis, CA, USA.

[#]Present address. Department of Medicine, Aab Cardiovascular Research Institute, University of Rochester School of Medicine and Dentistry, Rochester, NY, USA.

^{**}Present address. Department of Cardiology, University of Heidelberg, Heidelberg, Germany.

^{††}Present address. Department of Molecular Cell Biology, Leiden University Medical Center, Leiden, Netherlands.

Graphical Abstract



Keywords

Pim1 • Telomere • TGFβ • Cardiomyocyte • Smad2

1. Introduction

Critically short telomeres are associated with cellular senescence and cardiovascular disease.^{1–3} Cardiomyocyte-specific telomere attrition is a characteristic of heart failure in humans and has been associated with decreased contractile ability, hypertrophy, and senescence.^{4,5} In proliferative cells, multiple rounds of cell division promote telomeric shortening, activating DNA damage response that results in cellular senescence or apoptosis.⁶ Cardiomyocytes also exhibit telomere attrition as a result of biomechanical stress, oxidative stress, or inflammation.⁵ Unveiling molecular mechanisms responsible for preserving telomeres in cardiomyocytes could provide valuable insights into antagonizing cellular aging and preserving cardiac function.

Pim1 kinase plays a role in maintaining a youthful cellular phenotype as it regulates a myriad of physiological responses such as anti-apoptotic signalling,⁷ preservation of mitochondrial integrity, enhancing metabolic activity,⁸ and importantly, maintenance of telomere length.^{9,10} Pim1-mediated maintenance of telomeres was established in proliferating cardiac interstitial cells (CICs) derived from murine hearts⁹ or cardiac explants from heart failure patients.¹⁰ Additionally, telomere shortening was observed in response to knockout of Pim1 in mouse cardiomyocytes.¹¹ Although the role of Pim1 in maintaining telomeres has been established, the underlying mechanisms of Pim1-induced telomere maintenance are largely unknown.

Another pathway responsible for regulating telomere length is the transforming growth factor beta (TGF β) signalling pathway, where TGF β activation inhibits telomerase activity.^{12,13} TGF β is a cytokine that mediates a host of cellular responses through stimulation of TGF β receptors and downstream phosphorylation and activation of receptor-associated Smad proteins including Smad2. Activated Smad2 translocates into the nucleus and regulates expression of TGF β pathway target genes responsible for inducing hypertrophic effects on cardiomyocytes and promoting myocardial fibrosis in the failing heart.^{14,15} Although TGF β -mediated regulation of telomeres has been elucidated in proliferating cells,^{16,17} the impact of TGF β signalling inhibition upon telomere maintenance in post-mitotic cells is yet to be explored. Moreover, the intracellular molecular interactions between TGF β signalling and other pathways involved in telomere regulation have not been fully unravelled.

Independent studies showing opposing effects of Pim1 kinase and TGF β signalling upon telomere length point to a potential interaction between Pim1 and TGF β signalling cascades. This study was designed to assess the role of signalling interplay between Pim1 kinase and TGF β in regulating telomere length. Findings from this study demonstrate that both Pim1 overexpression and TGF β signalling inhibition maintains telomere length. Mechanistically, Pim1 maintains telomere length by inhibiting TGF β signalling as revealed by decreased phosphorylation of its downstream effector Smad2. TGF β signalling inhibition by Pim1 was further demonstrated by decreased expression of TGF β pathway target genes *Serpine1* and *FOXA2*. Signalling events were first established in A549 human lung carcinoma cells and recapitulated in post-mitotic adult murine cardiomyocytes, indicating that the Pim1/TGF β signalling axis extends to proliferating and non-proliferating cells. Overall, findings from this article introduce a novel molecular link between Pim1 and TGF β signalling cascades, suggesting a potential mechanism by which Pim1 maintains telomere length.

2. Methods

2.1 Cell culturing

About 200 000 A549 cells (ATCC; Manassas, VA, USA; CCL-185) were plated in 60 mm dishes and were cultured in high glucose Dulbecco's Modified Eagle Medium (ThermoFisher; Waltham, MA, USA; 11965092) supplemented with 10% foetal bovine serum (Omega Scientific; Tarzana, CA, USA; FB-11) and 1% Penicillin–Streptomycin–Glutamine (ThermoFisher; Waltham, MA, USA; 10378016) in a 37°C, 5% CO₂ humidified incubator.

2.2 Viral infections

Lentiviral plasmids were created as previously described.¹⁸ A549 cells were infected with lentivirus encoding enhanced green fluorescent protein (GFP) or human PIM1-GFP. GFP and PIM1 overexpression were confirmed via immunoblotting. Cardiomyocytes were infected with lentivirus harbouring sh-RNA against telomerase reverse transcriptase (TERT) (Lt-sh-TERT) (Dharmacon; Lafayette, CO) or a scramble (Lt-sh-Control).

2.3 Animal experiments

All animal protocols were approved by the Institutional Animal Care and Use Committee of San Diego State University and conform to the *Guide for the Care and Use of Laboratory Animals* published by the US National Institutes of Health. Transgenic mice with cardiac-specific overexpression of human PIM1 (PIM1) and global deletion of Pim1 (Pim1-KO) have been described before.⁷ Non-transgenic (NTg) age- and gender-matched mice of the same strain (FVB) were used as controls. Male adult mice aged 3 months and older were used for the study.

2.4 Isolation and culture of adult mouse cardiomyocytes

Cardiomyocytes were isolated from 3-month-old FVB mice as previously described.¹⁹ Briefly, mice were anaesthetized with ketamine (100 mg/kg) and injected with Heparin (100 U/kg) (Sigma–Aldrich; Burlington, MA, USA; H3393) intraperitoneally to prevent blood clots. After mice displayed no pedal reflex in response to a firm toe pinch, hearts were cannulated via the aorta and digested on a Langendorff apparatus using Collagenase II (230 U/mL) (Worthington; Lakewood, NJ, USA; LS004147). Hearts were finely minced and triturated until no large tissue remained. Following calcium introduction, 100 000 cardiomyocytes were pre-plated on laminin (ThermoFisher; Waltham, MA, USA; 23017015) -coated 35 mm dishes. After 2 h, cardiomyocytes were cultured in serum-free media overnight at 37°C following which pharmacological treatments were conducted.

2.5 Pharmacologic treatments

About 50 000 A549 cells were plated in 35 mm dishes. The following day, cells were treated with 10 μ M LY-364947 (TGFR-In) (Selleck Biochem; Houston, TX, USA; S2805) for 48 h or with 10 ng/mL TGF- β 1 (R&D Systems; Minneapolis, MN, USA; 766-MB) for 1 h. About 100 000 cardiomyocytes were plated in 35 mm dishes and were treated with 10 ng/mL TGF- β 1 10 μ M TGFR-In, 3 mM *N*-acetyl-cysteine (NAC) (Sigma–Aldrich; Burlington, MA; A9165), 200 μ M 6-hydroxy-2,5,7,8-tetramethylchroman-2-carboxylic acid (Trolox) (Sigma–Aldrich; Burlington, MA; 238813), or 25 μ M blebbistatin (Sigma–Aldrich; Burlington, MA; B0560) for 48 h.

2.6 Quantitative RT-PCR

RNA was isolated using Quick-RNA Miniprep Kit (Zymo Research; Irvine, CA, USA; R1054) following the manufacturer's instructions. RNA concentrations were determined using a Nanodrop 2000 spectrophotometer (ThermoFisher; Waltham, MA; ND-2000) and cDNA was synthesized using iScript cDNA synthesis kit (Bio-Rad; Hercules, CA, USA; 1708890). Reactions were prepared in triplicate using 6.5 ng cDNA per reaction and iQ SYBER Green (Bio-Rad; Hercules, CA, USA; 1708880) on a CFX Real-Time PCR thermocycler (Bio-Rad; Hercules, CA, USA; 1855201). Samples were normalized to 18S and data were analysed by $\Delta\Delta C_t$ method. Primers sequences are listed in Table 1.

2.7 Telomere measurement using quantitative RT-PCR

Telomere lengths were measured based on the monochrome multiplex qPCR method established by Cawthon *et al.*²⁰ Genomic DNA was isolated from A549 cells, myocardial tissue, or isolated adult myocytes using the NucleoSpin Tissue kit (Machery-Nagel; Bethlehem, PA, USA; 740952) following the manufacturer's instructions. Reactions were prepared in quintuplicate using 1.67 ng DNA and iQ SYBER Green (Bio-Rad; Hercules, CA, USA; 1708880) on a CFX Real-Time PCR thermocycler (Bio-Rad; Hercules, CA, USA; 1855201). Samples were normalized to albumin and data were analysed by $\Delta\Delta C_t$ method. The telomere length was calculated as $2^{-\Delta\Delta C_t}$ multiplied by the average telomere length for the mouse strain, determined from previous studies to be 75 000 bp for FVB mice.²¹ Primer sequences are listed in Table 1.

2.8 Immunoblot

Cells were lysed in ice-cold radioimmunoprecipitation assay buffer (ThermoFisher; Waltham, MA, USA; 89900) containing protease and phosphatase inhibitor cocktails (Sigma-Aldrich; Burlington, MA, USA; P8340, P5726, and P2850). Protein concentrations were analysed and normalized by Bradford Assay and lysates were prepared by addition of NuPAGE LDS Sample Buffer (ThermoFisher; Waltham, MA, USA; NP0007). Samples were sonicated and boiled then loaded onto a 4–12% NuPAGE Bis-Tris gel (ThermoFisher; Waltham, MA, USA; NP0321BOX). Proteins were transferred onto an Immobilon-FL Polyvinylidene difluoride membrane (EMD Millipore; Burlington, MA, USA; IPFL0010) and the

membrane was blocked with Odyssey Blocking Buffer (Li-Cor; Lincoln, NE, USA; 927-50000) for 1 h at room temperature and incubated with primary antibodies overnight at 4°C (Table 2). Secondary antibodies were applied for 2 h at room temperature. Fluorescent signal was detected using an Odyssey CLx imaging system (Li-Cor; Lincoln, NE, USA) and bands were quantified using Image Studio.

2.9 Immunofluorescence

Immunohistochemistry was performed on paraffin heart sections as previously described.^{22,23} Briefly, slides were deparaffinized in three changes of xylene for 5 min, followed by three changes of 100% ethanol for 3 min, two changes of 95% ethanol for 3 min, 70% ethanol for 3 min, then rehydrated in three changes of de-ionized water for 3 min. The sections were boiled in 10 mmol/L citrate buffer (pH 6.0) for 15 min then were blocked with Tris–NaCl-Blocking buffer (TNB) (Perkin Elmer; Waltham, MA, USA; FP1020) for 30 min. Primary antibody (Table 2) was applied overnight at 4°C in TNB. The following day, slides were washed with three changes of Tris–NaCl (1XTN), stained with secondary antibody for 2 h at room temperature, washed in three changes of 1XTN for 5 min, and then stained with 1 µg/mL 4',6-diamidino-2-phenylindole (DAPI; Sigma-Aldrich; Burlington, MA, USA; D8417). Slides were mounted with Vectashield (Vector Laboratories; Burlingame, CA, USA; H-1000), following which images were acquired on an SP8 confocal microscope (Leica; Buffalo Grove, IL, USA).

2.10 Quantitative-fluorescence in situ hybridization

Telomere lengths were measured in heart samples by quantitative-fluorescence in situ hybridization (qFISH) using a modified version of a previously published protocol.²⁴ Paraffin sections were subjected to deparaffinization and antigen retrieval as described previously.²³ The hybridization probe was warmed to 37°C for 5 min and the hybridization buffer (20 mM Tris at pH 7.4, 60% formamide, 0.1 µg/mL salmon sperm DNA) (Sigma-Aldrich; Burlington, MA, USA; D1626) was heated to 85°C for 5 min. The telomere probe (PNA Bio; Newbury Park, CA, USA; F1002) was mixed with hybridization buffer at a ratio of 1:12.5 by volume and heated to 85°C for 10 min. Simultaneously, the dried slides were preheated at 85°C for 5 min. Following application of the mix

Table 1 Primer list

Target	Forward	Reverse
Human <i>SERPINE1</i>	GCAACGTGGTTTTCTCACCC	GGCCATGCCCTTGTCATCAA
Human <i>N-CADHERIN</i>	ATTGATGCTGACGATCCCAATGCC	TCAAGTCCAGCTGCCACTGTGATCA
Human <i>18S</i>	CGAGCCGCCTGGATACC	CATGGCCTCAGTTCGGAAAA
Mouse <i>Serpine1</i>	TTCAGCCCTTGCTTGCCCTC	ACACTTTTACTCCGAAGTCGGT
Mouse <i>Foxa2</i>	CCCTACGCCAACATGAACTCG	GTTCTGCCGGTAGAAAGGGA
Mouse <i>Pim1</i>	ATCCGCGTCGCCGACAACTT	TCGGGTGCCATTGGGCAGTT
Mouse <i>18S</i>	CGAGCCGCCTGGATACC	CATGGCCTCAGTTCGGAAAA
Telomere	ACACTAAGGTTTGGGTTTGGGTTTGGGTTTGGGTTAGTGT	TGTTAGGTATCCCTATCCCTATCCCTATCCCTATCCCT AACA
Human <i>ALB</i>	CGGCGGCGGGCGGCGGGCTGGGCGGAAATGCTGCA CAGAATCCTTG	GCCCGGCCGCGCGCCCGTCCCGCCGAAAAGCATGG TCGCCTGTT
Mouse <i>Alb</i>	CGGCGGCGGGCGGCGGGCTGGGCGGTTGCATGAA GTTGCCAGAAG	GCCCGGCCGCGCGCCCGTCCCGCCGGTCAGGCAGCT TTCCTTGTC

Table 2 Antibody list

Antibody	Catalogue number	Dilution	Application
Smad2	Cell Signalling 5339	1:500	IB
Phospho-Smad2 (Ser465/467)	Cell Signalling 3108	1:500	IB
Smad2/3	Cell Signalling 3102	1:500	IB
Phospho-Smad3 (Ser423/425)	Abcam Ab52903	1:500	IB
Pim-1	ThermoFisher 39-4600	1:1000	IB
GFP	Invitrogen G10362	1:1000	IB
GFP	Rockland 600-101-215	1:100	IHC
TGF β	R&D Systems MAB240	1:500	IB
TGF β	Cell Signalling 3709	1:50	IHC
TERT	ThermoFisher PA5-80105	1:500	IB
Cardiac troponin T	ThermoFisher MA5-12960	1:100	IHC
GAPDH	Millipore AB2302	1:5000	IB
Myosin light chain 2	Santa Cruz Sc-34490	1:20	IHC

containing the telomere probe and hybridization buffer, slides were heated at 85°C for 10 min and subsequently incubated overnight at 37°C in a covered, humidified chamber. The next day, the slides were washed once with 37°C 2 \times saline-sodium citrate (2XSSC) (Sigma–Aldrich; Burlington, MA, USA; S6639), twice with 37°C 2XSSC containing 50% formamide for 5 min each, and twice with 37°C 2XSSC for 5 min each. Slides were then blocked with TNB (Perkin Elmer; Waltham, MA, USA; FP1020) for 1 h at room temperature, and incubated with primary antibody overnight at 4°C. The next day, slides were washed, incubated with secondary antibody for 1 h at room temperature, washed, stained with 1 μ g/mL DAPI (Sigma–Aldrich; Burlington, MA, USA; D8417), then mounted, and imaged.

To measure telomere length *in vitro*, slides of cultured adult cardiomyocytes were treated with fixative containing 3:1 ratio of ethanol: acetic acid for 30 min. Slides were treated with RNase solution (2XSSC containing 100 μ g/mL ribonuclease A) and incubated at 37°C for 1 h then washed with 2XSSC and phosphate-buffered saline for 5 min each. Slides were then treated with 37°C 0.005% pepsin (Sigma–Aldrich; Burlington, MA, USA; P7000) for 3 min and washed with phosphate-buffered saline for 5 min. The slides were dehydrated using cold 70%, 90%, and 100% ethanol for 2 min each then air dried. The PNA probe was warmed to 37°C for 5 min and mixed with hybridization buffer at a ratio of 1:12.5 by volume and heated to 85°C for 10 min. Simultaneously, the dried slides were preheated at 85°C for 5 min. Following application of the mix containing the telomere probe and hybridization buffer, slides were heated at 85°C for 5–10 min and subsequently incubated overnight at 37°C in a covered, humidified chamber. The next day, the slides were washed once with 37°C 2XSSC for 5 min, twice with 37°C 2XSSC containing 50% formamide for 5 min each, and twice with warm 2XSSC for 5 min each. Following addition of 1 μ g/mL DAPI, slides were mounted and cells were imaged.

2.11 Telomere length measurements using qFISH

qFISH images were acquired on the Leica SP8 confocal microscope. Mean values provided by the Leica software represent the pixel sum/sum processed pixel that accounts for the area of the region of interest, therefore, no additional normalizations to the nuclear area were done.

2.12 Statistical analyses

Statistical analyses were performed using paired or unpaired Student's *t*-test or one-way ANOVA with Tukey's multiple comparison test on GraphPad Prism v5.0. A *P*-value of <0.05 was considered statistically significant.

3. Results

3.1 Telomere length is elongated by PIM1 overexpression or TGF β receptor inhibition

Previous studies demonstrate that telomere length is maintained by PIM1 overexpression in mouse cardiac non-myocyte cells,⁹ and that telomeric shortening is associated with TGF β signalling.^{13,16,17,25} To recapitulate these results, A549 cells were used because these cells express low levels of endogenous PIM1. GFP or GFP-PIM1 (PIM1) was overexpressed via lentiviral infection in A549 cells and overexpression was confirmed via immunoblot (*Figure 1A*). PIM1 cells contained 1.55-fold longer telomeres than GFP cells after three passages as revealed by quantitative RT–PCR (*Figure 1B*). To determine basal level of telomere length, telomeres were measured in A549 cells before and after three passages in culture. There was no significant change in telomere length over three passages in culture suggesting that telomere length is already in equilibrium in A549 cells (see [Supplementary material](#) online, *Figure S1A*). Telomeres are dynamic in length as they are progressively eroded during rounds of cell division and elongated by telomerase.²⁶ PIM1 influences the equilibrium of telomeres in A549 cells, shifting it towards elongation. The role of TGF β signalling on telomere length was then evaluated by pharmacologic inhibition of TGF β receptor-1 (TGFR1) using LY-364947 (TGFR-In), a selective, ATP-competitive inhibitor of TGFR1. Inhibition of TGFR1 resulted in 1.22-fold longer telomeres compared to the vehicle control (*Figure 1C*). Collectively, these results demonstrate that telomeres can be elongated by either PIM1 overexpression or TGF β signalling inhibition in A549 cells.

3.2 Tgf β signalling is blunted by PIM1

The link between PIM1 and TGF β signalling was established by examining the TGF β pathway downstream targets, pSMAD2, *SERPINE1*,

and *N-CADHERIN* expression following treatment of 10 ng/mL TGF β 1 on GFP and PIM1 cells. In the presence of TGF β 1 agonist, PIM1 exhibited a 41% reduction in pSMAD2 levels compared to GFP as evaluated by immunoblotting (Figure 2A and B). Inhibitory effects of PIM1 on TGF β signalling were confirmed by expression of canonical downstream target *SERPINE1* and non-canonical target *N-CADHERIN* as measured via qRT-PCR. PIM1 displayed a 64% and 42% reduction in mRNA expression of *SERPINE1* and *N-CADHERIN*, respectively, compared to GFP (Figure 2C and D). These findings indicate that TGF β signalling is inhibited by PIM1 overexpression, suggesting a potential mechanism by which telomere length is maintained by PIM1 in proliferating A549 cells.



3.3 Telomere length is maintained by PIM1 in adult murine cardiomyocytes

Pim1 exerts cardioprotective effects in mouse myocardium.⁷ To determine if telomere length is maintained by Pim1 in cardiomyocytes, telomeres were measured in hearts of 3-month-old non-transgenic mice (NTg), transgenic mice with cardiac-specific overexpression of human PIM1 (PIM1),²⁷ and knockout mice with global deletion of Pim1 (Pim1-KO).⁷ Telomeres were 27% longer in PIM1 myocardium and 31% shorter in Pim1-KO hearts compared to NTg as revealed by qRT-PCR (Figure 3A). Telomere lengths were then analysed in cardiomyocytes from NTg, PIM1, and Pim1-KO heart sections via qFISH (Figure 3C). Consistent with findings from whole heart analysis, telomere fluorescence intensity was 28% greater in PIM1 cardiomyocytes and reduced by 30% in Pim1-KO cardiomyocytes compared to NTg controls (Figure 3B). Collectively, these results indicate that telomere length is maintained by PIM1 in adult mouse cardiomyocytes.

3.4 Telomere length is maintained by TGF β inhibition in cardiomyocytes

The link between TGF β signalling and telomere length maintenance has not been established in cardiomyocytes. Therefore, telomere lengths were measured by qRT-PCR in isolated adult NTg mouse cardiomyocytes following treatment with TGF β or TGFR-In for 48 h. Telomeres were 1.87-fold longer following treatment with TGFR-In, whereas no significant change to telomere length was observed with TGF β treatment compared to the vehicle control (Figure 4A).

Furthermore, treatment of Pim1-KO with TGFR-In rescued the telomere shortening observed (see Supplementary material online, Figure S2A). Addition of TGFR-In to PIM1 cardiomyocytes did not further maintain the telomeres, suggesting that PIM1 and TGFR-In act via similar or concurrent pathways to maintain the telomeres in cardiomyocytes (see Supplementary material online, Figure S2B). Telomeres rapidly eroded in isolated cardiomyocytes over 48 h in culture (see Supplementary material online, Figure S2A). Treatment with the antioxidant Trolox maintained telomere length in culture, whereas treatment with blebbistatin had no effect on telomere length, suggesting that telomere attrition in culture was due to oxidative stress and not biomechanical stress (see Supplementary material online, Figure S2C). To further confirm the effect of TGF β signalling on telomere length, qFISH analysis of telomeres was performed on isolated myocytes







Figure 3 Telomere length is maintained by PIM1 in adult murine cardiomyocytes. (A) Analysis of telomere lengths in isolated adult cardiomyocytes from PIM1 or Pim1-KO mice via qRT-PCR. $n = 4$, error bars represent SEM, $*P < 0.05$, $**P < 0.01$ vs. NTg, $***P < 0.001$ vs. PIM1 as measured by one-way ANOVA followed by Tukey's multiple comparison test. (B) Quantification of telomere fluorescence intensity indicated as mean fluorescence values from qFISH images. $n = 32$ nuclei per group, error bars represent SEM, $***P < 0.001$ vs. NTg and vs. PIM1 as measured by one-way ANOVA followed by Tukey's multiple comparison test. (C) Representative qFISH images of telomeres (white) from NTg, PIM1, and Pim1-KO mouse heart sections co-stained with myosin light chain 2 (MYLC2, red) and DAPI (blue). Scale bar represents 25 μm .

following treatment with TGF β or TGF β -In. Telomere fluorescence intensity was 3.84-fold greater following treatment with TGF β -In compared to the vehicle control (Figure 4B and C). Endogenous TGF β expression was measured to account for the lack of change in telomere length in TGF β -treated cells. Indeed, endogenous TGF β expression was detected in cardiomyocytes as determined by immunoblot (Figure 4D). TGF β is synthesized as a latent precursor protein that cannot interact with its receptor until it is cleaved to yield the active form.²⁸ There was no significant difference in latent and active TGF β expression between NTg and PIM1 cardiomyocytes (Figure 4E and F). Furthermore, TGF β expression was evaluated in NTg and PIM1 hearts 7-day post-myocardial infarction. Conditions of stress up-regulated TGF β expression in both NTg and PIM1, however, there was no significant difference in TGF β expression between NTg and PIM1 indicating that PIM1 does not directly affect ligand expression (see Supplementary material online, Figure S3B and C). Likewise, treatment of NTg cardiomyocytes with TGF β had no effect on Pim1 expression suggesting that Pim1 and TGF β do not exhibit feedback relationships (see Supplementary material online, Figure S3C). Overall, these results demonstrate that telomere length is maintained by inhibition of the TGF β signalling pathway in cardiomyocytes.

3.5 Expression of TGF β downstream targets is reduced by PIM1 in cardiomyocytes

To determine whether TGF β signalling inhibition is mediated by PIM1 in cardiomyocytes as revealed in A549 cells, phosphorylation of the downstream effector Smad2 was evaluated in PIM1 cardiomyocytes by immunoblot analysis. Cardiac-specific overexpression of PIM1 resulted in a 39% reduction in pSmad2 levels compared to NTg (Figure 5A and B). Pim1-KO cardiomyocytes also displayed an up-regulation of pSmad2 compared to PIM1 (see Supplementary material online, Figure S4A and B). However, there was no significant difference in pSmad2/Smad2 between the NTg and Pim1-KO (see Supplementary material online, Figure S4A and B). Adult cardiomyocytes already have very low basal levels of Pim1 which could account for the lack of difference observed between the NTg and Pim1-KO. Immunoblot analysis of pSmad3 also revealed a reduction of TGF β signalling in PIM1 compared to NTg and Pim1-KO (see Supplementary material online, Figure S4C and D). Additionally, PIM1 cardiomyocytes showed significant down-regulation of TGF β pathway target genes *Serpine1* and *FoxA2* (Figure 5C and D). Taken together, these results demonstrate the role of PIM1 in maintaining telomeres in cardiomyocytes through inhibition of the canonical TGF β signalling pathway.



3.6 Pim1-mediated maintenance of telomere length is TERT dependent

Upon TGFβ stimulation, Smad2/3 is phosphorylated and translocated into the nucleus where it binds to the TERT gene promoter, repressing transcriptional activation of TERT.¹³ To determine if TERT is involved in PIM1-mediated telomere maintenance in cardiomyocytes, TERT protein expression was analysed in PIM1 and Pim1-KO cardiomyocytes. Expression of TERT was coincident with telomere length such that PIM1 mice had increased TERT levels compared to NTg (Figure 6A and B). Conversely Pim1-KO mice had significant reduction in TERT protein expression relative to NTg (Figure 6A and B). The mechanism underlying PIM1-mediated telomere maintenance was further assessed by knocking down TERT expression in cardiomyocytes isolated from NTg and PIM1 mice. Knocking down TERT blunted PIM1-mediated increases in telomere length indicating that TERT is a critical intermediate in PIM1-mediated telomere maintenance in cardiomyocytes (Figure 6C).

4. Discussion

Telomere-associated cellular senescence of cardiomyocytes remains a hallmark of heart failure that contributes to long-term deterioration of the heart.⁴ Telomere attrition in mouse cardiomyocytes resulted in

increased cardiomyocyte apoptosis, cellular hypertrophy, up-regulation of the senescence marker, p16, and increased levels of the proapoptotic transcription factor, p53.^{5,29} Functionally, hearts with critically short telomeres in cardiomyocytes displayed cardiac dysfunction characterized by increased end diastolic left ventricular pressure and decreased contractility and relaxation of the left ventricle.³⁰ Delineating mechanisms of telomere length maintenance are key to antagonizing telomere-associated cellular aging and preserving cardiac function. Herein, we establish a novel link between two major signalling molecules that regulate telomere length. PIM1 inhibits phosphorylation of Smad2, the downstream effector of TGFβ signalling, which in turn maintains telomere length *in vitro*. These findings provide novel insights into PIM1-mediated mechanisms in maintaining telomere length and antagonizing cardiac cellular aging.



Previous reports have demonstrated PIM1-mediated telomere maintenance in CICs depends upon TERT activity.⁹ CICs overexpressing PIM1 showed no increase in telomere length following treatment with a TERT inhibitor, suggesting that PIM1 maintains telomeres through a TERT mediated pathway.⁹ Telomeres are maintained by telomerase enzyme that comprises telomerase RNA template (TER), and the protein component, TERT.¹³ TGFβ signalling has been shown to negatively regulate telomerase activity by repressing expression of the TERT gene.¹³ In the presence of TGFβ, Smad2/3 is phosphorylated and translocated into

the nucleus where it binds to the TERT gene promoter, repressing transcriptional activation of TERT.¹³ Inhibition of TGFβ signalling is likely influencing telomere length in a similar manner in cardiomyocytes. Findings from this article suggest PIM1-mediated telomere maintenance through inhibition of TGFβ signalling potentially occurs in a TERT-dependent pathway.

Collectively, findings from this study along with previously published reports emphasize the crucial role of Pim1 kinase in reversing aging phenotypes in different cardiac cell populations and antagonizing myocardial aging by preserving telomeres. Additionally, Pim1-mediated inhibition of TGFβ signalling further preserves cardiac function. Endogenous TGFβ

signalling in the heart contributes to the pathogenesis of cardiac fibrotic and hypertrophic re-modelling in the pressure-overloaded heart.¹⁵ TGFβ signalling drives pathological hypertrophy through suppression of antioxidant genes *Mt1/2*.³¹ PIM1-mediated inhibition of the TGFβ signalling pathway in cardiomyocytes (Figure 5) suggests a plausible mechanism by which Pim1 inhibits pathological re-modelling and preserves cardiac function with age and post-injury. On the other hand, a recent report documented the involvement of PIM1 in Smad2/3 activation in clear-cell renal-cell carcinoma,³² indicating differential regulation of Smad2/3 by Pim1 in different cell types. PIM1 has been demonstrated to play a dual role in cancers, acting as either an activator or a suppressor depending on the type of cancer.³² PIM1 is overexpressed in breast cancer, mesothelioma, glioblastoma, and clear-cell renal-cell carcinoma and contributes to increased Smad2 phosphorylation and EMT.³² Conversely, PIM1 is down-regulated in other cancers such as pancreatic ductal carcinoma and non-small cell lung cancer.^{32–34} Therefore, the interaction between PIM1 and Smad2 is highly dependent on the cell type and the associated signalling pathways. In this study A549 cells, a non-small cell lung cancer line, were utilized because these cells have very low basal expression of PIM1, similar to adult cardiomyocytes.

While the present study introduces a novel mechanistic link between Pim1 and TGFβ signalling pathways in telomere regulation, the mechanism in the signalling cascade remain to be elucidated. Pim1 is a serine/threonine kinase that phosphorylates a multitude of targets including Bad, p21, and Cdc25A.^{35–37} However, findings demonstrate that overexpression of Pim1 results in decreased levels of phosphorylated Smad2 (Figures 2 and 5). The antagonistic relationship between Pim1 and Smad phosphorylation supports previous reports demonstrating a repressive effect of Akt upon the TGFβ pathway. The mechanism of Pim1-mediated inhibition of SMAD2 phosphorylation has not been identified, but it hypothesized that Pim1 acts in a similar manner as Akt. Akt has been shown to inhibit TGFβ signalling through direct interaction with Smad3.^{38,39} Co-immunoprecipitation assays show that Akt physically binds unphosphorylated Smad3 to sequester it outside the nucleus and to prevent its phosphorylation.^{38,39} This interaction is constitutive but is inhibited by treatment with TGFβ.^{38,39} Furthermore, inhibition of Smad3 phosphorylation by Akt occurs in a kinase-independent manner. A catalytically inactive Akt mutant is effective at suppressing TGFβ-mediated



apoptosis.³⁹ Pim1 operates downstream of Akt and mediates the cardioprotective signalling of Akt.⁷ Akt is a serine-threonine kinase involved in cell proliferation and survival.⁷ Nuclear accumulation of Akt induces Pim1 expression, mediating the cardioprotective effects of Akt.⁴⁰ Ablation of Pim1 induces Akt expression and activation, but the increase in Akt activation does not mediate enhanced recovery or reduced apoptosis following infarction, indicating that the cardioprotective effects of Akt depend on Pim1.⁷ Future studies could assess whether Pim1-induced Smad inhibition is a consequence of physical interaction and sequestration of Smad proteins. Additionally, identifying downstream effectors of Pim1/TGF β signalling in telomere regulation remains a topic for future investigations.

In summary, findings from this article demonstrate a novel mechanism by which PIM1-mediated inhibition of the canonical TGF β signalling pathway contributes to telomere maintenance, consistent with cumulative evidence of Pim1-induced reversal of aging phenotypes, particularly in myocardial aging. Preservation of a youthful cellular phenotype is key to blunting damage in the injured myocardium and preserving cardiac function. Profound mechanistic understanding of telomere maintenance could provide valuable information towards developing novel therapeutic strategies for heart failure as well as a variety of age-related disorders.

Supplementary material

Supplementary material is available at *Cardiovascular Research* online.

Authors' contributions

M.S., N.G., and M.G. designed and directed the project. D.E., F.K., K.M., M.M., C.E., V.S., N.H., K.K., M.M., J.E., C.C., P.Q., J.N., R.A., M.V., M.K., B.W., F.F., J.N., performed the experiments. D.E., F.K., K.B., M.M., V.S., N.H., K.K., M.M., J.E., C.C., P.Q., J.N., R.A., M.V., M.K., B.W., F.F., N.G., and M.S. analysed and interpreted the data. D.E. and F.K. wrote the paper with input from all the authors.

Acknowledgements

The authors thank all former and current members of the Sussman lab for their critical feedback and review of the article.

Conflict of interest: M.S. is Chief Scientific Officer and a founding member of CardioCreate, Inc. N.H. is an employee of Roche Molecular Systems, Inc., where her work is unrelated to the content of this article.

Funding

This work was supported by National Institutes of Health (4R37HL091102, 5R01HL105759, 5R01HL067245, 1R01HL113647, 1R01HL117163, and 2P01HL085577 to M.A.S.), American Heart Association Scientist Development Grant (16SDG30970046 to N.H.), and Foundation Leducq Transatlantic Network Consortium.

References

- Lansdorf PM. Telomeres and disease. *EMBO J* 2009;**28**:2532–2540.
- Blasco MA. Telomeres and human disease: ageing, cancer and beyond. *Nat Rev Genet* 2005;**6**:611–622.
- Yeh J-K, Wang C-Y. Telomeres and telomerase in cardiovascular diseases. *Genes* 2016;**7**:58.

- Sharifi-Sanjani M, Oyster NM, Tichy ED, Bedi KC, Harel O, Margulies KB, Mourkioti F. Cardiomyocyte-specific telomere shortening is a distinct signature of heart failure in humans. *J Am Heart Assoc* 2017;**6**:e005086.
- Booth SA, Charchar FJ. Cardiac telomere length in heart development, function, and disease. *Physiol Genomics* 2017;**49**:368–384.
- Armanios M. Telomeres and age-related disease: how telomere biology informs clinical paradigms. *J Clin Invest* 2013;**123**:996–1002.
- Muraski JA, Rota M, Misao Y, Fransioli J, Cottage C, Gude N, Esposito G, Delucchi F, Arcarese M, Alvarez R, Siddiqi S, Emmanuel GN, Wu W, Fischer K, Martindale JJ, Glembotski CC, Leri A, Kajstura J, Magnuson N, Berns A, Beretta RM, Houser SR, Schaefer EM, Anversa P, Sussman MA. Pim-1 regulates cardiomyocyte survival downstream of Akt. *Nat Med* 2007;**13**:1467–1475.
- Borillo GA, Mason M, Quijada P, Völkers M, Cottage C, McGregor M, Din S, Fischer K, Gude N, Avitabile D, Barlow S, Alvarez R, Truffa S, Whittaker R, Glassy MS, Gustafsson AB, Miyamoto S, Glembotski CC, Gottlieb RA, Brown JH, Sussman MA. Pim-1 kinase protects mitochondrial integrity in cardiomyocytes. *Circ Res* 2010;**106**:1265–1274.
- Cottage CT, Neidig L, Sundararaman B, Din S, Joyo AY, Bailey B, Gude N, Hariharan N, Sussman MA. Increased mitotic rate coincident with transient telomere lengthening resulting from Pim-1 overexpression in cardiac progenitor cells. *Stem Cells* 2012;**30**:2512–2522.
- Mohsin S, Khan M, Nguyen J, Alkatib M, Siddiqi S, Hariharan N, Wallach K, Monsanto M, Gude N, Dembitsky W, Sussman MA. Rejuvenation of human cardiac progenitor cells with Pim-1 kinase. *Circ Res* 2013;**113**:1169–1179.
- Din S, Konstandin MH, Johnson B, Emathingier J, Völkers M, Toko H, Collins B, Ormachea L, Samse K, Kubli DA, La Torre AD, Kraft AS, Gustafsson AB, Kelly DP, Sussman MA. Metabolic dysfunction consistent with premature aging results from deletion of Pim kinases. *Circ Res* 2014;**115**:376–387.
- Yang H, Kyo S, Takatura M, Sun L. Autocrine transforming growth factor beta suppresses telomerase activity and transcription of human telomerase reverse transcriptase in human cancer cells. *Cell Growth Differ* 2001;**12**:119–127.
- Li H, Xu D, Li J, Berndt MC, Liu JP. Transforming growth factor β suppresses human telomerase reverse transcriptase (hTERT) by Smad3 interactions with c-Myc and the hTERT gene. *J Biol Chem* 2006;**281**:25588–25600.
- Shi Y, Massagué J. Mechanisms of TGF- β signaling from cell membrane to the nucleus. *Cell* 2003;**113**:685–700.
- Dobaczewski M, Chen W, Frangogiannis NG. Transforming growth factor (TGF)- β signaling in cardiac remodeling. *J Mol Cell Cardiol* 2011;**51**:600–606.
- Cassar L, Nicholls C, Pinto AR, Chen R, Wang L, Li H, Liu JP. TGF-beta receptor mediated telomerase inhibition, telomere shortening and breast cancer cell senescence. *Protein Cell* 2017;**8**:39–54.
- Lacerte A, Korah J, Roy M, Yang XJ, Lemay S, Lebrun JJ. Transforming growth factor- β inhibits telomerase through SMAD3 and E2F transcription factors. *Cell Signal* 2008;**20**:50–59.
- Fischer KM, Cottage CT, Wu W, Din S, Gude NA, Avitabile D, Quijada P, Collins BL, Fransioli J, Sussman MA. Enhancement of myocardial regeneration through genetic engineering of cardiac progenitor cells expressing pim-1 kinase. *Circulation* 2009;**120**:2077–2087.
- Kubli DA, Cortez MQ, Moyzis AG, Najor RH, Lee Y, Gustafsson ÅB. PINK1 is dispensable for mitochondrial recruitment of parkin and activation of mitophagy in cardiac myocytes. *PLoS One* 2015;**10**:e0130707.
- Cawthon RM. Telomere length measurement by a novel monochrome multiplex quantitative PCR method. *Nucleic Acids Res* 2009;**37**.
- Hemann MT. Wild-derived inbred mouse strains have short telomeres. *Nucleic Acids Res* 2000;**28**:4474–4478.
- Hariharan N, Quijada P, Mohsin S, Joyo A, Samse K, Monsanto M, La Torre AD, Avitabile D, Ormachea L, McGregor MJ, Tsai EJ, Sussman MA. Nucleostemin rejuvenates cardiac progenitor cells and antagonizes myocardial aging. *J Am Coll Cardiol* 2015;**65**:133–147.
- Mohsin S, Khan M, Toko H, Bailey B, Cottage CT, Wallach K, Nag D, Lee A, Siddiqi S, Lan F, Fischer KM, Gude N, Quijada P, Avitabile D, Truffa S, Collins B, Dembitsky W, Wu JC, Sussman MA. Human cardiac progenitor cells engineered with Pim-1 kinase enhance myocardial repair. *J Am Coll Cardiol* 2012;**60**:1278–1287.
- Zhao M-M, DePinho RA, Kost-Alimova M, Mourkioti F, Kraft P, Bernstein D, Protopopov A, Meeker AK, Blau HM, Day JW, Kustan J. Role of telomere dysfunction in cardiac failure in Duchenne muscular dystrophy. *Nat Cell Biol* 2013;**15**:895–904.
- Katakura Y, Nakata E, Miura T, Shirahata S. Transforming growth factor β triggers two independent-senescence programs in cancer cells. *Biochem Biophys Res Commun* 1999;**255**:110–115.
- Bernal A, Tusell L. Telomeres: implications for cancer development. *Int J Mol Sci* 2018;**19**:294.
- Muraski JA, Fischer KM, Wu W, Cottage CT, Quijada P, Mason M, Din S, Gude N, Alvarez R, Rota M, Kajstura J, Wang Z, Schaefer E, Chen X, MacDonnel S, Magnuson N, Houser SR, Anversa P, Sussman MA. Pim-1 kinase antagonizes aspects of myocardial hypertrophy and compensation to pathological pressure overload. *Proc Natl Acad Sci U S A* 2008;**105**:13889–13894.
- Khalil N. TGF- β : from latent to active. *Microbes Infect* 1999;**1**:1255–1263.
- Leri A, Franco S, Zacheo A, Barlucchi L, Chimenti S, Limana F, Nadal-Ginard B, Kajstura J, Anversa P, Blasco MA. Ablation of telomerase and telomere loss leads to

- cardiac dilatation and heart failure associated with p53 upregulation. *EMBO J* 2003; **22**:131–139.
30. Wong LSM, Oeseburg H, Boer RD, Gilst WV, Veldhuisen DV, Harst PVD. Telomere biology in cardiovascular disease: the TERC^{-/-} mouse as a model for heart failure and ageing. *Cardiovasc Res* 2009; **81**:244–252.
 31. Lighthouse JK, Burke RM, Velasquez LS, Dirx RA, Aiezza A, Moravec CS, Alexis JD, Rosenberg A, Small EM. Exercise promotes a cardioprotective gene program in resident cardiac fibroblasts. *JCI Insight* 2019; **4**:e92098.
 32. Zhao B, Liu L, Mao J, Zhang Z, Wang Q, Li Q. PIM1 mediates epithelial-mesenchymal transition by targeting Smads and c-Myc in the nucleus and potentiates clear-cell renal-cell carcinoma oncogenesis article. *Cell Death Dis* 2018; **9**: 307.
 33. Warnecke-Eberz U, Bollschweiler E, Drebber U, Pohl A, Baldus SE, Hoelscher AH, Metzger R. Frequent down-regulation of pim-1 mRNA expression in non-small cell lung cancer is associated with lymph node metastases. *Oncol Rep* 2008; **20**:619–624.
 34. Reiser-Erkan C, Erkan M, Pan Z, Bekasi S, Giese NA, Streit S, Michalski CW, Friess H, Kleeff J. Hypoxia-inducible proto-oncogene Pim-1 is a prognostic marker in pancreatic ductal adenocarcinoma. *Cancer Biol Ther* 2008; **7**:1353–1360.
 35. Macdonald A, Campbell DG, Toth R, McLauchlan H, Hastie CJ, Arthur J. Pim kinases phosphorylate multiple sites in Bad and promote 14-3-3 binding and dissociation from Bcl-XL. *BMC Cell Biol* 2006; **7**:1.
 36. Zhang Y, Wang Z, Magnuson NS. Pim-1 kinase-dependent phosphorylation of p21Cip1/WAF1 regulates its stability and cellular localization in H1299 cells. *Mol Cancer Res* 2007; **5**:909–922.
 37. Mochizuki T, Kitanaka C, Noguchi K, Muramatsu T, Asai A, Kuchino Y. Physical and functional interactions between pim-1 kinase and Cdc25A phosphatase. Implications for the Pim-1-mediated activation of the c-Myc signaling pathway. *J Biol Chem* 1999; **274**:18659–18666.
 38. Conery AR, Cao Y, Thompson EA, Townsend CM, Ko TC, Luo K. Akt interacts directly with Smad3 to regulate the sensitivity to TGF-β-induced apoptosis. *Nat Cell Biol* 2004; **6**:366–372.
 39. Remy I, Montmarquette A, Michnick SW. PKB/Akt modulates TGF-β signalling through a direct interaction with Smad3. *Nat Cell Biol* 2004; **6**:358–365.
 40. Sussman MA. Mitochondrial integrity: preservation through Akt/Pim-1 kinase signaling in the cardiomyocyte. *Expert Rev Cardiovasc Ther* 2009; **7**:929–938.

Translational perspective

Telomere maintenance is associated with preservation of cardiomyocyte functional competency and survival, with telomeric shortening linked to cardiomyopathic disease. Aging also contributes to telomere erosion that contributes to deterioration of myocardial performance. Pim1 kinase mediates beneficial effects to preserve cardiomyocyte survival and function and this report demonstrates a novel molecular mechanism of Pim1 action to mitigate telomeric shortening. Findings linking Pim1 to telomere biology introduce another facet of cardioprotection that can be developed as a molecular interventional approach to treat myocardial injury and heart failure.