



Universiteit  
Leiden  
The Netherlands

## Detection of OH at the evening terminator of the ultra-hot Jupiter WASP-76b

Landman, R.; Sánchez López, A.; Mollière, P.; Kesseli, A.Y.; Louca, A.J.; Snellen, I.A.G.

### Citation

Landman, R., Sánchez López, A., Mollière, P., Kesseli, A. Y., Louca, A. J., & Snellen, I. A. G. (2021). Detection of OH at the evening terminator of the ultra-hot Jupiter WASP-76b. *European Planetary Science Congress, EPSC2021-222*. doi:10.5194/epsc2021-222

Version: Publisher's Version

License: [Creative Commons CC BY 4.0 license](https://creativecommons.org/licenses/by/4.0/)

Downloaded from: <https://hdl.handle.net/1887/3276973>

**Note:** To cite this publication please use the final published version (if applicable).



## Detection of OH at the evening terminator of the ultra-hot Jupiter WASP-76b

Rico Landman<sup>1</sup>, Alejandro Sánchez-López<sup>1</sup>, Paul Mollière<sup>2</sup>, Aurora Kesseli<sup>1</sup>, Amy Louca<sup>1</sup>, and Ignas Snellen<sup>1</sup>

<sup>1</sup>Leiden University, Leiden Observatory, Netherlands (rlandman@strw.leidenuniv.nl)

<sup>2</sup>Max-Planck-Institut für Astronomie, Heidelberg, Germany

Ultra-hot Jupiters have dayside temperatures similar to those of M-dwarfs. While molecular absorption from the hydroxyl radical (OH) is easily observed in near-infrared spectra of M-dwarfs, it is often not considered when studying the atmospheres of (ultra-)hot Jupiters. We use high-resolution spectroscopic near-infrared observations of a transit of WASP-76b obtained using CARMENES to assess the presence of OH. After validating the OH line list, we generate model transit spectra of WASP-76b with petitRADTRANS. The data are corrected for telluric contamination and cross-correlated with the model spectra. After combining all cross-correlation functions from the transit, a detection map is constructed. OH is detected in the atmosphere of WASP-76b with a signal-to-noise ratio of 6.1. From a Markov Chain Monte Carlo retrieval we obtain  $K_p=234$  km/s and a blueshift of 13.9 km/s. Considering the fast spin-rotation of the planet, the OH signal is best explained with the signal mainly originating from the evening terminator and the presence of a strong day- to nightside wind. The signal appears to be broad, with a full width at half maximum of 16.2 km/s. The retrieval results in a weak constraint on the temperature of 2420-3150 K at the pressure of the OH signal. Our results demonstrate that OH is readily observable in the transit spectra of ultra-hot Jupiters. Studying this molecule can give new insights in the molecular dissociation processes in the atmospheres of such planets.