



Synthesis, in vitro antiproliferative activities, and Chk1 inhibitory properties of indolylpyrazolones and indolylpyridazinedione

Elisabeth Conchon, Fabrice Anizon, Roy Golsteyn, Stéphane Léonce, Bruno Pfeiffer, Michelle Prudhomme

► To cite this version:

Elisabeth Conchon, Fabrice Anizon, Roy Golsteyn, Stéphane Léonce, Bruno Pfeiffer, et al.. Synthesis, in vitro antiproliferative activities, and Chk1 inhibitory properties of indolylpyrazolones and indolylpyridazinedione. Eur. J. Med. Chem., 2006, 41, pp.1470-1477. <10.1016/j.ejmech.2006.06.012>. <hal-00121158v2>

HAL Id: hal-00121158

<https://hal.archives-ouvertes.fr/hal-00121158v2>

Submitted on 2 Mar 2007

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Synthesis, in vitro antiproliferative activities, and Chk1 inhibitory properties of dipyrrolo[3,4-*a*:3,4-*c*]carbazole-triones

Elisabeth Conchon,^a Fabrice Anizon,^a Roy M. Golsteyn,^b Stéphane Léonce,^b
Bruno Pfeiffer^b and Michelle Prudhomme^{a,*}

^aLaboratoire SEESIB, Université Blaise Pascal, Clermont Ferrand, UMR 6504 du CNRS, 63177 Aubière Cedex, France

^bInstitut de Recherches SERVIER, Division Recherche Cancérologie, 125 Chemin de ronde, 78290 Croissy sur Seine, France

Received 23 June 2006; revised 5 September 2006; accepted 12 September 2006

Available online 5 October 2006

Abstract—The syntheses of dipyrrolo[3,4-*a*:3,4-*c*]carbazole-1,4,6-triones and dipyrrolo[3,4-*a*:3,4-*c*]carbazole-3,4,6-triones are reported. These compounds can be considered as granulatinide analogues in which a maleimide replaces the imidazole moiety and a five-membered lactam ring replaces the upper maleimide. The Chk1 inhibitory properties of the more soluble compounds have been evaluated and their in vitro antiproliferative activities toward three tumor cell lines: murine leukemia L1210, and human colon carcinoma HT29 and HCT116. Due to their insolubility, the biological activities of the other compounds in this series could not be evaluated. All the tested compounds proved to be potent Chk1 inhibitors.

© 2006 Elsevier Ltd. All rights reserved.

1. Introduction

The carbazole framework is found in many biologically active compounds. Some of them, such as rebeccamycin, are topoisomerase I inhibitors. Others, such as staurosporine and UCN-01, are inhibitors of kinases.^{1–3} Granulatinide and isogranulatinide, natural compounds isolated from an ascidian, as well as staurosporine and UCN-01, indolocarbazole compounds isolated from cultures of *Streptomyces*, or synthetic compounds such as SB-218078 have triggered considerable interest as cell cycle G2 checkpoint inhibitors (Fig. 1).^{4–7} In the cell division cycle, the G2 checkpoint is activated in response to DNA damage. Its role consists in blocking the cell cycle to allow time for DNA repair. In more than 60% of cancer cells, the G1 checkpoint is lacking, due to mutations of the *p53* gene. In the *p53*-mutated cells, only the G2 checkpoint provides cancer cells with an opportunity to repair their DNA after damage. Accordingly, combining a DNA damaging agent with a G2 checkpoint inhibitor will force selectively cancer cells into a premature and lethal mitosis due to an accumulation of DNA lesions.^{8–10} The Chk1 kinase plays a major role in the G2 checkpoint regulation.^{11,12} Therefore, Chk1 inhibitors are relevant targets for the conception of agents that are able

to kill selectively cancer cells without causing damage to healthy cells. Granulatinide, isogranulatinide, staurosporine, UCN-01, and SB-218078 were found to be efficient Chk1 inhibitors. All of them possess a carbazole moiety, with an upper heterocycle containing an imide or a lactam function. Moreover, in staurosporine, UCN-01, and SB-218078, a carbohydrate-like heterocycle is linked to both indole nitrogens.

The crystal structures of SB-218078, staurosporine, UCN-01,¹³ and isogranulatinide¹⁴ in complex with Chk1 kinase, have been determined. These compounds are ATP-competitive Chk1 inhibitors. In the structures of the four complexes,




Figure 1.

Keywords: Granulatinide; Dipyrrolo[3,4-*a*:3,4-*c*]carbazole-1,4,6-trione; Dipyrrolo[3,4-*a*:3,4-*c*]carbazole-3,4,6-trione; Antitumor agents; Chk1 inhibitors.

* Corresponding author. Tel.: +33 4 73 40 7124; fax: +33 4 73 40 7717; e-mail: michelle.prudhomme@univ-bpclermont.fr




Figure 2.


two hydrogen bonds between the inhibitors and the ATP binding site of the enzyme are conserved: the first one between the NH of the upper heterocycle and the carbonyl oxygen of Glu⁸⁵, and the second one between the oxygen of the carbonyl group of the lactam or imide function of the drug and the amide nitrogen of Cys⁸⁷. Granulatimide and isogranulatimide isomers, and structurally related compounds bearing modified heterocycles, have been recently synthesized.^{15–24}

In this paper, we describe the syntheses of dipyrrolo[3,4-*a*:3,4-*c*]carbazole-1,4,6-triones and dipyrrolo[3,4-*a*:3,4-*c*]carbazole-3,4,6-triones (Fig. 2). Compared with granulatimide and isogranulatimide, the imidazole has been replaced by a maleimide and the upper heterocycle contains a lactam function, like in staurosporine, instead of the imide function present in granulatimide and isogranulatimide. Moreover, several substituents have been introduced in the 10-position of the indole moiety. Some compounds in this series proved to be extremely insoluble, therefore, their biological activities could not be evaluated. For the most soluble compounds, the Chk1 inhibitory activities and the cytotoxicities toward three tumor cell lines: murine leukemia L1210, and human colon carcinoma HT29 and HCT116 were evaluated.


2. Results and discussion

2.1. Chemistry

In previous works,^{22–24} we described the four-step synthesis of bis-imides granulatimide analogues (Scheme 1). A similar synthetic scheme was applied for the synthesis of



Scheme 1.



Scheme 2.

dipyrrolo[3,4-*a*:3,4-*c*]carbazole-triones. The reduction of the 3-(indol-3-yl)-maleimides intermediates led to lactams and hydroxylactams, from which the synthesis was completed.

Hydroxy and methyl substituents were introduced in the 10-position because in the bis-imide series, these substitutions led to the most efficient Chk1 inhibitors.


3-(Indol-3-yl)-maleimides **1a–d** (R=H, OBn, OH, and CH₃) were prepared as previously described^{22,24} in two steps from the corresponding substituted indoles via a Michael addition with maleimide followed by dehydrogenation of the Michael adduct using DDQ. The corresponding lactams and hydroxylactams were obtained by reduction of the 3-(indol-3-yl)-maleimides (Scheme 2).

Depending on the reducing agent (LiAlH₄, NaBH₄ or DIBAL-H) and the substituent on the indole moiety, important variations were observed in the yields of compounds **2–6** (Table 1). Indeed, Mase et al.²⁵ showed that the regioselectivity of the reduction of monosubstituted maleimides using NaBH₄ was due to the approach of the hydride anion from the less hindered carbonyl group. Therefore, the hydride anion attacks the more hindered carbonyl group. When using DIBAL-H, the inverted regioselectivity is explained by the complexation effect of the carbonyl group with an aluminum atom preferably coordinated to the less hindered carbonyl group. Therefore, the hydride anion approaches from the more hindered carbonyl group and attacks


Table 1. Percentages of compounds **2**, **3**, **4**, **5**, and **6** obtained by reduction of **1a–d** using LiAlH₄, NaBH₄ or DIBAL-H and ratio for the reductions on Cl/Cm (Cl: less hindered carbon, Cm: more hindered carbon)

Starting product	Reducing agent	Compounds (%)					Reduction ratio on Cl/Cm (%)	
		2	3	4	5	6		
1a	LiAlH ₄	9	6	11	23	19	29	71
	NaBH ₄	0	11	0	60	0	16	84
	DIBAL-H	0	17	33	15	0	77	23
1b	LiAlH ₄	1	0	11	17	21	22	78
	DIBAL-H	0	54	11	19	0	78	22
1c	LiAlH ₄	0	0	6	24	40	8	92
	DIBAL-H	0	42	13	13	0	81	19
1d	LiAlH ₄	0	0	12	33	0	27	73
	DIBAL-H	0	0	28	15	0	65	35

the less hindered carbonyl group (Scheme 3). In compounds **3** and **4**, the less hindered carbonyl group has been reduced, whereas in compounds **5** and **6**, the more hindered carbonyl group has been reduced. With the less bulky LiAlH₄, the complexation of both carbonyl groups with an aluminum atom may occur, but the hydride anion approaches very probably more quickly from the less hindered side. The structure of compounds **4a** was assigned from NMR NOESY correlations between the two methylene protons and the NH of the lactam function and the vicinal ethylenic proton (Scheme 4). In compound **6a**, no NOESY correlations were observed between the protons of the methylene group and the ethylenic proton. The H₄ of compound **4a** is shifted at 7.43 ppm, whereas the H₃ of compound **6a** is shifted at 6.26 ppm. Based on these NMR data, the structures of compounds **3a** and **5a** were assigned (**3a**: H₄ at 7.14 ppm, **5a**: H₃ at 6.17 ppm). By analogy with the unsubstituted compounds **3a–6a**, the structures of the analogues substituted in 5'-position on the indole moiety were assigned from the chemical shifts of the ethylenic protons.




Scheme 3.



Scheme 4.

The next step was a Diels–Alder cycloaddition with maleimide. In previous studies, it was observed that the Diels–Alder cycloaddition carried out between 3-indolyl-maleimide and maleimide could lead, according to the treatment, filtration or chromatography on silicagel, to indoline or indole isomers.^{22,26} When the Diels–Alder reaction was performed from **4a** and **4c**, the mixture of isomers could not be separated. The oxidation yielding to the final aromatic compounds **7** and **10** was carried out on the isomeric mixture. With lactams **4b** and **4d**, the indoline intermediate **8** and the indoline intermediate **11** were isolated. The position of the double bond was determined from ¹H NMR data. Indeed, an indole and an imide NH are usually shifted at about 11–12 ppm, whereas an indoline and a lactam NH are shifted at about 7–9 ppm. In compounds **8**, two exchangeable protons are shifted at 11.12, and 11.52 ppm whereas in compound **11**, only one exchangeable proton was shifted at 10.94 ppm. Oxidation of the intermediates in dioxane in



Scheme 5.

the presence of TFA gave lactams **7**, **9**, **10**, and **12** (Scheme 5).

Diels–Alder reactions performed from hydroxy-lactams **3a–d** did not lead to the cycloadducts, whereas from hydroxy-lactams **5a–d**, the cycloaddition occurred with the loss of a water molecule. With compounds **5c** and **5d**, the indole intermediates **15** and **17** could be isolated. Oxidation of the intermediates in dioxane, either in the presence of TFA or with DDQ, led to the required lactams **13**, **14**, and **16**. In spite of various modifications of the oxidation procedure of intermediate **17** (in dioxane in the presence of TFA from 6 to 20 equiv from 60 °C to 80 °C or with DDQ 2 equiv in dioxane at room temperature), the required aromatized compound could not be obtained. Concomitant with the aromatization, oxidation of the lactam heterocycle to imide was observed.

Cycloadditions between maleimide and lactams **6** did not lead to the required cycloadducts. With lactam **6** in which R=H, depending on the solvents used, either a double Diels–Alder reaction or a Diels–Alder reaction followed by a Michael addition with a second molecule of maleimide occurred.

2.2. Chk1 inhibitory activities

The Chk1 inhibitory activities could only be determined with compounds **7**, **10**, **12**, **13**, and intermediate **17** and were compared with those of granulatimide, isogranulatimide, and bis-imide analogue **A** (Fig. 2) (Table 2).²⁷ Due

Table 2. Percentages of Chk1 inhibition at a drug concentration of 10 μ M, IC₅₀ values (μ M) toward Chk1; in vitro antiproliferative activities against three tumor cell lines: murine leukemia L1210, and human HT29 and HCT116 colon carcinoma (IC₅₀ μ M)

Compound	% of Chk1 inhibition at 10 μ M	IC ₅₀ Chk1 (μ M)	L1210	HCT116	HT29
Granulatimide	93.9	0.08	2.8	6.1	5.7
Isogranulatimide	89.7	0.44	10	13	13.7
A	94.4	0.02	32.7	nd	9.7
7	85.4	0.05	nd	nd	nd
10	69.8	1	25.9	28.3	36.4
12	95.7	0.01	54.5	63.5	41.8
13	71.7	0.37	47.0	58.9	>100
17	78.6	2.83	>50	43.7	49.7

to the insolubility of compounds **9**, **14**, and **16**, their Chk1 inhibitory activities could not be evaluated. Compounds **7**, unsubstituted at the 10-position, and compound **12** bearing a hydroxy group, are stronger Chk1 inhibitors than granulatimide and isogranulatimide. Interestingly, when the carbonyl of the lactam heterocycle is oriented toward the indole moiety, the compounds seem to be more efficient Chk1 inhibitors than those in which the carbonyl of the lactam heterocycle is oriented toward the imide heterocycle (compare **7** and **13**). These results are not completely surprising since, in the crystal structures of staurosporine, UCN-01, and isogranulatimide in complex with Chk1, the carbonyl on the left of the upper heterocycle accepts a hydrogen bond from the amide nitrogen of Cys⁸⁷, whereas no hydrogen bond is formed with the carbonyl located on the right. However, the strong Chk1 inhibitory activity of compound **13** could be due to a different position of the molecule in the ATP binding site allowing the formation of the two fundamental hydrogen bonds with Glu⁸⁵ and Cys⁸⁷. In this orientation, the imide heterocycle would be positioned on the left and the indole moiety would lie on the right. No significant differences are observed between the Chk1 inhibitory activities of lactam **7** and imide **A**. Compared with unsubstituted compound **7**, compound **12** substituted with a hydroxy group is a stronger Chk1 inhibitor.

2.3. In vitro antiproliferative activities

The cytotoxicities of the soluble compounds were evaluated toward three tumor cell lines: murine leukemia L1210, and human colon carcinoma HT29 and HCT116 and compared with those of granulatimide, isogranulatimide, and compound **A** (Table 2). Compared with granulatimide and isogranulatimide, all the lactams tested are considerably less active, their cytotoxicities are in the same range as those of imide **A**. A checkpoint inhibitor is not expected to be cytotoxic by itself. However, the weak in vitro antiproliferative activities of the new compounds described in this paper suggests a possible instability of these compounds in the biological medium.

3. Conclusion

In conclusion, this work reports the synthesis of pyrrolo[3,4-*a*:3,4-*c*]carbazole-1,4,6-tetraones and pyrrolo[3,4-*a*:3,4-*c*]carbazole-3,4,6-tetraones. These compounds are structurally

related to the Chk1 inhibitor granulatimide. Their upper heterocycle contains a lactam function like in staurosporine. All the new compounds are potent Chk1 inhibitors suggesting that the orientation of the carbonyl group of the upper heterocycle, either toward the indole moiety or toward the maleimide unit, could modify the positioning of the drug in the active site of the kinase. Their weak cytotoxicity could be due to a limited penetration into the cells or to a degradation in the biological media. This hypothesis is currently under investigation.

4. Experimental

4.1. Chemistry

IR spectra were recorded on a Perkin–Elmer 881 spectrometer (ν in cm^{-1}). NMR spectra were performed on a Bruker AVANCE 400 and AVANCE 500 (chemical shifts δ in parts per million, the following abbreviations are used: singlet (s), broad singlet (br s), doublet (d), doubled doublet (dd), triplet (t), doubled triplet (dt), multiplet (m), pseudo quadruplet (pq), tertiary carbons (C tert), and quaternary carbons (C quat). The signals were assigned from ¹H–¹H COSY, HSQC, and HMBC NMR correlations. Low-resolution mass spectra (ESI+ and APCI+) and HRMS were determined on a MS Hewlett Packard instrument. Chromatographic purifications were performed by flash silicagel Geduran SI 60 (Merck) 0.040–0.063 mm column chromatography.

4.2. Typical procedure for the reduction using LiAlH₄

To a solution of 3-(indol-3-yl)-maleimide (200 mg, 0.94 mmol) in THF (40 mL) was added dropwise a 1 M solution of LiAlH₄ in Et₂O (6 mL) at room temperature. The mixture was stirred for 3 days. After cooling to 0 °C, water (14 mL) was added. The mixture was acidified to pH 2 with 2 M HCl (2 mL). After extraction with EtOAc, the organic phase was washed with saturated aqueous NaHCO₃. The organic phase was dried over MgSO₄ and the solvent was removed. The residue was purified by flash chromatography (eluent: from EtOAc/cyclohexane 1:1 to EtOAc/MeOH 9:1).

4.3. Typical procedure for the reduction using NaBH₄

To a solution of 3-(indol-3-yl)-maleimide (2.4 mmol) in THF (50 mL) was added sodium borohydride (90 mg, 2.4 mmol) in portions. The mixture was stirred at room temperature for 24 h. After cooling to 0 °C, water was added. After extraction with EtOAc, the organic phase was dried over MgSO₄, the solvent was removed and the residue was purified by flash chromatography (eluent: from cyclohexane/EtOAc 1:1 to EtOAc/MeOH 9:1).

4.4. Typical procedure for the reduction using DIBAL-H

To a solution of 3-(indol-3-yl)-maleimide (0.94 mmol) in THF (70 mL) at –78 °C was added dropwise a 1 M solution of DIBAL-H in toluene (2.3 mL). The mixture was stirred for 1.5 h at –78 °C, then a 1 M solution of DIBAL-H in toluene (2.3 mL) was added. The mixture was stirred at –78 °C for 6 h. Then the mixture was warmed to 0 °C and saturated aqueous NaHCO₃ was added dropwise. After extraction

with EtOAc, the organic phase was dried over MgSO₄, the solvent was removed and the residue was purified by flash chromatography (eluent: from EtOAc/cyclohexane 1:1 to EtOAc/MeOH 9:1).

4.4.1. 3-(1*H*-Indol-3-yl)-1*H*-pyrrole (2a). Yellow solid. Mp 65 °C. IR (KBr) ν_{NH} 2924 cm⁻¹. HRMS (ESI+) [M+H]⁺ calcd for C₁₂H₁₁N₂ 183.0922, found 183.0926.

¹H NMR (400 MHz, DMSO-*d*₆): 6.38 (1H, s), 6.79 (1H, s), 7.04 (1H, dt, $J_1=7.0$ Hz, $J_2=1.0$ Hz), 7.09–7.10 (2H, m), 7.37 (1H, d, $J=8.0$ Hz), 7.40 (1H, d, $J=2.5$ Hz), 7.77 (1H, d, $J=8.0$ Hz), 10.73 (1H, br s), 10.91 (1H, s).

¹³C NMR (100 MHz, DMSO-*d*₆): 106.1, 111.4, 113.4, 117.8, 118.5, 119.5, 120.6, 120.8 (C tert), 111.6, 117.4, 125.4, 136.5 (C quat).

4.4.2. 3-(1*H*-Indol-3-yl)-1*H*-5-hydroxy-2,5-dihydro-pyrrol-2-one (3a). Orange-brown solid. Mp >300 °C. IR (KBr) $\nu_{\text{C}=\text{C}}$ 1632 cm⁻¹, $\nu_{\text{C}=\text{O}}$ 1705 cm⁻¹, $\nu_{\text{NH-OH}}$ 3200–3500 cm⁻¹. Mass (ESI+) [M+Na]⁺ 237.

¹H NMR (400 MHz, DMSO-*d*₆): 5.59 (1H, d, $J=9.0$ Hz, H₅), 5.99 (1H, d, $J=9.0$ Hz, OH), 7.14 (1H, s, H₄), 7.18 (1H, dt, $J_1=8.0$ Hz, $J_2=1.0$ Hz), 7.21 (1H, dt, $J_1=8.0$ Hz, $J_2=1.0$ Hz), 7.51 (1H, d, $J=8.0$ Hz), 7.94 (1H, d, $J=8.0$ Hz), 8.28 (1H, d, $J=2.5$ Hz), 8.65 (1H, s), 11.46 (1H, s).

¹³C NMR (100 MHz, DMSO-*d*₆): 78.2 (CHOH), 106.0, 125.5, 130.6, 136.1 (C quat), 112.0, 119.7, 119.9, 121.7, 126.5, 133.8 (C tert), 171.7 (C=O).

4.4.3. 3-(1*H*-Indol-3-yl)-1*H*-2,5-dihydro-pyrrol-2-one (4a). Orange-brown solid. Mp 205–207 °C. IR (KBr) $\nu_{\text{C}=\text{C}}$ 1628 cm⁻¹, $\nu_{\text{C}=\text{O}}$ 1678 cm⁻¹, ν_{NH} 3284 cm⁻¹. HRMS (ESI+) [M+H]⁺ calcd for C₁₂H₁₁N₂O 199.0871, found 199.0880.

¹H NMR (400 MHz, DMSO-*d*₆): 4.06 (2H, s, CH₂), 7.15 (1H, dt, $J_1=8.0$ Hz, $J_2=1.0$ Hz, H₅'), 7.24 (1H, dt, $J_1=8.0$ Hz, $J_2=1.0$ Hz, H₆'), 7.43 (1H, d, $J=1.5$ Hz, H₄), 7.49 (1H, d, $J=8.0$ Hz, H₇'), 7.93 (1H, d, $J=8.0$ Hz, H₄'), 8.28 (1H, d, $J=2.5$ Hz, H₂'), 8.44 (1H, s, NH₁), 11.36 (1H, s, NH_{indole}).

¹³C NMR (100 MHz, DMSO-*d*₆): 46.0 (CH₂), 106.8, 125.5 (C quat), 130.6, 136.0 (C quat), 111.8, 119.3, 119.6, 121.5, 125.5 (C tert arom), 131.8 (CH lactam), 173.1 (C=O).

4.4.4. 4-(1*H*-Indol-3-yl)-1*H*-5-hydroxy-2,5-dihydro-pyrrol-2-one (5a). Orange-brown solid. Mp 207 °C. IR (KBr) $\nu_{\text{C}=\text{C}}$ 1608 cm⁻¹, $\nu_{\text{C}=\text{O}}$ 1662 cm⁻¹, ν_{NH} 3271 cm⁻¹. HRMS (ESI-) [M-H]⁻ calcd for C₁₂H₉N₂O₂ 213.0664, found 213.0667.

¹H NMR (400 MHz, DMSO-*d*₆): 5.80 (1H, dd, $J_1=9.5$ Hz, $J_2=1.0$ Hz, H₅), 6.17 (1H, d, $J=1.0$ Hz, H₃), 6.22 (1H, d, $J=9.5$ Hz, OH), 7.17 (1H, dt, $J_1=8.0$ Hz, $J_2=1.0$ Hz), 7.19 (1H, dt, $J_1=8.0$ Hz, $J_2=1.0$ Hz), 7.47 (1H, d, $J=8.0$ Hz), 7.84 (1H, d, $J=3.0$ Hz), 7.85 (1H, d, $J=8.0$ Hz), 8.16 (1H, s), 11.69 (1H, s).

¹³C NMR (100 MHz, DMSO-*d*₆): 80.5 (CHOH), 108.0, 125.3, 136.6, 154.4 (C quat), 112.1, 113.2, 120.1, 120.6, 122.1, 128.5 (C tert), 172.8 (C=O).

4.4.5. 4-(1*H*-Indol-3-yl)-1*H*-2,5-dihydro-pyrrol-2-one (6a). Brown solid. Mp 255 °C. IR (KBr) $\nu_{\text{C}=\text{C}}$ 1610 cm⁻¹, $\nu_{\text{C}=\text{O}}$ 1649 cm⁻¹, ν_{NH} 3255 cm⁻¹. HRMS (ESI-) [M-H]⁻ calcd for C₁₂H₉N₂O 197.0715, found 197.0722.

¹H NMR (400 MHz, DMSO-*d*₆): 4.42 (2H, s, CH₂), 6.26 (1H, d, $J=1.0$ Hz, H₃), 7.15 (1H, dt, $J_1=8.0$ Hz, $J_2=1.0$ Hz, H₅'), 7.19 (1H, dt, $J_1=8.0$ Hz, $J_2=1.0$ Hz, H₆'), 7.46 (1H, d, $J=8.0$ Hz, H₇'), 7.84 (3H, m, H₂', H₄', NH₁), 11.69 (1H, d, $J=1.0$ Hz, NH_{indole}).

¹³C NMR (100 MHz, DMSO-*d*₆): 48.3 (CH₂), 109.0, 124.8, 136.9, 152.7 (C quat), 112.1, 114.4, 119.9, 120.6, 122.1, 126.7 (C tert), 175.0 (C=O).

4.4.6. 3-(5-Methyl-1*H*-indol-3-yl)-1*H*-pyrrole (2b). Green solid. Mp 110 °C. IR (KBr) ν_{NH} 3423 cm⁻¹. HRMS (ESI+) [M+H]⁺ calcd for C₁₃H₁₃N₂ 197.1079, found 197.1084.

¹H NMR (400 MHz, DMSO-*d*₆): 2.44 (3H, s), 6.39 (1H, pq, $J=2.0$ Hz), 6.82 (1H, pq, $J=2.0$ Hz), 6.94 (1H, dd, $J_1=8.0$ Hz, $J_2=1.5$ Hz), 7.11 (1H, pq, $J=2.0$ Hz), 7.28 (1H, d, $J=8.0$ Hz), 7.37 (1H, d, $J=2.5$ Hz), 10.80 (1H, s, NH), 10.74 (1H, s, NH).

¹³C NMR (100 MHz, DMSO-*d*₆): 21.3 (CH₃), 106.1, 111.0, 113.3, 117.7, 119.1, 120.7, 122.3 (C tert), 111.1, 117.6, 125.7, 126.9, 134.9 (C quat).

4.4.7. 3-(5-Methyl-1*H*-indol-3-yl)-1*H*-5-hydroxy-2,5-dihydro-pyrrol-2-one (3b). Off-white solid. Mp >300 °C. IR (KBr) $\nu_{\text{C}=\text{C}}$ 1633 cm⁻¹, $\nu_{\text{C}=\text{O}}$ 1697 cm⁻¹, $\nu_{\text{NH,OH}}$ 3200–3500 cm⁻¹. HRMS (ESI+) [M+H-H₂O]⁺ calcd for C₁₃H₁₁N₂O 211.0871, found 211.0871.

¹H NMR (400 MHz, DMSO-*d*₆): 2.48 (3H, s), 5.59 (1H, d, $J=7.5$ Hz), 5.99 (1H, d, $J=8.5$ Hz, OH), 7.04 (1H, t, $J=8.0$ Hz), 7.14 (1H, s, H₄), 7.38 (1H, d, $J=8.0$ Hz), 7.73 (1H, s), 8.25 (1H, d, $J=3.0$ Hz), 8.65 (1H, s, NH), 11.32 (1H, s, NH).

¹³C NMR (100 MHz, DMSO-*d*₆): 21.3 (CH₃), 78.3 (CHOH), 105.6, 125.8, 128.7, 130.8, 134.5 (C quat), 111.6, 119.5, 123.3, 126.5, 133.5 (C tert), 171.8 (C=O).

4.4.8. 3-(5-Methyl-1*H*-indol-3-yl)-1*H*-2,5-dihydro-pyrrol-2-one (4b). Ochre solid. Mp >300 °C. IR (KBr) $\nu_{\text{C}=\text{C}}$ 1606 cm⁻¹, $\nu_{\text{C}=\text{O}}$ 1649 cm⁻¹, ν_{NH} 3100–3300 cm⁻¹. HRMS (ESI+) [M+Na]⁺ calcd for C₁₃H₁₂N₂ONa 235.0847, found 235.0853.

¹H NMR (400 MHz, DMSO-*d*₆): 2.47 (3H, s), 4.05 (2H, s), 7.02 (1H, dd, $J_1=8.0$ Hz, $J_2=1.0$ Hz), 7.36 (1H, d, $J=8.0$ Hz), 7.41 (1H, d, $J=1.5$ Hz, H₄), 7.71 (1H, br s), 8.22 (1H, d, $J=2.5$ Hz), 8.40 (1H, s, NH), 11.21 (1H, s, NH).

¹³C NMR (100 MHz, DMSO-*d*₆): 21.3 (CH₃), 45.9 (CH₂N), 106.3, 125.7, 128.3, 130.8, 134.4 (C quat), 111.5, 119.3, 123.1, 125.5, 131.5 (C tert), 173.2 (C=O).

4.4.9. 4-(5-Methyl-1*H*-indol-3-yl)-1*H*-5-hydroxy-2,5-dihydro-pyrrol-2-one (5b). Off-white solid. Mp 104 °C. IR (KBr) $\nu_{\text{C}=\text{C}}$ 1615 cm^{-1} , $\nu_{\text{C}=\text{O}}$ 1682 cm^{-1} , ν_{NH} 3274, 3398 cm^{-1} . HRMS (ESI+) $[\text{M}+\text{Na}]^+$ calcd for $\text{C}_{13}\text{H}_{12}\text{N}_2\text{O}_2\text{Na}$ 251.0796, found 251.0805.

^1H NMR (400 MHz, $\text{DMSO-}d_6$): 2.47 (3H, s), 5.83 (1H, dd, $J_1=9.5$ Hz, $J_2=1.5$ Hz), 6.21 (1H, d, $J=1.0$ Hz, H_3), 6.23 (1H, d, $J=9.5$ Hz), 7.05 (1H, dd, $J_1=8.0$ Hz, $J_2=1.0$ Hz), 7.39 (1H, d, $J=8.0$ Hz), 7.70 (1H, br s), 7.82 (1H, d, $J=2.5$ Hz), 8.16 (1H, s, NH), 11.59 (1H, s, NH).

^{13}C NMR (100 MHz, $\text{DMSO-}d_6$): 21.2 (CH_3), 80.5 (CHOH), 107.6, 125.6, 129.4, 134.9, 154.5 (C quat), 111.7, 113.0, 119.8, 123.6, 128.4 (C tert), 172.9 (C=O).

4.4.10. 4-(5-Methyl-1*H*-indol-3-yl)-1*H*-2,5-dihydro-pyrrol-2-one (6b). Off-white solid. Mp 140 °C. IR (KBr) $\nu_{\text{C}=\text{C}}$ 1605 cm^{-1} , $\nu_{\text{C}=\text{O}}$ 1649 cm^{-1} , ν_{NH} 3035–3430 cm^{-1} . HRMS (ESI+) $[\text{M}+\text{H}]^+$ calcd for $\text{C}_{13}\text{H}_{13}\text{N}_2\text{O}$ 213.1028, found 213.1027.

^1H NMR (400 MHz, $\text{DMSO-}d_6$): 2.46 (3H, s), 4.43 (2H, s), 6.28 (1H, d, $J=1.0$ Hz, H_3), 7.05 (1H, dd, $J_1=8.0$ Hz, $J_2=1.0$ Hz), 7.37 (1H, d, $J=8.0$ Hz), 7.69 (1H, s), 7.81 (1H, s), 7.82 (1H, s), 11.58 (1H, s, NH).

^{13}C NMR (100 MHz, $\text{DMSO-}d_6$): 21.2 (CH_3), 48.3 (CH_2N), 108.6, 125.1, 129.4, 135.1, 152.8 (C quat), 111.8, 114.1, 119.6, 123.6, 126.7 (C tert), 175.1 (C=O).

4.4.11. 3-(5-Benzyloxy-1*H*-indol-3-yl)-1*H*-5-hydroxy-2,5-dihydro-pyrrol-2-one (3c). Off-white solid. Mp >200 °C (decomposition). IR (KBr) $\nu_{\text{C}=\text{C}}$ 1631 cm^{-1} , $\nu_{\text{C}=\text{O}}$ 1706 cm^{-1} , $\nu_{\text{NH,OH}}$ 3100–3500 cm^{-1} . HRMS (ESI+) $[\text{M}+\text{H}]^+$ calcd for $\text{C}_{19}\text{H}_{17}\text{N}_2\text{O}_3$ 321.1239, found 321.1256.

^1H NMR (400 MHz, $\text{DMSO-}d_6$): 5.22 (2H, s, CH_2), 5.58 (1H, dt, $J_1=9.0$ Hz, $J_2=2.0$ Hz, H_5), 5.98 (1H, d, $J=9.0$ Hz, OH), 6.94 (1H, dd, $J_1=9.0$ Hz, $J_2=2.0$ Hz), 7.12 (1H, t, $J=1.5$ Hz), 7.36–7.46 (5H, m), 7.55 (2H, d, $J=7.0$ Hz), 8.25 (1H, d, $J=3.0$ Hz), 8.63 (1H, s, NH), 11.33 (1H, d, $J=2.5$ Hz, NH).

^{13}C NMR (100 MHz, $\text{DMSO-}d_6$): 69.9 (CH_2O), 78.3 (CHOH), 103.4, 112.3, 112.6, 127.1, 127.6 (3C), 128.3 (2C), 133.5 (C tert), 105.9, 125.9, 130.6, 131.3, 137.8, 153.2 (C quat), 171.8 (C=O).

4.4.12. 3-(5-Benzyloxy-1*H*-indol-3-yl)-1*H*-2,5-dihydro-pyrrol-2-one (4c). Brown solid. Mp 185 °C. IR (KBr) $\nu_{\text{C}=\text{C}}$ 1620 cm^{-1} , $\nu_{\text{C}=\text{O}}$ 1680 cm^{-1} , ν_{NH} 3100–3500 cm^{-1} . Mass (ESI+) $[\text{M}+\text{K}]^+$ 343, $[\text{M}+\text{Na}]^+$ 327.

^1H NMR (400 MHz, $\text{DMSO-}d_6$): 4.05 (2H, s), 5.20 (2H, s), 6.92 (1H, dd, $J_1=9.0$ Hz, $J_2=2.0$ Hz), 7.33–7.47 (6H, m), 7.54 (2H, d, $J=7.5$ Hz), 8.23 (1H, d, $J=2.5$ Hz), 8.41 (1H, s, NH), 11.22 (1H, s, NH).

^{13}C NMR (100 MHz, $\text{DMSO-}d_6$): 46.0 (CH_2N), 69.9 (CH_2OBn), 103.4, 112.0, 112.4, 126.1, 127.6, 127.7 (2C), 128.3 (3C), 131.3 (C tert), 106.7, 125.8, 130.6, 131.2, 137.8, 153.0 (C quat), 173.1 (C=O).

4.4.13. 4-(5-Benzyloxy-1*H*-indol-3-yl)-1*H*-5-hydroxy-2,5-dihydro-pyrrol-2-one (5c). Off-white solid. Mp 215–216 °C. IR (KBr) $\nu_{\text{C}=\text{C}}$ 1613 cm^{-1} , $\nu_{\text{C}=\text{O}}$ 1681 cm^{-1} , $\nu_{\text{NH,OH}}$ 3040–3663 cm^{-1} . HRMS (ESI+) $[\text{M}+\text{H}]^+$ calcd for $\text{C}_{19}\text{H}_{17}\text{N}_2\text{O}_3$ 321.1239, found 321.1242.

^1H NMR (400 MHz, $\text{DMSO-}d_6$): 5.23 (2H, s), 5.81 (1H, d, $J=9.5$ Hz), 6.22 (1H, s), 6.23 (1H, d, $J=9.0$ Hz), 6.95 (1H, d, $J=9.0$ Hz), 7.34–7.46 (5H, m), 7.54 (2H, d, $J=7.5$ Hz), 7.83 (1H, s), 8.17 (1H, s), 11.61 (1H, s, NH).

^{13}C NMR (100 MHz, $\text{DMSO-}d_6$): 69.7 (CH_2OBn), 80.5 (CHOH), 103.5, 111.4, 111.9, 112.8, 127.6 (3C), 128.3 (2C), 129.0 (C tert), 107.9, 125.8, 131.7, 137.7, 153.6, 154.3 (C quat), 173.0 (C=O).

4.4.14. 4-(5-Benzyloxy-1*H*-indol-3-yl)-1*H*-2,5-dihydro-pyrrol-2-one (6c). Off-white solid. Mp 235–237 °C. IR (KBr) $\nu_{\text{C}=\text{C}}$ 1606 cm^{-1} , $\nu_{\text{C}=\text{O}}$ 1651 cm^{-1} , ν_{NH} 3200–3430 cm^{-1} . HRMS (ESI+) $[\text{M}+\text{H}]^+$ calcd for $\text{C}_{19}\text{H}_{17}\text{N}_2\text{O}_2$ 305.1290, found 305.1297.

^1H NMR (400 MHz, $\text{DMSO-}d_6$): 4.41 (2H, s), 5.23 (2H, s), 6.29 (1H, s, H_3), 6.94 (1H, d, $J=9.0$ Hz), 7.34–7.46 (5H, m), 7.54 (2H, d, $J=7.5$ Hz), 7.82 (2H, s), 11.59 (1H, s, NH).

^{13}C NMR (100 MHz, $\text{DMSO-}d_6$): 48.3, 69.7 (CH_2), 103.2, 103.5, 112.9, 104.0, 114.0, 127.2, 127.6 (2C), 128.3 (2C), 129.0 (C tert), 108.9, 125.3, 132.0, 137.7, 152.7, 153.6 (C quat), 175.2 (CO).

4.4.15. 4-(5-Hydroxy-1*H*-indol-3-yl)-1*H*-5-hydroxy-2,5-dihydro-pyrrol-2-one (5d). Brown solid. Mp 102–104 °C. IR (KBr) $\nu_{\text{C}=\text{C}}$ 1614 cm^{-1} , $\nu_{\text{C}=\text{O}}$ 1678 cm^{-1} , $\nu_{\text{NH,OH}}$ 2981–3599 cm^{-1} . HRMS (ESI+) $[\text{M}+\text{Na}]^+$ calcd for $\text{C}_{12}\text{H}_{10}\text{N}_2\text{O}_3\text{Na}$ 253.0589, found 253.0595.

^1H NMR (400 MHz, $\text{DMSO-}d_6$): 5.81 (1H, d, $J=10.0$ Hz), 5.93 (1H, s), 6.23 (1H, d, $J=10.0$ Hz, OH), 6.75 (1H, d, $J=9.0$ Hz), 7.14 (1H, s), 7.31 (1H, dd, $J_1=9.0$ Hz, $J_2=1.0$ Hz), 7.78 (1H, s), 8.16 (1H, s), 8.97 (1H, s), 11.49 (1H, s, $\text{NH}_{\text{indole}}$).

^{13}C NMR (100 MHz, $\text{DMSO-}d_6$): 80.5 (CHOH), 104.2, 111.9, 112.0, 112.6, 128.8 (C tert), 107.2, 126.3, 130.9, 152.1, 154.9 (C quat), 172.8 (C=O).

4.4.16. 3-(5-Hydroxy-1*H*-indol-3-yl)-1*H*-2,5-dihydro-pyrrol-2-one (4d). Brown solid. Mp 180 °C. IR (KBr) $\nu_{\text{C}=\text{C}}$ 1618 cm^{-1} , $\nu_{\text{C}=\text{O}}$ 1672 cm^{-1} , ν_{NH} 3100–3550 cm^{-1} . Mass (APCI+) $[\text{M}+\text{H}]^+$ 215.

^1H NMR (400 MHz, $\text{DMSO-}d_6$): 4.04 (2H, s), 6.72 (1H, d, $J=9.0$ Hz), 7.20 (1H, s), 7.21 (1H, s), 7.27 (1H, d, $J=9.0$ Hz), 8.18 (1H, s), 8.39 (1H, s), 8.84 (1H, br s), 11.08 (1H, s, NH).

^{13}C NMR (100 MHz, $\text{DMSO-}d_6$): 45.9 (CH_2), 103.9, 111.6, 112.1, 125.8, 130.5 (C tert), 106.0, 126.3, 130.4, 131.0, 151.4 (C quat), 173.2 (C=O).

4.4.17. 2*H*,5*H*,7*H*-1,3,4,6-Tetrahydro-dipyrrolo[3,4-*a*:3,4-*c'*]carbazole-1,4,6-trione (7). A mixture of maleimide

(244 mg, 2.51 mmol) and compound **4a** (100 mg, 0.505 mmol) in xylene (14 mL) was refluxed for 3 days. After cooling, the yellow precipitate was filtered off washed with xylene and dried. The yellow solid (144 mg, 0.491 mmol, 97% yield) corresponds to an isomeric mixture of Diels–Alder adducts. The mixture of isomers (84 mg, 0.286 mmol) in dioxane (22 mL) was refluxed for 36 h in the presence of trifluoroacetic acid (293 μ L). After evaporation, EtOAc was added to the residue. The mixture was filtered off. The solid was successively washed with saturated aqueous NaHCO₃, brine, and EtOAc to give compound **7** as a yellow-brown solid (22 mg, 0.075 mmol, 27% yield).

Mp >310 °C. IR (KBr) $\nu_{\text{C=O}}$ 1690, 1700, 1720 cm^{-1} , ν_{NH} 3387 cm^{-1} . HRMS (ESI+) [M+H]⁺ calcd for C₁₆H₁₀N₃O₃ 292.0722, found 292.0745.

¹H NMR (400 MHz, DMSO-*d*₆): 4.77 (2H, s), 7.34 (1H, t, *J*=7.5 Hz), 7.61 (1H, t, *J*=8.0 Hz), 7.73 (1H, d, *J*=7.5 Hz), 9.08 (1H, s), 9.25 (1H, d, *J*=8.0 Hz), 11.40 (1H, s), 12.37 (1H, s).

Due to its insolubility, the ¹³C NMR spectrum could not be recorded.

4.4.18. 10-Methyl-2*H*,5*H*,7*H*-1,3,3*a*,3*b*,4,6,6*a*,11*c*-octahydro-dipyrrolo[3,4-*a*:3,4-*c*]carbazole-1,4,6-trione (8**).** A mixture of maleimide (179 mg, 1.84 mmol) and compound **4b** (78 mg, 0.37 mmol) in xylene (10 mL) was refluxed for 3 days. After cooling, the mixture was filtered off and the solid residue was washed with xylene then dried to give **8** as an off-white solid (116 mg, 0.37 mmol, 100% yield).

Mp >295 °C. IR (KBr) $\nu_{\text{C=O}}$ 1715, 1777 cm^{-1} , ν_{NH} 3061–3684 cm^{-1} . HRMS (ESI+) [M+H]⁺ calcd for C₁₇H₁₆N₃O₃ 310.1192, found 310.1178.

¹H NMR (400 MHz, DMSO-*d*₆): 2.37 (3H, s), 2.82 (1H, t, *J*=9.5 Hz), 3.11 (1H, t, *J*=9.0 Hz), 3.25 (1H, m), 3.51 (1H, t, *J*=7.0 Hz), 3.56 (1H, d, *J*=7.0 Hz), 4.38 (1H, d, *J*=8.5 Hz), 6.92 (1H, d, *J*=8.0 Hz), 7.28 (1H, d, *J*=8.5 Hz), 7.58 (1H, s, NH), 7.75 (1H, s), 11.12 (1H, s, NH), 11.52 (1H, s, NH).

¹³C NMR (100 MHz, DMSO-*d*₆): 21.3 (CH₃), 41.3 (CH₂N), 35.6, 38.6, 40.3, 40.7 (CH), 110.7, 120.2, 122.8 (C tert), 102.2, 125.9, 126.3, 126.7, 135.3 (C quat), 175.1, 177.0, 178.5 (C=O).

4.4.19. 10-Methyl-2*H*,5*H*,7*H*-1,3,4,6-tetrahydro-dipyrrolo[3,4-*a*:3,4-*c*]carbazole-1,4,6-trione (9**).** A solution of **8** (104 mg, 0.34 mmol) in dioxane (11 mL) and trifluoroacetic acid (266 μ L) was stirred at 80 °C for 48 h. After evaporation, water was added to the residue, the mixture was filtered off, the solid was washed with water and with small amounts of EtOAc to give **9** as an orange solid (80 mg, 0.263 mmol, 78% yield).

Mp >300 °C. IR (KBr) $\nu_{\text{C=O}}$ 1716, 1758 cm^{-1} , ν_{NH} 3208–3664 cm^{-1} . HRMS (ESI+) [M+H]⁺ calcd for C₁₇H₁₂N₃O₃ 306.0879, found 306.0894.

¹H NMR (400 MHz, DMSO-*d*₆): 2.50 (3H, s), 4.75 (2H, s), 7.43 (1H, d, *J*=8.0 Hz), 7.61 (1H, d, *J*=8.0 Hz), 9.05 (2H, s), 11.36 (1H, s), 12.23 (1H, s).

Due to its insolubility, the ¹³C NMR spectrum could not be recorded.

4.4.20. 10-Benzyloxy-2*H*,5*H*,7*H*-1,3,4,6-tetrahydro-dipyrrolo[3,4-*a*:3,4-*c*]carbazole-1,4,6-trione (10**).** A mixture of maleimide (122 mg, 1.25 mmol) and compound **4c** (74 mg, 0.25 mmol) in xylene (6 mL) was refluxed for 18 h. After cooling, the mixture was filtered off and the solid residue was washed with CH₂Cl₂ then was dried to give an orange solid as a mixture of isomers (97 mg, 0.25 mmol, quantitative yield).

The mixture of isomers (90 mg, 0.23 mmol), dioxane (6 mL) and trifluoroacetic acid (2.2 mmol, 172 μ L) was stirred at 80 °C for 48 h. After evaporation, water was added to the residue, the mixture was filtered off and the solid residue was washed repeatedly with water and small amounts of EtOAc to give **10** (61 mg, 0.15 mmol, 65% yield) as a dark red solid.

Mp >300 °C. IR (KBr) $\nu_{\text{C=C}}$ 1617 cm^{-1} , $\nu_{\text{C=O}}$ 1722, 1774 cm^{-1} , ν_{NH} 3100–3550 cm^{-1} . HRMS (ESI+) [M+Na]⁺ calcd for C₂₃H₁₅N₃O₄Na 420.0960, found 420.0974.

¹H NMR (400 MHz, DMSO-*d*₆): 4.76 (2H, s), 5.21 (2H, s), 7.33–7.65 (7H, m), 8.95 (1H, d, *J*=2.0 Hz), 9.07 (1H, s), 11.37 (1H, s), 12.21 (1H, s).

Due to its insolubility, the ¹³C NMR spectrum could not be recorded.

4.4.21. 10-Hydroxy-2*H*,5*H*,7*H*-1,3,3*a*,3*b*,4,6,6*a*,11*c*-octahydro-dipyrrolo[3,4-*a*:3,4-*c*]carbazole-1,4,6-trione (11**).** A mixture of maleimide (40 mg, 0.48 mmol) and **4d** (92 mg, 0.43 mmol) in xylene (5 mL) was refluxed for 24 h. After cooling, the mixture was filtered off and the solid residue was washed with CH₂Cl₂ then was dried to give **11** (40 mg, 0.13 mmol, 30% yield) as a dark orange solid.

Mp >300 °C. IR (KBr) $\nu_{\text{C=O}}$ 1691, 1763 cm^{-1} , $\nu_{\text{NH,OH}}$ 3300–3550 cm^{-1} . Mass (ESI+) [M+H]⁺ 312.

¹H NMR (400 MHz, DMSO-*d*₆): 2.94 (1H, m), 3.27 (1H, t, *J*=8.0 Hz), 3.47 (1H, t, *J*=9.0 Hz), 3.61 (1H, t, *J*=10.0 Hz), 4.23–4.29 (2H, m), 6.28 (1H, s), 6.59 (1H, d, *J*=8.0 Hz), 6.63 (1H, d, *J*=8.0 Hz), 7.83 (1H, s), 8.06 (1H, s), 8.67 (1H, s), 10.94 (1H, s).

¹³C NMR (100 MHz, DMSO-*d*₆): 41.1 (CH₂N), 36.2, 42.7, 60.5, 70.0 (CH), 110.4, 113.2, 119.8 (C tert arom), 119.5, 122.8, 143.1, 148.8, 149.8 (C quat arom), 168.4, 175.7, 178.4 (C=O).

4.4.22. 10-Hydroxy-2*H*,5*H*,7*H*-1,3,4,6-tetrahydro-dipyrrolo[3,4-*a*:3,4-*c*]carbazole-1,4,6-trione (12**).** A solution of **11** (80 mg, 0.257 mmol) in dioxane (8 mL) was stirred at 80 °C for 48 h in the presence of trifluoroacetic acid (2.56 mmol, 200 μ L). After evaporation, water was added to the residue, the mixture was filtered off and the solid residue was washed successively with water and small amounts

of EtOAc to give **12** (52 mg, 0.169 mmol, 66% yield) as a red solid.

Mp >300 °C. IR (KBr) $\nu_{\text{C=O}}$ 1711 cm^{-1} , $\nu_{\text{NH,OH}}$ 3200–3672 cm^{-1} . HRMS (ESI+) $[\text{M}+\text{H}]^+$ calcd for $\text{C}_{16}\text{H}_{10}\text{N}_3\text{O}_4$ 308.0671, found 308.0672.

^1H NMR (400 MHz, $\text{DMSO-}d_6$): 4.75 (2H, s), 7.11 (1H, dd, $J_1=9.0$ Hz, $J_2=2.0$ Hz), 7.54 (1H, d, $J=9.0$ Hz), 8.66 (1H, d, $J=2.0$ Hz), 9.02 (1H, s), 9.27 (1H, s), 11.33 (1H, s), 12.06 (1H, s).

Due to its insolubility, the ^{13}C NMR spectrum could not be recorded.

4.4.23. 2H,5H,7H-1,3,4,6-Tetrahydro-dipyrrolo[3,4-*a*:3,4-*c*]carbazole-3,4,6-trione (13). A mixture of **5a** (241 mg, 1.126 mmol) and maleimide (131 mg, 1.35 mmol) in xylene (20 mL) was refluxed for 4 days. After filtration, the solid residue was washed with water to give an isomeric mixture of the Diels–Alder adducts as a brown solid (254 mg, 0.86 mmol, 77% yield).

A solution of the isomeric mixture (60 mg, 0.205 mmol) in dioxane (7 mL) was refluxed for 21 days in the presence of trifluoroacetic acid (900 μL). After evaporation, EtOAc was added to the residue. The mixture was filtered off, and the solid was successively washed with saturated aqueous NaHCO_3 , water, and EtOAc to give **13** as a brown solid (32 mg, 0.110 mmol, 54% yield).

Mp >300 °C. IR (KBr) $\nu_{\text{C=O}}$ 1710, 1720, 1780 cm^{-1} , $\nu_{\text{N-H}}$ 3000–3500 cm^{-1} . Mass (ESI+) $[\text{M}+\text{H}]^+$ 292. HRMS (ESI+) $[\text{M}+\text{Na}]^+$ calcd for $\text{C}_{16}\text{H}_9\text{N}_3\text{O}_3\text{Na}$ 314.0542, found 314.0556.

^1H NMR (400 MHz, DMSO): 5.03 (2H, s), 7.41 (1H, t, $J=7.5$ Hz), 7.63 (1H, t, $J=7.5$ Hz), 7.78 (1H, d, $J=8.0$ Hz), 8.12 (1H, d, $J=8.0$ Hz), 8.76 (1H, s, NH), 11.21 (1H, s, NH), 12.46 (1H, s, NH). Due to its insolubility, the ^{13}C NMR spectrum could not be recorded.

4.4.24. 10-Methyl-2H,5H,7H-1,3,4,6-tetrahydro-dipyrrolo[3,4-*a*:3,4-*c*]carbazole-3,4,6-trione (14). A mixture of compound **5b** (120 mg, 0.53 mmol) and maleimide (61 mg, 0.62 mmol) in xylene (5 mL) was refluxed for 3 days. After filtration, the solid was washed with water and dried to give an isomeric mixture of the Diels–Alder adducts (161 mg, 0.52 mmol, 100% yield) as a brown solid.

A solution of the isomeric mixture (150 mg, 0.49 mmol) in dioxane (16 mL) was refluxed for 48 h in the presence of trifluoroacetic acid (2.2 mL). After evaporation, EtOAc was added to the residue. The mixture was filtered off to give **14** (75 mg, 0.24 mmol, 46% yield) as a brown solid.

Mp >300 °C. IR (KBr) $\nu_{\text{C=O}}$ 1719, 1770 cm^{-1} , ν_{NH} 3100–3550 cm^{-1} . HRMS (ESI+) $[\text{M}+\text{Na}]^+$ calcd for $\text{C}_{17}\text{H}_{11}\text{N}_3\text{O}_3\text{Na}$ 328.0698, found 328.0681.

^1H NMR (400 MHz, $\text{DMSO-}d_6$): 5.01 (2H, s), 7.45 (1H, d, $J=8.5$ Hz), 7.66 (1H, d, $J=8.0$ Hz), 7.92 (1H, s), 8.76 (1H, s, NH), 11.18 (1H, s, NH), 12.34 (1H, s, NH).

Due to its insolubility, the ^{13}C NMR spectrum could not be recorded.

4.4.25. 10-Benzyloxy-2H,5H,7H-1,3,3a,3b,4,6,6a,11c-octahydro-dipyrrolo[3,4-*a*:3,4-*c*]carbazole-3,4,6-trione (15). A mixture of **5c** (90 mg, 0.28 mmol) and maleimide (34 mg, 0.35 mmol) in xylene (7 mL) was refluxed for 48 h. After filtration, the solid was washed with water and dried to give **15** (102 mg, 0.26 mmol, 91% yield) as a brown-orange solid.

Mp >300 °C. IR (KBr) $\nu_{\text{C=O}}$ 1719, 1778 cm^{-1} , ν_{NH} 3100–3550 cm^{-1} . HRMS (ESI+) $[\text{M}+\text{H}]^+$ calcd for $\text{C}_{23}\text{H}_{18}\text{N}_3\text{O}_4$ 400.1297, found 400.1285.

^1H NMR (400 MHz, $\text{DMSO-}d_6$): 4.33–4.39 (2H, m), 4.59 (1H, d, $J=11.0$ Hz), 4.64 (1H, d, $J=17.5$ Hz), 5.16 (2H, s), 6.89 (1H, dd, $J_1=9.0$ Hz, $J_2=2.0$ Hz), 7.13 (1H, d, $J=2.0$ Hz), 7.36–7.52 (6H, m), 7.97 (1H, s, NH), 11.72 (1H, s, NH), 11.73 (1H, s, NH).

^{13}C NMR (100 MHz, $\text{DMSO-}d_6$): 45.2, 69.6 (CH_2), 42.3, 46.7 (CH), 102.4, 111.2, 113.1, 127.6 (2C), 127.7, 128.3 (2C) (C tert), 105.2, 115.7, 123.1, 132.1, 133.0, 137.4, 148.3, 153.4 (C quat), 172.3, 177.3, 177.5 (CO).

4.4.26. 10-Benzyloxy-2H,5H,7H-1,3,4,6-tetrahydro-dipyrrolo[3,4-*a*:3,4-*c*]carbazole-3,4,6-trione (16). A mixture of compound **15** (50 mg, 0.12 mmol) and DDQ (28 mg, 0.12 mmol) in dioxane (5 mL) was stirred overnight. After removal of the solvent, the residue was washed successively with water, EtOAc, and small amounts of THF. Compound **16** (19 mg, 0.05 mmol, 40% yield) was isolated as a brown solid.

Mp >300 °C. IR (KBr) $\nu_{\text{C=C}}$ 1617 cm^{-1} , $\nu_{\text{C=O}}$ 1722, 1774 cm^{-1} , ν_{NH} 3100–3500 cm^{-1} . Mass (ESI+) $[\text{M}+\text{H}]^+$ 398. HRMS (ESI+) $[\text{M}+\text{Na}]^+$ calcd for $\text{C}_{23}\text{H}_{15}\text{N}_3\text{O}_4\text{Na}$ 420.0960, found 420.0955.

^1H NMR (400 MHz, $\text{DMSO-}d_6$): 5.03 (2H, s), 5.29 (2H, s), 7.36–7.70 (8H, m), 8.79 (1H, s, NH), 11.19 (1H, s, NH), 12.33 (1H, s, NH).

Due to its insolubility, the ^{13}C NMR spectrum could not be recorded.

4.4.27. 10-Hydroxy-2H,5H,7H-1,3,3b,4,6,6a-hexahydro-dipyrrolo[3,4-*a*:3,4-*c*]carbazole-3,4,6-trione (17). A mixture of **5d** (120 mg, 0.52 mmol) and maleimide (61 mg, 0.62 mmol) in xylene (14 mL) was refluxed for 48 h. After filtration, the solid was washed with water and dried to give **17** (129 mg, 0.42 mmol, 81% yield) as an off-white solid.

Mp >300 °C. IR (KBr) $\nu_{\text{C=O}}$ 1708, 1778 cm^{-1} , $\nu_{\text{NH,OH}}$ 3200–3550 cm^{-1} . Mass (ESI+) $[\text{M}+\text{H}]^+$ 310, $[\text{M}+\text{Na}]^+$ 332. HRMS (ESI+) $[\text{M}+\text{Na}]^+$ calcd for $\text{C}_{16}\text{H}_{11}\text{N}_3\text{O}_4\text{Na}$ 332.0647, found 332.0664.

^1H NMR (400 MHz, $\text{DMSO-}d_6$): 4.32 (1H, d, $J=18.0$ Hz), 4.36 (1H, d, $J=11.0$ Hz), 4.54 (1H, d, $J=19.0$ Hz), 4.57 (1H, d, $J=11.0$ Hz), 6.69 (1H, dd, $J_1=9.0$ Hz, $J_2=2.0$ Hz),

6.84 (1H, d, $J=2.0$ Hz), 7.29 (1H, d, $J=9.0$ Hz), 7.88 (1H, s, NH), 8.96 (1H, s), 11.57 (1H, s, NH), 11.71 (1H, s, NH).

Due to its insolubility, the ^{13}C NMR spectrum could not be recorded.

4.5. Chk1 inhibitory assays

Human Chk1 full-length enzyme with an N-terminal GST sequence was either purchased from Upstate Biochemicals (No. 14-346) or purified from extracts of Sf9 cells infected with a baculovirus encoding GST-Chk1. Assays for compound testing were based upon the method described by Davies et al.²⁸

4.6. Growth inhibition assays

Tumor cells were provided by American Type Culture Collection (Frederik, MD, USA). They were cultivated in RPMI 1640 medium (Life Science technologies, Cergy-Pontoise, France) supplemented with 10% fetal calf serum, 2 mM L-glutamine, 100 units/mL penicillin, 100 $\mu\text{g}/\text{mL}$ streptomycin, and 10 mM HEPES buffer (pH=7.4). Cytotoxicity was measured by the microculture tetrazolium assay as described.²⁹ Cells were continuously exposed to graded concentrations of the compounds for four doubling times, then 15 μL of 5 mg/mL 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide was added to each well and the plates were incubated for 4 h at 37 °C. The medium was then aspirated and the formazan solubilized by 100 μL of DMSO. Results are expressed as IC_{50} , concentration, which reduced by 50% the optical density of treated cells with respect to untreated controls.

Acknowledgements

The authors are grateful to Bertrand Légeret, University Blaise Pascal, Clermont-Ferrand, for recording the mass spectra.

References and notes

1. Prudhomme, M. *Curr. Pharm. Des.* **1997**, *3*, 265–290.
2. Prudhomme, M. *Curr. Med. Chem.: Anti-Cancer Agents* **2004**, *4*, 509–521.
3. Knölker, H.-J.; Reddy, K. R. *Chem. Rev.* **2002**, *102*, 4303–4427.
4. Anderson, H. J.; Andersen, R. J.; Roberge, M. *Progress in the Cell Cycle Research*; Meijer, L., Jézéquel, A., Roberge, M., Eds.; Editions: Roscoff, France, 2003; Vol. 5, pp 423–430.
5. Roberge, M.; Berlinck, R. G. S.; Xu, L.; Anderson, H. J.; Lim, L. Y.; Curman, D.; Stringer, C. M.; Friend, S. H.; Davies, P.; Vincent, I.; Haggarty, S. J.; Kelly, M. T.; Britton, R.; Piers, E.; Andersen, R. J. *Cancer Res.* **1998**, *58*, 5701–5706.
6. Busby, E. C.; Leistritz, R. T.; Karnitz, L. M.; Sarkaria, J. N. *Cancer Res.* **2000**, *60*, 2108–2112.
7. Jackson, J. R.; Gilmartin, A.; Imburgia, C.; Winkler, J. D.; Marshall, L. A.; Roshak, A. *Cancer Res.* **2000**, *60*, 566–572.
8. Vermeulen, K.; Van Bockstaele, D. R.; Berneman, Z. N. *Cell Prolif.* **2003**, *36*, 131–149.
9. Cuddihy, A. R.; O'Connell, M. J. *Cytology* **2003**, *222*, 99–140.
10. McGowan, C. H. *Mutat. Res.* **2003**, *532*, 75–84.
11. Bartek, J.; Lukas, J. *Cancer Cell* **2003**, *3*, 421–429.
12. Carrassa, L.; Brogini, M.; Erba, E.; Damia, G. *Cell Cycle* **2004**, *3*, 1177–1181.
13. Zhao, B.; Bower, M. J.; McDevitt, P. J.; Zhao, H.; Davis, S. T.; Johanson, K. O.; Green, S. M.; Concha, N. O.; Zhou, B. B. S. *J. Biol. Chem.* **2002**, *277*, 46609–46615.
14. Jiang, X.; Zhao, B.; Britton, R.; Lim, L. Y.; Leong, D.; Sanghera, J. S.; Zhou, B.-B. S.; Piers, E.; Andersen, R. J.; Roberge, M. *Mol. Cancer Ther.* **2004**, *3*, 1221–1227.
15. Berlinck, R. G. S.; Britton, R.; Piers, E.; Lim, L.; Roberge, M.; Moreira da Rocha, R.; Andersen, R. J. *J. Org. Chem.* **1998**, *63*, 9850–9856.
16. Piers, E.; Britton, R.; Andersen, R. J. *J. Org. Chem.* **2000**, *65*, 530–535.
17. Yoshida, T.; Nishiyachi, M.; Nakashima, N.; Murase, M.; Kotani, E. *Chem. Pharm. Bull.* **2002**, *50*, 872–876.
18. Yoshida, T.; Nishiyachi, M.; Nakashima, N.; Murase, M.; Kotani, E. *Chem. Pharm. Bull.* **2003**, *51*, 209–214.
19. Terpin, A.; Winklhofer, C.; Schumann, S.; Steglich, W. *Tetrahedron* **1998**, *54*, 1745–1752.
20. Hugon, B.; Pfeiffer, B.; Renard, P.; Prudhomme, M. *Tetrahedron Lett.* **2003**, *44*, 3927–3930.
21. Hugon, B.; Pfeiffer, B.; Renard, P.; Prudhomme, M. *Tetrahedron Lett.* **2003**, *44*, 4607–4611.
22. Hugon, B.; Pfeiffer, B.; Renard, P.; Prudhomme, M. *Tetrahedron Lett.* **2003**, *44*, 3935–3937.
23. Hénon, H.; Messaoudi, S.; Hugon, B.; Anizon, F.; Pfeiffer, B.; Prudhomme, M. *Tetrahedron* **2005**, *61*, 5599–5614.
24. Hénon, H.; Anizon, F.; Kucharczyk, N.; Loynel, A.; Casara, P.; Pfeiffer, B.; Prudhomme, M. *Synthesis* **2006**, 711–715.
25. Mase, N.; Nishi, T.; Hiyoshi, M.; Ichihara, K.; Bessho, J.; Yoda, H.; Takabe, K. *J. Chem. Soc., Perkin Trans. 1* **2002**, 707–709.
26. Bergman, J.; Janosik, T.; Koch, E.; Pelcman, B. *J. Chem. Soc., Perkin Trans. 1* **2000**, 2615–2621.
27. Hénon, H.; Anizon, F.; Golsteyn, R. M.; Léonce, S.; Hofmann, R.; Pfeiffer, B.; Prudhomme, M. *Bioorg. Med. Chem.* **2006**, *14*, 3825–3834.
28. Davies, S. P.; Reddy, H.; Caivano, M.; Cohen, P. *Biochem. J.* **2000**, *351*, 95–105.
29. Léonce, S.; Pérez, V.; Casabianca-Pignède, M. R.; Anstett, M.; Bisagni, E.; Atassi, G. *Invest. New Drugs* **1996**, *14*, 169–180.