



(Neutrino indirect) detection of neutralino dark matter in (non) universals SUSY GUT models

E. Nezri

► To cite this version:

E. Nezri. (Neutrino indirect) detection of neutralino dark matter in (non) universals SUSY GUT models. COSMO-02 International Workshop on Particle Physics and the Early Universe, Sep 2002, Chicago, United States. pp.1-15, 2002. <in2p3-00021538>

HAL Id: in2p3-00021538

<http://hal.in2p3.fr/in2p3-00021538>

Submitted on 20 Feb 2003

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

(Neutrino Indirect) Detection of Neutralino Dark Matter in (non-)Universals SUSY GUT Models.

Emmanuel Nezri

Laboratoire de Physique Corpusculaire de Clermont-Ferrand : Theory
Centre de Physique des Particules de Marseille : Antares

- ⇒ [V.Bertin, E.N., J.Orloff, non-Universal models, hep-ph/0210034](#)
- ⇒ [V.Bertin, E.N., J.Orloff, hep-ph/0204135 accepted in EPJ C \(mSugra\)](#)

see also: J.L.Feng, K.T. Matchev, F. Wilczek, PRD63(01)040524
V. Barger, F. Halzen, D. Hooper, C. Kao, PRD65(02)075022
and: L. Bergström, J. Edsjö, P. Gondolo; PRD58(98)103519
G. Jungman, M. Kamionkowski, K.Griest, Phys. Rep. 267 (96)

Contents

Detecting cold dark matter (WIMPS) in neutrino telescopes

mSugra summary

non-Universality :

 Scalar sector

 Gaugino sector

Conclusion

DM indirect detection with a neutrino telescope: ingredients


- ★ A cold dark matter candidate: choose χ , lightest neutralino in Constrained MSSM
- ★ A relic density: depends on (co-)annihilation processes σ_A
- ★ A cosmic storage ring: to re-start annihilation, need to concentrate n_χ ; halo, clumps? too small for ν 's! Need big, nearby, heavy body with large capture rate (C) \Rightarrow depends on $\sigma_{\chi p}^{el}$. Sun: OK! Earth: small.
- ★ χ = Majorana fermion: can self-annihilate, limiting the total population N_χ :

$$\dot{N}_\chi = C - C_A N_\chi^2$$

Annihilation rate: $\Gamma_A = \frac{1}{2} C_A N_\chi^2 = \frac{C}{2} \tanh^2 \sqrt{C C_A t} \stackrel{eq}{\approx} \frac{C}{2}$ can be insensitive to σ_A

- ★ among decay products $\chi\chi \rightarrow b\bar{b}, t\bar{t}, WW, ZZ, \dots \rightarrow \nu + \dots$, only ν 's can escape the sun and reach a detector. $\Phi(E_\nu)$ depends on dominant annihilation channel
- ★ Cerenkov detector watches for ν 's converted into μ

DM indirect detection with a neutrino telescope: picture



Neutralino

SM \xrightarrow{SUSY} **MSSM**

- group $SU(3) \times SU(2) \times U(1)$
- 2 Higgs doublets : $\tan \beta = \frac{v_u}{v_d}$, 5 scalars : h, A, H, H^\pm
- R-parity conservation \rightarrow **stable LSP**
- $m_p \neq m_{\tilde{p}} \Rightarrow$ Soft breaking terms $\mathcal{L}_{\text{soft}}$

In the basis $(-i\tilde{B}, -i\tilde{W}^3, \tilde{H}_1^0, \tilde{H}_2^0)$:

$$M_\chi = \begin{pmatrix} M_1 & 0 & -m_Z c\beta sW & m_Z s\beta sW \\ 0 & M_2 & m_Z c\beta cW & -m_Z s\beta cW \\ -m_Z c\beta sW & m_Z c\beta cW & 0 & -\mu \\ m_Z s\beta sW & -m_Z s\beta cW & -\mu & 0 \end{pmatrix}$$

$$\chi = z_{11}\tilde{b} + z_{12}\tilde{W}^3 + z_{13}\tilde{H}_1^0 + z_{14}\tilde{H}_2^0$$

$$\text{gaugino fraction : } f_G = z_{11}^2 + z_{12}^2$$

$$\text{higgsino fraction : } f_H = z_{13}^2 + z_{14}^2$$

Parameters at GUT scale $\sim 2 \cdot 10^{16}$ GeV:

- ★ a common gaugino mass $m_{1/2}$
- ★ a common scalar mass m_0
- ★ a common trilinear coupling A_0
- ★ a common bilinear coupling B_0
- ★ Higgs parameter μ_0

+ Renormalization group equations and radiative ElectroWeak Symmetry Breaking:

$$\frac{1}{2} m_Z^2 = \frac{m_{H_d}^2|_{Q_{EW\text{SB}}} - m_{H_u}^2|_{Q_{EW\text{SB}}} \tan^2 \beta}{\tan^2 \beta - 1} - \mu^2|_{Q_{EW\text{SB}}}$$

achieved at $Q_{EW\text{SB}} \sim \sqrt{m_{\tilde{t}_1} m_{\tilde{t}_2}}$

⇒ Input parameters :

$$m_0, m_{1/2}, A_0, \tan \beta, \text{sgn}(\mu)$$

Advantages: REWSB, less free parameters, contact with acc. analyses, but also addressing CCB, Landau poles, high energy extrapoll.

Thanks to **Suspect** authors

<http://www.lpm.univ-montp2.fr:7082/kneur/suspect.html>

(study of the CMSSM with Suspect :

hep-ph/0107316, A. Djouadi, M. Drees, J.L. Kneur)

mSugra/CMSSM models

Composition of the lightest neutralino :

- ★ bino χ : for low m_0 , **RGE** drive

$$\begin{aligned} M_1|_{Q_{EW\text{SB}}} &= \frac{M_2|_{Q_{EW\text{SB}}}}{2} \\ &= 0.41 m_{1/2} \ll |\mu|_{Q_{EW\text{SB}}} \end{aligned}$$

- ★ mixed bino-higgsino χ : **EWSB**

$$\sim -m_{H_u}^2|_{Q_{EW\text{SB}}} - \mu^2|_{Q_{EW\text{SB}}} \quad \text{if} \quad \tan \beta \geq 5$$



for m_0 large (> 1000),

increasing $m_0 \Rightarrow m_{H_u}^2$ less negative $\Rightarrow \mu$ decreases $\Rightarrow \mu \sim M_1$

“Focus point region” hep-ph/9909334 Feng, Matchev, Moroi

Experiment sensitivities in the $(m_0, m_{1/2})$ plane

$A_0=0 ; \tan(\beta)=45 ; \mu > 0$



$$A_0 = 0 ; \tan \beta = 45 ; \mu > 0$$

Calculation with Suspect + Darksusy package :


<http://www.lpm.univ-montp2.fr:7082/kneur/suspect.html>

<http://www.physto.se/edsjo/darksusy/>

mSugra : Direct Detection Experiments vs Neutrino Telescopes


$$\mu \text{ flux}_{\odot} : m_{\chi}$$

mSugra with 5 GeV threshold vs neutrino Indirect Detection



$$\sigma_{\chi-p}^{scal} : m_{\chi}$$

mSugra models vs Direct Detection





Non-universal scalar soft terms

- ★ Sfermions : non-universality in sfermions matrices can lead to light third generation sfermions
→ **coannihilations** $\chi\tilde{\tau}$, $\chi\tilde{t}$ can modify the relic density but happen in region out of reach of detectors.
Detection imply “real” quarks mainly on u and d valence and due to their low Yukawa couplings RGE evolutions of the first and second generation squarks depend on gaugino soft masses → their masses can not be lowered by changing scalar soft terms to enhance $\sigma_{\chi-q}^{scal}$ and $\sigma_{\chi-q}^{spin}$ through the process $\chi q \xrightarrow{\tilde{q}} \chi q$.
- ★ Higgses : relax universality $m_{H_i}|_{M_{GUT}} = (1 + \delta_i)m_0$ → modify REWSB relation parameters , m_A , μ can change life.

non universal Higgs masses effects in the $(m_0'', m_{1/2})$ plane

$$m_{H_2}=m_0 ; m_{H_1}=0.5*m_0 ; A_0=0 ; \tan(\beta)=45 ; \mu > 0$$



$$m_{H_2} = m_0 ; m_{H_1} = 0.5m_0 ; A_0 = 0 ; \tan \beta = 45 ; \mu > 0$$

★ wider noREWSB region ★ $m_{A(H)} < m_{A(H)}|_{mSugra} \rightarrow$ good for relic density ($\chi\chi \xrightarrow{A} b\bar{b}$) and direct detection at low m_0 ($\chi q \xrightarrow{H} \chi q$) ★ but lower indirect detection \neq Barger et al PRD65 2002, due to the low Isajet value of μ at high m_0 (Allanach, Kraml, Porod, susy 02, hep-ph/0207314) which increases the higgsino fraction (?).

Free relations in gaugino mass parameters

$M_2|_{GUT}$ parameter

- ★ essentially modify neutralino composition (and slightly low energy values of fields without $SU(3)$ charge through RGE).

Increasing the **wino** component of the neutralino favours $\rightarrow \chi\chi \xrightarrow{x_1^+, x_2^0} W^+W^-, ZZ$ and enhance the annihilation cross section $\sigma_{\chi-\chi}^A$. the strong $\chi\chi_2^0$ and $\chi\chi_1^+$ coannihilations become active and $\Omega_\chi h^2$ **strongly** decreases.

- ★ larger coupling entering in $\sigma_{\chi-p}^{scal} \rightarrow$ increases the direct detection
- ★ favours neutralino annihilations into the hard W^+W^- spectrum \rightarrow increases the indirect detection muon fluxes
- ★ However, the relevant value of M_2 is very critical : = **fine-tuning** (except moduli decays giving wino in AMSB models. Moroi, Randall hep-ph/9906527).

$M_3|_{GUT}$ parameter

- ★ 1-loop RGE analyse $\rightarrow M_3|_{GUT}$ is the main parameter for non-universality in the MSSM (Kazakov, Moultaqa hep-ph/9912271).

$$(M_{soft}^{scal}|_{low})^2 = (M_{soft}^{scal}|_{GUT})^2 + c_3 f_3 + c_2 f_2 + c_1 f_1 + corrections$$

$$\text{with } f_i = \frac{(M_i^{GUT})^2}{b_i} \left(1 - \frac{1}{(1+b_i \alpha_0 t)^2} \right)$$

and c_3 **strongly dominant**.

$\rightarrow M_3|_{GUT} < m_{1/2}$ decreases $m_A, m_{H_2}^2, \mu$ (and $m_{\tilde{q}}$)

- ★ increase annihilation

$$\chi\chi \xrightarrow{A} b\bar{b} \quad (\chi\chi \xrightarrow{\tilde{f}} f\bar{f})$$

$\chi\chi \xrightarrow{Z} t\bar{t}, \chi\chi \xrightarrow{x_i^+, x_i} W^+W^-, ZZ \propto$ higgsino fraction


\rightarrow very **favourable** for $\Omega_\chi h^2$

Gain for detection :

- ★ direct detection : increase $\sigma_{\chi-p}^{scal}$ via $\chi q \xrightarrow{H} \chi q$ (m_H lower and $z_{11(2)} z_{13(4)}$ \nearrow)



- ★ indirect detection : increase capture $\sigma_{\chi-p}^{spin}$ via $\chi q \xrightarrow{Z} \chi q$ (higgsino fraction \nearrow) and annihilation $\chi\chi \rightarrow W^+W^-, ZZ, t\bar{t}$ with energetic neutrinos

Effects of $x = \frac{M_{3(2)}|_{GUT}}{m_{1/2}}$ on relic density and detection rate



$M_3|_{GUT}$ effect on experiment sensitivities in the $(m_0, m''_{1/2})$ plane



$A_0=0 ; \tan(\beta)=45 ; \mu > 0 ; M_3/m_{1/2} = 0.63$



$$\frac{M_3|_{GUT}}{m_{1/2}} = 0.63 ; \tan \beta = 45$$

$M_3|_{GUT}$ effect on experiment sensitivities in the $(m_0, m''_{1/2})$ plane

$A_0=0$; $\tan(\beta)=10$; $\mu > 0$; $M_3/m_{1/2} = 0.55$



$$\frac{M_3|_{GUT}}{m_{1/2}} = 0.55 ; \tan \beta = 10$$

Conclusion

RGE models OK with neutrino indirect detection :

- ★ very heavy scalars (beyond reach of LHC)
- ★ neutrino telescope signal in mSugra $\rightarrow m_{\chi^+} < 300 - 350$ GeV
- ★ $\Omega h^2 \sim 0.15$ can be accommodated for all “mSugra” points using $x = M_3|_{GUT}/m_{1/2} \sim 0.6(\pm 0.1) + \text{corrections}(m_0, \tan \beta, m_b)$ with a strong enhancement of detection rates improving the experiment possibilities.
- ★ Earth out of reach neutrino telescopes in those framework.
- ★ These models are compatible with the Standard Model value of $(g - 2)_\mu : a_\mu^{SUSY} \simeq 0$.