



Efficient basis selection for the computation of vibrational spectrum

Romain Garnier, Vincent Le Bris, Marc Odunlami, Didier Bégué, Isabelle Baraille, Olivier Coulaud

► To cite this version:

Romain Garnier, Vincent Le Bris, Marc Odunlami, Didier Bégué, Isabelle Baraille, et al.. Efficient basis selection for the computation of vibrational spectrum. 10th European Conference on Computational Chemistry, Aug 2015, Fulda, Germany. hal-01200748

HAL Id: hal-01200748

<https://hal.archives-ouvertes.fr/hal-01200748>

Submitted on 17 Sep 2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Efficient basis selection for the computation of vibrational spectrum

Romain Garnier^{1*}, Vincent Le Bris², Marc Odunlami², Didier Bégue², Isabelle Baraille², Olivier Coulaud¹

¹ INRIA Bordeaux, 33405 France

² IPREM Pau, 64000 France

*romain.garnier@inria.fr

We propose here an efficient method to define an approximation subspace to compute the first vibrational frequencies of the molecular Hamiltonian which are those of interest in the experimental results. In the Born approximation the Hamiltonian can be written as the sum of the kinetic energy and the potential energy function in normal coordinates [1] reflecting the relative movements of atoms within the molecule. When we keep the kinetic energy and the terms of order 2 of the potential energy, it brings up what is commonly called the Harmonic part of the Hamiltonian.

The eigen-states of this part are completely described by the multivariable Hermite functions that are the traditional solutions of the quantum harmonic oscillator [2]. These are a relevant choice to provide a basis for the calculation of the eigen-states of the Hamiltonian by the variational method. However the curse of dimensionality appears when processing of large molecules. For example, for a molecule to 8 atoms we have 18 normal coordinates and if we vary the degree of the basis functions from 0 to 6 in each direction the total number of basis functions is 7^{18} . Select a minimal basis to solve the eigen system is an important issue. Many current means such that the variation-perturbation theory [2] can target a relevant subspace for calculating a given number of vibrational frequencies of the Hamiltonian, but generally results are not as good as those we obtain with classical full Vibrational Configuration Interaction method [3]. Based on the same approach, we propose a new method to obtain comparable results to the full VCI method [3] with a much smaller number of basis vectors. For example we were able to divide by two the number of basis vectors needed to calculate the first 40 vibrational frequencies of the molecule H_2CO . Results on larger molecules that have unstable behavior will be presented.

[1] Wilson, E. B., Decius, J. C., & Cross, P. C. (1955). *Molecular Vibrations: The Theory of Infrared and Raman Vibrational Spectra*. Dover Publications.

[2] Sinano O. (1961). Variation-Perturbation Method for Excited States. *Phys. Rev.*, 122(2), 491–492.

[3] D. Bégue, N. Gohaud, C. Pouchan, P. Cassam-Chenai, and J. Liévin, “A comparison of two methods for selecting vibrational configuration interaction spaces on a

heptatomic system: Ethylene oxide,” *Journal of Chemical Physics*, vol. 127, 2007.