

Purdue University
Purdue e-Pubs

LARS Symposia

Laboratory for Applications of Remote Sensing

1-1-1977

Classifying Vegetative Cover with Landsat Digital Data, Great Dismal Swamp, Virginia and North Carolina

Patricia T. Gammon

Lurie J. Shima

Virginia Carter

Follow this and additional works at: http://docs.lib.purdue.edu/lars_symp

Gammon, Patricia T.; Shima, Lurie J.; and Carter, Virginia, "Classifying Vegetative Cover with Landsat Digital Data, Great Dismal Swamp, Virginia and North Carolina" (1977). *LARS Symposia*. Paper 233.
http://docs.lib.purdue.edu/lars_symp/233

This document has been made available through Purdue e-Pubs, a service of the Purdue University Libraries. Please contact epubs@purdue.edu for additional information.

Reprinted from

**Symposium on
Machine Processing of
Remotely Sensed Data**

June 21 - 23, 1977

The Laboratory for Applications of
Remote Sensing

Purdue University
West Lafayette
Indiana

IEEE Catalog No.
77CH1218-7 MPRSD

Copyright © 1977 IEEE
The Institute of Electrical and Electronics Engineers, Inc.

Copyright © 2004 IEEE. This material is provided with permission of the IEEE. Such permission of the IEEE does not in any way imply IEEE endorsement of any of the products or services of the Purdue Research Foundation/University. Internal or personal use of this material is permitted. However, permission to reprint/republish this material for advertising or promotional purposes or for creating new collective works for resale or redistribution must be obtained from the IEEE by writing to pubs-permissions@ieee.org.

By choosing to view this document, you agree to all provisions of the copyright laws protecting it.

CLASSIFYING VEGETATIVE COVER WITH LANDSAT DIGITAL DATA, GREAT DISMAL SWAMP, VIRGINIA AND NORTH CAROLINA

PATRICIA T. GAMMON

U.S. Geological Survey, Suffolk, Virginia

LURIE J. SHIMA

Goddard Space Flight Center, Greenbelt,
Maryland

VIRGINIA CARTER

U.S. Geological Survey, Reston, Virginia

Geometrically corrected and temporally registered Landsat digital data from April and February 1974 have been used to classify and map vegetative cover in the Great Dismal Swamp in Virginia and North Carolina. Cover classes were grouped into dominance types and subclasses according to the new U.S. Fish and Wildlife Service wetland classification system. Four data sets were analyzed on the Interactive Digital Image Manipulation System (IDIMS): (1) April data (2) February data (3) bands 5 and 7 from April and February and (4) all 8 bands from April and February.

Eighteen dominance types were established for the swamp on the basis of season, dominant canopy species and, in some deciduous classes, on type of understory. The subclasses and dominance types vary in number and composition according to season of data acquisition. Percentage estimates for the swamp subclasses were as follows: shallow water/benthos 2.2 percent, broad-leaved evergreen forested wetland 6.5 percent, needle-leaved evergreen forested wetland 16 percent, broad-leaved deciduous forested wetland 67.7 percent, narrow-leaved deciduous forested wetland 1 percent, broad-leaved evergreen shrub wetland 5 percent, and altered 1.6 percent.

IDIMS algorithms were used to choose 65 random, 10x10 pixel, sample plots and to generate the geometric coordinates (line/column; latitude/longitude) for these samples. The plots were transferred to 1:65,000-scale color infrared photographs of the swamp and each sample plot was photointerpreted for dominance type, subclass, and acreage. The photointerpretation results are being compared with IDIMS classification for each data set to measure the accuracy of classification. In addition, we are using the same sample plots to assess the accuracy of previous

classifications made with Landsat digital data from the same data sets and from different dates using IDIMS, Image 100, and Laboratory for Applications of Remote Sensing System (LARSYS).