

Purdue University Purdue e-Pubs

LARS Symposia

Laboratory for Applications of Remote Sensing

1-1-1980

Determination of Potentially Arable Land and Measurements of Non-Agricultural Uses for Nine Selected Areas in Africa

Irvin A. Goldblatt

Richard F. Hyde

Follow this and additional works at: http://docs.lib.purdue.edu/lars_symp

Goldblatt, Irvin A. and Hyde, Richard F., "Determination of Potentially Arable Land and Measurements of Non-Agricultural Uses for Nine Selected Areas in Africa" (1980). *LARS Symposia*. Paper 371. http://docs.lib.purdue.edu/lars_symp/371

This document has been made available through Purdue e-Pubs, a service of the Purdue University Libraries. Please contact epubs@purdue.edu for additional information.

Reprinted from

Symposium on

Machine Processing of

Remotely Sensed Data

and

Soil Information Systems

and

Remote Sensing and Soil Survey

June 3-6, 1980

Proceedings

The Laboratory for Applications of Remote Sensing

Purdue University West Lafayette Indiana 47907 USA

IEEE Catalog No. 80CH1533-9 MPRSD

Copyright © 1980 IEEE The Institute of Electrical and Electronics Engineers, Inc.

Copyright © 2004 IEEE. This material is provided with permission of the IEEE. Such permission of the IEEE does not in any way imply IEEE endorsement of any of the products or services of the Purdue Research Foundation/University. Internal or personal use of this material is permitted. However, permission to reprint/republish this material for advertising or promotional purposes or for creating new collective works for resale or redistribution must be obtained from the IEEE by writing to pubs-permissions@ieee.org.

By choosing to view this document, you agree to all provisions of the copyright laws protecting it.

DETERMINATION OF POTENTIALLY ARABLE LAND AND MEASUREMENTS OF NON-AGRICULTURAL USES FOR NINE SELECTED AREAS IN AFRICA

IRVIN A. GOLDBLATT AND RICHARD F. HYDE Butler University

The purpose of this study was to determine the percentage of total potentially arable land currently committed to non-agricultural human settlement uses in selected African agro-climatic zones.

Nine study areas equal in size to Landsat scenes, were selected by climatic zones as specified by the Food and Agriculture Organization, Rome, Italy. Soils information were fundamental to this project and were combined with climatic and crop suitability information for each study area. This information was manually converted to digital form for input to a computer-aided geographic information system, thereby creating nine independent data bases.

Landsat CCT data for these study areas were processed by computer to identify and map the extent of human settlement. These data were automatically input as files into the appropriate data bases using the registration and aggregation program LEVI, developed at the Holcomb Research Institute.

Potentially arable lands were identified within each Landsat scene, using FAO criteria, based upon soil and crop suitability within each climatic zone. Landsat digital data identified the extent and location of human settlement, and file manipulation techniques identified settlements occurring upon these potentially arable lands. The results of this study indicate that the amount of potentially arable land varied dramatically between the nine study areas, ranging from 778 km² in South Africa (.19% committed to human settlements) to 29,017 km² in Nigeria (2.1% committed to human settlement). ¹The research described in this paper was supported by the Food and Agriculture Organization of the United Nations under contract No. UNFPA/INT/75/P13-2/AGL.

CH1533-9/80/0000-0245 \$00.75 (c) 1980 IEEE

1980 Machine Processing of Remotely Sensed Data Symposium