

Purdue University
Purdue e-Pubs

LARS Symposia

Laboratory for Applications of Remote Sensing

1-1-1981

Biotic Contributions to the Global Carbon Cycle: The Role of Remote Sensing

R. A. Houghton

G. M. Woodwell

Follow this and additional works at: http://docs.lib.purdue.edu/lars_symp

Houghton, R. A. and Woodwell, G. M., "Biotic Contributions to the Global Carbon Cycle: The Role of Remote Sensing" (1981).
LARS Symposia. Paper 473.
http://docs.lib.purdue.edu/lars_symp/473

This document has been made available through Purdue e-Pubs, a service of the Purdue University Libraries. Please contact epubs@purdue.edu for additional information.

Reprinted from

Seventh International Symposium

Machine Processing of

Remotely Sensed Data

with special emphasis on

Range, Forest and Wetlands Assessment

June 23 - 26, 1981

Proceedings

Purdue University
The Laboratory for Applications of Remote Sensing
West Lafayette, Indiana 47907 USA

Copyright © 1981

by Purdue Research Foundation, West Lafayette, Indiana 47907. All Rights Reserved.

This paper is provided for personal educational use only,
under permission from Purdue Research Foundation.

Purdue Research Foundation

BIOTIC CONTRIBUTIONS TO THE GLOBAL CARBON CYCLE: THE ROLE OF REMOTE SENSING

R.A. HOUGHTON, G.M. WOODWELL

The Ecosystems Center
Marine Biological Laboratory
Woods Hole, Massachusetts

ABSTRACT

The CO₂ content of the atmosphere is increasing currently as a result of the combustion of fossil fuels and the oxidation of vegetation and soils associated with changes in the use of land. Prediction of the atmospheric CO₂ concentration in the future requires a better understanding of how important these land-use changes are currently and how important they have been in the past. In this paper we present an analysis of past changes in the terrestrial biota and soils of the earth. The analysis is based on rates of forest harvest and regrowth, rates of land conversion to agriculture, and on the changes in biomass and soil carbon that accompany these uses of land. The results of the analysis show that changes in land use have caused a net release of carbon to the atmosphere that until recently was larger than the release from combustion of fossil fuels. There is still a large uncertainty in the analysis, however, largely because of conflicting reports as to the current rate of disappearance of tropical forests. We outline the kinds of information needed to improve the analysis and believe that remote sensing is of use immediately in reducing the range of uncertainty by a factor of two to four.

I. INTRODUCTION: THE GLOBAL CO₂ PROBLEM

The atmospheric CO₂ concentration is rising as a result of the combustion of fossil fuels and the destruction of forests. The importance of the destruction of forests as a source of CO₂ is uncertain within a factor of five or more. The amount of CO₂ from fossil fuels is known within less than 50%. Existing tabulations of data on the harvest and transformation of forests to agricultural and grazing land, changes

that are important in the global CO₂ flux, could be improved greatly by use of satellite imagery. In this paper we examine methods currently being developed for this purpose.

The atmosphere contains about 0.03% CO₂, little more than a trace. The concentration has been increasing at an accelerating rate since the beginning of the century, possibly much longer. The best evidence for the increase has been produced at Mauna Loa in the Hawaiian Islands beginning in 1958 (Fig. 1), when C. D. Keeling started his famous program of measurements there and in the Antarctic.^{9,10} These measurements and many other more recent data show an increase in CO₂ concentration that appears to assure a doubling of the preindustrial concentration by 2040, possibly earlier.

The increase in CO₂ is important because CO₂, in contrast to the other major gases of the atmosphere, O₂ and N₂, absorbs infra-red energy. An increase in the amount of CO₂ will result in greater retention in the atmosphere of radiant heat from the sun and a higher equilibrium temperature for the earth as a whole. The rise in temperature is expected to be important, 2-4° C averaged for the earth as a whole for a doubling of the CO₂ in the atmosphere.^{19,22} The increase at the poles will be greater, as much as 10° C, or more.^{12,13} The transition threatens rapid changes in climate that will affect agricultural productivity globally. The problem is serious. A clearer definition of the problem requires better data and more detailed analysis, especially of the biotic segments of the cycle, which appear to be less well known than other segments.

The release of carbon annually from the combustion of fossil fuels is about 6×10^{15} g C. The accumulation in the

atmosphere. Decay of organic matter and wood products also introduce time lags into the analysis. In this analysis we have attempted to document changes in the use of land since 1700. We have used the reports of agricultural historians, geographers, and economists for the years prior to 1945 and regional and global surveys conducted by various organizations in more recent years. A more thorough discussion of the model, the data and sources used, and the results are given in Moore et al.¹⁴ We report here some of the major results.

The results show that there has been a net release of carbon from terrestrial systems worldwide for as long as we have documented changes in land use (Fig. 4). Individual regions, Europe and North America, for example, do not show this sustained release, but the global results are what concern us here. The major reason for the release of carbon through history has been the clearing of lands for permanent agriculture (Fig. 4). Although the area of forests harvested has been greater than the area permanently cleared during this time interval, the regrowth of harvested forests stores carbon, and the net flux from both harvest and regrowth is less than from harvest itself.




Figure 4. Net annual flux of carbon to the atmosphere as a result of clearing and cultivation of agricultural land (DLTAG), harvest and regrowth of forests (DLTFOR), oxidation of wood harvested or cleared from a site (DLTPOL), and clearing of land for pasture (DLTGRZ).

When the net fluxes of carbon from all land-use changes are summed, the net release in 1980 was 2.6×10^{15} g C. The carbon released from the combustion of fossil fuels was about 6×10^{15} g C in the same year, but appears until recently to have been less of an annual release than the release from the biota and soils (Fig. 5).




Figure 5. Net annual flux of carbon to the atmosphere as a result of changes in land use (solid curve) and as a result of the combustion of fossil fuels and production of cement.

The net flux of 2.6×10^{15} g C from terrestrial systems in 1980 is an estimate based on what appear to us to be the most reasonable sources of information. It may be more appropriate, however, to express the 1980 flux of carbon as a range of $1-5 \times 10^{15}$ g C. This range results from use of different assumptions and different sources of information in alternative analyses with the model. The widest range occurred as a result of one uncertainty: the rate at which tropical forests were converted to agriculture between 1950 and 1980. The low value of the range (Fig. 6) (1.0×10^{15} g C released in 1980) was obtained using changes in agricultural area reported by the Production Yearbook of the Food and Agricultural Organization of the United Nations. Several authors have pointed out the problems inherent in such an approach¹⁷ and it may be that the rates of agricultural expansion are greater than reported. For the high end of the range we used rates of tropical




Figure 6. Net annual flux of carbon to the atmosphere calculated from agricultural information obtained from FAO Production Yearbooks.

forest destruction reported by Myers (Fig. 7).^{15,16} It is not always clear from Myers' reports whether or not the forests grow back.




Figure 7. Net annual flux of carbon to the atmosphere calculated with information from Myers (1980a, b) on deforestation rates in the tropics.

If we are to understand the global

carbon cycle, we must continue to narrow the range of uncertainty in the magnitudes of fluxes. We have reached the point in our analyses where the largest uncertainty ($1-5 \times 10^{15} \text{ g C yr}^{-1}$) is caused by conflicting reports as to the area of tropical forests cleared for agriculture between 1950 and 1980. Were these areas converted to permanent agriculture or to shifting cultivation? How large an area was cleared? We need an objective method for measuring changes in area. Can remote sensing supply that need?

III. THE CHALLENGE FOR REMOTE SENSING

It is important to recall that the emphasis is on change; change in carbon stocks, change in area, and change in land use. Is it likely that we can use remote sensing to obtain these changes? Before answering the question it is useful to distinguish two different kinds of change: change in carbon per unit area and change in area. The annual rate of change we are interested in is of the order of $1.0 \times 10^{15} \text{ g C}$. The carbon content of the world's vegetation and soils is about $1500 \times 10^{15} \text{ g C}$. Thus we are looking for a change of 1 part in 1500, or 1-2 parts per thousand. The average carbon content of forests worldwide, including soils, is about 300 metric tons/ha (Table 1). If the annual change of 1 in 1500 were distributed equally over the forests of the earth, we would be looking for a change of about 0.2 mt C/ha in 300. There is no hope that remote sensing would detect such a change in less than 100 years. Direct sampling of forest biomass on the ground is unlikely to reveal the change either; the variability of plots measured within the same forest type is of the order of tens of mt/ha.^{7,21}

On the other hand if the change in the storage of terrestrial carbon is concentrated in areas where, say, all of the carbon in a forested hectare is released to the atmosphere, 1 part in 1500 amounts to an annual change of about $3.5 \times 10^6 \text{ ha}$ of forest (world forests occupy about $5000 \times 10^6 \text{ ha}$). Actually the minimum area is $5-7 \times 10^6 \text{ ha}$ since the carbon lost in the transformation of forests to cultivated land is about 50% of the carbon originally held in the forest. Thus the change in area sufficient to release $1 \times 10^{15} \text{ g C}$ is a change that can be measured with remote sensing. In fact the transformation of forests to agriculture and the reverse transformation are the kinds of change responsible for the major

Table 1. Average amount of carbon (mt/ha) in different forest types.

	Tropical moist forests	Tropical seasonal forests	Temperate evergreen forests	Temperate deciduous forests	Boreal forests
Vegetation ²³	200	160	160	135	90
Soil ¹⁸	117	117	134	134	206
Total	317	277	294	269	296

uncertainty in our current analysis of net carbon flux. The changes include the clearing of forest for agriculture, abandonment of agricultural land, and reforestation.

The current resolution of the carbon problem is such that a direct measure of the transformations of forests is sufficient to decrease our uncertainty by a factor of 2 to 4 - from a range of $1-5 \times 10^{15} \text{ g C yr}^{-1}$ to one of $1.5-3.0 \times 10^{15} \text{ g C yr}^{-1}$. At this point in the analyses, therefore, we need not classify cover types or land uses into more than two categories. Differences in carbon content between forest types, for example, is of the order of 10-50 mt/ha (Table 1) as opposed to 100-200 mt C/ha for forest to non-forest differences. The former differences are of secondary importance at this stage of our analysis. Furthermore, to the extent the techniques exist for detecting change between two satellite images without prior classification, the task of classification will be made still easier. Only those pixels, or groups of pixels, that register a change need classification. This approach requires that the registration of two images be to within several pixels at least; but to the extent that registration is a simpler problem to deal with than classification, the approach seems justified.

If LANDSAT is used in the detection of change, there are several kinds of information it will provide. As long as images are analyzed frequently enough to detect a clearing before it is covered again, imagery will yield when a change has occurred. As long as the registration of images is to within several pixels, the approach will provide where a change has occurred and how much area was changed. But will image analysis reveal what kind of change took place? Hopefully forest to non-forest and non-forest to forest transformations will be evident. But the response of vegetation and soils to logging is different from their response to clearing and cultivation. The analysis of two

images may reveal a cleared forest, but without a third image at a later date the analysis will not reveal whether the clearing was permanent or not. Neither will it be clear what exchanges of carbon should be assigned to that area of change. Our analysis, thus, has a further requirement.

We need to know and to remember where a change has occurred. There are two reasons why we need to be able to return to the same group of pixels with a subsequent image. First, we need to establish what the change was. Was it a harvest with regrowth or was it a more permanent clearing? Second, we want to know the carbon content of a forest when it is harvested or cleared a second time in the record of the imagery. If satellite imagery could be used to obtain values of biomass and soil carbon directly, the model would be unnecessary. But the development of a forest canopy, or leaf area index, (spectral response) is not proportional to the development of biomass and soil carbon (carbon content).^{6,4} Seven year-old pine stands look spectrally similar to 50 year-old stands (Barker, this symposium), yet their carbon contents are clearly different. As long as remote sensing cannot distinguish carbon differences associated with succession (Hicks and Park, this symposium), our analysis requires an independent source of data. We do not require that satellite imagery yield information on the biomass and soil carbon of a forest but only that it detect when and where a change has occurred. The model, based on ecological information, predicts how carbon in a hectare of regrowing forest will change with time. Imagery will provide the information to start the regrowing process; the model will calculate the changes in carbon per unit area. To be useful for our purpose, remote sensing must indicate when and where a change has occurred, and the methodology must provide a means of returning to that changed area on subsequent dates.

As this discussion represents work

in progress it is appropriate to end with a question. The global carbon problem requires a global approach and, hence, a global sampling scheme. The area of forests is approximately 5000×10^6 ha; the total area of land, about three times that. One satellite image covers about 2.2×10^6 ha, so complete coverage of the total land area once would require over 6000 frames. Including the frames required to cover land-water boundaries and the frames necessary to account for equipment failures, cloudiness, and other problems, the total number of frames is probably about 24,000 for one look. Change detection requires a minimum of two images for each time interval. Clearly there is a need for stratification, but on what basis? How many scenes must be analyzed to reduce our uncertainty to 1×10^{15} g C yr⁻¹? The questions are difficult, the task of sampling is formidable. But the development and practice of such a scheme would have countless uses in managing the resources of the earth.

IV. REFERENCES

1. Adams, J. A. S., M. S. M. Mantovani, and L. L. Lundell. 1977. Wood versus fossil fuel as a source of excess carbon dioxide in the atmosphere: A preliminary report. *Science* 196: 54-56.
2. Bacastow, R. and C. D. Keeling. 1973. Atmospheric carbon dioxide and radiocarbon in the natural carbon cycle: II. Changes from A.D. 1700 to 2070 as deduced from a geochemical model. Pp. 86-135 in: G. M. Woodwell and E. V. Pecan (eds.), *Carbon and the Biosphere*. National Technical Information Service, Springfield, VA.
3. Bolin, B. 1977. Changes of land biota and their importance for the carbon cycle. *Science* 196: 613-615.
4. Bormann, F. H., and G. E. Likens. 1979. *Pattern and process in a forested ecosystem*. Springer-Verlag, New York.
5. Broecker, W. S., T. Takahashi, H. J. Simpson and T. -H. Peng. 1979. Fate of fossil fuel carbon dioxide and the global carbon budget. *Science* 206: 409-418.
6. Covington, W. W., and J. D. Aber. 1980. Leaf production during secondary succession in northern hardwoods. *Ecology* 61: 200-204.
7. Edwards, P. J., and P. J. Grubb. 1977. Studies of mineral cycling in a montane rain forest in New Guinea: I. The distribution of organic matter in the vegetation and soil. *J. Ecol.* 65: 943-970.
8. Hampicke, U. 1979. Net transfer of carbon between the land biota and the atmosphere, induced by man. Pp. 219-236 in: B. Bolin, E. J. Degens, S. Kempe, and P. Ketner (eds.), *The Global Carbon Cycle*. SCOPE 13. John Wiley & Sons, New York.
9. Keeling, C. D., R. B. Bacastow, A. E. Bainbridge, C. A. Ekdahl Jr., P. R. Guenther, and L. S. Waterman. 1976 a. Atmospheric carbon dioxide variations at Mauna Loa Observatory, Hawaii. *Tellus* 28: 538-551.
10. Keeling, C. D., J. A. Adams, Jr., C. A. Ekdahl, Jr., and P. R. Guenther. 1976 b. Atmospheric carbon dioxide variations at the South Pole. *Tellus* 28: 552-564.
11. Machta, L. 1973. Prediction of CO₂ in the atmosphere. Pp. 21-31 in: G. M. Woodwell and E. V. Pecan (eds.), *Carbon in the Biosphere*. National Technical Information Service, Springfield, VA.
12. Manabe, S., and R. J. Stouffer. 1979. A CO₂-climate sensitivity study with a mathematical model of the global climate. *Nature* 282: 491-493.
13. Manabe, S., and R. T. Wetherald. 1980. On the distribution of climate change resulting from an increase in CO₂ content of the atmosphere. *J. Atmos. Sci.* 37: 99-118.
14. Moore, B., R. D. Boone, J. E. Hobbie, R. A. Houghton, J. M. Melillo, B. J. Peterson, G. R. Shaver, C. J. Vorosmarty and G. M. Woodwell. 1981. A simple model for analysis of the role of terrestrial ecosystems in the global carbon budget. In: B. Bolin (ed.), *Modelling the Global Carbon Cycle*, SCOPE 16. John Wiley & Sons, New York. (In press).
15. Myers, N. 1980a. Report of survey of conversion rates in tropical moist forests. National Research Council, Washington, D.C.

16. Myers, N. 1980b. The present and future prospects of tropical moist forests. *Environ. Conservation* 7: 101-114.
17. Persson, R. 1974. World Forest Resources: Review of the World's Forest Resources in the Early 1970's. Royal College of Forestry. Research Notes No. 17, Stockholm, Sweden.
18. Schlesinger, W. H. 1977. Carbon balance in terrestrial detritus. *Ann. Rev. Ecol. Syst.* 8: 51-81.
19. Schneider, S. H. 1975. On the carbon dioxide - climate confusion. *J. Atmos. Sci.* 32: 2060-2066.
20. Siegenthaler, U. and H. Oeschger. 1978. Predicting future atmospheric carbon dioxide levels. *Science* 199: 388-394.
21. Tanner, E. V. J. 1980. Studies on the biomass and productivity in a series of Montane Rain forests in Jamaica. *J. Ecol.* 68: 573-588.
22. Watts, R. G. 1980. Climate models and CO₂-induced climatic changes. *Climatic Change* 2: 387-408.
23. Whittaker, R. H. and G. E. Likens. 1973. Carbon in the biota. Pp. 281-302 in: G. M. Woodwell and E. V. Pecan (eds.), *Carbon and the Biosphere*. National Technical Information Service, Springfield, VA.
24. Wong, C. S. 1978. Atmospheric input of carbon dioxide from burning wood. *Science* 200: 197-200.
25. Woodwell, G. M., R. H. Whittaker, W. A. Reiners, G. E. Likens, C. C. Delwiche and D. B. Botkin. 1978. The biota and the world carbon budget. *Science* 199: 141-146.