

Tropical Fourier–Motzkin elimination, with an application to real-time verification

Xavier Allamigeon, Axel Legay, Uli Fahrenberg, Ricardo Katz, Stéphane Gaubert

► To cite this version:

Xavier Allamigeon, Axel Legay, Uli Fahrenberg, Ricardo Katz, Stéphane Gaubert. Tropical Fourier–Motzkin elimination, with an application to real-time verification. International Journal of Algebra and Computation, World Scientific Publishing, 2014, 24 (5), pp.569 - 607. 10.1142/S0218196714500258. hal-01087367

HAL Id: hal-01087367 https://hal.inria.fr/hal-01087367

Submitted on 26 Nov 2014

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Tropical Fourier-Motzkin Elimination, with an Application to Real-Time Verification*

Xavier Allamigeon

INRIA and CMAP, École Polytechnique, CNRS, 91128 Palaiseau Cédex, France xavier.allamigeon@inria.fr

> Uli Fahrenberg Irisa / INRIA, Campus Beaulieu, 35042 Rennes Cédex, France ulrich.fahrenberg@inria.fr

> > Stéphane Gaubert

INRIA and CMAP, École Polytechnique, CNRS, 91128 Palaiseau Cédex, France stephane.gaubert@inria.fr

Ricardo D. Katz CONICET-CIFASIS, Bv. 27 de Febrero 210 bis, 2000 Rosario, Argentina katz@cifasis-conicet.gov.ar

Axel Legay Irisa / INRIA, Campus Beaulieu, 35042 Rennes Cédex, France axel.legay@inria.fr

> Received (Day Month Year) Accepted (Day Month Year)

Communicated by [editor]

We introduce a generalization of tropical polyhedra able to express both strict and nonstrict inequalities. Such inequalities are handled by means of a semiring of germs (encoding infinitesimal perturbations). We develop a tropical analogue of Fourier-Motzkin elimination from which we derive geometrical properties of these polyhedra. In particular, we show that they coincide with the tropically convex union of (non-necessarily closed) cells that are convex both classically and tropically. We also prove that the

*The five authors were partially supported by the programme "Ingénierie Numérique & Sécurité" of the French National Agency of Research (ANR), project "MALTHY", number ANR-13-INSE-0003. Xavier Allamigeon was partially supported by the same programme, project "CAFEIN", number ANR-12-INSE-0007. Ricardo D. Katz was partially supported by CONICET Grant PIP 112-201101-01026. Xavier Allamigeon, Ricardo D. Katz and Stéphane Gaubert were partially supported by the PGMO program of EDF and Fondation Mathématique Jacques Hadamard. Uli Fahrenberg and Axel Legay were partially supported by the EU IP project DANSE and by the regional CREATE Estase initiative.

redundant inequalities produced when performing successive elimination steps can be dynamically deleted by reduction to mean payoff game problems. As a complement, we provide a coarser (polynomial time) deletion procedure which is enough to arrive at a simply exponential bound for the total execution time. These algorithms are illustrated by an application to real-time systems (reachability analysis of timed automata).

 $Keywords\colon$ Tropical polyhedra, strict inequalities, Fourier-Motzkin elimination, mean payoff games, real-time verification, timed automata

Mathematics Subject Classification 2010: 14T05, 52A01, 52B55

1. Introduction

1.1. Tropical convexity

Tropical or max-plus algebra refers to the set $\mathbb{R}_{\max} := \mathbb{R} \cup \{-\infty\}$ equipped with $x \oplus y := \max(x, y)$ as addition and $x \otimes y := x + y$ as multiplication (the latter will be also denoted by concatenation xy). In this setting, an inequality constraint on variables x_1, \ldots, x_n is said to be *(tropically) affine* if it is of the form:

$$a_0 \oplus a_1 \boldsymbol{x}_1 \oplus \dots \oplus a_n \boldsymbol{x}_n \leqslant b_0 \oplus b_1 \boldsymbol{x}_1 \oplus \dots \oplus b_n \boldsymbol{x}_n \quad (1.1)$$

or equivalently, with usual notation,

$$\max(a_0, a_1 + x_1, \dots, a_n + x_n) \leqslant \max(b_0, b_1 + x_1, \dots, b_n + x_n) , \qquad (1.2)$$

where $a_i, b_i \in \mathbb{R}_{\max}$ for i = 0, 1, ..., n. By analogy with the terminology of usual convex geometry, a *tropical (convex) polyhedron* is defined as a set composed of all the vectors $\boldsymbol{x} \in \mathbb{R}_{\max}^n$ satisfying finitely many such inequalities. An example is depicted on the left-hand side of Figure 1.

Tropical polyhedra and, more generally, tropically convex sets, have been introduced and studied in various contexts, including optimization [49], control theory [19], idempotent functional analysis [46], or combinatorics [23]. Several basic results of convex analysis and geometry have been shown to have tropical analogues. These include Hahn-Banach [49,46,20,23], Minkowski [29,15,39], and Carathéodory/Helly-type [13,34,32] theorems. Some algorithmic aspects have also been studied (*e.g.* [16,40,3]). We refer the reader to [3] for further references.

1.2. Motivation

The present work is motivated by a specific application of tropical algebra to the verification of real-time systems. Indeed, a remarkable property of tropical polyhedra is their ability to concisely encode possibly non-convex sets expressed as disjunctions of closed zones. A *closed zone*, also known in the literature as *polytrope*, is a set of vectors $\boldsymbol{x} \in \mathbb{R}^n$ defined by inequalities of the form $\boldsymbol{x}_i \ge m_i, \, \boldsymbol{x}_i \le M_i$, and $\boldsymbol{x}_i \le k_{ij} + \boldsymbol{x}_j$, for certain constants $m_i, M_i, k_{ij} \in \mathbb{R}$. More generally, zones are obtained by replacing some of the previous inequalities by strict ones. See the right-hand side of Figure 1 for an illustration.



Fig. 1. Left: a tropical polyhedron (including the black border). Right: a (non-closed) zone defined by the inequalities $1 \leq x_1 < 7$, $1 < x_2 \leq 5$, $-2 < x_1 - x_2 \leq 3$.

Zones are extensively used in the area of verification of real-time systems, where these systems are modelled by formalisms such as *e.g.* timed automata [8] or timed Petri nets [12]. More precisely, zones are used by model checking tools as *symbolic states*, typically representing infinitely many states of the system. They can be represented using difference-bound matrices (DBM), which are essentially adjacency matrices of weighted graphs. This allows for efficient algorithms for the manipulation of zones during the verification process.

An inherent drawback of zones is that they are convex sets, and consequently they are not closed under set union. This means that during the analysis process, symbolic states cannot generally be combined, which potentially leads to statespace explosion. Due to this, tropical polyhedra have been proposed in [47] as a replacement for zones. However, an important drawback in this approach is that the analysis of timed automata often requires to express strict constraints, for instance in the analysis of communication protocols [22,44], while tropical polyhedra are by definition topologically closed. An example illustrating these drawbacks will be given in Section 5.

1.3. Contributions

In this paper we first introduce (Section 2) a class of non-necessarily closed tropically convex sets. This class is called *tropical polyhedra with mixed constraints*. It can express not only inequalities of the form (1.2) in which the relation \leq has been replaced by <, but also finer constraints exploiting the disjunctive character of tropical inequalities. For instance, the inequality

$$\boldsymbol{x}_1 \leqslant \max(1 + \boldsymbol{x}_2, 0^- + \boldsymbol{x}_3)$$

is going to represent the disjunction of the inequalities $x_1 \leq 1 + x_2$ and $x_1 < x_3$. These *mixed inequalities* are defined using coefficients in a semiring of affine germs, which represent infinitesimal perturbations of reals.

In the second place, we present a tropical counterpart of Fourier-Motzkin elimination (Section 3). It provides a constructive method to show that the projection in \mathbb{R}_{\max}^{n-1} of a tropical polyhedron with mixed constraints $\mathcal{P} \subset \mathbb{R}_{\max}^n$ is a polyhedron with mixed constraints (Theorem 3.4). It computes a representation by mixed inequalities of the projection by combining the defining inequalities of \mathcal{P} . Actually, this approach handles more generally systems of inequalities with coefficients in a totally ordered idempotent semiring, modulo some assumptions. Note that such an analogue of Fourier-Motzkin algorithm has not been considered previously in the tropical setting, even in the case of standard (closed) tropical polyhedra. Fourier-Motzkin elimination also appears as a useful tool to show the polyhedral character of some non-closed tropically convex sets. As an application, we indeed prove that tropical polyhedra with mixed constraints are precisely the tropically convex union of finitely many zones, and the intersection of finitely many tropical hemispaces (*i.e.* tropically convex sets whose complements are also tropically convex, which were studied in *e.g.* [14,42]), see Theorem 3.6 and Corollary 3.10.

Superfluous inequalities may be produced by Fourier-Motzkin algorithm, so that the size of the constraint systems can grow in a double exponential way during consecutive applications of the method. In order to eliminate such redundant inequalities, in Section 4 we extend to mixed inequalities a result of [1] and its subsequent refinement in [5], building on techniques of these two papers. The result of [1] shows that deciding the feasibility of a system of tropical linear inequalities is (Karp) polynomial-time equivalent to solving mean payoff games. The result of [5] shows that deciding logical implications over tropical linear inequalities is also equivalent to solving mean payoff games. Theorem 4.7 generalizes these two results to mixed inequalities. We note that the present approach (through germs) also yields an alternative, simpler derivation of the result of [5]. Indeed, deciding whether a given inequality of the form (1.2) is logically implied by a system of other inequalities of the same kind amounts to checking if the intersection of a tropical polyhedron with the complement of a closed half-space is empty or not. Such an intersection is obviously a tropical polyhedron with mixed constraints.

Experimentally efficient algorithms have been developed to solve mean payoff games, but no polynomial time algorithm is known. Hence we also provide a weak criterion which allows to eliminate some of the superfluous inequalities in polynomial time (Section 4.2). We prove that, in the case of non-strict inequalities, this weak elimination is sufficient to obtain a *single*-exponential bound for Fourier-Motzkin elimination (Section 4.3).

Finally, Section 5 illustrates the application of tropical polyhedra with mixed constraints to the verification (reachability analysis) of timed automata. We show that the operations necessary for forward exploration of timed automata can be defined on tropical polyhedra with mixed constraints, using Fourier-Motzkin elimination and the algorithms developed to eliminate redundant inequalities.

Tropical Fourier-Motzkin Elimination, with an Application to Real-Time Verification 5

1.4. Related work

The algorithms developed so far for tropical polyhedra usually benefit from the fact that these can be represented either *externally*, using inequalities (as in (1.1)), or *internally*, as sets generated by finitely many points and rays (see [30] for details). In contrast, non-closed tropically convex sets may not be finitely generated. Generating representations of (non-necessarily closed) tropical convex cones have been studied in [15], and in [42] in the particular case of tropical hemispaces. A certain class of possibly infinite generating representations was treated in [28], however, the associated algorithms rely on the expensive Presburger arithmetic. Defining non-closed polyhedra using infinitesimal perturbations of generators also presents some difficulties, see Remark 2.6 below. Moreover, we should warn the reader that some geometric aspects of tropical polyhedra, in particular the notion of faces, are still not yet understood [24]. Thus it does not seem easy to manipulate non-closed polyhedra from closed ones by excluding some "facets" or "edges".

The present tropical Fourier-Motzkin algorithm may be thought of as a dual of the tropical double description method [3], in which one successively eliminates inequalities rather than variables. In both algorithms, redundant intermediate data (inequalities or generators) are produced, and the key to the efficiency of the algorithm lies in the dynamic elimination of such data. Redundant generators can be eliminated in almost linear time using a combinatorial hypergraph algorithm, however the hypergraph criterion appears to have no natural dual analogue which can detect redundant inequalities.

As mentioned above, the equivalence between mean payoff games and the emptiness problem for tropical polyhedra with mixed constraints generalizes a result of [1]. Moreover, it generalizes an earlier result [48] concerning finite solutions of a class of non-strict disjunctive constraints appearing in scheduling. The non-strict inequality satisfiability problem has also been studied under the name of "max-atom problem" in [10], with motivations from SMT solving. Note here a fundamental difference between strict and non-strict constraints: in the latter case, for inequalities with integer coefficients, it is shown in [1] that emptiness over the integers is equivalent to emptiness over the reals. The same is not true for strict inequalities (consider for example the open hypercube $]0, 1[^n$ which is non-empty, but contains no integer points), so that the present result for mixed inequalities cannot be deduced from earlier ones.

The infinitesimal perturbation of reals used in mixed inequalities is based on a semiring of affine germs, which was used in [27,25] to provide policy iteration based methods to solve mean payoff games. It also appeared in the context of tropical linear programming, see $[31, \S 3.7]$. The idea here is that germs allow one to determine algebraically the value of a perturbed game. Related perturbation or parametric game ideas were used in [5,35].

2. Tropical Polyhedra with Mixed Constraints

In the semiring $(\mathbb{R}_{\max}, \oplus, \otimes)$, addition and multiplication admit neutral elements, namely $-\infty$ and 0. Addition does not generally admit inverses. In contrast, any nonzero (in the tropical sense) element x admits a multiplicative inverse, which is given by -x and will be denoted x^{-1} . The semiring operations are extended to vectors and matrices in the usual way, *i.e.* $(A \oplus B)_{ij} := A_{ij} \oplus B_{ij}$ and $(A \otimes B)_{ij} := \bigoplus_k (A_{ik} \otimes B_{kj})$. We will work in the semimodule \mathbb{R}^n_{\max} , for $n \in \mathbb{N}$. Its elements can be seen as points or vectors and are denoted x, y, etc. The order \leq on \mathbb{R}_{\max} is extended to vectors entry-wise. We equip \mathbb{R}_{\max} with the topology induced by the metric $(x, y) \mapsto |\exp x - \exp y|$, and \mathbb{R}^n_{\max} with the product topology. In the sequel, we also use the completed max-plus semiring $\overline{\mathbb{R}}_{\max} := \mathbb{R}_{\max} \cup \{+\infty\}$, with the conventions $x < +\infty$ for all $x \in \mathbb{R}_{\max}, x \otimes (+\infty) = +\infty$ if $x \neq -\infty$, and $(-\infty) \otimes (+\infty) = -\infty$. Finally, given a positive integer n, we denote by [n] the set $\{1, \ldots, n\}$.

The notion of convexity can be transposed to tropical algebra. A subset C of \mathbb{R}^n_{\max} is said to be *(tropically) convex* if it contains the *tropical segment*

$$\{\lambda \boldsymbol{x} \oplus \mu \boldsymbol{y} \mid \lambda, \mu \in \mathbb{R}_{\max}, \ \lambda \oplus \mu = 0\}$$

joining any two points x and y of C. This is analogous to the usual definition of convexity, except that in the tropical setting the non-negativity constraint on λ and μ is implicit (any scalar $x \in \mathbb{R}_{\max}$ satisfies $x \ge -\infty$).

We now introduce the algebraic structure which will allow us to handle possibly strict tropical inequalities. We use a disjoint copy \mathbb{R}^- of \mathbb{R} composed of elements denoted α^- for $\alpha \in \mathbb{R}$, and we set $\mathbb{G} := \overline{\mathbb{R}}_{\max} \cup \mathbb{R}^-$. The *modulus* |x| of an element $x \in \mathbb{G}$ is defined by:

$$|x| := \begin{cases} x & \text{if } x \in \overline{\mathbb{R}}_{\max} ;\\ \alpha & \text{if } x = \alpha^- \in \mathbb{R}^- \end{cases}$$

The set \mathbb{G} is totally ordered by the order relation $\leq_{\mathbb{G}}$ defined by:

$$x \leq_{\mathbb{G}} y \Longleftrightarrow \begin{cases} |x| < |y| & \text{if } x \in \overline{\mathbb{R}}_{\max} \text{ and } y \in \mathbb{R}^- \\ |x| \leq |y| & \text{otherwise.} \end{cases}$$

We use the notation $x \leq_{\mathbb{G}} y$ when $x \leq_{\mathbb{G}} y$ and $x \neq y$. As an illustration, the Hasse diagram of $\leq_{\mathbb{G}}$ over the elements with modulus in $\mathbb{Z} \cup \{\pm \infty\}$ is given in Figure 2.

The element α^- can be interpreted as an infinitesimal perturbation of α of the form $\alpha - \varepsilon$ with $\varepsilon > 0$. Formally, given $x \in \mathbb{G}$ and $\varepsilon > 0$, the valuation of x at ε , denoted by $x(\varepsilon)$, is the element of $\overline{\mathbb{R}}_{\max}$ defined as follows:

$$x(\varepsilon) := \begin{cases} x & \text{if } x \in \overline{\mathbb{R}}_{\max} ;\\ |x| - \varepsilon & \text{if } x \in \mathbb{R}^- . \end{cases}$$

The valuation is extended to vectors and matrices entry-wise.

The set \mathbb{G} has a semiring structure when equipped with the sum of two elements $x, y \in \mathbb{G}$ defined as the greatest element among them, and the multiplication given



Fig. 2. Hasse diagram of the order $\leq_{\mathbb{G}}$ over the elements of \mathbb{G} with modulus in $\mathbb{Z} \cup \{\pm \infty\}$.

by:

$$\begin{cases} x \otimes y & \text{if } x, y \in \mathbb{R} ; \\ (|x| \otimes |y|)^{-} & \text{if } x \in \mathbb{R}^{-} \text{ or } y \in \mathbb{R}^{-}, \text{ and } x, y \neq \pm \infty ; \\ -\infty & \text{if } x = -\infty \text{ or } y = -\infty ; \\ +\infty & \text{if } x, y \neq -\infty, \text{ and } x = +\infty \text{ or } y = +\infty . \end{cases}$$

By abuse of notation, the multiplication in \mathbb{G} will be simply denoted by concatenation and the sum by \oplus , as in the case of \mathbb{R}_{\max} . Observe that in \mathbb{G} the neutral elements are still $-\infty$ and 0, and that only the elements $x \in \mathbb{R}$ are invertible with respect to multiplication. Also note that the modulus map is a semiring morphism.

We begin with a technical lemma on the arithmetic operations in the semiring $\mathbb{G}.$

Lemma 2.1. The following properties hold:

- (i) for any $x, y \in \mathbb{G}$, $x \leq_{\mathbb{G}} y$ if, and only if, $x(\varepsilon) \leq y(\varepsilon)$ for $\varepsilon > 0$ sufficiently small;
- (ii) for any $x, y \in \mathbb{G}$ and $\varepsilon > 0$ sufficiently small, $(x \oplus y)(\varepsilon) = x(\varepsilon) \oplus y(\varepsilon)$;
- (iii) for any $x \in \mathbb{R}_{\max}$, $y \in \mathbb{G}$, and $\varepsilon > 0$, $(xy)(\varepsilon) = xy(\varepsilon)$;
- (iv) for any $x, y, z \in \mathbb{G}$, $x \leq_{\mathbb{G}} y$ implies $xz \leq_{\mathbb{G}} yz$, and the converse holds if $z \in \mathbb{R}$;
- (v) for any $x, y \in \mathbb{R}_{\max}$, x < y is equivalent to $x \leq_{\mathbb{G}} 0^- y$ when $x \in \mathbb{R}$, and to $0 \leq_{\mathbb{G}} (+\infty)y$ when $x = -\infty$.

Proof. (i) The only non-trivial case is when $x \in \mathbb{R}$ and $y \in \mathbb{R}^-$, so assume we are in this case. Then, $x \leq_{\mathbb{G}} y$ amounts to x < |y|. This is equivalent to $x(\varepsilon) = x \leq |y| - \varepsilon = y(\varepsilon)$ for $\varepsilon > 0$ sufficiently small.

(ii) Straightforward from Property (i).

(iii) This property readily follows from the definition of the multiplication.

(iv) If $z = \pm \infty$, the first property is straightforward, so assume $|z| \in \mathbb{R}$. Let $x <_{\mathbb{G}} y$ (the implication is trivial when x = y). If |x| < |y|, we have |xz| < |yz| because $z \neq \pm \infty$. Moreover, if $|x| = |y| =: \alpha$, then $\alpha \neq \pm \infty$ (as x and y are distinct), and thus necessarily $x = \alpha^-$ and $y = \alpha$ (because $x <_{\mathbb{G}} y$). Hence $xz \in \mathbb{R}^-$ and |xz| = |yz|. In both cases, we conclude that $xz \leq_{\mathbb{G}} yz$.

Conversely, assume that $z \in \mathbb{R}$. Using the first part of the proof, $xz \leq_{\mathbb{G}} yz$ implies $x \leq_{\mathbb{G}} y$ by multiplying both sides of the inequality $xz \leq_{\mathbb{G}} yz$ by z^{-1} .

(v) In the first place, we suppose that $x \in \mathbb{R}$. If x < y, then $y \in \mathbb{R}$ and $0^-y = y^-$. Thus, x < y implies $x \leq_{\mathbb{G}} y^-$, *i.e.* $x \leq_{\mathbb{G}} 0^-y$. Conversely, if the latter inequality holds, then y is distinct from $-\infty$. Thus, $0^-y = y^-$, and $x \leq_{\mathbb{G}} y^-$ ensures that x < y.

Assume now that $x = -\infty$. Note that $(+\infty)y$ is equal to $+\infty$ if $y \neq -\infty$, and to $-\infty$ otherwise. Thus, we have $0 \leq_{\mathbb{G}} (+\infty)y$ if, and only if, $y > -\infty = x$.

A mixed tropical affine inequality is defined as a constraint of the form

$$a_0 \oplus a_1 \boldsymbol{x}_1 \oplus \cdots \oplus a_n \boldsymbol{x}_n \leqslant_{\mathbb{G}} b_0 \oplus b_1 \boldsymbol{x}_1 \oplus \cdots \oplus b_n \boldsymbol{x}_n \quad (2.1)$$

where the coefficients a_i on the left-hand side belong \mathbb{R}_{\max} , while the coefficients b_i on the right-hand side are in \mathbb{G} . When the set of \boldsymbol{x} in \mathbb{R}^n_{\max} satisfying a mixed inequality is a non-empty proper subset of \mathbb{R}^n_{\max} , it is called *mixed half-space*.

Lemma 2.2. A vector $\mathbf{x} \in \mathbb{R}^n_{\max}$ satisfies (2.1) if, and only if, there exists $\varepsilon > 0$ such that

$$a_0 \oplus a_1 \boldsymbol{x}_1 \oplus \cdots \oplus a_n \boldsymbol{x}_n \leqslant b_0(\varepsilon) \oplus b_1(\varepsilon) \boldsymbol{x}_1 \oplus \cdots \oplus b_n(\varepsilon) \boldsymbol{x}_n$$
 . (2.2)

Proof. If (2.1) is satisfied, then by Property (i) of Lemma 2.1 we have

 $a_0 \oplus a_1 \boldsymbol{x}_1 \oplus \cdots \oplus a_n \boldsymbol{x}_n \leq (b_0 \oplus b_1 \boldsymbol{x}_1 \oplus \cdots \oplus b_n \boldsymbol{x}_n)(\varepsilon)$

for $\varepsilon > 0$ sufficiently small, since the left-hand side of (2.1) belongs to \mathbb{R}^n_{\max} . Besides, by Properties (ii) and (iii) of Lemma 2.1, it follows that

$$(b_0 \oplus b_1 \boldsymbol{x}_1 \oplus \cdots \oplus b_n \boldsymbol{x}_n)(\varepsilon) = b_0(\varepsilon) \oplus b_1(\varepsilon) \boldsymbol{x}_1 \oplus \cdots \oplus b_n(\varepsilon) \boldsymbol{x}_n$$

for $\varepsilon > 0$ sufficiently small, which shows that (2.2) holds.

Conversely, suppose that (2.2) is satisfied for some $\varepsilon > 0$. Then

$$a_0 \oplus a_1 \boldsymbol{x}_1 \oplus \cdots \oplus a_n \boldsymbol{x}_n \leqslant b_0(arepsilon') \oplus b_1(arepsilon') \boldsymbol{x}_1 \oplus \cdots \oplus b_n(arepsilon') \boldsymbol{x}_n$$

for any $\varepsilon' < \varepsilon$ (the map $\varepsilon' \mapsto b_0(\varepsilon') \oplus b_1(\varepsilon') \boldsymbol{x}_1 \oplus \cdots \oplus b_n(\varepsilon') \boldsymbol{x}_n$ is non-increasing). It follows that (2.1) holds, by Properties (i), (ii) and (iii) of Lemma 2.1 and the fact that the left-hand side of (2.1) belongs to \mathbb{R}_{\max} .

A tropical polyhedron with mixed constraints is defined as a set composed of the vectors $\boldsymbol{x} \in \mathbb{R}^n_{\max}$ which satisfy finitely many mixed tropical affine inequalities.

To contrast with, we use the term *closed tropical polyhedron* when the defining mixed inequalities only involve coefficients in \mathbb{R}_{\max} , *i.e.* they are of the form (1.1). The following proposition establishes that polyhedra with mixed constraints are (possibly non-closed) tropically convex sets.

Proposition 2.3. Any tropical polyhedron with mixed constraints is a tropically convex set.

Proof. Let x, y be two solutions of (2.1), and $\lambda, \mu \in \mathbb{R}_{\max}$ be such that $\lambda \oplus \mu = 0$. By Lemma 2.2, there exist $\varepsilon, \varepsilon' > 0$ such that:

$$a_0 \oplus a_1 \boldsymbol{x}_1 \oplus \cdots \oplus a_n \boldsymbol{x}_n \leqslant b_0(\varepsilon) \oplus b_1(\varepsilon) \boldsymbol{x}_1 \oplus \cdots \oplus b_n(\varepsilon) \boldsymbol{x}_n ,$$

$$a_0 \oplus a_1 \boldsymbol{y}_1 \oplus \cdots \oplus a_n \boldsymbol{y}_n \leqslant b_0(\varepsilon') \oplus b_1(\varepsilon') \boldsymbol{y}_1 \oplus \cdots \oplus b_n(\varepsilon') \boldsymbol{y}_n .$$

These inequalities are still valid if we replace ε and ε' by min $(\varepsilon, \varepsilon')$. Hence, we can assume, without loss of generality, that $\varepsilon = \varepsilon'$. Then, $\boldsymbol{z} = \lambda \boldsymbol{x} \oplus \mu \boldsymbol{y}$ satisfies

 $a_0 \oplus a_1 \boldsymbol{z}_1 \oplus \cdots \oplus a_n \boldsymbol{z}_n \leqslant b_0(\varepsilon) \oplus b_1(\varepsilon) \boldsymbol{z}_1 \oplus \cdots \oplus b_n(\varepsilon) \boldsymbol{z}_n$,

which proves that z is a solution of (2.1) by Lemma 2.2. Thus, any mixed halfspace is tropically convex. We conclude that every tropical polyhedron with mixed constraints is tropically convex, as the intersection of finitely many tropically convex sets.

Example 2.4. The vectors $\boldsymbol{x} \in \mathbb{R}^2$ satisfying the strict inequality $\boldsymbol{x}_1 < \max(-1 + \boldsymbol{x}_2, 0)$, depicted on the left-hand side of Figure 3, are obtained as the real solutions of the mixed affine inequality $\boldsymbol{x}_1 \leq_{\mathbb{G}} (-1)^- \boldsymbol{x}_2 \oplus 0^-$. Similarly, the solutions of $\boldsymbol{x}_1 \leq_{\mathbb{G}} (-1)^- \boldsymbol{x}_2 \oplus 0$ correspond to the previous set in which the half-line $\{(0, \lambda) \mid \lambda \leq 1\}$ is added (middle of Figure 3).

The set depicted on the right-hand side of Figure 3 is the tropical polyhedron with mixed constraints defined by the following mixed inequalities:

$$(-2)\mathbf{x}_{2} \leq_{\mathbb{G}} 0^{-} \oplus 0^{-} \mathbf{x}_{1} \qquad -2 \leq_{\mathbb{G}} \mathbf{x}_{2}$$

$$-3 \leq_{\mathbb{G}} \mathbf{x}_{1} \qquad \mathbf{x}_{1} \leq_{\mathbb{G}} 3^{-} \mathbf{x}_{2} \qquad (2.3)$$

$$0 \leq_{\mathbb{G}} 1\mathbf{x}_{1} \oplus 0^{-} \mathbf{x}_{2} \qquad (-2)\mathbf{x}_{1} \leq_{\mathbb{G}} 0^{-} \oplus (-1)\mathbf{x}_{2}$$

Observe that the inequalities in (2.3) have the property that no variable (or constant term) ever appears on both the left- and right-hand sides. The following lemma ensures that this situation is not restrictive, and that in any inequality of the form (2.1), we can always assume $a_i = -\infty$ or $b_i = -\infty$ for all $i \in \{0, \ldots, n\}$.

Lemma 2.5. Let $a, b \in \mathbb{R}_{\max}$ and $c, d \in \mathbb{G}$. The set of solutions in \mathbb{R}_{\max} of the mixed tropical affine inequality $ax \oplus b \leq_{\mathbb{G}} cx \oplus d$ is given by $\{x \in \mathbb{R}_{\max} \mid b \leq_{\mathbb{G}} cx \oplus d\}$ if $a \leq_{\mathbb{G}} c$, and by $\{x \in \mathbb{R}_{\max} \mid ax \oplus b \leq_{\mathbb{G}} d\}$ otherwise.

Proof. In the first place, suppose that $a \leq_{\mathbb{G}} c$. Then, $ax \leq_{\mathbb{G}} cx$ for all $x \in \mathbb{R}_{\max}$ by Property (iv) of Lemma 2.1. It follows that $ax \leq_{\mathbb{G}} cx \oplus d$ is always satisfied.



Fig. 3. Tropical polyhedra with mixed constraints (the ends of the black segments marked by points are included in the polyhedra).

Suppose now that $a >_{\mathbb{G}} c$. If $x \in \mathbb{R}$, we have $ax \oplus b \ge_{\mathbb{G}} ax >_{\mathbb{G}} cx$ by Property (iv) of Lemma 2.1. As a consequence, any $x \in \mathbb{R}_{\max}$ such that $ax \oplus b \leqslant_{\mathbb{G}} cx \oplus d$ also satisfies $ax \oplus b \leqslant_{\mathbb{G}} d$. This completes the proof.

Remark 2.6. Following the analogy with closed tropical polyhedra, we could also consider subsets of \mathbb{G}^n generated by finitely many points and rays (given by vectors with entries in \mathbb{G}), having in mind to encode non-closed subsets of \mathbb{R}^n_{\max} thanks to infinitesimal perturbation of generators. More precisely, we could consider that a subset \mathcal{C} of \mathbb{G}^n encodes the subset $\widetilde{\mathcal{C}}$ of \mathbb{R}^n_{\max} given by the points $\boldsymbol{y} \in \mathbb{R}^n_{\max}$ such that for all $\varepsilon > 0$ there exist $\varepsilon' \in]0, \varepsilon[$ and $\boldsymbol{x} \in \mathcal{C}$ verifying $\boldsymbol{y} = \boldsymbol{x}(\varepsilon')$. However, this approach presents several difficulties. For example, consider the segment \mathcal{S} of \mathbb{G}^2 joining the points $\boldsymbol{v} = (0^-, 1)$ and $\boldsymbol{w} = (1^-, 1)$, *i.e.*

$$\mathcal{S} = \{\lambda \boldsymbol{v} \oplus \mu \boldsymbol{w} \in \mathbb{G}^2 \mid \lambda, \mu \in \mathbb{G}, \ \lambda \oplus \mu = 0\}$$

Then, it can be checked that $\widetilde{\mathcal{S}} = \{(\alpha, 1) \in \mathbb{R}^2_{\max} \mid \alpha \in [0, 1]\}$. Now, observe that if we consider the segment \mathcal{S}' joining $\boldsymbol{v}' = (1^-, 0)$ and $\boldsymbol{w}' = (0, 1)$, we have $\widetilde{\mathcal{S}}' = \widetilde{\mathcal{S}}$. Noticing that $\boldsymbol{v}' \notin \mathcal{S}$ and $\boldsymbol{v} \notin \mathcal{S}'$, we see that different polyhedra of \mathbb{G}^n can encode the same subset of \mathbb{R}^n_{\max} . Consequently, it seems far from trivial to determine the equality of two subsets of \mathbb{R}^n_{\max} when they are encoded as polyhedra of \mathbb{G}^n .

Remark 2.7. As a complement of Remark 2.6, we point out that the closure of a polyhedron with mixed constraints is apparently harder to compute than in the case of usual convex polyhedra. Indeed, in the latter case, the closure can be simply obtained by substituting $\langle by \leq in$ the defining inequalities. In contrast, in the current setting, replacing the coefficients b_i of the form β_i^- by β_i on the right-hand side of mixed inequalities provides a closed tropical polyhedron which may be larger than the closure. For example, consider the polyhedron with mixed constraints defined by the inequalities:

$$0 \leqslant_{\mathbb{G}} \boldsymbol{x}_1 \qquad \boldsymbol{x}_1 \oplus \boldsymbol{x}_2 \leqslant_{\mathbb{G}} 1 \qquad 0 \leqslant_{\mathbb{G}} (-1)^- \boldsymbol{x}_1 \oplus 0^- \boldsymbol{x}_2 \;,$$



Fig. 4. Left: a tropical polyhedron with mixed constraints, together with an open mixed halfspace (in blue) defining it. Right: the closed polyhedron defined by the corresponding non-strict inequalities.

which is depicted on the left-hand side of Figure 4. The mixed half-space defined by the last inequality is represented in blue. The closure of the polyhedron is the usual unit square. However, the closed polyhedron defined by the inequalities:

 $0 \leqslant \boldsymbol{x}_1 \qquad \boldsymbol{x}_1 \oplus \boldsymbol{x}_2 \leqslant 1 \qquad 0 \leqslant (-1) \boldsymbol{x}_1 \oplus \boldsymbol{x}_2 \; ,$

contains additionally the half-line $\{(1, \lambda) \mid \lambda \leq 0\}$ (right-hand side of Figure 4).

3. Tropical Fourier-Motzkin Elimination

In this section, we first present a tropical Fourier-Motzkin elimination method, which allows to eliminate a variable in a finite system of mixed inequalities. Then we apply this method to establish relationships between tropical polyhedra with mixed constraints, zones and tropical hemispaces.

3.1. The algorithm

We first illustrate the algorithm on an example.

Example 3.1. Consider the system given in (2.3), and assume we want to eliminate the variable x_1 . From the last (rightmost) two inequalities of (2.3), we know that:

$$\boldsymbol{x}_1 \leqslant_{\mathbb{G}} 3^- \boldsymbol{x}_2 \qquad \boldsymbol{x}_1 \leqslant_{\mathbb{G}} 2^- \oplus 1 \boldsymbol{x}_2 \tag{3.1}$$

In each inequality involving the variable x_1 on the right-hand side (*i.e.* the leftmost three inequalities of (2.3)), we propose to replace x_1 by the two upper bounds provided by (3.1). This produces the following six inequalities:

$$(-2)\boldsymbol{x}_{2} \leqslant_{\mathbb{G}} 0^{-} \oplus 3^{-}\boldsymbol{x}_{2} \qquad -3 \leqslant_{\mathbb{G}} 3^{-}\boldsymbol{x}_{2} \qquad 0 \leqslant_{\mathbb{G}} 4^{-}\boldsymbol{x}_{2} (-2)\boldsymbol{x}_{2} \leqslant_{\mathbb{G}} 2^{-} \oplus 1^{-}\boldsymbol{x}_{2} \qquad -3 \leqslant_{\mathbb{G}} 2^{-} \oplus 1\boldsymbol{x}_{2} \qquad 0 \leqslant_{\mathbb{G}} 3^{-} \oplus 2\boldsymbol{x}_{2}$$
(3.2)

Besides, in each inequality not involving x_1 on the right-hand side, we remove the term in x_1 from the left-hand side, if any. From the rightmost inequalities of (2.3), we obtain the following three inequalities:

$$-2 \leqslant_{\mathbb{G}} \boldsymbol{x}_2 \qquad -\infty \leqslant_{\mathbb{G}} 3^- \boldsymbol{x}_2 \qquad -\infty \leqslant_{\mathbb{G}} 0^- \oplus (-1) \boldsymbol{x}_2 \quad . \tag{3.3}$$

We claim that the inequalities in (3.2) and (3.3) precisely describe the projection on the x_2 axis of the polyhedron with mixed constraints defined by (2.3). Note that the collection of inequalities obtained in this way is redundant: one inequality, $-2 \leq_{\mathbb{G}} x_2$, suffices.

We now formalize the approach sketched in Example 3.1. Under some assumptions (which are specified in Theorem 3.3 below), it applies more generally to systems of constraints over a totally ordered idempotent semiring.

Recall that a semiring $(\mathbb{S}, \oplus, \otimes, \mathbb{O}, \mathbb{1})$ is said to be *totally ordered* if there exists a total order $\leq_{\mathbb{S}}$ on \mathbb{S} such that:

- (i) $0 \leq a$ for all $a \in S$,
- (ii) for all $a, b, c \in \mathbb{S}$, $a \leq_{\mathbb{S}} b$ implies $a \oplus c \leq_{\mathbb{S}} b \oplus c$, $a \otimes c \leq_{\mathbb{S}} b \otimes c$, and $c \otimes a \leq_{\mathbb{S}} c \otimes b$.

The semiring S is said to be *idempotent* if $a \oplus a = a$ for all $a \in S$. The next lemma shows that such a semiring is *naturally ordered*, meaning that $a \oplus b$ is equal to the maximal element among a and b.

Lemma 3.2. Let $(\mathbb{S}, \oplus, \otimes, \mathbb{O}, \mathbb{1}, \leq_{\mathbb{S}})$ be a totally ordered idempotent semiring. Then, for all $a, b \in \mathbb{S}$, $a \oplus b = a$ if $a \geq_{\mathbb{S}} b$, and $a \oplus b = b$ otherwise.

Proof. First, observe that $b \ge_{\mathbb{S}} 0$ implies $a \oplus b \ge_{\mathbb{S}} a$. Analogously, we have $a \oplus b \ge_{\mathbb{S}} b$. Now suppose that $a \ge_{\mathbb{S}} b$. Then, $a = a \oplus b$ since $a = a \oplus a \ge_{\mathbb{S}} a \oplus b$.

Now we explain how to eliminate \boldsymbol{x}_n in a linear system of inequalities over S in the variables $\boldsymbol{x}_1, \ldots, \boldsymbol{x}_n$. For the sake of simplicity, we extend the operations of S to matrices and vectors in the usual way and represent the multiplication \otimes by concatenation.

Theorem 3.3 (Fourier-Motzkin elimination for systems over totally ordered idempotent semirings). Let $(\mathbb{S}, \oplus, \otimes, \mathbb{O}, \mathbb{1}, \leq_{\mathbb{S}})$ be a totally ordered idempotent semiring and $\mathcal{P} \subset \mathbb{S}^n$ be the solution set of the system $A\mathbf{x} \oplus \mathbf{c} \leq_{\mathbb{S}} B\mathbf{x} \oplus \mathbf{d}$, where $A, B \in \mathbb{S}^{p \times n}$ and $\mathbf{c}, \mathbf{d} \in \mathbb{S}^p$ satisfy the following conditions:

- (i) A_{in} is (left-)invertible with respect to \otimes if $A_{in} \neq 0$ (we denote its inverse by A_{in}^{-1}),^a
- (ii) for any $\alpha \in \mathbb{S}$ there exists $\beta \in \mathbb{S}$ such that $\alpha \leq_{\mathbb{S}} B_{in}\beta$ if $B_{in} \neq \mathbb{O}$,
- (iii) either $A_{ij} = 0$ or $B_{ij} = 0$ for $i \in [p]$ and $j \in [n]$.

Let $\mathcal{Q} \subset \mathbb{S}^{n-1}$ be the set defined by the following inequalities in the variables x_1, \ldots, x_{n-1} :

$$(\oplus_{j \neq n} A_{ij} \boldsymbol{x}_j) \oplus \boldsymbol{c}_i \leqslant_{\mathbb{S}} (\oplus_{j \neq n} B_{ij} \boldsymbol{x}_j) \oplus \boldsymbol{d}_i \quad , \tag{3.4}$$

^aHere and below, M_{in} denotes the (i, n)-entry of matrix M, and should not be confused with any abbreviation.

Tropical Fourier-Motzkin Elimination, with an Application to Real-Time Verification 13

for all $i \in [p]$ such that $B_{in} = 0$, and

$$(\oplus_{j\neq n}A_{ij}\boldsymbol{x}_j)\oplus\boldsymbol{c}_i\leqslant_{\mathbb{S}}(\oplus_{j\neq n}(B_{ij}\oplus B_{in}A_{kn}^{-1}B_{kj})\boldsymbol{x}_j)\oplus\boldsymbol{d}_i\oplus B_{in}A_{kn}^{-1}\boldsymbol{d}_k \quad , \quad (3.5)$$

for all $i, k \in [p]$ such that $B_{in} \neq 0$ and $A_{kn} \neq 0$.

Then $x \in \mathcal{Q}$ if, and only if, there exists $\lambda \in \mathbb{S}$ such that $(x, \lambda) \in \mathcal{P}$.

Proof. Assume that $(\boldsymbol{x}, \lambda) \in \mathcal{P}$ for some $\lambda \in S$. Then for all $i \in [p]$ such that $B_{in} = 0$, we have:

$$(\oplus_{j\neq n}A_{ij}\boldsymbol{x}_j)\oplus\boldsymbol{c}_i\leqslant_{\mathbb{S}}(\oplus_{j\neq n}A_{ij}\boldsymbol{x}_j)\oplus A_{in}\lambda\oplus\boldsymbol{c}_i\leqslant_{\mathbb{S}}(\oplus_{j\neq n}B_{ij}\boldsymbol{x}_j)\oplus\boldsymbol{d}_i$$

which proves that \boldsymbol{x} satisfies the inequalities of the form (3.4). Now consider $i, k \in [p]$ such that $B_{in} \neq 0$ and $A_{kn} \neq 0$. Then, $A_{in} = B_{kn} = 0$ by Condition (iii), and A_{kn} is left-invertible by Condition (i). Since $A_{kn}\lambda \leq_{\mathbb{S}} (\oplus_{j\neq n}B_{kj}\boldsymbol{x}_j) \oplus \boldsymbol{d}_k$, we know that

$$\lambda \leqslant_{\mathbb{S}} A_{kn}^{-1}(\oplus_{j
eq n} B_{kj} \boldsymbol{x}_j) \oplus A_{kn}^{-1} \boldsymbol{d}_k$$

Replacing λ by the latter upper bound in the inequality

$$(\oplus_{j\neq n}A_{ij}\boldsymbol{x}_j)\oplus\boldsymbol{c}_i=(\oplus_{j\neq n}A_{ij}\boldsymbol{x}_j)\oplus A_{in}\lambda\oplus\boldsymbol{c}_i\leqslant_{\mathbb{S}}(\oplus_{j\neq n}B_{ij}\boldsymbol{x}_j)\oplus B_{in}\lambda\oplus\boldsymbol{d}_i$$

precisely yields inequality (3.5), using distributivity of \otimes and commutativity of \oplus . Thus, we conclude that $x \in Q$.

Conversely, let $\boldsymbol{x} \in \mathcal{Q}$. In the first place, assume that $A_{in} = 0$ for all $i \in [p]$. Define

$$\lambda := \oplus_{i \in I} \beta_i \;\;,$$

where $I := \{i \in [p] \mid B_{in} \neq 0\}$ and $\beta_i \in \mathbb{S}$ is such that $(\bigoplus_{j \neq n} A_{ij} \boldsymbol{x}_j) \oplus \boldsymbol{c}_i \leq \mathbb{S} B_{in} \beta_i$ for $i \in I$ (β_i exists thanks to Condition (ii)). Then, if $B_{in} \neq 0$, we have

$$(\oplus_{j\neq n}A_{ij}\boldsymbol{x}_j)\oplus A_{in}\lambda\oplus\boldsymbol{c}_i=(\oplus_{j\neq n}A_{ij}\boldsymbol{x}_j)\oplus\boldsymbol{c}_i\leqslant_{\mathbb{S}}B_{in}\lambda\leqslant_{\mathbb{S}}(\oplus_{j\neq n}B_{ij}\boldsymbol{x}_j)\oplus B_{in}\lambda\oplus\boldsymbol{d}_i.$$

The same inequality is trivially satisfied if $B_{in} = 0$ due to (3.4). It follows that $(\boldsymbol{x}, \lambda) \in \mathcal{P}$.

Now assume that $A_{kn} \neq 0$ for some $k \in [p]$. Define

$$\lambda := \min_{\substack{k \in [p] \\ A_{kn} \neq 0}} \left(A_{kn}^{-1}(\oplus_{j \neq n} B_{kj} \boldsymbol{x}_j) \oplus A_{kn}^{-1} \boldsymbol{d}_k \right) , \qquad (3.6)$$

where the operator min is understood as providing the minimum of its operands with respect to the order $\leq_{\mathbb{S}}$. As a consequence, for all *i* such that $A_{in} \neq 0$, we have $A_{in}\lambda \leq_{\mathbb{S}} (\bigoplus_{j\neq n} B_{ij}\boldsymbol{x}_j) \oplus \boldsymbol{d}_i$. The fact that $B_{in} = 0$ (by Condition (iii)) and the conjunction with (3.4) yield:

$$(\oplus_{j\neq n}A_{ij}\boldsymbol{x}_j)\oplus A_{in}\lambda\oplus\boldsymbol{c}_i\leqslant_{\mathbb{S}}(\oplus_{j\neq n}B_{ij}\boldsymbol{x}_j)\oplus\boldsymbol{d}_i=(\oplus_{j\neq n}B_{ij}\boldsymbol{x}_j)\oplus B_{in}\lambda\oplus\boldsymbol{d}_i$$

Note that the latter inequality also holds for all $i \in [p]$ such that A_{in} and B_{in} are both equal to \mathbb{O} .

Now suppose that $i \in [p]$ satisfies $B_{in} \neq 0$ and $k \in [p]$ attains the minimum in (3.6). Since \boldsymbol{x} satisfies (3.5), *i.e.*

$$(\oplus_{j\neq n}A_{ij}\boldsymbol{x}_j)\oplus\boldsymbol{c}_i\leqslant_{\mathbb{S}}(\oplus_{j\neq n}B_{ij}\boldsymbol{x}_j)\oplus\boldsymbol{d}_i\oplus B_{in}A_{kn}^{-1}(\oplus_{j\neq n}B_{kj}\boldsymbol{x}_j\oplus\boldsymbol{d}_k)$$

it follows that

$$(\oplus_{j\neq n}A_{ij}\boldsymbol{x}_j)\oplus A_{in}\lambda\oplus\boldsymbol{c}_i=(\oplus_{j\neq n}A_{ij}\boldsymbol{x}_j)\oplus\boldsymbol{c}_i\leqslant_{\mathbb{S}}(\oplus_{j\neq n}B_{ij}\boldsymbol{x}_j)\oplus B_{in}\lambda\oplus\boldsymbol{d}_i$$

because $B_{in} \neq 0$ implies $A_{in} = 0$ by Condition (iii). This shows that $(\boldsymbol{x}, \lambda)$ belongs to \mathcal{P} .

The case of tropical polyhedra with mixed constraints is obtained by setting $\mathbb{S} = \mathbb{G}$. Note that the conditions of Theorem 3.3 are satisfied when $\mathbb{S} = \mathbb{G}$, due in particular to Lemma 2.5 and the fact that any non-zero coefficient on the left-hand side of a mixed inequality is invertible (as an element of \mathbb{R}). However, a tropical polyhedron with mixed constraints consists of the solutions belonging to \mathbb{R}_{\max}^n , while Theorem 3.3 applies to the solutions in \mathbb{G}^n . The following result shows that the projection algorithm is still valid when restricted to \mathbb{R}_{\max}^n .

Theorem 3.4 (Fourier-Motzkin elimination for systems of mixed inequalities). Assume \mathcal{P} and \mathcal{Q} are defined as in Theorem 3.3 with $\mathbb{S} = \mathbb{G}$. Then, for all $\boldsymbol{x} \in \mathbb{R}_{\max}^{n-1}$, $\boldsymbol{x} \in \mathcal{Q}$ if, and only if, there exists $\lambda \in \mathbb{R}_{\max}$ such that $(\boldsymbol{x}, \lambda) \in \mathcal{P}$.

Proof. Observe that to prove the theorem, it is enough to show that if $\boldsymbol{y} := (\boldsymbol{x}, \lambda) \in \mathcal{P}$ for some $\boldsymbol{x} \in \mathbb{R}_{\max}^{n-1}$ and $\lambda \in \mathbb{G}$, then there exists $\lambda' \in \mathbb{R}_{\max}$ such that $(\boldsymbol{x}, \lambda') \in \mathcal{P}$.

If $\lambda \in \mathbb{R}_{\max}$, there is nothing to prove, so assume $\lambda \in \mathbb{R}^- \cup \{+\infty\}$. We will show that for a certain choice of $\lambda' \in \mathbb{R}$ verifying $\lambda' \leq_{\mathbb{G}} \lambda$, the vector $\mathbf{y}' := (\mathbf{x}, \lambda') \in \mathbb{R}_{\max}^n$ belongs to \mathcal{P} . Note that for any such choice of λ' , the vector \mathbf{y}' satisfies the inequalities in $A\mathbf{x} \oplus \mathbf{c} \leq_{\mathbb{G}} B\mathbf{x} \oplus \mathbf{d}$ indexed by $i \in [p]$ such that $B_{in} = \pm \infty$. Indeed, in this case, we have

$$(\oplus_j A_{ij} \boldsymbol{y}'_j) \oplus \boldsymbol{c}_i \leqslant_{\mathbb{G}} (\oplus_j A_{ij} \boldsymbol{y}_j) \oplus \boldsymbol{c}_i \leqslant_{\mathbb{G}} (\oplus_j B_{ij} \boldsymbol{y}_j) \oplus \boldsymbol{d}_i = (\oplus_j B_{ij} \boldsymbol{y}'_j) \oplus \boldsymbol{d}_i . \quad (3.7)$$

Thus, we next focus on the inequalities indexed by elements of the set $I := \{i \in [p] \mid B_{in} \in \mathbb{R} \cup \mathbb{R}^-\}$. In consequence, $A_{in} = -\infty$ for all $i \in I$. It is convenient to split the rest of the proof into two cases.

In the first place, assume $\lambda = +\infty$. Define $\lambda' := \delta \epsilon \in \mathbb{R}$, where $\epsilon > 0$ and

$$\delta = \begin{cases} \bigoplus_{i \in I} |B_{in}|^{-1} (\bigoplus_{j \neq n} A_{ij} \boldsymbol{x}_j \oplus \boldsymbol{c}_i) & \text{if } I \neq \emptyset, \\ 0 & \text{otherwise.} \end{cases}$$

Obviously, $\lambda' \leq_{\mathbb{G}} \lambda$. Besides, for all $i \in I$ we have

$$(\oplus_{j\neq n}A_{ij}\boldsymbol{x}_{j})\oplus A_{in}\lambda'\oplus\boldsymbol{c}_{i}\leqslant_{\mathbb{G}}(\oplus_{j\neq n}B_{ij}\boldsymbol{x}_{j})\oplus B_{in}\lambda'\oplus\boldsymbol{d}_{i},$$

since $\oplus_{j\neq n} A_{ij} \boldsymbol{x}_j \oplus \boldsymbol{c}_i < |B_{in}| \lambda'$ and $A_{in} = -\infty$.

Now assume $\lambda \in \mathbb{R}^-$. Let I' be the set of indices $i \in I$ such that $B_{in}\lambda >_{\mathbb{G}} \oplus_{i \neq n} B_{ij} \boldsymbol{x}_i \oplus \boldsymbol{d}_i$. Then,

$$(\oplus_j A_{ij} \boldsymbol{y}_j) \oplus \boldsymbol{c}_i = (\oplus_{j \neq n} A_{ij} \boldsymbol{x}_j) \oplus \boldsymbol{c}_i \leqslant_{\mathbb{G}} B_{in} \lambda$$

for all $i \in I'$. For $i \in I'$, let $\nu_i = \bigoplus_{j \neq n} A_{ij} \boldsymbol{x}_j \oplus \boldsymbol{c}_i$. Since ν_i belongs to \mathbb{R}_{\max} $(A_{ij}, \boldsymbol{x}_j)$ and \boldsymbol{c}_i belong to \mathbb{R}_{\max}) and $B_{in}\lambda$ belongs to \mathbb{R}^- , we necessarily have $\nu_i < |B_{in}\lambda|$. Define $\lambda' := \kappa |\lambda| \in \mathbb{R}$, where

$$\kappa = \begin{cases} \max_{i \in I'} (\nu_i - |B_{in}\lambda|)/2 & \text{if } \{i \in I' \mid \nu_i \in \mathbb{R}\} \neq \emptyset, \\ -1 & \text{otherwise.} \end{cases}$$

As $\kappa < 0$, we have $\lambda' \leq_{\mathbb{G}} \lambda$. Moreover, if $i \in I'$, then $\nu_i < \kappa |B_{in}\lambda|$ and so

$$(\oplus_j A_{ij} \boldsymbol{y}'_j) \oplus \boldsymbol{c}_i \leqslant_{\mathbb{G}} (\oplus_j A_{ij} \boldsymbol{y}_j) \oplus \boldsymbol{c}_i = (\oplus_{j \neq n} A_{ij} \boldsymbol{x}_j) \oplus \boldsymbol{c}_i = = \nu_i \leqslant_{\mathbb{G}} \kappa B_{in} \lambda \leqslant_{\mathbb{G}} \kappa B_{in} |\lambda| = B_{in} \lambda' \leqslant_{\mathbb{G}} (\oplus_j B_{ij} \boldsymbol{y}'_j) \oplus \boldsymbol{d}_i .$$

Finally, note that for $i \in I \setminus I'$ the relations in (3.7) are still valid because in that case we have $\bigoplus_{j \neq n} B_{ij} \boldsymbol{x}_j \oplus \boldsymbol{d}_i \geq_{\mathbb{G}} B_{in} \lambda \geq_{\mathbb{G}} B_{in} \lambda'$. This completes the proof. \Box

3.2. Characterization of tropical polyhedra with mixed constraints in terms of zones and tropical hemispaces

In this subsection we discuss some consequences of tropical Fourier-Motzkin elimination. In the first place, we establish that the tropical convex hull of the union of two tropical polyhedra with mixed constraints is a tropical polyhedron with mixed constraints. Recall that the *tropical convex hull* tconv(G) of a set $G \subset \mathbb{R}^n_{\max}$ is defined as the set of the vectors of the form

$$\lambda_1 x^1 \oplus \cdots \oplus \lambda_m x^n$$

where m is a positive integer, $x^i \in G$, $\lambda_i \in \mathbb{R}_{\max}$ $(i \in [m])$, and $\bigoplus_{i \in [m]} \lambda_i = 0$.

Proposition 3.5. Let $\mathcal{P}, \mathcal{P}' \subset \mathbb{R}^n_{\max}$ be polyhedra with mixed constraints, respectively defined by the systems $A\mathbf{x} \oplus \mathbf{c} \leq_{\mathbb{G}} B\mathbf{x} \oplus \mathbf{d}$ and $A'\mathbf{x} \oplus \mathbf{c}' \leq_{\mathbb{G}} B'\mathbf{x} \oplus \mathbf{d}'$, and let \mathcal{Q} be the polyhedron defined by the inequalities which are obtained eliminating $y_1, \ldots, y_n, y'_1, \ldots, y'_n, \lambda$ and μ in the following system:

$oldsymbol{x} = oldsymbol{y} \oplus oldsymbol{y}'$	$\lambda \oplus \mu = 0$
$Aoldsymbol{y} \oplus \lambdaoldsymbol{c} \leqslant_{\mathbb{G}} Boldsymbol{y} \oplus \lambdaoldsymbol{d}$	$\boldsymbol{y}_1 \oplus \cdots \oplus \boldsymbol{y}_n \leqslant_{\mathbb{G}} (+\infty)\lambda$ (3.8)
$A'oldsymbol{y}'\oplus\muoldsymbol{c}'\leqslant_{\mathbb{G}}B'oldsymbol{y}'\oplus\muoldsymbol{d}'$	$oldsymbol{y}_1^\prime\oplus\cdots\oplusoldsymbol{y}_n^\prime\leqslant_{\mathbb{G}}(+\infty)\mu$

Then, we have $\operatorname{tconv}(\mathcal{P} \cup \mathcal{P}') = \mathcal{Q}$.

Proof. In the first place, note that $\mathcal{P} \subset \mathcal{Q}$. Indeed, given $\boldsymbol{x} \in \mathcal{P}$, let $\lambda = 0$, $\mu = -\infty$, $\boldsymbol{y} = \boldsymbol{x}$ and $\boldsymbol{y}'_i = -\infty$ for $i \in [n]$. Then, (3.8) is clearly satisfied, and so from Theorem 3.4 we deduce that $\boldsymbol{x} \in \mathcal{Q}$. Similarly, $\mathcal{P}' \subset \mathcal{Q}$. Since \mathcal{Q} is tropically convex, we deduce that $\operatorname{tconv}(\mathcal{P} \cup \mathcal{P}') \subset \mathcal{Q}$.

Conversely, let $\boldsymbol{x} \in \mathcal{Q}$, and consider $\boldsymbol{y}, \boldsymbol{y}', \lambda, \mu$ as in (3.8). If both λ, μ are distinct from $-\infty$, then $\lambda^{-1}\boldsymbol{y} \in \mathcal{P}, \mu^{-1}\boldsymbol{y}' \in \mathcal{P}'$, and $\boldsymbol{x} = \lambda(\lambda^{-1}\boldsymbol{y}) \oplus \mu(\mu^{-1}\boldsymbol{y}')$, which ensures that $\boldsymbol{x} \in \operatorname{tconv}(\mathcal{P} \cup \mathcal{P}')$. Otherwise, if for instance $\mu = -\infty$, then we necessarily have $\lambda = 0$, and so $\boldsymbol{y} \in \mathcal{P}$. Moreover, from $\boldsymbol{y}'_1 \oplus \cdots \oplus \boldsymbol{y}'_n \leq_{\mathbb{G}} (+\infty)\mu$ we deduce that $\boldsymbol{y}'_i = -\infty$ for all $i \in [n]$, and in consequence $\boldsymbol{x} = \boldsymbol{y}$. This completes the proof, because again we have $\boldsymbol{x} = \boldsymbol{y} \in \mathcal{P} \subset \operatorname{tconv}(\mathcal{P} \cup \mathcal{P}')$, and so $\mathcal{Q} \subset \operatorname{tconv}(\mathcal{P} \cup \mathcal{P}')$.

In order to characterize tropical polyhedra with mixed constraints in terms of zones, we first need to extend the definition of zones to \mathbb{R}_{\max} : a zone (of \mathbb{R}_{\max}^n) is a set of vectors $\boldsymbol{x} \in \mathbb{R}_{\max}^n$ defined by inequalities of the form

$$m_i \triangleleft \boldsymbol{x}_i \qquad \boldsymbol{x}_i \triangleleft M_i \qquad \boldsymbol{x}_i \triangleleft k_{ij} \boldsymbol{x}_j$$

$$(3.9)$$

where $\triangleleft \in \{\leq, <\}$ and $m_i, M_i, k_{ij} \in \mathbb{R}_{\max}$.

Theorem 3.6. A subset \mathcal{P} of \mathbb{R}^n_{\max} is a tropical polyhedron with mixed constraints if, and only if, it is a tropically convex union of finitely many zones.

Proof. Observe that when \triangleleft is equal to \leq in (3.9), these inequalities are equivalent to the ones obtained replacing \triangleleft by $\leq_{\mathbb{G}}$. Moreover, we have

$$\begin{split} m_i < \boldsymbol{x}_i &\iff \begin{cases} m_i \leqslant_{\mathbb{G}} 0^- \boldsymbol{x}_i & \text{if } m_i \in \mathbb{R} \\ 0 \leqslant_{\mathbb{G}} (+\infty) \boldsymbol{x}_i & \text{if } m_i = -\infty \end{cases} \\ \boldsymbol{x}_i < M_i &\iff \boldsymbol{x}_i \leqslant_{\mathbb{G}} 0^- M_i & \text{if } M_i \in \mathbb{R} \end{cases} \\ \boldsymbol{x}_i < k_{ij} \boldsymbol{x}_j &\iff (\boldsymbol{x}_i \leqslant_{\mathbb{G}} (0^-) k_{ij} \boldsymbol{x}_j \text{ and } 0 \leqslant_{\mathbb{G}} (+\infty) \boldsymbol{x}_j) & \text{if } k_{ij} \in \mathbb{R} \end{cases}$$

In consequence, any zone is a tropical polyhedron with mixed constraints. Then, by Proposition 3.5, so is the tropical convex hull of the union of finitely many zones. The "if" part of the theorem follows from the fact that if the union of finitely many zones is tropically convex, then it coincides with its tropical convex hull.

Suppose now that Q is a polyhedron defined by mixed inequalities of the form (2.1), in which only one of the coefficients b_j is distinct from $-\infty$. If this coefficient is b_0 , as we can assume that $b_0 \neq +\infty$, it follows that (2.1) can be rewritten as a system of inequalities of the form $\mathbf{x}_i \leq_{\mathbb{G}} \beta M_i$, with $\beta \in \{0, 0^-\}$ and $M_i \in \mathbb{R}$. If the considered coefficient is b_j for $j \in [n]$, then (2.1) can be rewritten as a system of inequalities of the form $m_j \leq_{\mathbb{G}} \beta \mathbf{x}_j$ and $\mathbf{x}_i \leq_{\mathbb{G}} \beta k_{ij} \mathbf{x}_j$, with $\beta \in \{0, 0^-, +\infty\}$ and $m_j, k_{ij} \in \mathbb{R}$. Since

$$\begin{aligned} \boldsymbol{x}_i \leqslant_{\mathbb{G}} \beta k_{ij} \boldsymbol{x}_j & \Leftrightarrow \begin{cases} \boldsymbol{x}_i \leqslant k_{ij} \boldsymbol{x}_j & \text{if } \beta = 0\\ \boldsymbol{x}_i < k_{ij} \boldsymbol{x}_j \text{ or } (\boldsymbol{x}_i \leqslant -\infty \text{ and } \boldsymbol{x}_j \leqslant -\infty) & \text{if } \beta = 0^-\\ -\infty < \boldsymbol{x}_j \text{ or } (\boldsymbol{x}_i \leqslant -\infty \text{ and } \boldsymbol{x}_j \leqslant -\infty) & \text{if } \beta = +\infty \end{cases} \\ \boldsymbol{x}_i \leqslant_{\mathbb{G}} \beta M_i & \Longleftrightarrow \begin{cases} \boldsymbol{x}_i \leqslant M_i & \text{if } \beta = 0\\ \boldsymbol{x}_i < M_i & \text{if } \beta = 0^- \end{cases} \end{aligned}$$

Tropical Fourier-Motzkin Elimination, with an Application to Real-Time Verification 17

$$m_j \leqslant_{\mathbb{G}} \beta \boldsymbol{x}_j \iff \begin{cases} m_j \leqslant \boldsymbol{x}_j & \text{if } \beta = 0\\ m_j < \boldsymbol{x}_j & \text{if } \beta = 0^-\\ -\infty < \boldsymbol{x}_j & \text{if } \beta = +\infty \end{cases}$$

it follows that \mathcal{Q} is a finite union of zones.

The "only if" part of the theorem follows from the fact that any polyhedron with mixed constraints can be written as a finite union of polyhedra of the form considered in the previous paragraph (it suffices to choose, in each inequality, one term in the right-hand side to be the maximizing one). \Box

Theorem 3.6 raises the problem, of independent interest, of determining whether a given union of finitely many zones is tropically convex. To our knowledge, this problem has not been studied so far, even in the case of closed zones. A naive solution consists in computing the tropical convex hull of the union of zones (using Proposition 3.5), and checking whether it intersects the complement of the union of zones (the latter can be expanded into a union of zones, and the intersection test requires the techniques developed in Section 4, see Theorem 4.7). This approach would be particularly expensive. Yet, it is similar to the technique implemented in the UPPAAL DBM library [21] to test if a union of zones is a zone. Whether there exists a more efficient method is left for future work.

Now we study the relationship between tropical polyhedra with mixed constraints and tropical hemispaces. Recall that a *tropical hemispace* is a tropically convex set whose complement is also tropically convex. Tropical hemispaces and mixed half-spaces share the property that their closure is a closed half-space (in the case of hemispaces, this is proved in [14]). In fact, Proposition 3.7 and Example 3.8 below show that mixed half-spaces are a proper subclass of hemispaces.

Proposition 3.7. Mixed half-spaces are tropical hemispaces.

Proof. We already proved in Proposition 2.3 that mixed half-spaces are tropically convex.

The complement of the mixed half-space defined by (2.1) consists of the vectors $\boldsymbol{x} \in \mathbb{R}^n_{\max}$ such that

$$a_0 \oplus a_1 \boldsymbol{x}_1 \oplus \cdots \oplus a_n \boldsymbol{x}_n >_{\mathbb{G}} b_0 \oplus b_1 \boldsymbol{x}_1 \oplus \cdots \oplus b_n \boldsymbol{x}_n . \tag{3.10}$$

Let $\boldsymbol{x}, \boldsymbol{y} \in \mathbb{R}^n_{\max}$ be in this complement, and $\lambda, \mu \in \mathbb{R}_{\max}$ be such that $\lambda \oplus \mu = 0$. Without loss of generality, assume that $\lambda = 0$. If $\mu = -\infty$, then $\lambda \boldsymbol{x} \oplus \mu \boldsymbol{y} = \boldsymbol{x}$ trivially satisfies (3.10). Otherwise, *i.e.* if $\mu \in \mathbb{R}$ and $\mu \leq 0$, then by Property (iv) of Lemma 2.1 we have $\mu(a_0 \oplus a_1\boldsymbol{y}_1 \oplus \cdots \oplus a_n\boldsymbol{y}_n) >_{\mathbb{G}} \mu(b_0 \oplus b_1\boldsymbol{y}_1 \oplus \cdots \oplus b_n\boldsymbol{y}_n)$. It readily follows that $a_0 \oplus a_1(\lambda \boldsymbol{x}_1 \oplus \mu \boldsymbol{y}_1) \oplus \cdots \oplus a_n(\lambda \boldsymbol{x}_n \oplus \mu \boldsymbol{y}_n) >_{\mathbb{G}} b_0 \oplus b_1(\lambda \boldsymbol{x}_1 \oplus \mu \boldsymbol{y}_1) \oplus \cdots \oplus b_n(\lambda \boldsymbol{x}_n \oplus \mu \boldsymbol{y}_n)$. This shows that the complement of a mixed half-space is tropically convex.

Example 3.8. Consider the tropical hemispace \mathcal{H} defined as the set of vectors $\boldsymbol{x} \in \mathbb{R}^4_{\max}$ such that

$$(\boldsymbol{x}_3 \leqslant \boldsymbol{x}_1 ext{ and } \boldsymbol{x}_4 \leqslant \boldsymbol{x}_1) \quad ext{or} \quad (\boldsymbol{x}_3 \leqslant \boldsymbol{x}_2 ext{ and } \boldsymbol{x}_4 < \boldsymbol{x}_2)$$
 .

We claim that \mathcal{H} is not a mixed half-space. Its closure is the tropical half-space defined by the inequality

$$\boldsymbol{x}_3 \oplus \boldsymbol{x}_4 \leqslant 0 \boldsymbol{x}_1 \oplus 0 \boldsymbol{x}_2 \;. \tag{3.11}$$

Any mixed half-space whose closure is given by (3.11) is defined by a mixed inequality obtained from (3.11) by replacing some of the coefficients 0 on the right-hand side by 0^- . However, it can be easily verified that none of these mixed half-spaces is equal to \mathcal{H} .

Similarly, it can be checked that the complement of \mathcal{H} , which is given by the set of vectors $\boldsymbol{x} \in \mathbb{R}^4_{\max}$ such that

$$\boldsymbol{x}_1 < \boldsymbol{x}_3 \oplus \boldsymbol{x}_4 \quad ext{and} \quad (\boldsymbol{x}_2 < \boldsymbol{x}_3 ext{ or } \boldsymbol{x}_2 \leqslant \boldsymbol{x}_4) \; ,$$

is not a mixed half-space either.

To prove the following proposition, we shall use the characterization of tropical hemispaces in terms of (P, R)-decompositions established in [42].

Proposition 3.9. Tropical hemispaces are tropical polyhedra with mixed constraints.

Proof. In the first place, let us recall that the *tropical conic hull* of a subset G of \mathbb{R}^n_{\max} is defined as

$$\operatorname{tcone}(G) := \{\mu_1 \boldsymbol{x}^1 \oplus \cdots \oplus \mu_m \boldsymbol{x}^m \mid m \in \mathbb{N}, \ \boldsymbol{x}^i \in G, \ \mu_i \in \mathbb{R}_{\max}\}.$$

Moreover, given two subsets G and G' of \mathbb{R}^n_{\max} , their tropical Minkowski sum $G \oplus G'$ is defined as $\{ \boldsymbol{x} \oplus \boldsymbol{x}' \mid \boldsymbol{x} \in G, \ \boldsymbol{x}' \in G' \}$. In this proof, we denote by $\boldsymbol{u}^i \in \mathbb{R}^n_{\max}$, for $i \in [n]$, the vector defined by $\boldsymbol{u}^i_j := 0$ if j = i and $\boldsymbol{u}^i_j := -\infty$ otherwise.

Let $\mathcal{H}_1, \mathcal{H}_2$ be two complementary hemispaces. Suppose that the vector all of whose entries are equal to $-\infty$ belongs to \mathcal{H}_1 . Then, by [42, Theorem 4.22] there exist a partition I, J of [n] and interval sets $\Lambda^1_{ij}, \Lambda^2_{ij} \subset \mathbb{R}_{\max}$ for $i \in I \cup \{0\}$ and $j \in J$ such that

$$\begin{aligned} \mathcal{H}_1 &= \operatorname{tconv}(\{\lambda \boldsymbol{u}^j \mid j \in J, \lambda \in \Lambda_{0j}^1\}) \oplus \operatorname{tcone}(\{\boldsymbol{u}^i \oplus \lambda \boldsymbol{u}^j \mid i \in I, \ j \in J, \ \lambda \in \Lambda_{ij}^1\}) \\ \mathcal{H}_2 &= \operatorname{tconv}(\{\lambda \boldsymbol{u}^j \mid j \in J, \lambda \neq +\infty, \ \lambda \in \Lambda_{0j}^2\}) \\ &\oplus \operatorname{tcone}(\{\lambda \boldsymbol{u}^i \oplus \boldsymbol{u}^j \mid i \in I, \ j \in J, \ \lambda \in -\Lambda_{ij}^2\}) \end{aligned}$$

where each couple of intervals $\Lambda_{ij}^1, \Lambda_{ij}^2$ has one of the following forms:

$$(\Lambda_{ij}^1, \Lambda_{ij}^2) = \begin{cases} ([-\infty, \beta],]\beta, +\infty]), & \beta \in \mathbb{R} \cup \{-\infty\} \\ ([-\infty, \alpha[, [\alpha, +\infty]), & \alpha \in \mathbb{R} \cup \{+\infty\} \end{cases}$$
(3.12)

Thus, note that $x \in \mathcal{H}_1$ if, and only if, there exist λ_{ij} and μ_{ij} in \mathbb{R}_{\max} for $i \in I \cup \{0\}$ and $j \in J$ such that

$$\begin{aligned} \boldsymbol{x}_{i} &= \bigoplus_{j \in J} \mu_{ij} & \text{for } i \in I \\ \boldsymbol{x}_{j} &= \bigoplus_{i \in I \cup \{0\}} \lambda_{ij} \mu_{ij} & \text{for } j \in J \\ \oplus_{j \in J} \mu_{0j} &= 0 \\ \lambda_{ij} &\in \Lambda^{1}_{ij} & \text{for } i \in I \cup \{0\} \text{ and } j \in J \end{aligned}$$

Equivalently, $x \in \mathcal{H}_1$ if, and only if, there exist μ_{ij} and ν_{ij} in \mathbb{R}_{\max} for $i \in I \cup \{0\}$ and $j \in J$ such that

$$\begin{aligned} \boldsymbol{x}_{i} &= \bigoplus_{j \in J} \mu_{ij} & \text{for } i \in I \\ \boldsymbol{x}_{j} &= \bigoplus_{i \in I \cup \{0\}} \nu_{ij} & \text{for } j \in J \\ \oplus_{j \in J} \mu_{0j} &= 0 \end{aligned}$$
$$\nu_{ij} \leq_{\mathbb{G}} \begin{cases} \beta \mu_{ij} & \text{if } \Lambda^{1}_{ij} = [-\infty, \beta], \ \beta \in \mathbb{R}_{\max} \\ \alpha^{-} \mu_{ij} & \text{if } \Lambda^{1}_{ij} = [-\infty, \alpha[, \ \alpha \in \mathbb{R} \\ (+\infty) \mu_{ij} & \text{if } \Lambda^{1}_{ij} = [-\infty, +\infty[\end{cases} & \text{for } i \in I \cup \{0\} \text{ and } j \in J \end{cases}$$
$$(3.13)$$

By Theorem 3.4, we conclude that \mathcal{H}_1 is a tropical polyhedron with mixed constraints, since it is defined by the system of mixed inequalities obtained by eliminating μ_{ij} and ν_{ij} for $i \in I \cup \{0\}$ and $j \in J$ in (3.13).

The same result can be obtained for \mathcal{H}_2 using a symmetric argument. \Box

The following result is a straightforward consequence of Propositions 3.7 and 3.9.

Corollary 3.10. Tropical polyhedra with mixed constraints are precisely the intersections of finitely many tropical hemispaces.

Remark 3.11. We have previously defined a closed tropical polyhedron as the solution set of inequalities of the form (1.1), *i.e.* mixed inequalities with coefficients in \mathbb{R}_{max} . We point out that this definition is consistent with the fact that such polyhedra are precisely the tropical polyhedra with mixed constraints which are closed. Indeed, by Theorem 3.6 a tropical polyhedron with mixed constraints \mathcal{P} can be written as a finite union of zones. If \mathcal{P} is closed, then it is equal to the union of the closure of these zones (*i.e.* closed zones, which can be generated by finitely many points and rays by the tropical Minkowski-Weyl Theorem, see [30, Theorem 2]). Since \mathcal{P} is tropically convex, it is even equal to the tropical convex hull of this union. It follows that it is generated by finitely many points and rays. By the tropical Minkowski-Weyl Theorem, we deduce that it is the solution set of finitely many inequalities of the form (1.1).

4. Eliminating Redundant Mixed Inequalities

Like in the classical setting, Fourier-Motzkin elimination generates a system of $O(p^2)$ inequalities from an input with p constraints. Consequently, the number of inequalities may grow in $O(p^{2^k})$ after k successive applications. To avoid the explosion of the size of the constraint system, superfluous inequalities must be eliminated. With this aim, we present a decision procedure for implications of the form:

$$A \boldsymbol{x} \oplus \boldsymbol{c} \leqslant_{\mathbb{G}} B \boldsymbol{x} \oplus \boldsymbol{d} \implies \boldsymbol{e} \boldsymbol{x} \oplus g \leqslant_{\mathbb{G}} \boldsymbol{f} \boldsymbol{x} \oplus h \quad \text{for all } \boldsymbol{x} \in \mathbb{R}^n_{\max},$$
 (4.1)

where $A \in \mathbb{R}_{\max}^{p \times n}$, $B \in \mathbb{G}^{p \times n}$, $c \in \mathbb{R}_{\max}^{p}$, $d \in \mathbb{G}^{p}$, $g \in \mathbb{R}_{\max}$, $h \in \mathbb{G}$, and e and f are *n*-dimensional row vectors with entries in \mathbb{R}_{\max} and \mathbb{G} respectively. We assume that e and g are not identically null (in the tropical sense) and that $h \neq +\infty$, because otherwise deciding implication (4.1) is trivial.

4.1. Equivalence with mean payoff games

We first show that deciding an implication of the form (4.1) is equivalent to an emptiness problem for tropical polyhedra with mixed constraints.

Proposition 4.1. Let Q be the tropical polyhedron with mixed constraints defined by the system $Ax \oplus c \leq_{\mathbb{G}} Bx \oplus d$ and the following inequalities:

$$\begin{cases} \boldsymbol{f}_{i}\boldsymbol{x}_{i} \leq_{\mathbb{G}} (0^{-}\boldsymbol{e})\boldsymbol{x} \oplus 0^{-}g & \text{if } \boldsymbol{f}_{i} \in \mathbb{R}_{\max} \\ |\boldsymbol{f}_{i}|\boldsymbol{x}_{i} \leq_{\mathbb{G}} \boldsymbol{e}\boldsymbol{x} \oplus g & \text{if } \boldsymbol{f}_{i} \in \mathbb{R}^{-} \\ \boldsymbol{x}_{i} \leq_{\mathbb{G}} -\infty & \text{if } \boldsymbol{f}_{i} = +\infty \end{cases}$$

$$(4.2)$$

for all $i \in [n]$,

$$\begin{cases} h \leqslant_{\mathbb{G}} (0^{-}\boldsymbol{e})\boldsymbol{x} \oplus 0^{-}g & \text{if } h \in \mathbb{R}_{\max} \\ |h| \leqslant_{\mathbb{G}} \boldsymbol{e}\boldsymbol{x} \oplus g & \text{if } h \in \mathbb{R}^{-} \end{cases}$$
(4.3)

and

 $0 \leq_{\mathbb{G}} \oplus_{e_i \neq -\infty} (+\infty) \boldsymbol{x}_i \quad if \ g = -\infty.$

Then, implication (4.1) holds if, and only if, Q is empty.

Proof. Implication (4.1) is false if, and only if, there exists $\boldsymbol{x} \in \mathbb{R}^n_{\max}$ such that $A\boldsymbol{x} \oplus \boldsymbol{c} \leq_{\mathbb{G}} B\boldsymbol{x} \oplus \boldsymbol{d}$ and $\boldsymbol{e}\boldsymbol{x} \oplus g >_{\mathbb{G}} \boldsymbol{f}\boldsymbol{x} \oplus h$. Observe that $\boldsymbol{e}\boldsymbol{x} \oplus g >_{\mathbb{G}} \boldsymbol{f}\boldsymbol{x} \oplus h$ holds if, and only if, $\boldsymbol{e}\boldsymbol{x} \oplus g >_{\mathbb{G}} h$ and $\boldsymbol{e}\boldsymbol{x} \oplus g >_{\mathbb{G}} \boldsymbol{f}\boldsymbol{x} \oplus h$. Observe that $\boldsymbol{e}\boldsymbol{x} \oplus g >_{\mathbb{G}} \boldsymbol{f}\boldsymbol{x} \oplus h$ holds if, and only if, $\boldsymbol{e}\boldsymbol{x} \oplus g >_{\mathbb{G}} h$ and $\boldsymbol{e}\boldsymbol{x} \oplus g >_{\mathbb{G}} \boldsymbol{f}_i\boldsymbol{x}_i$ for all $i \in [n]$. This implies $\boldsymbol{e}\boldsymbol{x} \oplus g > -\infty$, *i.e.* $0 \leq_{\mathbb{G}} (+\infty)(\boldsymbol{e}\boldsymbol{x} \oplus g)$. The latter inequality is trivially satisfied if $g \neq -\infty$. When $g = -\infty$, it is equivalent to $0 \leq_{\mathbb{G}} \oplus_{\boldsymbol{e}_i \neq -\infty} (+\infty)\boldsymbol{x}_i$. Finally, assuming $\boldsymbol{e}\boldsymbol{x} \oplus \boldsymbol{g} > -\infty$, note that $\boldsymbol{e}\boldsymbol{x} \oplus \boldsymbol{g} >_{\mathbb{G}} \boldsymbol{f}_i\boldsymbol{x}_i$ is equivalent to (4.2), and $\boldsymbol{e}\boldsymbol{x} \oplus \boldsymbol{g} >_{\mathbb{G}} h$ is equivalent to (4.3). This completes the proof.

Tropical Fourier-Motzkin Elimination, with an Application to Real-Time Verification 21

In light of Proposition 4.1, it is enough to develop a decision procedure to determine whether a tropical polyhedron with mixed constraints is empty. Our approach relies on *parametric mean payoff games*, following the lines of [1,5].

Let \mathcal{R} be a tropical polyhedron with mixed constraints defined by the system

$$M\boldsymbol{x} \oplus \boldsymbol{p} \leqslant_{\mathbb{G}} N\boldsymbol{x} \oplus \boldsymbol{q} , \qquad (4.4)$$

where $M \in \mathbb{R}_{\max}^{r \times n}$, $N \in \mathbb{G}^{r \times n}$, $p \in \mathbb{R}_{\max}^{r}$ and $q \in \mathbb{G}^{r}$. It is convenient to consider two different cases, depending on whether $+\infty$ coefficients appear or not in (4.4).

4.1.1. Polyhedra defined by systems with $no +\infty$ coefficients.

In the first place, we restrict to the following case:

Assumption A. No coefficient is equal to $+\infty$ on the right-hand side of mixed inequalities.

In other words, in this subsection we assume $N \in (\mathbb{G} \setminus \{+\infty\})^{r \times n}$ and $q \in (\mathbb{G} \setminus \{+\infty\})^r$.

In this case, with each $\varepsilon \ge 0$ we associate a closed tropical polyhedron $\mathcal{R}_{\varepsilon}$, given by the system $M \boldsymbol{x} \oplus \boldsymbol{p} \leq_{\mathbb{G}} N(\varepsilon) \boldsymbol{x} \oplus \boldsymbol{q}(\varepsilon)$, and a mean payoff game involving two players, "Max" and "Min", playing on a weighted bipartite digraph $\mathcal{G}_{\varepsilon}$ composed of two kinds of nodes: row nodes $i \in [r]$, and column nodes $j \in [n + 1]$. This digraph contains an arc from row node i to column node j with weight $N_{ij}(\varepsilon)$ when $N_{ij} \neq -\infty$, and an arc from j to i with weight $-M_{ij}$ when $M_{ij} \neq -\infty$. Similarly, it contains an arc from row node i to column node n+1 with weight $\boldsymbol{q}_i(\varepsilon)$ if $\boldsymbol{q}_i \neq -\infty$, and an arc from column node n+1 to row node i with weight $-\boldsymbol{p}_i$ when $\boldsymbol{p}_i \neq -\infty$.

Example 4.2. Figure 5 provides an illustration of the digraph $\mathcal{G}_{\varepsilon}$ corresponding to the tropical polyhedron with mixed constraints defined by the system:

$$\begin{array}{rrrr} 1: & -3 \leqslant_{\mathbb{G}} \boldsymbol{x}_1 & & 3: & -2 \leqslant_{\mathbb{G}} \boldsymbol{x}_2 \\ 2: & & 0 \leqslant_{\mathbb{G}} 1 \boldsymbol{x}_1 \oplus 0^- \boldsymbol{x}_2 & & 4: & (-2) \boldsymbol{x}_1 \leqslant_{\mathbb{G}} 0^- \oplus (-1) \boldsymbol{x}_2 \end{array}$$

The principle of the game is the following. Players Min and Max alternatively move a pawn over the nodes of $\mathcal{G}_{\varepsilon}$. When it is placed on a column node, Player Min selects an outgoing arc, moves the pawn to the corresponding row node, and pays to Player Max the weight of the selected arc. Once the pawn is on a row node, Player Max similarly selects an outgoing arc, moves the pawn along it, and receives from Player Min a payment equal to the weight of the selected arc.

In the sequel, we suppose that Players Max and Min always have at least one possible action in each node:

Assumption B. Each node of $\mathcal{G}_{\varepsilon}$ has at least one outgoing arc.



Fig. 5. The digraph associated with a parametric mean payoff game (column and row nodes are respectively represented by circles and squares).

This technical property can be assumed without loss of generality, up to adding trivial inequalities or removing non-relevant unknowns in the system. We refer to the discussion of Assumptions 2.1 and 2.2 in [1] for further details.

We consider infinite runs of the game, in which case the payoff is defined as the mean of the payments of Player Min to Player Max. Player Min wants to minimize this mean of payments while Player Max wants to maximize it. We denote by $v(\varepsilon)$ the value of the game associated with $\mathcal{G}_{\varepsilon}$ when it starts at column node n + 1. It is shown in [1, Theorem 3.5] that the tropical polyhedron $\mathcal{R}_{\varepsilon}$ is non-empty if, and only if, $v(\varepsilon) \ge 0$, *i.e.* column node n + 1 is a winning initial node (for Player Max). Then, the following result immediately follows from Lemma 2.2.

Proposition 4.3. The tropical polyhedron with mixed constraints \mathcal{R} is non-empty if, and only if, there exists $\varepsilon > 0$ such that $v(\varepsilon) \ge 0$.

Let \tilde{M} (resp. \tilde{N}) be the matrix of size $r \times (n+1)$ obtained by concatenating matrix M and column vector \boldsymbol{p} (resp. N and \boldsymbol{q}). The dynamic programming operator g_{ε} associated with the game over $\mathcal{G}_{\varepsilon}$ is the function from \mathbb{R}^{n+1}_{\max} to itself defined by

$$(g_{\varepsilon}(\boldsymbol{x}))_{j} := \min_{\substack{i \in [r]\\ \tilde{M}_{ij} \neq -\infty}} \left(-\tilde{M}_{ij} + \max_{\substack{k \in [n+1]\\ \tilde{N}_{ik} \neq -\infty}} \left(\tilde{N}_{ik}(\varepsilon) + \boldsymbol{x}_{k} \right) \right) , \qquad (4.5)$$

for $j \in [n+1]$. This function satisfies the following properties:

- (i) it is order preserving, *i.e.* $\boldsymbol{x} \leq \boldsymbol{y}$ implies $g_{\varepsilon}(\boldsymbol{x}) \leq g_{\varepsilon}(\boldsymbol{y})$ for all $\boldsymbol{x}, \boldsymbol{y} \in \mathbb{R}_{\max}^{n+1}$,
- (ii) it is additively homogeneous, *i.e.* $g_{\varepsilon}(\lambda \boldsymbol{x}) = \lambda g_{\varepsilon}(\boldsymbol{x})$ for all $\lambda \in \mathbb{R}_{\max}$ and $\boldsymbol{x} \in \mathbb{R}_{\max}^{n+1}$,
- (iii) it preserves \mathbb{R}^{n+1} , thanks to Assumption B above.

Such a function can be shown to be non-expansive for the sup-norm. Since it is also piecewise affine, a theorem due to Kohlberg [43] implies the following vector $\chi(g_{\varepsilon})$,

Tropical Fourier-Motzkin Elimination, with an Application to Real-Time Verification 23

referred to as the *cycle-time vector* of g_{ε} , exists and has finite entries:

$$\chi(g_{\varepsilon}) := \lim_{h \to +\infty} g_{\varepsilon}^{h}(0)/h \; \; .$$

Kohlberg's theorem also implies the *j*-th entry of $\chi(g_{\varepsilon})$, which we denote by $\chi_j(g_{\varepsilon})$, corresponds to the value of the game when it starts at column node *j*. We refer to [1] for further details. Following the notation above, we consequently have

$$v(\varepsilon) = \chi_{n+1}(g_{\varepsilon})$$

for $\varepsilon \ge 0$.

The cycle-time vector $\chi(g_{\varepsilon})$ can be expressed in terms of the cycle-time vectors of dynamic programming operators associated with certain one-player games. More precisely, a *(positional) strategy* for Player Min is a function $\sigma : [n + 1] \rightarrow [r]$ associating with each column node j a row node $\sigma(j)$ such that $\tilde{M}_{\sigma(j)j} \neq -\infty$. Such a strategy defines a one-player game (played by Player Max) over the digraph $\mathcal{G}_{\varepsilon}^{\sigma}$ obtained from $\mathcal{G}_{\varepsilon}$ by deleting the arcs connecting column nodes j with row nodes isuch that $i \neq \sigma(j)$. Its dynamic programming operator g_{ε}^{σ} is given by:

$$(g_{\varepsilon}^{\sigma}(\boldsymbol{x}))_{j} = -\tilde{M}_{\sigma(j)j} + \max_{\substack{k \in [n+1] \ \tilde{N}_{\sigma(j)k} \neq -\infty}} \left(\tilde{N}_{\sigma(j)k}(\varepsilon) + \boldsymbol{x}_{k} \right) ,$$

for $j \in [n + 1]$. Observe that this operator is linear in the tropical semiring \mathbb{R}_{\max} . If we denote by Σ the (finite) set of strategies for Player Min, then as another consequence of Kohlberg's theorem, it can be shown that

$$\chi(g_{\varepsilon}) = \min_{\sigma \in \Sigma} \chi(g_{\varepsilon}^{\sigma}) .$$
(4.6)

A dual result based on strategies for Player Max can also be established.

Given $\sigma \in \Sigma$, the (n+1)-th entry of the vector $\chi(g_{\varepsilon}^{\sigma})$ can be similarly interpreted as the value of the one-player game associated with the digraph $\mathcal{G}_{\varepsilon}^{\sigma}$ when it starts at column node n + 1. As the function g_{ε}^{σ} is linear in the tropical semiring, it is known [18] that $\chi_{n+1}(g_{\varepsilon}^{\sigma})$ is equal to the maximal weight-to-length ratio of the elementary circuits of $\mathcal{G}_{\varepsilon}^{\sigma}$ reachable from column node n+1. A circuit in this digraph is referred to as a sequence of column nodes $j_0, \ldots, j_{l-1}, j_l = j_0$, where $l \ge 1$, and so l is considered to be its length. Note that the reachability relation in $\mathcal{G}_{\varepsilon}^{\sigma}$ does not depend on ε . Let C_{σ} be the set of the elementary circuits of \mathcal{G}_{0}^{σ} reachable from column node n + 1. By (4.6), we readily obtain

$$v(\varepsilon) = \chi_{n+1}(g_{\varepsilon}) = \min_{\sigma \in \mathcal{L}} \max_{(j_0, \dots, j_{l-1}, j_l) \in C_{\sigma}} \frac{1}{l} \left(\sum_{0 \leq s \leq l-1} -\tilde{M}_{\sigma(j_s)j_s} + \tilde{N}_{\sigma(j_s)j_{s+1}}(\varepsilon) \right)$$
(4.7)

for $\varepsilon \ge 0$. We deduce that:

Lemma 4.4. The function $\varepsilon \mapsto v(\varepsilon)$ is non-increasing and piecewise affine.

Proof. The fact that $\varepsilon \mapsto v(\varepsilon)$ is piecewise affine straightforwardly follows from (4.7).

We next prove by induction on h that the function $\varepsilon \mapsto (g_{\varepsilon}^{h}(\boldsymbol{x}))_{j}$ is non-increasing for any $\boldsymbol{x} \in \mathbb{R}^{n}$ and $j \in [n]$. In the first place, observe that ε appears only negatively in the coefficients $\tilde{N}_{ik}(\varepsilon)$ of (4.5), so the function $\varepsilon \mapsto (g_{\varepsilon}(\boldsymbol{x}))_{j}$ is non-increasing. Now consider $h \in \mathbb{N}$ and $\varepsilon \leqslant \varepsilon'$. By induction, we know that $g_{\varepsilon'}^{h}(\boldsymbol{x}) \leqslant g_{\varepsilon}^{h}(\boldsymbol{x})$ for any $\boldsymbol{x} \in \mathbb{R}^{n}$. Since the function $\boldsymbol{x} \mapsto g_{\varepsilon}(\boldsymbol{x})$ is order preserving, we have $g_{\varepsilon}(g_{\varepsilon'}^{h}(\boldsymbol{x})) \leqslant$ $g_{\varepsilon}^{h+1}(\boldsymbol{x})$. Besides, $g_{\varepsilon}(g_{\varepsilon'}^{h}(\boldsymbol{x})) \geqslant g_{\varepsilon'}(g_{\varepsilon'}^{h}(\boldsymbol{x}))) = g_{\varepsilon'}^{h+1}(\boldsymbol{x})$. It follows that $(g_{\varepsilon'}^{h+1}(\boldsymbol{x}))_{j} \leqslant$ $(g_{\varepsilon}^{h+1}(\boldsymbol{x}))_{j}$ for all $j \in [n]$.

We conclude that $\varepsilon \mapsto v(\varepsilon)$ is non-increasing as the limit of non-increasing functions.

If we assume that the numerical parts of the non-zero entries of M, N, p and q are integers (this assumption is obviously satisfied in the application to timed automata in Section 5), the criterion of Proposition 4.3 can be determined by evaluating $v(\varepsilon)$ at $\varepsilon = 0$ and at a small positive value.

Proposition 4.5. The tropical polyhedron with mixed constraints \mathcal{R} is non-empty if, and only if, $v(0) \ge 1/(n+1)$ or $v(1/(n+1)^2) \ge 0$.

Proof. Each linear piece of the function $\varepsilon \mapsto v(\varepsilon)$ corresponds to an affine map given by the weight-to-length ratio of an elementary circuit of $\mathcal{G}_{\varepsilon}^{\sigma}$ for some strategy σ for Player Min. In consequence, this affine map is of the form $\frac{\lambda - k\varepsilon}{l}$, where:

- (i) l is the length of the circuit $(l \leq n+1)$,
- (ii) λ is the sum of 2*l* integers given by the modulus, or their opposite, of some entries of the matrices *M* and *N* and of the vectors *p* and *q*,
- (iii) k is the number of occurrences of $-\varepsilon$ in the weight $N_{ij}(\varepsilon)$ or $q_i(\varepsilon)$ of some arcs of the circuit (so $k \leq l$).

Therefore, any non-differentiability point $\tilde{\varepsilon}$ of the map $\varepsilon \mapsto v(\varepsilon)$ satisfies $(\lambda - k\tilde{\varepsilon})/l = (\lambda' - k'\tilde{\varepsilon})/l'$, where l', λ' , and k' have the same properties as l, λ , and k above respectively. It follows that any positive non-differentiability point (if any) is of the form:

$$\tilde{\varepsilon} = rac{\lambda' l - \lambda l'}{k' l - l' k}$$

Assume, without loss of generality, that k/l < k'/l'. Since $\tilde{\varepsilon} > 0$, the numerator $\lambda' l - \lambda l'$ is a positive integer. This implies that $\tilde{\varepsilon} \ge \varepsilon^*$, where $\varepsilon^* = 1/(n+1)^2$.

Consequently, the function $\varepsilon \mapsto v(\varepsilon)$ is affine on the interval $[0, \epsilon^*]$. Since it is non-increasing by Lemma 4.4, we have $v(\varepsilon) < 0$ for all $\varepsilon > 0$ if, and only if, $v(0) \leq 0$ (by continuity) and $v(\varepsilon^*) < 0$. The application of Proposition 4.3 shows that \mathcal{R} is non-empty if, and only if, v(0) > 0 or $v(\varepsilon^*) \geq 0$. Since v(0) is equal to the weightto-length ratio of an elementary circuit of \mathcal{G}_0^{σ} , for some $\sigma \in \Sigma$, it can be written as λ''/l'' , where λ'' and l'' have the same properties as λ and l above. It follows that v(0) is positive if, and only if, it is greater than or equal to 1/(n+1). This completes the proof. Let \mathcal{G}'_0 be the digraph obtained from \mathcal{G}_0 by subtracting 1/(n+1) from the weight of each arc connecting a row node with a column node. Then, v(0) - (1/(n+1)) is the value of the mean payoff game associated with \mathcal{G}'_0 when it starts from column node n + 1. It follows that the condition of Proposition 4.5 holds if, and only if, column node n + 1 is winning (for Player Max) in one of the two games associated with \mathcal{G}'_0 and $\mathcal{G}_{1/(n+1)^2}$. Let \mathcal{G}^* be the digraph obtained as the disjoint union of \mathcal{G}'_0 and $\mathcal{G}_{1/(n+1)^2}$, adding a special row node 0 and two arcs, with zero weight, connecting it with column nodes n + 1 of \mathcal{G}'_0 and $\mathcal{G}_{1/(n+1)^2}$. The criterion of Proposition 4.5 can be restated as follows:

Proposition 4.6. The tropical polyhedron with mixed constraints \mathcal{R} is non-empty if, and only if, row node 0 is a winning initial node (for Player Max) in the mean payoff game associated with \mathcal{G}^* .

As an immediate consequence, we obtain the following complexity result, in which the equivalence (ii) \Leftrightarrow (iii) extends Theorem 3.2 of [1], whereas the equivalence (i) \Leftrightarrow (iii) extends Theorem 18 of [5] (only non-strict constraints are considered there).

Theorem 4.7. Under Assumption A, the following problems are (Karp) polynomial-time equivalent:

- (i) deciding whether a mixed tropical affine inequality is implied by a system of such inequalities;
- (ii) deciding whether a tropical polyhedron with mixed constraints is empty;
- (iii) determining whether a given initial node in a mean payoff game is winning.

Problem (iii) is known to be in NP \cap coNP, see [50]. We deduce from Theorem 4.7 that Problems (i) and (ii) both belong to the same complexity class (NP and coNP are closed under Karp reductions). Whether Problem (iii) can be solved in polynomial time has been an open question since the first combinatorial algorithm [36]. Value iteration leads to a pseudo-polynomial algorithm [50]. Several algorithms rely on the idea of strategy iteration [38], applying various strategy improvement rules, see in particular [11,18,25,41]. A remarkable example has recently been constructed [26] in which some common rules lead to an exponential number of iterations. However, many algorithms, including the one of [18,25], are known to have experimentally a small average case number of iterations (growing sublinearly with the dimension), see the benchmarks in [17].

The support of a tropically convex set $C \subset \mathbb{R}^n_{\max}$ is defined as the set supp(C) of indices $j \in [n]$ such that there exists $\boldsymbol{x} \in C$ verifying $\boldsymbol{x}_j \neq -\infty$. Note that supp(C) is the greatest subset $J \subset [n]$ such that $J = \{j \in [n] \mid \boldsymbol{x}_j \neq -\infty\}$ for a certain $\boldsymbol{x} \in C$.

Mean payoff games can be used to compute the support of the tropical polyhedron with mixed constraints \mathcal{R} . Indeed, j belongs to the support of \mathcal{R} if, and only if, $j \in \text{supp}(\mathcal{R}_{\varepsilon})$ for some $\varepsilon > 0$. By [1, Theorem 3.2], the fact that $j \in \text{supp}(\mathcal{R}_{\varepsilon})$ is equivalent to $\chi_j(g_{\varepsilon}) \ge 0$ and $\chi_{n+1}(g_{\varepsilon}) \ge 0$, *i.e.* to the fact that column nodes j and

n+1 are both winning initial nodes for Player Max in the game associated with $\mathcal{G}_{\varepsilon}$. Using the same arguments as above, it can be shown that there exists $\varepsilon > 0$ such that $\chi_j(g_{\varepsilon}) \ge 0$ if, and only if, $\chi_j(g_0) \ge 1/(n+1)$ or $\chi_j(g_{1/(n+1)^2}) \ge 0$.

In consequence, the support of \mathcal{R} can be computed by determining the winning initial nodes in the games associated with \mathcal{G}'_0 and $\mathcal{G}_{1/(n+1)^2}$. We point out that some policy iteration based algorithms, such as the one of [18,25], directly provide the cycle-time vector $\chi(g)$ of the dynamic programming operator g of a mean payoff game (and so all the winning initial nodes).

Remark 4.8. Positional strategies for Player Max are defined symmetrically to the ones for Player Min, *i.e.* as functions τ from row nodes to column nodes, such that for each row node *i* there is an arc in $\mathcal{G}_{\varepsilon}$ connecting it with column node $\tau(i)$. Such a strategy τ induces a one-player game (now played by Player Min) whose associated digraph $\mathcal{G}_{\varepsilon}^{\tau}$ is obtained by removing from $\mathcal{G}_{\varepsilon}$ the arcs connecting row nodes *i* with columns nodes *j* such that $j \neq \tau(i)$.

Positional strategies can be used as certificates to ensure that a mean payoff game is winning for one of the players, and these certificates can be checked in polynomial time. For instance, given a column node $j \in [n + 1]$, a strategy σ for Player Min satisfying $\chi_j(g_{\varepsilon}^{\sigma}) \leq 0$ ensures that $\chi_j(g_{\varepsilon}) \leq 0$ by (4.6). In other words, column node j is a winning initial node for Player Min in the game associated with $\mathcal{G}_{\varepsilon}$. Since as explained above $\chi_j(g_{\varepsilon}^{\sigma})$ is given by the maximal weight-to-length ratio of the circuits reachable from column node j in $\mathcal{G}_{\varepsilon}^{\sigma}$, it can be checked that $\chi_j(g_{\varepsilon}^{\sigma})$ is less than or equal to 0 in polynomial time using Karp's algorithm.

4.1.2. Polyhedra defined by systems with $+\infty$ coefficients.

We now deal with the case in which Assumption A is not satisfied. Suppose that the tropical polyhedron with mixed constraints \mathcal{R} is defined by (4.4), where now $N \in \mathbb{G}^{r \times n}$. Observe that we can still assume that $\mathbf{q} \in (\mathbb{G} \setminus \{+\infty\})^r$, because any inequality in which \mathbf{q}_i is equal to $+\infty$ is trivial.

Given $I \subset [r]$, we denote by \mathcal{R}_I the polyhedron defined by the inequalities

$$M_{i1}\boldsymbol{x}_1 \oplus \cdots \oplus M_{in}\boldsymbol{x}_n \oplus \boldsymbol{p}_i \leq_{\mathbb{G}} (\oplus_{N_{ij} \neq +\infty} N_{ij}\boldsymbol{x}_j) \oplus \boldsymbol{q}_i \quad \text{for } i \in I.$$

The algorithm in Figure 6 determines whether \mathcal{R} is empty by evaluating the emptiness of polyhedra of the form \mathcal{R}_I . To prove the correctness of this algorithm, we shall use the following lemma.

Lemma 4.9. At each iteration of the loop at Line 4, we have $\mathcal{R} \subset \mathcal{R}_{I \cup I'}$.

Proof. We prove the lemma by induction on the number of iterations of the loop. Before the first iteration, we have J = [n] and so $I \cup I' = \{i \in [r] \mid N_{ij} \neq +\infty$ for all $j \in [n]\}$. Thus, the polyhedron $\mathcal{R}_{I \cup I'}$ is defined by a subsystem of $M \boldsymbol{x} \oplus \boldsymbol{p} \leq_{\mathbb{G}} N \boldsymbol{x} \oplus \boldsymbol{q}$, and the inclusion $\mathcal{R} \subset \mathcal{R}_{I \cup I'}$ is immediate. Input: polyhedron with mixed constraints \mathcal{R} defined by the system $M \boldsymbol{x} \oplus \boldsymbol{p} \leq_{\mathbb{G}} N \boldsymbol{x} \oplus \boldsymbol{q}$ Output: true if \mathcal{R} is empty, false otherwise

1: J := [n] $2{:}\ I:=\emptyset$ 3: $I' := \{i \in [r] \mid N_{ij} \neq +\infty \text{ for all } j \in J\}$ 4: while $I' \neq \emptyset$ do $I := I \cup I'$ 5:if \mathcal{R}_I is empty then 6: 7: return true 8: else $J := \operatorname{supp}(\mathcal{R}_I)$ 9: $I' := \{ i \notin I \mid N_{ij} \neq +\infty \text{ for all } j \in J \}$ 10: end if 11:12: end while 13: return false

Fig. 6. Determining whether a polyhedron with mixed constraints is empty in the general case.

Now suppose that at iteration k we have $\mathcal{R} \subset \mathcal{R}_{I_k \cup I'_k}$, and let $x \in \mathcal{R}$. If the loop is iterated again, then the sets J, I and I' are respectively given by

· --

$$J_{k+1} = \operatorname{supp}(\mathcal{R}_{I_k \cup I'_k})$$
$$I_{k+1} = I_k \cup I'_k$$
$$I'_{k+1} = \{i \notin I_{k+1} \mid N_{ij} \neq +\infty \text{ for all } j \in J_{k+1}\}$$

In consequence, we have $supp(\mathcal{R}) \subset J_{k+1}$, and so $x_j = -\infty$ for $j \in [n] \setminus J_{k+1}$. Then, we deduce that x satisfies the inequality

 $M_{i1}\boldsymbol{x}_1 \oplus \cdots \oplus M_{in}\boldsymbol{x}_n \oplus \boldsymbol{p}_i \leqslant_{\mathbb{G}} (\oplus_{j \in J_{k+1}} N_{ij}\boldsymbol{x}_j) \oplus \boldsymbol{q}_i \leqslant_{\mathbb{G}} (\oplus_{N_{ij} \neq +\infty} N_{ij}\boldsymbol{x}_j) \oplus \boldsymbol{q}_i$

for any $i \in I'_{k+1}$. Thus, we have $\boldsymbol{x} \in \mathcal{R}_{I'_{k+1}}$. Since $\boldsymbol{x} \in \mathcal{R} \subset \mathcal{R}_{I_k \cup I'_k} = \mathcal{R}_{I_{k+1}}$, we readily obtain $\boldsymbol{x} \in \mathcal{R}_{I_{k+1} \cup I'_{k+1}}$, which completes the proof. \Box

Proposition 4.10. The algorithm of Figure 6 is correct, and the number of iterations of the loop at Line 4 is bounded by $\min(n, r)$.

Proof. Suppose that the algorithm returns **true**. Then $\mathcal{R}_I = \emptyset$, and by Lemma 4.9 we have $\mathcal{R} = \emptyset$.

Now assume that **false** is returned. Let $\boldsymbol{x} \in \mathcal{R}_I$ be such that $\{j \in [n] \mid \boldsymbol{x}_j \neq -\infty\} = \operatorname{supp}(\mathcal{R}_I) = J$. We claim that $\boldsymbol{x} \in \mathcal{R}$. Indeed, since $I' = \emptyset$ (which is the condition to reach Line 13 and return **false**), we know that for all $i \notin I$ there exists $j \in J$ such that $N_{ij} = +\infty$. As $\boldsymbol{x}_j \neq -\infty$ for $j \in J$, this means that the *i*-th inequality of the system $M\boldsymbol{x} \oplus \boldsymbol{p} \leq_{\mathbb{G}} N\boldsymbol{x} \oplus \boldsymbol{q}$ is satisfied (the right-hand side is equal to $+\infty$). As \boldsymbol{x} also satisfies the inequalities of the system indexed by $i \in I$, the claim is proved.

Finally, observe that at each iteration of the loop, the set I is strictly increased at Line 5. Similarly, if at Line 10 the set I' is not empty, then necessarily the set J has been strictly decreased at Line 9. We deduce that the number of iterations is indeed bounded by $\min(n, r)$.

The idea behind the algorithm of Figure 6 can be used to build certificates.

Proposition 4.11. The problem of determining whether a polyhedron with mixed constraints is empty belongs to NP \cap coNP.

Proof. A certificate that $\mathcal{R} \neq \emptyset$ can be provided by two sets $J \subset [n]$ and $I \subset [r]$ such that $\{i \notin I \mid N_{ij} \neq +\infty$ for all $j \in J\} = \emptyset$, together with positional strategies ensuring that $\mathcal{R}_I \neq \emptyset$ and $\mathsf{supp}(\mathcal{R}_I) = J$ (see Remark 4.8 and the discussion on supports which precedes it). The first property of the sets I and J can be checked in polynomial time, the same as the properties $\mathcal{R}_I \neq \emptyset$ and $\mathsf{supp}(\mathcal{R}_I) = J$ thanks to the positional strategies for the players. As shown in the proof of Proposition 4.10, this ensures that \mathcal{R} is not empty. In consequence, the problem is in coNP.

To certify that $\mathcal{R} = \emptyset$, we use a decreasing sequence $J_1 = [n], J_2, \ldots, J_k$ of subsets of [n] and an increasing sequence I_1, \ldots, I_k of subsets of [r] such that

$$I_l = \{i \in [r] \mid N_{ij} \neq +\infty \text{ for all } j \in J_l\},\$$

for all $l \in [k]$, together with positional strategies ensuring that $\mathcal{R}_{I_l} \neq \emptyset$ and $J_{l+1} = \sup (\mathcal{R}_{I_l})$ for $l \in [k-1]$, and that $\mathcal{R}_{I_k} = \emptyset$. Since $I_{l+1} = I_l \cup \{i \notin I_l \mid N_{ij} \neq +\infty$ for all $j \in J_{l+1}\}$ for $l \in [k-1]$, it can be shown by induction on $l \in [k]$ that $\mathcal{R} \subset \mathcal{R}_{I_l}$, using the same technique as in the proof of Lemma 4.9. Thus, these certificates allow to prove that $\mathcal{R} = \emptyset$, and they can be checked in polynomial time. This completes the proof.

4.2. Polynomial-time weak redundancy elimination

Since no polynomial-time algorithm is known to evaluate the criteria given in Section 4.1, we also develop a sufficient criterion for which a potentially faster algorithm exists. It consists in checking whether $e x \oplus g \leq_{\mathbb{G}} f x \oplus h$ is a linear combination of the inequalities in $Ax \oplus c \leq_{\mathbb{G}} Bx \oplus d$. Note that in general, this condition is not necessary for (4.1) to hold (the tropical analogue of Farkas' lemma given in [5] shows that taking tropical linear combinations does not suffice to deduce all valid inequalities).

Now we see the constraint $\boldsymbol{ex} \oplus \boldsymbol{g} \leq_{\mathbb{G}} \boldsymbol{fx} \oplus h$ as the (2n+2)-dimensional row vector $\boldsymbol{v} := (\boldsymbol{e}, \boldsymbol{g}, \boldsymbol{f}, h)$. Similarly, we introduce the matrix $R \in \mathbb{G}^{p \times (2n+2)}$, whose rows are given by the vectors $(A_i, \boldsymbol{c}_i, B_i, \boldsymbol{d}_i)$, for $i \in [p]$, where A_i and B_i denote the *i*-th rows of A and B respectively. We are reduced to the problem of determining whether there exists a p-dimensional row vector \boldsymbol{w} (with entries in \mathbb{G}) such that $\boldsymbol{v} = \boldsymbol{w}R$. Without loss of generality, we assume that no row of R is identically zero. We propose a method based on residuation theory (see *e.g.* [33]). Given $\boldsymbol{x} \in \mathbb{G}$, the self-map $\boldsymbol{z} \mapsto \boldsymbol{zx}$ on \mathbb{G} can be shown to be residuated, meaning that for each $y \in \mathbb{G}$ there exists a maximal element of the set $\{z \in \mathbb{G} \mid zx \leq_{\mathbb{G}} y\}$, denoted by y/x. Indeed, the later element is given by:

$$y \mapsto y/x := \begin{cases} (|y| - |x|)^- & \text{if } x \in \mathbb{R}, \ y \in \mathbb{R}^-, \\ |y| - |x| & \text{otherwise,} \end{cases}$$

where it is used the conventions $\alpha - (-\infty) = +\infty$ for $\alpha \in \mathbb{R}_{\max}$, $\alpha - (+\infty) = -\infty$ for $\alpha \in \mathbb{R}_{\max}$, and $(+\infty) - (+\infty) = +\infty$.

Proposition 4.12. Define the p-dimensional row vector w^* by

$$\boldsymbol{w}_i^* := \min_{j \in [2n+2]} (\boldsymbol{v}_j / R_{ij})$$

for $i \in [p]$. Then, the inequality $\mathbf{ex} \oplus g \leq_{\mathbb{G}} \mathbf{fx} \oplus h$ is a linear combination of the inequalities in the system $A\mathbf{x} \oplus \mathbf{c} \leq_{\mathbb{G}} B\mathbf{x} \oplus \mathbf{d}$ if, and only if, $\mathbf{v} = \mathbf{w}^* R$.

The principle of Proposition 4.12 is that \boldsymbol{w}^* can be shown to be the greatest solution of $\boldsymbol{w}R \leq_{\mathbb{G}} \boldsymbol{v}$. Thus, there exists a solution to $\boldsymbol{v} = \boldsymbol{w}R$ if, and only if, the equality is satisfied for $\boldsymbol{w} = \boldsymbol{w}^*$. It follows that the criterion of Proposition 4.12 can be checked efficiently, in time $O(n \times p)$.

4.3. Complexity of successive Fourier-Motzkin eliminations

As discussed in the beginning of Section 4, propagating redundant inequalities may produce $O(p^{2^k})$ constraints after k calls to Fourier-Motzkin elimination method (recall that p refers to the number of constraints defining the initial polyhedron). We claim that, in the case of closed tropical polyhedra, the number of constraints remains simply exponential at every Fourier-Motzkin elimination when the weak redundancy criterion is used. To see this, let $\mathcal{P}_0 = \mathcal{P}, \mathcal{P}_1, \ldots, \mathcal{P}_k$ be the closed tropical polyhedra arising during a sequence of k calls to Fourier-Motzkin elimination followed by the weak inequality redundancy elimination of Proposition 4.12.

Proposition 4.13. There exists a constant K (independent from k, n, and p) such that for all $l \in [k]$, the number of inequalities describing \mathcal{P}_l obtained using the weak inequality redundancy elimination is bounded by

$$K(n-l+1)p^{\lfloor n/2 \rfloor \lfloor (n-l)/2 \rfloor} .$$

$$(4.8)$$

Proof. By the tropical analogue of McMullen's upper bound theorem [4], we know that the number q of extreme generators (points and rays) of \mathcal{P} is bounded by U(p+n+1,n), where U(p,n) is the number of facets of (classical) cyclic polytopes with p extreme points in dimension n. In particular, q is bounded by $p^{\lfloor n/2 \rfloor}$ when n and p are sufficiently large (see Appendix Appendix A).

It is straightforward to see that the extreme generators of every polyhedron \mathcal{P}_l $(l \in [k])$ arise as the projection of some of the extreme generators of \mathcal{P} . In consequence, the number of extreme generators of each \mathcal{P}_l is bounded by q.

Dually, the set of inequalities satisfied by all the points of a tropical polyhedron \mathcal{Q} forms a tropical polyhedral cone called the *polar cone* of \mathcal{Q} , see [5]. As proved in [4], if \mathcal{Q} is generated by q points and rays in dimension n-l, the number of (non-trivial) extreme rays of its polar is bounded by (n-l+1)(U(q+n-l+1,n-l)-n+l+2). In particular, this also bounds the size of any description of \mathcal{Q} by linearly independent inequalities. It can be shown that there exists a constant K' > 0 such that the latter quantity is bounded by $K'(n-l+1)q^{\lfloor (n-l)/2 \rfloor}$ for all values of q and n-l (see Appendix Appendix A). It follows that the representation by inequalities obtained after the application of the weak redundancy criterion is bounded by a quantity of the form (4.8).

As a consequence, after the (l + 1)-th call to Fourier-Moztkin elimination, the number of inequalities (defining \mathcal{P}_{l+1}) is bounded by the square of (4.8), and the weak redundancy criterion can be applied in time $O((n - l)^3 p^{n(n-l)/2})$. It follows that the complexity of k successive calls to tropical Fourier-Motzkin elimination using the weak redundancy criterion can be bounded by

$$O(kn^3p^{n^2/2})$$

Therefore, the time complexity is only exponential in the worst case.

Since the criterion provided by Theorem 4.7 may eliminate further inequalities, the number of inequalities describing \mathcal{P}_l obtained using this criterion instead of the weak one is also bounded by (4.8). However, this has to be balanced with the potentially greater cost of the associated algorithm for solving mean payoff games.

5. Tropical Forward Exploration for Timed Automata

Timed automata is one of the formalisms used for modelling and verification of realtime systems. As an application of the methods developed in this paper, we show how the forward exploration algorithm for timed automata [8] can be implemented using tropical polyhedra with mixed constraints as symbolic states. This algorithm is used to solve the reachability problem, to which most verification problems for timed automata can be reduced, see [37,7].

We first recall some notions concerning timed automata, and illustrate with an example the drawbacks (mentioned in the introduction) of using zones or closed tropical polyhedra as symbolic states. Note that zones are also used as symbolic states in the verification of other real-time models such as e.g. timed Petri nets [12, 45], hence the corresponding algorithms can potentially benefit from our results too.

We consider a timed automaton over the set of clocks $C = \{x_1, \ldots, x_n\}$. It is represented by a directed graph whose nodes correspond to locations (l_0, l_1, \ldots) . We denote by l_0 the initial location and by L the set of locations. An edge from location l to l' is denoted by $l \xrightarrow{\phi, r} l'$, where ϕ represents a clock constraint and r a set of reset operations. More precisely, ϕ is a (possibly empty) conjunction of atomic clock constraints of the form $x_i \bowtie k$ and $x_i \bowtie k + x_j$, where $x_i, x_j \in C$,



Fig. 7. Left: Timed automaton. Right: Symbolic states accumulated during forward exploration, shown after delays (black borders are included in the depicted gray regions).

 $k \in \mathbb{Z}$, and $\bowtie \in \{<, \leq, =, \geq, >\}$. Besides, r is defined as a partial function from C to \mathbb{N} , meaning that \boldsymbol{x}_i is mapped to k when the clock \boldsymbol{x}_i is reset to the value k. The set of edges is denoted by E. Each location l can be additionally labeled by a clock constraint $\theta(l)$. Then, the states of the automaton are of the form (l, v), where $l \in L$ and $v : C \to \mathbb{R}_{\geq 0}$ is such that $(v(\boldsymbol{x}_1), \ldots, v(\boldsymbol{x}_n))$ satisfies the constraint $\theta(l)$. The evolution of the system (*i.e.* the *semantics* of the automaton) is expressed as a transition relation on these states, denoted by \sim . Transitions can be of two kinds:

Delays: where clock values increase synchronously at a given location. More precisely, $(l, v) \rightsquigarrow (l, v')$ if there exists $t \ge 0$ such that $v'(\boldsymbol{x}_i) := v(\boldsymbol{x}_i) + t$ for all $i \in [n]$, and $(v(\boldsymbol{x}_1) + t', \ldots, v(\boldsymbol{x}_n) + t')$ satisfies $\theta(l)$ for all $t' \in [0, t]$.

Switches: which are governed by the edges $l \xrightarrow{\phi,r} l'$. In this case, we have $(l, v) \rightsquigarrow (l', v')$ if $(v(\boldsymbol{x}_1), \ldots, v(\boldsymbol{x}_n))$ satisfies the constraints ϕ , and $v'(\boldsymbol{x}_i) := r(\boldsymbol{x}_i)$ if r is defined on $\boldsymbol{x}_i, v'(\boldsymbol{x}_i) := v(\boldsymbol{x}_i)$ otherwise.

The basic problem in the verification of timed automata is (untimed) reachability: is a final location l_f of the automaton reachable? More precisely, the reachability problem consists in determining whether there exist clock values $v_f : C \to \mathbb{R}_{\geq 0}$ and a finite path $(l_0, v_0) \rightsquigarrow^* (l_f, v_f)$ of transitions in the automata, where v_0 is the function which maps every \boldsymbol{x}_i to 0.

Example 5.1. Consider the timed-automaton fragment depicted in Figure 7, which involves two clocks x_1 and x_2 . Its edges are labeled by the reset operations (for instance, $x_1 := 0$) and/or the constraints on clocks (for instance, $x_2 > 1$). As mentioned above, the initial location is l_0 , and the two clocks are initialized to 0.

The diagrams on the right-hand side of Figure 7 depict the symbolic states of this automaton, *i.e.* the sets of states (clock values) which can arise at each location. For instance, at location l_0 we recover the initial state $\mathbf{x}_1 = \mathbf{x}_2 = 0$, and all the other states $\mathbf{x}_1 = \mathbf{x}_2 = t > 0$ arise as time goes by while staying at the same

location. Similarly, the states at location l_1 are obtained from the ones at location l_0 by resetting clock \boldsymbol{x}_1 , *i.e.* setting $\boldsymbol{x}_1 = 0$, while $\boldsymbol{x}_2 \ge 0$ is not affected. Then, as time goes by, we get all the states satisfying $0 \le \boldsymbol{x}_1 \le \boldsymbol{x}_2$. Note that the final location l_f is not reachable in this example.

The symbolic states at locations l_0 , l_1 and l_2 can be represented exactly by zones, but the one at location l_3 cannot. Hence, the symbolic state at l_3 has to be split, potentially doubling the number of symbolic states to be visited after. If several such timed-automaton fragments are concatenated, it is easy to see that this splitting of symbolic states may lead to a situation where an exponential number of zones have to be used to determine that the final location is not reachable. Alternatively, the symbolic state at location l_3 could be over-approximated by a single zone [, but in this case we cannot certify anymore that the final location l_f is not reachable. On the other hand, the use of closed tropical polyhedra to over-approximate the union of the sets of states arising from l_1 and l_2 provides the closure [of this union, which contains the point (1, 1). Then, the final location l_f becomes reachable, while it should not if strict constraints were correctly handled.

The reachability problem can be solved using a symbolic forward exploration algorithm, first given in [37], which is still used in state-of-the-art tools. This algorithm, shown in Figure 8, is usually implemented using zones (or DBMs). The algorithm explores sets of reachable states, representing them as symbolic states using zones, and performing symbolic delay and switch operations on them. We do not discuss this algorithm further, but we point out that any class of symbolic states can be used, provided that it supports the operations is_empty, is_included, intersect, reset, and delay used in the above algorithm. We now detail the definition of these operations, and show how to implement them over tropical polyhedra with mixed constraints.

The operation $is_empty(\mathcal{P})$ determines whether the polyhedron \mathcal{P} is empty. It is implemented using the methods described in Section 4.1, *i.e.* through a reduction to mean payoff games. The operation $is_included(\mathcal{P}_1, \mathcal{P}_2)$ checks if $\mathcal{P}_1 \subset \mathcal{P}_2$. This can be performed using the procedure for deciding implications of Section 4.1, by determining whether the system of inequalities defining \mathcal{P}_1 implies each defining inequality of the polyhedron \mathcal{P}_2 . The operation $intersect_{\psi}(\mathcal{P})$ computes the intersection of \mathcal{P} with the constraints in ψ . It is simply defined by appending the inequalities in ψ to the system defining \mathcal{P} , where any strict constraint in ψ is encoded as an inequality over \mathbb{G} using elements of the form λ^- . The operation $reset_{x_i:=k}(\mathcal{P})$ consists in computing the polyhedron

$$\{oldsymbol{y} \in \mathbb{R}^n_{ ext{max}} \mid oldsymbol{x} \in \mathcal{P}, \ oldsymbol{y}_j = oldsymbol{x}_j ext{ if } j
eq i, \ oldsymbol{y}_i = k \}$$
 .

It can be obtained by eliminating \boldsymbol{x}_i in the system of constraints defining \mathcal{P} using Fourier-Motzkin elimination, and then intersecting the resulting polyhedron with the constraint $\boldsymbol{x}_i = k$ (encoded as two inequalities $\boldsymbol{x}_i \leq_{\mathbb{G}} k$ and $k \leq_{\mathbb{G}} \boldsymbol{x}_i$). Finally, the operation delay(\mathcal{P}) consists in converting the polyhedron \mathcal{P} into the set $\{\lambda \boldsymbol{x} \mid$ Tropical Fourier-Motzkin Elimination, with an Application to Real-Time Verification 33

Input: timed automaton $(L, l_0, C, \theta, E), l_f \in L$ **Output:** true if $\exists v_f : C \to \mathbb{R}_{\geq 0} : (l_0, v_0) \rightsquigarrow^* (l_f, v_f)$, false otherwise 1: $Waiting := \{(l_0, \texttt{intersect}_{\theta(l_0)}(\texttt{delay}(\{v_0\})))\}; Passed := \emptyset$ 2: while $Waiting \neq \emptyset$ do Choose and remove (l, V) from Waiting 3: if $l = l_f$ then 4:return true 5: 6: end if if (not is_included(V, V')) for all $(l, V') \in Passed$ then 7: $Passed := Passed \cup \{(l, V)\}$ 8: for all $l \xrightarrow{\phi, r} l'$ do 9: $V' := \texttt{intersect}_{\theta(l')}(\texttt{delay}(\texttt{reset}_r(\texttt{intersect}_{\phi}(V)))$ 10: if not is_empty(V') then 11: Waiting := Waiting $\cup \{(l', V')\}$ 12:end if 13:14: end for 15: end if 16: end while 17: return false

Fig. 8. The symbolic forward reachability algorithm for timed automata.

 $x \in \mathcal{P}, \lambda \ge 0$ (recall that λx corresponds to the vector with entries $\lambda + x_i$). Assuming that \mathcal{P} is given by the system $Ax \oplus c \leq_{\mathbb{G}} Bx \oplus d$, we first let \mathcal{Q} be the polyhedron defined by $Ax \oplus \lambda c \leq_{\mathbb{G}} Bx \oplus \lambda d$ and $0 \leq_{\mathbb{G}} \lambda$, and then apply Fourier-Motzkin elimination on λ to get $delay(\mathcal{P})$. To prove this algorithm is correct, observe that:

$$\begin{aligned} \mathsf{delay}(\mathcal{P}) &= \{ \lambda \boldsymbol{x} \in \mathbb{R}^n_{\max} \mid 0 \leqslant_{\mathbb{G}} \lambda, \ A \boldsymbol{x} \oplus \boldsymbol{c} \leqslant_{\mathbb{G}} B \boldsymbol{x} \oplus \boldsymbol{d} \} \\ &= \{ \boldsymbol{x} \in \mathbb{R}^n_{\max} \mid 0 \leqslant_{\mathbb{G}} \lambda, \ A(\lambda^{-1}\boldsymbol{x}) \oplus \boldsymbol{c} \leqslant_{\mathbb{G}} B(\lambda^{-1}\boldsymbol{x}) \oplus \boldsymbol{d} \} \\ &= \{ \boldsymbol{x} \in \mathbb{R}^n_{\max} \mid 0 \leqslant_{\mathbb{G}} \lambda, \ A \boldsymbol{x} \oplus \lambda \boldsymbol{c} \leqslant_{\mathbb{G}} B \boldsymbol{x} \oplus \lambda \boldsymbol{d} \} \end{aligned}$$

To combat state space explosion, symbolic states are also equipped with an over-approximating union operator. In this way, symbolic states which are reached through different paths may be recombined, leading to a significant reduction in the number of symbolic states the forward exploration algorithm has to consider. Given two polyhedra with mixed constraints $\mathcal{P}, \mathcal{P}' \subset \mathbb{R}^n_{\max}$, the over-approximation union operator over_approx($\mathcal{P}, \mathcal{P}'$) is defined as the tropical convex hull of $\mathcal{P} \cup \mathcal{P}'$. Note that given systems of mixed inequalities describing \mathcal{P} and \mathcal{P}' , a system describing over_approx($\mathcal{P}, \mathcal{P}'$) can be computed by means of Proposition 3.5. As this involves 2n+2 calls to tropical Fourier-Motzkin elimination, it is crucial to implement some redundancy elimination.

Observe that the error introduced using the operation over_approx over polyhedra with mixed constraints is smaller than using zone-based over-approximation. Indeed, zones are tropically convex, and thus any zone containing $\mathcal{P} \cup \mathcal{P}'$ also contains the tropical convex hull of $\mathcal{P} \cup \mathcal{P}'$. In the example of Figure 7, the over-

approximating union by polyhedra with mixed constraints of the two sets of states arising at l_3 is exact, and given by the polyhedron defined by $\boldsymbol{x}_1, \boldsymbol{x}_2 \geq_{\mathbb{G}} 0$ and $1 \leq_{\mathbb{G}} 0^- \boldsymbol{x}_1 \oplus 0^- \boldsymbol{x}_2$, or equivalently, $1 < \max(\boldsymbol{x}_1, \boldsymbol{x}_2)$.

We have implemented a prototype of the forward exploration algorithm based on tropical polyhedra with mixed constraints. The algorithms of Section 3 and the operations described above have been implemented within the OCaml library TPLib [2], whose purpose is to provide algorithms for tropical polyhedra. It relies on the library MPGLib [6], which implements the algorithm in [25] for solving mean payoff games by policy iteration. Our prototype successfully checks that location l_f is not reachable in the timed automaton of Figure 7. In future works, we plan to apply our method on more representative examples taken from real case studies, and to compare it in terms of performance/precision with state-of-the-art tools such as UPPAAL [9].

Acknowledgments

The authors warmly thank Alexandre David for helpful discussions on the algorithms involved in the verification of timed automata. The authors are also grateful to the anonymous referee for his comments and suggestions which helped to improve the paper.

References

- M. Akian, S. Gaubert and A. Guterman, Tropical polyhedra are equivalent to mean payoff games, *Int. J. Alg. Comp.* 22(1) (2012) p. 125001 (43 pp.).
- [2] X. Allamigeon, TPLib: Tropical polyhedra library in OCaml, Available at https: //gforge.inria.fr/projects/tplib, (2012).
- [3] X. Allamigeon, S. Gaubert and E. Goubault, Computing the vertices of tropical polyhedra using directed hypergraphs, *Discrete & Computational Geometry* 49(2) (2013) 247–279.
- [4] X. Allamigeon, S. Gaubert and R. D. Katz, The number of extreme points of tropical polyhedra, J. Combin. Theory, Ser. A 118(1) (2011) 162 – 189.
- [5] X. Allamigeon, S. Gaubert and R. D. Katz, Tropical polar cones, hypergraph transversals, and mean payoff games, *Lin. Alg. Appl.* 435(7) (2011) 1549 1574.
- [6] X. Allamigeon and S. Sergeev, MPGLib: Mean payoff games library, Available at https://gforge.inria.fr/projects/mpglib, (2011).
- [7] R. Alur, C. Courcoubetis and D. Dill, Model-checking in dense real-time, *Inf. Comput.* 104(1) (1993) 2–34.
- [8] R. Alur and D. Dill, A theory of timed automata, Theor. Comput. Sci. 126 (1994) 183–235.
- [9] G. Behrmann, A. David, K. G. Larsen, J. Håkansson, P. Pettersson, W. Yi and M. Hendriks, UPPAAL 4.0, in *QEST* IEEE Computer Society, (2006) 125–126.
- [10] M. Bezem, R. Nieuwenhuis and E. Rodríguez-Carbonell, The max-atom problem and its relevance, in I. Cervesato, H. Veith and A. Voronkov (eds.), *LPAR*, **5330** of *Lect. Notes in Comput. Sci.* Springer-Verlag, (2008) 47–61.
- [11] H. Björklund and S. Vorobyov, A combinatorial strongly subexponential strategy improvement algorithm for mean payoff games, *Discrete Appl. Math.* 155 (2007) 210–229.

Tropical Fourier-Motzkin Elimination, with an Application to Real-Time Verification 35

- [12] F. D. J. Bowden, Modelling time in Petri nets, in Second Australian-Japan Workshop on Stochastic Models (1996)
- [13] W. Briec and C. Horvath, B-convexity, Optimization 53 (2004) 103–127.
- [14] W. Briec and C. Horvath, Halfspaces and Hahn-Banach like properties in B-convexity and max-plus convexity, *Pacific J. Optim.* 4(2) (2008) 293–317.
- [15] P. Butkovič, H. Schneider and S. Sergeev, Generators, extremals and bases of max cones, *Lin. Alg. Appl.* **421**(2-3) (2007) 394–406.
- [16] P. Butkovič and G. Hegedüs, An elimination method for finding all solutions of the system of linear equations over an extremal algebra, *Ekonom.-Mat. Obzor* 20(2) (1984) 203–215.
- [17] J. Chaloupka, Parallel algorithms for mean-payoff games: An experimental evaluation, in A. Fiat and P. Sanders (eds.), ESA, Lect. Notes in Comput. Sci. 5757 (Springer-Verlag, 2009) pp. 599–610.
- [18] J. Cochet-Terrasson, S. Gaubert and J. Gunawardena, A constructive fixed point theorem for min-max functions, *Dyn. and Stability of Syst.* 14(4) (1999) 407–433.
- [19] G. Cohen, S. Gaubert and J. P. Quadrat, Max-plus algebra and system theory: where we are and where to go now, *Annual Reviews in Control* 23 (1999) 207–219.
- [20] G. Cohen, S. Gaubert, J. P. Quadrat and I. Singer, Max-plus convex sets and functions, in G. L. Litvinov and V. P. Maslov (eds.), *Idempotent Mathematics and Mathematical Physics, Contemporary Mathematics* (American Mathematical Society, 2005) pp. 105–129.
- [21] A. David, UPPAAL DBM library, Available at http://people.cs.aau.dk/~adavid/ UDBM/, (2012).
- [22] A. David, J. I. Rasmussen, K. G. Larsen and A. Skou, Model-based framework for schedulability analysis using UPPAAL 4.1, in G. Nicolescu and P. J. Mosterman (eds.), *Model-Based Design for Embedded Systems* (CRC Press, 2010) pp. 93–119.
- [23] M. Develin and B. Sturmfels, Tropical convexity, Doc. Math. 9 (2004) 1–27 (electronic).
- [24] M. Develin and J. Yu, Tropical polytopes and cellular resolutions, *Exper. Math.* 16(3) (2007) 277–292.
- [25] V. Dhingra and S. Gaubert, How to solve large scale deterministic games with mean payoff by policy iteration, in *VALUETOOLS* ACM Press, (2006).
- [26] O. Friedmann, An exponential lower bound for the parity game strategy improvement algorithm as we know it, in *LICS* IEEE (August, 2009) 145–156.
- [27] S. Gaubert and J. Gunawardena, The duality theorem for min-max functions, C. R. Acad. Sci. Paris. 326, Série I (1998) 43–48.
- [28] S. Gaubert and R. D. Katz, Rational semimodules over the max-plus semiring and geometric approach to discrete event systems, Kybernetika 40(2) (2004) 153–180.
- [29] S. Gaubert and R. D. Katz, The Minkowski theorem for max-plus convex sets, *Lin. Alg. Appl.* 421(2-3) (2007) 356–369.
- [30] S. Gaubert and R. D. Katz, Minimal half-spaces and external representation of tropical polyhedra, J. Algebraic Combin. 33(3) (2011) 325–348.
- [31] S. Gaubert, R. D. Katz and S. Sergeev, Tropical linear-fractional programming and parametric mean payoff games, *Journal of Symbolic Computation* 47(12) (2012) 1447– 1478.
- [32] S. Gaubert and F. Meunier, Carathéodory, Helly and the Others in the Max-Plus World, Discrete & Computational Geometry 43(3) (2010) 648–662.
- [33] S. Gaubert and M. Plus, Methods and applications of (max, +) linear algebra, in STACS, 1200 of Lect. Notes in Comput. Sci. Springer-Verlag, (1997) 261–282.
- [34] S. Gaubert and S. Sergeev, Cyclic projectors and separation theorems in idempotent

convex geometry, Journal of Mathematical Sciences 155(6) (2008) 815-829.

- [35] S. Gaubert and S. Sergeev, The level set method for the two-sided max-plus eigenproblem, J. Disc. Event Dyn. Syst. 23(2) (2013) 105–134.
- [36] V. A. Gurvich, A. V. Karzanov and L. G. Khachivan, Cyclic games and an algorithm to find minimax cycle means in directed graphs, USSR Computational Mathematics and Mathematical Physics 28(5) (1988) 85–91.
- [37] T. A. Henzinger, X. Nicollin, J. Sifakis and S. Yovine, Symbolic model checking for real-time systems, *Inf. Comput.* 111(2) (1994) 193–244.
- [38] R. A. Howard, Dynamic Programming and Markov Processes (MIT Press, 1960)
- [39] M. Joswig, Tropical halfspaces, in Combinatorial and computational geometry, Math. Sci. Res. Inst. Publ. 52 (Cambridge Univ. Press, Cambridge, 2005) pp. 409–431.
- [40] M. Joswig, Tropical convex hull computations, in G. L. Litvinov and S. N. Sergeev (eds.), Proceedings of the International Conference on Tropical and Idempotent Mathematics, Contemporary Mathematics 495 (American Mathematical Society, 2009) pp. 193–212.
- [41] M. Jurdziński, M. Paterson and U. Zwick, A deterministic subexponential algorithm for solving parity games, in SODA ACM, (2006) 117–123.
- [42] R. D. Katz, V. Nitica and S. Sergeev, Characterization of tropical hemispaces by (P, R)-decompositions, Lin. Alg. Appl. 440(1) (2013) 131–163.
- [43] E. Kohlberg, Invariant half-lines of nonexpansive piecewise-linear transformations, Math. Oper. Res. 5(3) (1980) 366–372.
- [44] M. Kot, Modeling selected real-time database concurrency control protocols in Uppaal, ISSE 5(2) (2009) 129–138.
- [45] D. Lime, O. H. Roux, C. Seidner and L.-M. Traonouez, Romeo: A parametric modelchecker for Petri nets with stopwatches, in S. Kowalewski and A. Philippou (eds.), *TACAS*, **5505** of *Lect. Notes in Comput. Sci.* Springer-Verlag, (2009) 54–57.
- [46] G. L. Litvinov, V. P. Maslov and G. B. Shpiz, Idempotent functional analysis: an algebraic approach, *Math. Notes* 69(5) (2001) 696–729.
- [47] Q. Lu, M. Madsen, M. Milata, S. Ravn, U. Fahrenberg and K. G. Larsen, Reachability analysis for timed automata using max-plus algebra, J. Logic Alg. Prog. 81(3) (2012) 298–313.
- [48] R. H. Möhring, M. Skutella and F. Stork, Scheduling with AND/OR precedence constraints, SIAM J. Comput. 33(2) (2004) 393–415.
- [49] K. Zimmermann, A general separation theorem in extremal algebras, Ekonom.-Mat. Obzor 13(2) (1977) 179–201.
- [50] U. Zwick and M. Paterson, The complexity of mean payoff games on graphs, *Theor. Comput. Sci.* 158(1-2) (1996) 343–359.

Appendix A. Additional Details for the Proof of Proposition 4.13

We first recall that:

$$U(p,n) = \begin{cases} \binom{p-\lfloor n/2 \rfloor}{\lfloor n/2 \rfloor} + \binom{p-\lfloor n/2 \rfloor-1}{\lfloor n/2 \rfloor-1} & \text{for } n \text{ even,} \\ 2\binom{p-\lfloor n/2 \rfloor-1}{\lfloor n/2 \rfloor} & \text{for } n \text{ odd.} \end{cases}$$

Thus, if n is even,

$$U(p+n+1,n) = \binom{p+1+\lfloor n/2 \rfloor}{\lfloor n/2 \rfloor} + \binom{p+\lfloor n/2 \rfloor}{\lfloor n/2 \rfloor-1},$$

and if n is odd,

$$U(p+n+1,n) = 2 \binom{p+1+\lfloor n/2 \rfloor}{\lfloor n/2 \rfloor} .$$

In both cases, it can be easily checked that

$$U(p+n+1,n) \leq 2 \binom{p+1+\lfloor n/2 \rfloor}{\lfloor n/2 \rfloor}.$$

We claim that for p and n sufficiently large,

$$U(p+n+1,n) \leqslant p^{\lfloor n/2 \rfloor} . \tag{A.1}$$

Let $m := \lfloor n/2 \rfloor$. By Stirling approximation formulas, we know that for all positive integer h,

$$\sqrt{2\pi h}(h/e)^h \leqslant h! \leqslant e\sqrt{h}(h/e)^h$$

As a result,

$$\binom{p+1+m}{m} \leqslant \frac{e}{2\pi} \sqrt{\frac{p+1+m}{(p+1)m}} \left(1+\frac{p+1}{m}\right)^m \left(1+\frac{m}{p+1}\right)^{p+1}$$
$$\leqslant \frac{1}{2} \left(1+\frac{p+1}{m}\right)^m \left(1+\frac{m}{p+1}\right)^{p+1}$$

when $p \ge 1$ and $m \ge 2$. We next show that $\left(1 + \frac{p+1}{m}\right)^m \left(1 + \frac{m}{p+1}\right)^{p+1}$ is bounded by p^m by considering their logarithm:

$$m \ln\left(1 + \frac{p+1}{m}\right) + (p+1)\ln\left(1 + \frac{m}{p+1}\right) \leqslant$$

$$m \ln\left(1 + \frac{p+1}{m}\right) + m = m \ln\left(\frac{e(p+1+m)}{m}\right) \leqslant m \ln p$$

as soon as $e(p + m + 1) \leq pm$. The latter condition is satisfied if $p \geq 6$ and $m \geq 6$. This shows that (A.1) holds when $p \geq 6$ and $n \geq 12$.

Using the same arguments, we obtain that:

$$U(q+n-l+1,n-l)\leqslant q^{\lfloor (n-l)/2\rfloor}$$

as soon as $q \ge 6$ and $n - l \ge 12$. In any case, we can find a constant K' > 0 such that for all values of q and n - l,

$$U(q+n-l+1,n-l) \leqslant K' q^{\lfloor (n-l)/2 \rfloor}$$

Assuming that $q \leq U(p+n+1,n)$, it follows that for a certain constant K'' > 0,

$$(n-l+1)(U(q+n-l+1,n-l)-n+l+2) \leqslant K''(n-l+1)p^{\lfloor n/2 \rfloor \lfloor (n-l)/2 \rfloor}$$

if $p \ge 6$ and $n \ge 12$. This allows to show that there exists K > 0 such that for all $p, n, and l \in [n-1]$,

$$(n-l+1)(U(q+n-l+1,n-l)-n+l+2) \leq K(n-l+1)p^{\lfloor n/2 \rfloor \lfloor (n-l)/2 \rfloor}$$