



# Hölder equivalence of the value function for control-affine systems

Dario Prandi

► **To cite this version:**

Dario Prandi. Hölder equivalence of the value function for control-affine systems. ESAIM: Control, Optimisation and Calculus of Variations, EDP Sciences, 2014, 20 (4), pp.1224 - 1248. 10.1051/cocv/2014014 . hal-00817300v2

**HAL Id: hal-00817300**

**<https://hal.archives-ouvertes.fr/hal-00817300v2>**

Submitted on 26 Nov 2013

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# HÖLDER CONTINUITY OF THE VALUE FUNCTION FOR CONTROL-AFFINE SYSTEMS

DARIO PRANDI<sup>†‡</sup>

**Abstract** We prove the continuity and we give a Hölder estimate for the value function associated with the  $L^1$  cost of the control-affine system  $\dot{q} = f_0(q) + \sum_{j=1}^m u_j f_j(q)$ , satisfying the strong Hörmander condition. This is done by proving a result in the same spirit as the Ball-Box theorem for driftless (or sub-Riemannian) systems. The techniques used are based on a reduction of the control-affine system to a linear but time-dependent one, for which we are able to define a generalization of the nilpotent approximation. Finally, we also prove the continuity of the value function associated with the  $L^1$  cost of time-dependent systems of the form  $\dot{q} = \sum_{j=1}^m u_j f_j^t(q)$ .

**Key words:** control-affine systems, time-dependent systems, sub-Riemannian geometry, value function, Ball-Box theorem, nilpotent approximation.

**2010 AMS subject classifications:** 53C17, 93C15.

## 1. INTRODUCTION

A sub-Riemannian control system on a smooth manifold  $M$  is a control system in the form

$$(1) \quad \dot{\gamma}(t) = \sum_{i=1}^m u_i(t) f_i(\gamma(t)), \quad \text{a.e. } t \in [0, T],$$

where  $u : [0, T] \rightarrow \mathbb{R}^m$  is an integrable control function and  $\{f_1, \dots, f_m\}$  is a family of smooth vector fields satisfying the Hörmander condition, i.e., such that its iterated Lie brackets generate the whole tangent space at any point. The length of a curve  $\gamma$  solving (1), is then defined as the  $\text{length}(\gamma) = \min \int_0^T |u| dt$ , where the minimum is taken over all the possible  $u(\cdot)$  satisfying the above ODE. Due to the linearity of the system w.r.t.  $u$ , this length will be independent of the parametrization of  $\gamma$ . Finally, we define

$$d_{\text{SR}}(q, q') = \inf \{ \text{length}(\gamma) : \gamma : [0, 1] \rightarrow M \text{ is a solution of (1), } \gamma(0) = q \text{ and } \gamma(1) = q' \}.$$

By the Hörmander condition,  $d_{\text{SR}}$  is a distance, called Carnot-Carathéodory distance, endowing  $M$  with a natural metric space structure. A manifold considered together with a sub-Riemannian control system is called a sub-Riemannian manifold.

Define  $\Delta^1 = \text{span}\{f_1, \dots, f_m\}$  and  $\Delta^{s+1} = \Delta^s + [\Delta^s, \Delta^1]$ , for every  $s \in \mathbb{N}$ . Under the hypothesis that  $\{f_1, \dots, f_m\}$  is equiregular, i.e., that, for each  $s \in \mathbb{N}$ , the dimension of  $\Delta^s(q)$  is independent of  $q \in M$ , the Hörmander condition implies that there exists a (minimal)  $r \in \mathbb{N}$  such that  $\Delta^r(q) = T_q M$  for all  $q \in M$ . Such  $r$  is called the *degree of non-holonomy* of the sub-Riemannian control system.

---

<sup>†</sup>CENTRE NATIONAL DE RECHERCHE SCIENTIFIQUE (CNRS), CMAP, ÉCOLE POLYTECHNIQUE, ROUTE DE SACLAY, 91128 PALAISEAU CEDEX, FRANCE AND TEAM GECO, INRIA-CENTRE DE RECHERCHE SACLAY,

<sup>‡</sup>SISSA, TRIESTE, ITALY

*E-mail address:* prandi@cmap.polytechnique.fr.

This work was supported by the European Research Council, ERC StG 2009 “GeCoMethods”, contract number 239748, and by the ANR project *GCM*, program “Blanche”, project number NT09\_504490.

A fundamental result in the theory of sub-Riemannian manifolds is the celebrated Ball-Box theorem (see for example [8]). This theorem gives a rough description of the infinitesimal shape of the sub-Riemannian balls. Namely, at any point  $q$  of an equiregular sub-Riemannian manifold, the sub-Riemannian ball of small radius  $\varepsilon$  is equivalent, in privileged coordinates, to the box

$$\underbrace{[-\varepsilon, \varepsilon] \times \dots \times [-\varepsilon, \varepsilon]}_{\dim \Delta^1} \times \dots \times \underbrace{[-\varepsilon^s, \varepsilon^s] \times \dots \times [-\varepsilon^s, \varepsilon^s]}_{\dim \Delta^s - \dim \Delta^{s-1}} \times \dots \times \underbrace{[-\varepsilon^r, \varepsilon^r] \times \dots \times [-\varepsilon^r, \varepsilon^r]}_{\dim \Delta^r - \dim \Delta^{r-1}}.$$

By this we mean that, for some constant  $C > 0$ , the sub-Riemannian ball is contained in a box of side  $C\varepsilon$ , and contains a box of side  $\varepsilon/C$ .

This fact has a plethora of applications. First, it allows to prove a Hölder regularity estimate with respect to the Euclidean distance in coordinates, namely that, for  $q'$  sufficiently close to  $q$ , it holds

$$(2) \quad |q'| \lesssim d_{\text{SR}}(q, q') \lesssim |q'|^{1/r}.$$

Here we use “ $\lesssim$ ” to denote an inequality up to a multiplicative constant, independent of  $q'$ . Then, among many others, it is a fundamental step in the computation of the Hausdorff dimension of the manifold (see [21]), and it is used to obtain asymptotic estimates on the heat kernel (see e.g., [23, 20, 14, 2]). Moreover, it is the main tool in computing the asymptotic equivalents of the entropy and the complexity of curves (see e.g., [18, 19, 22, 15, 16]).

In this paper, we focus on a very important generalization of the control system (1), namely on control-affine systems. These systems are obtained by adding to (1) an uncontrolled vector field  $f_0$ , called the *drift*, and are in the form

$$(3) \quad \dot{\gamma}(t) = f_0(\gamma(t)) + \sum_{i=1}^m u_i(t) f_i(\gamma(t)), \quad \text{a.e. } t \in [0, T].$$

These kind of systems appears in plenty of applications. As an example we cite, mechanical systems with controls on the acceleration (see e.g., [10], [7]), where the drift is the velocity, or quantum control (see e.g., [12], [9]), where the drift is the free Hamiltonian. We always assume the strong Hörmander condition, i.e., that the family  $\{f_1, \dots, f_m\}$  satisfies the Hörmander condition.

The cost of a curve  $\gamma$  solving (3) is  $c_{f_0}(\gamma) = \int_0^T |u| dt$ . Unlike the sub-Riemannian length, due to the presence of the drift, the cost depends on the parametrization of the curve. Finally, the value function, between  $q, q' \in M$ , of the control system at time  $T > 0$ , is defined as

$$\rho_T^{f_0}(q, q') = \inf \{c_{f_0}(\gamma) : \gamma : [0, T'] \rightarrow M \text{ solves (3), } \gamma(0) = q, \gamma(T') = q', \text{ and } T' \leq T\}.$$

Assume now that the drift is regular, in the sense that there exists  $s \in \mathbb{N}$  such that  $f_0(q) \in \Delta^s(q) \setminus \Delta^{s-1}(q)$ , for any  $q \in M$ , where  $\Delta^s$  is defined through the vector fields  $\{f_1, \dots, f_m\}$  as before. Our main result is, then, a generalization of (2) to this context.

**Theorem 1.1.** *Let  $z = (z_1, \dots, z_n)$  be a system of privileged coordinates at  $q$  for  $\{f_1, \dots, f_m\}$ , rectifying  $f_0$  as the  $k$ -th coordinate vector field  $\partial_{z_k}$ , for some  $1 \leq k \leq n$ . Then, for sufficiently small  $T$  and  $\varepsilon$ , if  $\rho_T^{f_0}(q, q') \leq \varepsilon$  then in  $z$  coordinates it holds*

$$\text{dist} \left( z(q'), z(e^{[0, T]f_0} q) \right) \lesssim \rho_T^{f_0}(q, q') \lesssim \text{dist} \left( z(q'), z(e^{[0, T]f_0} q) \right)^{1/r}.$$

Here for any  $x \in \mathbb{R}^n$  and  $A \subset \mathbb{R}^n$ ,  $\text{dist}(x, A) = \inf_{y \in A} |x - y|$  denotes the Euclidean distance between them,  $e^{tf_0}$  denotes the flow of  $f_0$ , and  $r$  is the degree of non-holonomy of the sub-Riemannian control system defined by  $\{f_1, \dots, f_m\}$ .

In this result, instead of the Euclidean distance from the origin that appeared in (2), we have the distance from the integral curve of the drift. This is due to the fact that moving in this direction has null cost. As in the sub-Riemannian case, Theorem 1.1 is a consequence of an

estimate on the shape of the reachable sets, contained in Theorem 4.27. Moreover, although Theorem 1.1 seems a natural generalization of (2), the shape of the reachable sets described in Theorem 4.27 is much more complicated than the boxes of the sub-Riemannian case, yielding a more difficult proof. Theorem 1.1 and 4.27 represent the key step for generalizing the estimates on the complexity of curves from sub-Riemannian control systems to control-affine systems.

It is worth to mention that these results regarding control-affine systems are obtained by reducing them, as in [3], to time-dependent control systems in the form

$$(4) \quad \dot{\gamma}(t) = \sum_{i=1}^m u_i(t) f_i^t(\gamma(t)), \quad \text{a.e. } t \in [0, T],$$

where  $f_i^t = (e^{-tf_0})_* f_i$  is the pull-back of  $f_i$  through the flow of the drift. On these systems, that are linear in the control, we are able to define a good notion of approximation of the control vector fields. Namely, in Section 3.3 we will define a generalization of the nilpotent approximation, used in the sub-Riemannian context, taking into account the fact that in the system (4), exploiting the time, we can generate the direction of the brackets between  $f_0$  and the  $f_j$ s. This approximation and an iterated integral method yield the correct estimates on the reachable set, contained in Theorem 3.19.

The paper is divided in three sections. In Section 2 we recall some generalities and definitions regarding sub-Riemannian control systems, used in the following sections. In Section 3 we consider control systems in the form (4), and we prove the continuity of the value function for general time-dependent vector fields. Then, in Theorem 3.19, restricting then to the case where the time dependency is explicitly given as  $f_i^t = (e^{-tf_0})_* f_i$ , we establish some estimates on the reachable sets, in the same spirit as the Ball-Box theorem. Finally, in Section 4 we consider control-affine systems. After proving the relation between control-affine systems and time-dependent systems, we prove the continuity of the value function. Then, in Lemma 4.28, exploiting the affine nature of the control system, we give an upper bound on the time needed to join two points  $q$  and  $q'$  as a function of  $\rho_T^{f_0}(q, q')$ . From this fact and the estimates of Section 3, Theorems 4.26 and 4.27 follow. Theorem 1.1 is then a particular case of Theorem 4.26, that holds under slightly milder assumptions on  $f_0$  and  $\{f_1, \dots, f_m\}$ .

## 2. SUB-RIEMANNIAN GEOMETRY

Throughout this paper,  $M$  is an  $n$ -dimensional connected smooth manifold. In this section we recall some classical notions and results of sub-Riemannian geometry.

**2.1. Sub-Riemannian control systems.** A sub-Riemannian (or non-holonomic) control system on  $M$  is a control system in the form

$$(SR) \quad \dot{q} = \sum_{i=1}^m u_i f_i(q), \quad q \in M, \quad u = (u_1, \dots, u_m) \in \mathbb{R}^m,$$

where  $\{f_1, \dots, f_m\}$  is a family of smooth vector fields on  $M$ . We let  $f_u = \sum_{i=1}^m u_i f_i$ .

An absolutely continuous curve  $\gamma : [0, T] \rightarrow M$  is (SR)-*admissible* if there exists a control  $u \in L^1([0, T], \mathbb{R}^m)$  such that  $\dot{\gamma}(t) = f_{u(t)}(\gamma(t))$ , for a.e.  $t \in [0, T]$ . The curve is said to be *associated* to any such control. The length of  $\gamma$  is defined as

$$(5) \quad \text{length}(\gamma) = \min \|u\|_{L^1([0, T], \mathbb{R}^m)},$$

where the minimum is taken over all controls  $u$  such that  $\gamma$  is associated with  $u$ . It is attained, due to convexity reasons. Notice that, by definition,  $\text{length}(\gamma)$  is invariant under time

reparametrization of the curve. The *distance* induced by the sub-Riemannian system on  $M$  is then defined as

$$d_{\text{SR}}(q, q') = \inf \{ \text{length}(\gamma) : \gamma \text{ (SR)-admissible and } \gamma : q \rightsquigarrow q' \},$$

where  $\gamma : q \rightsquigarrow q'$  stands for  $\gamma : [0, T] \rightarrow M$ , for some  $T > 0$ ,  $\gamma(0) = q$  and  $\gamma(T) = q'$ .

Let  $\Delta$  be the  $C^\infty$ -module generated by the vector fields  $\{f_1, \dots, f_m\}$  (in particular, it is closed under multiplication by  $C^\infty(M)$  functions and summation). Let  $\Delta^1 = \Delta$ , and define recursively  $\Delta^{s+1} = \Delta^s + [\Delta^s, \Delta]$ , for every  $s \in \mathbb{N}$ . Due to the Jacobi identity  $\Delta^s$  is the  $C^\infty$ -module of linear combinations of all commutators of  $f_1, \dots, f_m$  with length  $\leq s$ . For  $q \in M$ , let  $\Delta^s(q) = \{f(q) : f \in \Delta^s\} \subset T_q M$ . We say that  $\{f_1, \dots, f_m\}$  satisfies the *Hörmander condition* (or that it is a bracket-generating family of vector fields) if  $\bigcup_{s \geq 1} \Delta^s(q) = T_q M$  for any  $q \in M$ . In the following we will always assume this condition to be satisfied.

By the Chow–Rashevsky theorem (see for instance [1]), the hypothesis of connectedness of  $M$  and the Hörmander condition guarantee the finiteness and continuity of  $d_{\text{SR}}$  with respect to the topology of  $M$ . Hence, the function  $d_{\text{SR}}$ , called *sub-Riemannian* or *Carnot-Carathéodory distance*, induces on  $M$  a metric space structure. The open balls of radius  $\varepsilon > 0$  and centered at  $q \in M$ , with respect to  $d_{\text{SR}}$ , are denoted by  $B_{\text{SR}}(q, \varepsilon)$ .

We say that a (SR)-admissible curve  $\gamma$  is a minimizer of the sub-Riemannian distance between  $q, q' \in M$  if  $\gamma : q \rightsquigarrow q'$  and  $\text{length}(\gamma) = d_{\text{SR}}(q, q')$ . Equivalently,  $\gamma$  is a minimizer between  $q, q' \in M$  if it is a solution of the free-time optimal control problem, associated with (SR),

$$(6) \quad \|u\|_{L^1(0,T)} = \int_0^T \sqrt{\sum_{j=1}^m u_j^2(t)} dt \rightarrow \min, \quad \gamma(0) = q, \quad \gamma(T) = q', \quad T > 0.$$

Indeed, the sub-Riemannian distance is the value function associated with this problem. It is a classical result that, for any couple of points  $q, q' \in M$  sufficiently close, there exists at least one minimizer.

**Remark 2.2.** The optimal control problem (6) is equivalent to the following, with  $p \geq 1$  and  $T > 0$  fixed,

$$(7) \quad \|u\|_{L^p(0,T)} = \left( \int_0^T |u|^p dt \right)^{1/p} \rightarrow \min, \quad \gamma(0) = q, \quad \gamma(T) = q',$$

In fact, due to the invariance under time reparametrization of system (SR), in (6) we can fix either  $T > 0$  or the Euclidean norm of  $u$ . Moreover, by the Hölder inequality, for any  $p > 1$ , letting  $p'$  be the conjugated exponent to  $p$  (i.e.,  $1/p + 1/p' = 1$ ), we get  $\|u\|_{L^1(0,T)} \leq T^{1/p'} \|u\|_{L^p(0,T)}$ , with the equality holding if and only if  $|u|$  is constant. From these two facts, it is easy to check that minimizers of the optimal control problem (7) coincide with the minimizers of (6) with constant norm. Thus the two optimal control problems are equivalent.

**Remark 2.3.** This control theoretical setting can be stated in purely geometric terms. Indeed, it is equivalent to a *generalized sub-Riemannian* structure. Such a structure is defined by a rank-varying smooth distribution and a Riemannian metric on it (see [1] for a precise definition). In a sub-Riemannian control system, in fact, the map  $q \mapsto \text{span}\{f_1(q), \dots, f_m(q)\} \subset T_q M$  defines a rank-varying smooth distribution, which is naturally endowed with the Riemannian norm defined, for  $v \in \Delta(q)$ , by

$$\mathbf{g}(q, v) = \inf \left\{ |u| = \sqrt{u_1^2 + \dots + u_m^2} : f_u(q) = v \right\}.$$

The pair  $(\Delta, \mathbf{g})$  is thus a generalized sub-Riemannian structure on  $M$ . Conversely, every rank-varying distribution is finitely generated, see [6, 1, 4, 13], and thus a sub-Riemannian distance can be written, globally, as the value function of a control system of the type (SR).

**2.2. Privileged coordinates and nilpotent approximation.** We now introduce the equivalent, in the sub-Riemannian context, of the linearization of a vector field. This classical procedure, called *nilpotent approximation*, is possible only in a carefully chosen set of coordinates, called *privileged coordinates*.

Since  $\{f_1, \dots, f_m\}$  is bracket-generating, the values of the sets  $\Delta^s$  at  $q$  form a flag of subspaces of  $T_qM$ ,

$$\Delta^1(q) \subset \Delta^2(q) \subset \dots \subset \Delta^r(q) = T_qM.$$

The integer  $r = r(q)$ , which is the minimum number of brackets required to recover the whole  $T_qM$ , is called *degree of non-holonomy* (or *step*) of the family  $\{f_1, \dots, f_m\}$  at  $q$ . Set  $n_s(q) = \dim \Delta^s(q)$ . The integer list  $(n_1(q), \dots, n_r(q))$  is called the *growth vector* at  $q$ . From now on we fix  $q \in M$ , and denote by  $r$  and  $(n_1, \dots, n_r)$  its degree of non-holonomy and its growth vector, respectively. Finally, let  $w_1 \leq \dots \leq w_n$  be the *weights* associated with the flag, defined by  $w_i = s$  if  $n_{s-1} < i \leq n_s$ , setting  $n_0 = 0$ .

For any smooth vector field  $f$ , we denote its action, as a derivation on smooth functions, by  $f : a \in C^\infty(M) \mapsto fa \in C^\infty(M)$ . For any smooth function  $a$  and every vector field  $f$  with  $f \neq 0$  near  $q$ , their (*non-holonomic*) *order* at  $q$  is

$$\begin{aligned} \text{ord}_q(a) &= \min\{s \in \mathbb{N} : \exists i_1, \dots, i_s \in \{1, \dots, m\} \text{ s.t. } (f_{i_1} \dots f_{i_s} a)(q) \neq 0\}, \\ \text{ord}_q(f) &= \max\{\sigma \in \mathbb{Z} : \text{ord}_q(fa) \geq \sigma + \text{ord}_q(a) \text{ for any } a \in C^\infty(M)\}. \end{aligned}$$

In particular it can be proved that  $\text{ord}_q(a) \geq s$  if and only if  $a(q') = \mathcal{O}(d_{\text{SR}}(q', q))^s$ .

**Definition 2.4.** A *system of privileged coordinates* at  $q$  for  $\{f_1, \dots, f_m\}$  is a system of local coordinates  $z = (z_1, \dots, z_n)$  centered at  $q$  and such that  $\text{ord}_q(z_i) = w_i$ ,  $1 \leq i \leq n$ .

For any point  $q \in M$  there always exists a system of privileged coordinates around  $q$ . Consider such a system  $z = (z_1, \dots, z_n)$ . We now show that this allows to compute the order of functions or vector fields in a purely algebraic way. Given a multiindex  $\alpha = (\alpha_1, \dots, \alpha_n)$  we define the weighted degree of the monomial  $z^\alpha = z_1^{\alpha_1} \dots z_n^{\alpha_n}$  as  $w(\alpha) = w_1\alpha_1 + \dots + w_n\alpha_n$  and the weighted degree of the monomial vector field  $z^\alpha \partial_{z_j}$  as  $w(\alpha) - w_j$ . Then one can prove that, given  $a \in C^\infty(M)$  and a smooth vector field  $f$ , with Taylor expansion

$$a(z) \sim \sum_{\alpha} a_{\alpha} z^{\alpha} \quad \text{and} \quad f(z) \sim \sum_{\alpha, j} f_{\alpha, j} z^{\alpha} \partial_{z_j},$$

their orders at  $q$  can be computed as

$$\text{ord}_q(a) = \min\{w(\alpha) : a_{\alpha} \neq 0\} \quad \text{and} \quad \text{ord}_q(f) = \min\{w(\alpha) - w_j : f_{\alpha, j} \neq 0\}.$$

A function or a vector field are said to be *homogeneous* if all the nonzero terms of the expansion have the same weighted degree.

We recall that, for any  $a, b \in C^\infty(M)$  and any smooth vector fields  $f, g$ , the order satisfies the following properties

$$(8) \quad \begin{aligned} \text{ord}_q(a + b) &= \min\{\text{ord}_q(a), \text{ord}_q(b)\}, & \text{ord}_q(ab) &= \text{ord}_q(a) + \text{ord}_q(b), \\ \text{ord}_q(f + g) &= \min\{\text{ord}_q(f), \text{ord}_q(g)\}, & \text{ord}_q([f, g]) &\geq \text{ord}_q(f) + \text{ord}_q(g). \end{aligned}$$

Consider  $f_i$ ,  $1 \leq i \leq m$ . By the definition of order, it follows that  $\text{ord}_q(f_i) \geq -1$ . Then we can express  $f_i$  in coordinates as

$$z_* f_i = \sum_{j=1}^n (h_{ij} + r_{ij}) \partial_{z_j},$$

where  $z_*$  is the push-forward operator on vector fields associated with the coordinates, defined as  $z_*f = dz \circ f \circ z^{-1}$ ,  $h_{ij}$  are homogeneous polynomials of weighted degree  $w_j - 1$ , and  $r_{ij}$  are functions of order larger than or equal to  $w_j$ .

**Definition 2.5.** The nilpotent approximation at  $q$  of  $f_i$ ,  $1 \leq i \leq m$ , associated with the privileged coordinates  $z$  is the vector field with coordinate representation

$$z_*\hat{f}_i = \sum_{j=1}^n h_{ij} \partial_{z_j}.$$

The nilpotentized sub-Riemannian control system is then defined as

$$(NSR) \quad \dot{q} = \sum_{j=1}^m u_j(t) \hat{f}_j(q).$$

The family of vector fields  $\{\hat{f}_1, \dots, \hat{f}_m\}$  is bracket-generating and nilpotent of step  $r$  (i.e., every iterated bracket  $[f_{i_1}, [\dots, [f_{i_{k-1}}, f_{i_k}]]]$  of length larger than  $r$  is zero).

The main consequence of the nilpotent approximation is the following (see for example [8, Proposition 7.29]).

**Proposition 2.6.** *Let  $z = (z_1, \dots, z_n)$  be a system of privileged coordinates at  $q \in M$  for  $\{f_1, \dots, f_m\}$ . For  $T > 0$  and  $u \in L^1([0, T]; \mathbb{R}^m)$ , with  $|u| \equiv 1$ , let  $\gamma(\cdot)$  and  $\hat{\gamma}(\cdot)$  be the trajectories associated with  $u$  in (SR) and (NSR), respectively, and such that  $\gamma(0) = \hat{\gamma}(0) = q$ . Then, there exist  $C, T_0 > 0$ , independent of  $u$ , such that, for any  $t < T_0$ , it holds*

$$(9) \quad |z_i(\gamma(t)) - z_i(\hat{\gamma}(t))| \leq Ct^{w_i+1}, \quad i = 1, \dots, n.$$

We recall, finally, the celebrated Ball-Box Theorem, that gives a rough description of the shape of small sub-Riemannian balls.

**Theorem 2.7** (Ball-Box Theorem). *Let  $z = (z_1, \dots, z_n)$  be a system of privileged coordinates at  $q \in M$  for  $\{f_1, \dots, f_m\}$ . Then there exist  $C, \varepsilon_0 > 0$  such that for any  $\varepsilon < \varepsilon_0$ , it holds*

$$\text{Box}\left(\frac{1}{C}\varepsilon\right) \subset B_{\text{SR}}(q, \varepsilon) \subset \text{Box}(C\varepsilon),$$

where,  $B_{\text{SR}}(q, \varepsilon)$  is identified with its coordinate representation  $z(B_{\text{SR}}(q, \varepsilon))$  and, for any  $\eta > 0$ , we let

$$(10) \quad \text{Box}(\eta) = \{z \in \mathbb{R}^n : |z_i| \leq \eta^{w_i}\},$$

Observe that the first inclusion follows directly from the definition of privileged coordinates.

As a corollary of the Ball-Box Theorem, we get the following result on the regularity of the distance.

**Corollary 2.8.** *Let  $z = (z_1, \dots, z_n)$  be a system of privileged coordinates at  $q \in M$  for  $\{f_1, \dots, f_m\}$ . Then there exists  $C, \varepsilon > 0$  such that*

$$\frac{1}{C}|z(q')| \leq d_{\text{SR}}(q, q') \leq C|z(q')|^{1/r}, \quad q' \in B_{\text{SR}}(q, \varepsilon).$$

### 3. TIME-DEPENDENT SYSTEMS

**3.1. Time-dependent control systems.** We now consider a more general situation. Namely, we consider on  $M$  the time-dependent non-holonomic control system

$$(TD) \quad \dot{q} = \sum_{i=1}^m u_i f_i^t(q), \quad q \in M, \quad u = (u_1, \dots, u_m) \in \mathbb{R}^m, \quad t \in I,$$

where  $I = [0, b)$  for some  $b \leq +\infty$  and  $\{f_1^t, \dots, f_m^t\}$  is a family of non-autonomous smooth vector fields, with smooth dependence on the time parameter. We let  $f_u^t = \sum_{i=1}^m u_i f_i^t$ .

In analogy with the autonomous case, we define (TD)-admissible curves as absolutely continuous curves  $\gamma : [0, T] \subset I \rightarrow M$  such that  $\dot{\gamma}(t) = f_{u(t)}^t(\gamma(t))$  for a.e.  $t \in [0, T]$ , for some control  $u \in L^1([0, T], \mathbb{R}^m)$ . Observe, however, that contrary to what happens in the sub-Riemannian case, the (TD)-admissibility property is not invariant under time reparametrization, e.g., a time reversal. Thus, we define the cost (and not the length) of  $\gamma$  to be

$$c(\gamma) = \min \|u\|_{L^1([0, T], \mathbb{R}^m)},$$

where the minimum is taken over all controls  $u$  such that  $\gamma$  is associated with  $u$  and is attained due to convexity. The *value function* induced by the time-dependent system is then defined as

$$\rho(q, q') = \inf \{c(\gamma) : \gamma \text{ is (TD)-admissible and } \gamma : q \rightsquigarrow q'\}.$$

Clearly, the value function is non-negative. It is not a metric since, in general, it fails both to be symmetric and to satisfy the triangular inequality. Moreover, as the following example shows,  $\rho$  could be degenerate. Namely, it could happen that  $q \neq q'$  but  $\rho(q, q') = 0$ .

**Example 3.9.** Let  $M = \mathbb{R}$ , with coordinate  $x$  and consider the vector field  $f^t = (1-t)^{-2} \partial_x$  defined on  $[0, 1)$ . For any  $x_0 \in \mathbb{R}$ ,  $x_0 \neq 0$ , and for any sequence  $t_n \uparrow 1$ , let  $u_n \in L^1([0, t_n])$  be defined as  $u_n \equiv (1-t_n)x_0$ . By definition, each  $u_n$  steers the system from 0 to  $x_0$ . Hence,

$$\rho_1(0, x_0) \leq \inf_{n \in \mathbb{N}} \|u_n\|_{L^1([0, t_n])} = \inf_{n \in \mathbb{N}} \int_0^{t_n} (1-t_n)x_0 dt = x_0 \inf_{n \in \mathbb{N}} t_n(1-t_n) = 0.$$

This proves that, for any  $x_0 \in \mathbb{R}$ ,  $\rho_1(0, x_0) = 0$ .

For  $T > 0$ ,  $q \in M$  and  $\varepsilon > 0$ , we denote the reachable set from  $q$  with cost less than  $\varepsilon$  by

$$\mathcal{R}(q, \varepsilon) = \{q' \in M : \rho(q, q') < \varepsilon\}.$$

We will also consider the reachable set from  $q$  in time less than  $T > 0$  and cost less than  $\varepsilon$ , and denote it by  $\mathcal{R}_T(q, \varepsilon)$ . Clearly  $\mathcal{R}_T(q, \varepsilon) \subset \mathcal{R}(q, \varepsilon)$ .

In general, the existence of minimizers for the optimal control problem associated with (TD) is not guaranteed. We conclude this section with an example of this fact.

**Example 3.10.** Let  $M = \mathbb{R}$ , with coordinate  $x$ , and consider the vector field  $f^t = e^{-t} \partial_x$  for  $t \in [0, 1)$ . Fix  $x_0 \in \mathbb{R}$ ,  $x_0 \neq 0$ . Observe that, for any  $T > 0$  and any control  $u \in L^1([0, T])$  steering the system from 0 to  $x_0$ , it holds

$$(11) \quad |x_0| = \left| \int_0^T u(t) e^{-t} dt \right| \leq \int_0^T |u(t)| e^{-t} dt < \|u\|_{L^1([0, T])}.$$

This implies  $\rho(0, x_0) \geq |x_0|$ . Let now  $u_n \in L^1([0, 1/n])$  be defined as  $u_n(t) = x_0 n e^t$ . Clearly  $u_n$  steers the system from 0 to  $x_0$ . Moreover,

$$\rho(0, x_0) \leq \inf_{n \in \mathbb{N}} \|u_n\|_{L^1([0, 1/n])} = |x_0| \inf_{n \in \mathbb{N}} \frac{e^{\frac{1}{n}} - 1}{\frac{1}{n}} = |x_0|.$$

This proves that  $\rho(0, x_0) = |x_0|$ . Hence, the non-existence of minimizers follows from (11).

**3.2. Finiteness and continuity of the value function.** In this section, we extend the Chow-Rashevsky Theorem to time-dependent non-holonomic systems, under the strong Hörmander condition, whose definition follows.

**Definition 3.11.** We say that a family of time-dependent vector fields  $\{f_1^t, \dots, f_m^t\}_{t \in I}$  satisfies the *strong Hörmander condition* if  $\{f_1^{t_0}, \dots, f_m^{t_0}\}$  satisfies the Hörmander condition for any  $t_0 \in I$ .



As we will see later on in Section 4, when considering families of time-dependent vector fields of the form  $f_i^t = (e^{-tf_0})_* f_i$  this condition is equivalent to the strong Hörmander condition for the affine control system with drift  $f_0$  and control vector fields  $\{f_1, \dots, f_m\}$ .

From now on we will assume that the following holds.

(H<sub>0</sub>) The family of smooth vector fields  $\{f_1^t, \dots, f_m^t\}_{t \in I}$ , depends smoothly on  $t$  and satisfies the strong Hörmander condition.

This section will be devoted to the proof of the following.

**Theorem 3.12.** *Assume that  $\{f_1^t, \dots, f_m^t\}_{t \in I}$  satisfies (H<sub>0</sub>). Then, the function  $\rho : M \times M \rightarrow [0, +\infty)$  is continuous. Moreover, for any  $t_0 \in I$  and any  $q, q' \in M$ , letting  $d_{\text{SR}}$  be the sub-Riemannian distance induced by  $\{f_1^{t_0}, \dots, f_m^{t_0}\}$ , it holds  $\rho(q, q') \leq d_{\text{SR}}(q, q')$ .*

Now, we need to introduce some notation. Following [5], the flows between times  $s, t \in \mathbb{R}$  of an autonomous vector field  $f$  and of a non-autonomous vector field  $\tau \mapsto f^\tau$  will be denoted by, respectively,

$$e^{(t-s)f} : M \rightarrow M \quad \text{and} \quad \overrightarrow{\text{exp}} \int_s^t f^\tau d\tau : M \rightarrow M.$$

Fix  $q \in M$  and assume, for the moment, that  $t_0 = 0$ . Let  $\ell \in \mathbb{N}$  and  $\mathcal{F} = (i_1, \dots, i_\ell) \in \{1, \dots, m\}^\ell$ . For any  $\mathcal{T} \in I$ ,  $\mathcal{T} > 0$ , we define the *switching end-point map* at time  $\mathcal{T}$  and associated with  $\mathcal{F}$  to be the function  $E_{\mathcal{T}, \mathcal{F}} : \mathbb{R}^\ell \rightarrow M$  defined as

$$\begin{aligned} E_{\mathcal{T}, \mathcal{F}}(\xi) &= \overrightarrow{\text{exp}} \int_{\frac{\ell-1}{\mathcal{T}}}^{\mathcal{T}} \frac{1}{\mathcal{T}} \xi_\ell f_{i_\ell}^\tau d\tau \circ \dots \circ \overrightarrow{\text{exp}} \int_0^{\frac{\mathcal{T}}{\ell}} \frac{1}{\mathcal{T}} \xi_1 f_{i_1}^\tau d\tau (q) \\ (12) \quad &= \overrightarrow{\text{exp}} \int_{\frac{\ell-1}{\mathcal{T}}}^1 \xi_\ell f_{i_\ell}^{\tau \mathcal{T}} d\tau \circ \dots \circ \overrightarrow{\text{exp}} \int_0^{\frac{1}{\ell}} \xi_1 f_{i_1}^{\tau \mathcal{T}} d\tau (q). \end{aligned}$$

Here we applied a standard change of variables formula for non-autonomous flows. Let then

$$(13) \quad g_{\mathcal{T}, \mathcal{F}}^\tau = \begin{cases} \xi_1 f_{i_1}^{\tau \mathcal{T}} & \text{if } 0 \leq \tau < 1/\ell, \\ \xi_2 f_{i_2}^{(\tau-1/\ell)\mathcal{T}} & \text{if } 1/\ell \leq \tau < 2/\ell, \\ \vdots & \\ \xi_\ell f_{i_\ell}^{(\tau-(\ell-1)/\ell)\mathcal{T}} & \text{if } (\ell-1)/\ell \leq \tau < 1, \end{cases}$$

so that we can write

$$E_{\mathcal{T}, \mathcal{F}}(\xi) = \overrightarrow{\text{exp}} \int_0^1 g_{\mathcal{T}, \mathcal{F}}^\tau(\xi) d\tau (q).$$

Clearly,  $t \mapsto \overrightarrow{\text{exp}} \int_0^t g_{\mathcal{T}, \mathcal{F}}^\tau(\xi) d\tau (q)$ ,  $t \in [0, 1]$ , is a (TD)-admissible trajectory. Thus,  $E_{\mathcal{T}, \mathcal{F}}(\xi)$ ,  $\mathcal{T} > 0$ , is the end-point of a piecewise smooth (TD)-admissible curve.

We recall that, by the series expansion of  $\overrightarrow{\text{exp}}$  (see [5]), for any non-autonomous smooth vector field  $f^\tau$ , it holds  $\overrightarrow{\text{exp}} \int_0^t f^\tau d\tau (q) = e^{t f^0} (q) + \mathcal{O}(t^2)$ . Thus, we can define

$$E_{0, \mathcal{F}}(\xi) = \lim_{\mathcal{T} \downarrow 0} E_{\mathcal{T}, \mathcal{F}}(\xi) = e^{\xi_\ell f_{i_\ell}^0} \circ \dots \circ e^{\xi_1 f_{i_1}^0} (q) = \overrightarrow{\text{exp}} \int_0^1 g_{0, \mathcal{F}}^\tau(\xi) d\tau (q),$$

where,  $g_{0, \mathcal{F}}^\tau(\xi)$  is defined in (13). Then  $t \mapsto \overrightarrow{\text{exp}} \int_0^t g_{0, \mathcal{F}}^\tau(\xi) d\tau (q)$ ,  $t \in [0, 1]$ , is an (SR)-admissible curve for the sub-Riemannian structure defined by  $\{f_1^0, \dots, f_m^0\}$  and  $E_{0, \mathcal{F}}(\xi)$  is the end-point of a piecewise smooth trajectory in (SR).

After [24], we say that a point  $q' \in M$  is (TD)-*reachable* from  $q$  at time  $t_0 = 0$ , if there exist  $\ell \in \mathbb{N}$ ,  $\mathcal{F} \in \{1, \dots, m\}^\ell$ ,  $\mathcal{T} > 0$  and  $\xi \in \mathbb{R}^\ell$ , such that  $E_{\mathcal{T}, \mathcal{F}}(\xi) = q'$ . In this case it is clear that  $\rho(q, q') \leq \sum_i |\xi_i|$ . Moreover, if  $\xi' \mapsto E_{\mathcal{T}, \mathcal{F}}(\xi')$  has rank  $n$  at  $\xi$ , the point  $q'$  is said to be

(TD)-normally reachable at time  $t_0 = 0$ . Finally, the point  $q'$  is said to be (SR)-reachable or (SR)-normally reachable for the vector fields  $\{f_1^0, \dots, f_m^0\}$ , if these properties holds for  $\mathcal{T} = 0$ .

In the case  $t_0 > 0$ , taking  $\mathcal{T} > 0$  such that  $\mathcal{T} + t_0 \in I$  and changing the interval of integration in (12) from  $[0, \mathcal{T}]$  to  $[t_0, t_0 + \mathcal{T}]$ , it is clear how to define (TD)-reachable and (TD)-normally reachable points from  $q$  at time  $t_0$ , and (SR)-reachable and (SR)-normally reachable points for the vector fields  $\{f_1^{t_0}, \dots, f_m^{t_0}\}$ .

The proof of the following lemma is an adaptation of [24, Lemma 3.1].

**Lemma 3.13.** *Let  $q' \in M$  be (SR)-normally reachable for the vector fields  $\{f_1^{t_0}, \dots, f_m^{t_0}\}$  from  $q$ , by some  $\ell \in \mathbb{N}$ ,  $\xi \in \mathbb{R}^\ell$  and  $\mathcal{F} \in \{1, \dots, m\}^\ell$ . Then, there exist  $\varepsilon_0, \mathcal{T}_0 > 0$  such that, for any  $\varepsilon < \varepsilon_0$ , the point  $q'$  is (TD)-normally reachable at time  $t_0$ , by the same  $\ell$  and  $\mathcal{F}$ , and some  $\xi' \in \mathbb{R}^\ell$ , with  $\sum_j |\xi_j - \xi'_j| \leq \varepsilon$ , and any  $\mathcal{T} < \mathcal{T}_0$ .*

*Proof.* Without loss of generality, we assume  $t_0 = 0$ .

Let  $U \subset \mathbb{R}^\ell$  be a neighborhood of  $\xi$  such that  $E_{0, \mathcal{F}}$  has still rank  $n$  when restricted to it. Then, there exists  $B = \{x: \sum_j |x_j - \xi_j| \leq \varepsilon\} \subset U$  such that  $E_{0, \mathcal{F}}$  maps diffeomorphically a neighborhood of  $B$  in  $U$  onto a neighborhood of  $q$ . It follows, from standard properties of differential equations, that, for  $\mathcal{T} > 0$  sufficiently small, the map  $E_{\mathcal{T}, \mathcal{F}}$  is well defined on  $B$  and that  $E_{\mathcal{T}, \mathcal{F}} \rightarrow E_{0, \mathcal{F}}$  as  $\mathcal{T} \downarrow 0$  in the  $C^1$ -topology over  $B$ . Thus, there exists  $\mathcal{T}_1 > 0$  such that, for  $\mathcal{T} < \mathcal{T}_1$ ,  $E_{\mathcal{T}, \mathcal{F}}$  has rank  $n$  at every point of  $B$ .

Since the map  $E_{0, \mathcal{F}}$  is an homeomorphism from  $B$  onto a neighborhood of  $q$ , and  $E_{\mathcal{T}, \mathcal{F}} \rightarrow E_{0, \mathcal{F}}$  uniformly as  $\mathcal{T} \downarrow 0$ , it follows that there exists a fixed neighborhood  $V$  of  $q$  and  $\mathcal{T}_2 > 0$  such that  $V \subset E_{\mathcal{T}, \mathcal{F}}(B)$ , for any  $\mathcal{T} < \mathcal{T}_2$ . Then, for any  $\mathcal{T} < \min\{\mathcal{T}_1, \mathcal{T}_2\}$ , there exists  $\xi' \in B$  such that the point  $q' = E_{\mathcal{T}, \mathcal{F}}(\xi')$  is (TD)-normally reachable.  $\square$

We will use the following consequence of Lemma 3.13. We remark that the result holds even if  $\{f_1^t, \dots, f_m^t\}_{t \in I}$  satisfies the Hörmander condition only at time  $t_0 \in I$ .

**Lemma 3.14.** *Let  $d_{\text{SR}}$  be the sub-Riemannian distance induced by  $\{f_1^{t_0}, \dots, f_m^{t_0}\}$ , then for any  $t_1 \in I$ , such that  $t_1 - t_0 > 0$  is sufficiently small, and for any  $q, q' \in M$  it holds*

$$\inf\{c(\gamma): \gamma: [t_0, t_1] \rightarrow M \text{ is (TD)-admissible, } \gamma(t_0) = q \text{ and } \gamma(t_1) = q'\} \leq d_{\text{SR}}(q, q').$$

*In particular,  $\rho(q, q') \leq d_{\text{SR}}(q, q')$ .*

*Proof.* Fix  $\varepsilon > 0$ . By Chow's theorem it is clear that  $q'$  is (SR)-reacheable from  $q$ . Moreover, since there exist (SR)-normally reachable points from  $q'$  arbitrarily close to  $q'$  (see e.g., [1, Lemma 3.21]), follows that  $q'$  is always (SR)-normally reacheable from  $q$  by  $\xi$  such that  $\sum_j |\xi_j| \leq d_{\text{SR}}(q, q') + \varepsilon/2$ . Hence, by Lemma 3.13, if  $\varepsilon$  and  $\eta > 0$  are sufficiently small, we have that  $q'$  is (TD)-normally reachable from  $q$  at time  $t_0$  by  $\xi'$  such that  $\sum_j |\xi'_j| \leq d_{\text{SR}}(q, q') + \varepsilon$  and  $T < t_1$ . This clearly implies that

$$\inf\{c(\gamma): \gamma \text{ is (TD)-admissible, } \gamma(t_0) = q \text{ and } \gamma(t_1) = q'\} \leq d_{\text{SR}}(q, q') + \varepsilon.$$

Finally, the lemma follows letting  $\varepsilon \downarrow 0$ .  $\square$

We now prove the main theorem of the section.

*Proof of Theorem 3.12.* By Lemma 3.14, we only need to prove the continuity of  $\rho$ . We will prove only the lower semicontinuity, sinche the upper semicontinuity follows by similar arguments.

We start by proving the lower semicontinuity of  $\rho(q, \cdot)$  at  $q'$ . Consider a sequence  $q_k \rightarrow q'$  and let  $u_k \in L^1([0, T_k], \mathbb{R}^m)$  be controls such that each one steers system (TD) from  $q$  to  $q_k$  and  $\liminf_n \rho^{f_0}(q, q_k) = \liminf_n \|u_k\|_{L^1}$ . Then, by Lemma 3.14, for any  $\varepsilon > 0$  there exists a sequence

of  $\tilde{T}_k > 0$  and a sequence of controls  $v_k \in L^1([T_k, \tilde{T}_k], \mathbb{R}^m)$  all steering system (TD) from  $q_k$  to  $q'$  and such that  $\|v_k\|_{L^1([T_k, \tilde{T}_k], \mathbb{R}^m)} \leq d_{\text{SR}}(q_k, q') + \varepsilon$ . Since  $d_{\text{SR}}(q_k, q') \rightarrow 0$ , this implies that

$$\rho(q, q') \leq \lim_{n \rightarrow \infty} \left( \|u_k\|_{L^1([0, T_k], \mathbb{R}^m)} + \|v_k\|_{L^1([T_k, \tilde{T}_k], \mathbb{R}^m)} \right) = \liminf_n \rho(q, q_k) + \varepsilon.$$

Letting  $\varepsilon \downarrow 0$  proves that  $\rho(q, \cdot)$  is lower semicontinuous at  $q'$ .

In order to prove the lower semicontinuity of  $\rho(\cdot, q')$  at  $q$ , let us define

$$\varphi_\varepsilon(p) = \inf\{c(\gamma) : \gamma : [\varepsilon, T] \subset I \rightarrow M \text{ is (TD)-admissible and } \gamma : p \rightsquigarrow q'\}.$$

We claim that for any  $p \in M$  it holds that  $\varphi_\varepsilon(p) \rightarrow \rho(p, q')$  as  $\varepsilon \downarrow 0$ . Since it is clear that  $\varphi_\varepsilon(\cdot) \geq \rho(\cdot, q')$ , it suffices to prove that

$$(14) \quad \lim_{\varepsilon \downarrow 0} \varphi_\varepsilon(p) \leq \rho(p, q') \quad \text{for any } p \in M.$$

To this aim, fix  $p \in M$  and  $\eta > 0$  and let  $\gamma : [0, T] \rightarrow M$  be such that  $c(\gamma) \leq \rho(p, q') + \eta$ . It is clear that  $\gamma(2\varepsilon) \rightarrow p$  as  $\varepsilon \downarrow 0$ , and hence that  $\rho(p, \gamma(2\varepsilon)) \rightarrow 0$  as  $\varepsilon \downarrow 0$ , by the first part of the proof. Thus, for any  $\varepsilon > 0$  sufficiently small, there exists a (TD)-admissible curve  $\gamma_\varepsilon : [\varepsilon, 2\varepsilon] \rightarrow M$  such that  $\gamma_\varepsilon : p \rightsquigarrow \gamma(2\varepsilon)$  and  $c(\gamma_\varepsilon) \leq \rho(p, \gamma(2\varepsilon)) + \eta$ . By concatenating  $\gamma_\varepsilon$  with  $\gamma|_{[2\varepsilon, T]}$ , we get that

$$\varphi_\varepsilon(p) \leq c(\gamma_\varepsilon) + c(\gamma) \leq \rho(p, \gamma(2\varepsilon)) + \rho(p, q') + 2\eta.$$

Letting  $\varepsilon \downarrow 0$  and then  $\eta \downarrow 0$ , this proves (14) and thus the claim.

Let now  $q_k \rightarrow q$  and fix  $\eta > 0$ . By Lemma 3.14 this implies that  $\rho(q_k, q) \rightarrow 0$  and that for any  $\varepsilon > 0$  sufficiently small, there exists a (TD)-admissible curve  $\gamma_\varepsilon : [0, \varepsilon] \rightarrow M$  such that  $\gamma_\varepsilon : q_k \rightsquigarrow q$  and  $c(\gamma_\varepsilon) \leq \rho(q_k, q) + \eta$ . Hence

$$\rho(q_k, q') \leq c(\gamma_\varepsilon) + \varphi_\varepsilon(q) \leq \rho(q_k, q) + \varphi_\varepsilon(q) + \eta.$$

By the previous claim, letting  $\varepsilon, \eta \downarrow 0$ , this implies that  $\rho(q_k, q') \leq \rho(q_k, q) + \rho(q, q')$ . Since  $\rho(q_k, q) \rightarrow 0$ , taking the liminf as  $k \rightarrow +\infty$ , this proves the lower semicontinuity of  $\rho(\cdot, q')$  at  $q$ , completing the proof.  $\square$

**Remark 3.15.** From the proof of Theorem 3.12, it follows that hypothesis (H<sub>0</sub>) is not sharp. Indeed, the following is sufficient to prove the theorem.

- The family of smooth vector fields  $\{f_1^t, \dots, f_m^t\}_{t \in I}$ , depends smoothly on  $t$ ,  
(H<sub>1</sub>) and satisfies the strong Hörmander condition at  $t = 0$  and in an open neighborhood of  $\sup I$ .

We will conclude this section by showing that, in our framework, it is necessary to assume the Hörmander condition on both ends of  $I$ . Although outside the scope of the present work, we remark that stronger assumptions on the regularity of the vector fields, i.e., that they are uniformly Lipschitz, would allow to prove Theorem 3.12 assuming only that  $\{f_1^t, \dots, f_m^t\}_{t \in I}$  satisfies the Hörmander condition at one time  $t_0 \in I$ .

The following example proves that if the family  $\{f_1^t, \dots, f_m^t\}_{t \in I}$  satisfies the Hörmander condition only near  $t = 0$ , then the value function is in general not continuous. Through a slight modification, the same argument can also be used to prove that the same holds if the Hörmander condition is satisfied only at a neighborhood of  $\sup I$  or of any  $t_0 \in I$ .

**Example 3.16.** Let  $M = (-2, 2) \times (-1, +\infty)$ , with coordinates  $(x, y)$ , and consider the vector fields

$$f(x, y) = \frac{((y+1)(1-x^2), -x)}{\sqrt{(y+1)^2(1-x^2)^2 + x^2}}, \quad g(x, y) = \frac{(x, h(x)(y+2))}{\sqrt{x^2 + h(x)^2(y+2)^2}},$$




FIGURE 1. The two vector fields of Example 3.16 with  $h(x) = ce^{-\frac{1}{1-x^2}}$  for  $x \in [-1, 1]$ .

where  $h : [-2, 2] \rightarrow \mathbb{R}$  is a smooth cutoff function such that  $\text{supp } h \subset [-1, 1]$ ,  $h \geq 0$  and  $h(0) = 1$  (see Figure 1). Fix  $0 < \varepsilon < 1$ ,  $C \geq 16$  and let  $\phi, \psi : [0, 1] \rightarrow \mathbb{R}$  be two smooth functions such that

$$\phi(t) = \begin{cases} 1 & \text{if } 0 \leq t \leq \varepsilon, \\ 0 & \text{if } 2\varepsilon \leq t \leq 1, \end{cases} \quad \psi(t) = \begin{cases} 1 & \text{if } 0 \leq t \leq 2\varepsilon, \\ C & \text{if } 3\varepsilon \leq t \leq 1, \end{cases}$$

and such that  $\phi$  is nonincreasing while  $\psi$  is nondecreasing. Finally, consider the time-dependent system on  $M$  specified by the vector fields  $f^t(x, y) = \phi(t)f(x, y)$  and  $g^t(x, y) = \psi(t)g(x, y)$ ,  $t \in [0, 1]$ . We will show that  $\{f^t, g^t\}$  satisfies the Hörmander condition for  $t \in [0, \varepsilon]$ , but that the value function associated with the family  $\{f^t, g^t\}_{t \in [0, 1]}$  is not lower semicontinuous.

We start by showing that  $f(p)$  and  $g(p)$  are transversal for any  $p = (x, y) \in M$ , implying the Hörmander condition for  $\{f^t, g^t\}$ ,  $t \in [0, \varepsilon]$ . If  $x \in (-2, -1] \cup [1, 2)$ , then, by definition of  $h$ ,  $g(p) = (1, 0)$  is clearly transversal to  $f(p)$ . On the other hand, if  $x \in (-1, 1) \setminus \{0\}$  and  $g(p)$  is parallel to  $f(p)$ , a simple computation shows that  $h(x) < 0$ , which is a contradiction. Finally, for  $x = 0$ , it is clear that  $g(p) = (0, y + 2)$  and  $f(p) = (y + 1, 0)$  are never parallel. We remark that this implies also that the value function  $\rho_\varepsilon$ , induced by controls defined on  $[0, \varepsilon]$ , is a distance equivalent to the Euclidean one. In particular,  $|p_1 - p_2| \leq 2\rho_\varepsilon(p_1, p_2)$  for any  $p_1, p_2 \in M$ .

Fix now  $q' = (1, 0)$ . The set of points from which  $q'$  is reachable using only  $f$  is exactly  $\mathcal{O}_{q'} = \{(1, y) : y > -1\}$ . Let then  $q_0 \in (-1, 0) \times \{0\}$  be such that  $\rho_\varepsilon(q_0, (-1, 0)) \leq \frac{1}{4} \min_{p \in \mathcal{O}_{q'}} \rho_\varepsilon(q_0, p)$ . In order to show that  $\rho_1(q_0, \cdot)$  is not lower semicontinuous at  $q'$ , consider any sequence  $\{q_n\}_{n \in \mathbb{N}} \subset (1/2, 1) \times \{0\}$  such that  $q_n \rightarrow q'$ . By continuity of  $\rho_\varepsilon$  and the fact that  $-q_n \rightarrow (-1, 0)$ , we can always assume that, up to subsequences,  $\rho_\varepsilon(q_0, -q_n) \leq \frac{1}{2} \min_{p \in \mathcal{O}_{q'}} \rho_\varepsilon(q_0, p)$ .

Since  $g^t \equiv 0$  for  $t \geq 2\varepsilon$ , if  $u \in L^1([0, 1], \mathbb{R}^2)$  is a control steering the system from  $q_0$  to  $q'$ , the control  $u|_{[0, 2\varepsilon]}$  steers the system from  $q_0$  to some  $p \in \mathcal{O}_{q'}$ . Exploiting the fact that  $\rho_{2\varepsilon} \geq \rho_\varepsilon$  by monotonicity of  $\psi$ , this implies that

$$(15) \quad \rho_1(q_0, q') \geq \min_{p \in \mathcal{O}_{q'}} \rho_\varepsilon(q_0, p) \geq 2\rho_\varepsilon(q_0, -q_n).$$

Let now  $u \in L^1([0, 1], \mathbb{R}^m)$  be the control constructed as follows. From time 0 to  $\varepsilon$ ,  $u|_{[0, \varepsilon]}$  is the minimizer of  $\rho_\varepsilon$  steering the system from  $q_0$  to  $-q_n$ . Then,  $u|_{(\varepsilon, 3\varepsilon)} \equiv 0$  and, after this, the control acts only on  $f^t$  for time  $t \in [3\varepsilon, 1]$ , steering the system from  $-q_n$  to  $q_n$ . Hence,

$$(16) \quad |q_n - (-q_n)| = \left| \int_{3\varepsilon}^1 u(t) f^t(x(t), y(t)) dt \right| = C \int_{3\varepsilon}^1 |u(t)| dt.$$

Since  $|q_n - (-q_n)| < 2$ ,  $C \geq 16/|q_0 - q'| \geq 8/\rho_\varepsilon(q_0, q')$ , and by (16), it holds that

$$\rho_1(q_0, q_n) \leq \int_0^1 |u(t)| dt = \rho_\varepsilon(q_0, -q_n) + \frac{1}{C} |q_n - (-q_n)| \leq \frac{3}{4} \rho_1(q_0, q').$$

Taking the  $\liminf$  as  $n \rightarrow \infty$  shows that  $\rho_1(q_0, \cdot)$  is not l.s.c. at  $q'$ .

**3.3. Estimates on reachable sets.** In this section, we concentrate on a particular class of time-dependent systems. Namely, let  $\{f_1, \dots, f_m\}$  be a bracket-generating family of smooth vector fields,  $f_0$  be a smooth vector field, and consider the time-dependent system

$$(17) \quad \dot{q} = \sum_{i=1}^m u_i f_i^t, \quad f_i^t = (e^{-tf_0})_* f_i(q), \quad q \in M, \quad u = (u_1, \dots, u_m) \in \mathbb{R}^m.$$

Here,  $(e^{-tf_0})_*$  is the push-forward operator associated with the flow of  $f_0$ .

As we will see in the next section, this class of systems arises naturally when dealing with control systems that are affine with respect to the control. Observe, in particular, that from the bracket-generating property of  $\{f_1, \dots, f_m\}$  it follows immediately that the time-dependent family  $\{(e^{-tf_0})_* f_1, \dots, (e^{-tf_0})_* f_m\}$  satisfies the strong Hörmander condition, as per Definition 3.11.

Before proceeding with the estimates of the reachable sets, we need to define a suitable approximation of system (17). Namely, fix a system of privileged coordinates (in the sub-Riemannian sense) at  $q$  for  $\{f_1, \dots, f_m\}$ . Assume that  $f_0(q) \neq 0$ , and let  $s \in \{1, \dots, r\}$  be such that  $\text{ord}_q(f_0) = -s$ . In this case, there exist, in coordinates, an homogeneous vector field  $f_0^{-s}$ , of weighted degree  $-s$ , and a vector field  $f_0^{>-s}$ , of weighted degree  $\geq -s + 1$ , such that

$$(18) \quad z_* f_0 = f_0^{-s} + f_0^{>-s}.$$

In particular, it holds that  $f_0^{-s} \neq 0$  near  $z(q) = 0$ .

**Remark 3.17.** The fact that  $\text{ord}_q(f_0) = -s$  is not equivalent, in general, to  $f_0 \in \Delta^s$  near  $q$ . In particular, if the growth vector is non-constant around  $q$ , from  $\text{ord}_q(f_0) \geq -s$  it does not follow that  $f_0 \in \Delta^s$ . For example, consider the sub-Riemannian control system on  $\mathbb{R}^2$  with (privileged) coordinates  $(x, y)$ , defined by the vector fields  $\partial_x$  and  $x\partial_y$ , called the Grushin plane. Outside  $\{x = 0\}$ , the non-holonomic degree of these vector fields is 1, while, on  $\{x = 0\}$ , we need one bracket to generate the  $y$  direction, and thus it is 2. Hence, if  $\bar{y} \neq 0$  the vector field  $y\partial_y$  is never in  $\Delta$  near  $(0, \bar{y})$ , but  $\text{ord}_{(0, \bar{y})}(y\partial_y) = \text{ord}_{(0, \bar{y})}(y) + \text{ord}_{(0, \bar{y})}(\partial_y) = 0$ .

However, due to the properties (8) and the fact that  $\Delta^s$  is a module, the converse is always true. Namely, if  $f_0 \in \Delta^s$ , then  $\text{ord}_q(f_0) \geq -s$ .

For any smooth vector field  $f$ , let  $(\text{ad}^1 f_0)f = [f_0, f]$  and  $(\text{ad}^\ell f_0)f = [f_0, (\text{ad}^{\ell-1} f_0)f]$ , for any  $\ell \in \mathbb{N}$ . We recall (see for example [17]) that the following Taylor expansion holds

$$(19) \quad (e^{-tf_0})_* f \sim \sum_{\ell=0}^{\infty} \frac{t^\ell}{\ell!} (\text{ad}^\ell f_0)f.$$

Since  $\text{ord}_q(f_j) \geq -1$ , by (8) we have that  $\text{ord}_q((\text{ad}^\ell f_0)f_j) \geq -\ell s - 1$ . Then, using the decomposition (18), for any  $\ell \geq 0$ , there exists, in coordinates, an homogeneous vector field  $F_j^\ell$  of weighted

degree  $-\ell s$ , and a remainder  $r^\ell$  of order  $\geq -\ell s - 1$ , such that

$$(20) \quad z_*[(\text{ad}^\ell f_0)f_j] = F_j^\ell + r^\ell.$$

**Definition 3.18.** The *homogeneous series approximation* at  $q$  of  $f_j^t$ ,  $1 \leq j \leq m$ , associated with the privileged coordinates  $z$ , is the vector field with coordinate representation

$$(21) \quad \widehat{f}_j^t = \sum_{\ell=0}^{\varrho} \frac{t^\ell}{\ell!} F_j^\ell,$$

where  $\varrho = \lfloor r^{-1}/s \rfloor$  and  $r$  is the non-holonomic degree of  $\{f_1, \dots, f_m\}$  at  $q$ . The approximated time-dependent control system is then defined as

$$(ATD) \quad \dot{q} = \sum_{j=1}^m u_j(t) \widehat{f}_j^t(q).$$

If a system, in some system of privileged coordinates, coincides with its homogeneous series approximation, we will say that it is series homogeneous.

The homogeneous series approximation encodes the idea that the time  $t$  is of weight  $s = -\text{ord}_q(f_0)$ . This is a consequence of the fact that, due to the expansion (19),  $t$  allows to build brackets of  $f_0$  with the  $f_j$ s. In this sense, the homogeneous series approximation is a generalization of the nilpotent approximation.

We are now ready to state the main theorem of this section.

**Theorem 3.19.** *Let  $\{f_1, \dots, f_m\}$  satisfy the Hörmander condition, let  $z = (z_1, \dots, z_n)$  be a system of privileged coordinates at  $q \in M$  for  $\{f_1, \dots, f_m\}$ . Then there exist  $C, T, \varepsilon_0 > 0$  such that, for any  $\varepsilon < \varepsilon_0$  and any  $q' \in \mathcal{R}_T(q, \varepsilon)$ , setting  $s = -\text{ord}_q(f_0)$  it holds*

$$(22) \quad |z_i(q')| \leq C \left( \varepsilon^{w_i} + \varepsilon T^{\frac{w_i}{s}} \right) \quad \text{if } w_i \leq s,$$

$$(23) \quad |z_i(q')| \leq C \varepsilon \left( \varepsilon + T^{\frac{1}{s}} \right)^{w_i - 1} \quad \text{if } w_i > s.$$

Moreover, if the system is series homogeneous, then it holds the stronger estimate

$$(24) \quad |z_i(q')| \leq C \varepsilon^{w_i} \quad \text{if } w_i \leq s.$$

To prove this theorem we need the following proposition, estimating the difference between (17) and (ATD).

**Proposition 3.20.** *Let  $\{f_1, \dots, f_m\}$  satisfy the Hörmander condition, and let  $z = (z_1, \dots, z_n)$  be a system of privileged coordinates at  $q \in M$  for  $\{f_1, \dots, f_m\}$ . For  $T > 0$  and  $u \in L^1([0, T]; \mathbb{R}^m)$ , let  $\gamma(\cdot)$  and  $\widehat{\gamma}(\cdot)$  be the trajectories associated with  $u$  in (17) and (ATD), respectively, and such that  $\gamma(0) = \widehat{\gamma}(0) = q$ . Then there exist  $C, \varepsilon_0, T_0 > 0$ , independent of  $u$ , such that, if  $t < T_0$  and  $\int_0^t |u| ds = \varepsilon < \varepsilon_0$ , and setting  $s = -\text{ord}_q(f_0)$  it holds*

$$(25) \quad |z_i(\gamma(t)) - z_i(\widehat{\gamma}(t))| \leq C \varepsilon \left( \varepsilon + t^{\frac{1}{s}} \right)^{w_i}, \quad i = 1, \dots, n.$$

**Remark 3.21.** This proposition generalizes Proposition 2.6. In fact, in the sub-Riemannian case, since  $f_0 \equiv 0$ , if, for  $t > 0$ , the curve  $\gamma$  is associated to  $u \in L^1([0, t], \mathbb{R}^m)$ , it is associated also to  $u_\tau(\cdot) = \frac{\tau}{t} u(\frac{\cdot}{t})$ , for any  $\tau > 0$ . Thus, since  $\int_0^\tau |u_\tau| ds = \int_0^t |u| ds = \varepsilon$ , (25) reduces to

$$|z_i(\gamma(t)) - z_i(\widehat{\gamma}(t))| \leq \lim_{\tau \downarrow 0} C (\varepsilon^{w_i+1} + \tau \varepsilon^\tau) = C \varepsilon^{w_i+1}.$$

Finally, assuming that  $u$  satisfies the hypotheses of Proposition 2.6, i.e., that  $|u| = 1$ , we get  $t = \varepsilon$ .

*Proof.* Let  $z(\gamma(\cdot)) = x(\cdot)$ ,  $z(\hat{\gamma}(\cdot)) = y(\cdot)$ , and  $\|z\| = \sum_{\ell=1}^n |z_\ell|^{1/w_i}$ . We mimic the proof of Proposition 7.29 in [8]. The first step is to prove that there exists a constant  $C > 0$  such that  $\|x(t)\|, \|y(t)\| \leq C\varepsilon$  for  $t$  and  $\varepsilon = \int_0^t |u| ds$  small enough. We prove this for  $\|x(t)\|$ , the same argument works also for  $\|y(t)\|$ .

In  $z$  coordinates, the equation of the control system (17) is,

$$\dot{x}_i(t) = \sum_{j=1}^m u_j(t) (z_i)_* f_j^t(\gamma(t)), \quad i = 1, \dots, n.$$

Due to the fact that  $z_* f_j^t = z_* f_j + \mathcal{O}(t)$ , uniformly in a neighborhood of  $q$  that  $\text{ord}_q(z_i) = w_i$  and that  $\text{ord}_q(f_j) \geq -1$ , we have that there exist  $T_0$  and  $C > 0$  such that  $|(z_i)_* f_j^t(q)| \leq \frac{C}{2} |(z_i)_* f_j(q)| \leq C \|x(t)\|^{w_i-1}$ , for any  $t < T_0$ . Thus we get

$$(26) \quad |\dot{x}_i(t)| \leq C \sum_{j=1}^m |u_j(t)| \|x_i(t)\|^{w_j-1}.$$

As in the proof for the sub-Riemannian case, choosing  $N$  sufficiently large, so that all  $N/w_i$  are even integers, and setting  $\|z\| = (\sum_{\ell=1}^n |z_\ell|^{N/w_i})^{1/N}$  we get a norm equivalent to  $\|z\|$ . Deriving with respect to time and using (26) we get  $\frac{d}{dt} \|x(t)\| \leq C \sum_{j=1}^n |u_j(t)|$ . Finally, by integration, equivalence of the norms, and the fact that  $x(0) = z(q) = 0$ , we conclude that  $\|x(t)\| \leq C\varepsilon$ .

Now we move to proving (25). By construction of (ATD) and the Taylor expansion of  $f_j^t$ , for any  $\ell \leq \varrho = \lfloor r-1/s \rfloor$ , there exist homogeneous polynomials  $h_{ji}^\ell$  of order  $w_i - \ell s - 1$  and remainders  $r_{ji}^\ell$  of order larger than or equal to  $w_i - \ell s$ , such that we can write

$$(z_i)_* f_j^t = \sum_{\ell=0}^{\varrho} \frac{t^\ell}{\ell!} (h_{ji}^\ell + r_{ji}^\ell) + \mathcal{O}(t^{\varrho+1}),$$

$$(z_i)_* \hat{f}_j^t = \sum_{\ell=0}^{\varrho} \frac{t^\ell}{\ell!} h_{ji}^\ell.$$

Here, the  $\mathcal{O}$  is intended as  $t \downarrow 0$  and is uniform in a compact neighborhood of the origin. Then,

$$\begin{aligned} \dot{x}_i(t) - \dot{y}_i(t) &= \sum_{j=1}^m u_j(t) \left( \sum_{\ell=0}^{\varrho} \frac{t^\ell}{\ell!} (h_{ji}^\ell(x) - h_{ji}^\ell(y) + r_{ji}^\ell(x)) + \mathcal{O}(t^{\varrho+1}) \right) \\ &= \sum_{j=1}^m u_j(t) \left( \sum_{\ell=0}^{\varrho} \frac{t^\ell}{\ell!} \left( \sum_{w_k < w_i - \ell s} (x_k(t) - y_k(t)) Q_{jik}^\ell(x, y) + r_{ji}^\ell(x) \right) + \mathcal{O}(t^{\varrho+1}) \right), \end{aligned}$$

where  $Q_{jik}^\ell$  are homogeneous polynomial in  $x$  and  $y$ , of order  $w_i - w_k - \ell s - 1$ . We observe that, if  $w_i - w_k - \ell s - 1 < 0$ , then  $Q_{jik}^\ell \equiv 0$ . Thus, for sufficiently small  $\|x\|$  and  $\|y\|$ , we have

$$|Q_{jik}^\ell(x, y)| \leq C (\|x\|^{(w_i - w_k - \ell s - 1)^+} + \|y\|^{(w_i - w_k - \ell s - 1)^+}), \quad |r_{ji}^\ell(x)| \leq C \|x\|^{(w_i - \ell s)^+}.$$

Here we let  $(\xi)^+ = \max\{\xi, 0\}$ , for any  $\xi \in \mathbb{R}$ . Using the inequalities of the first step, taking  $t < T$  sufficiently small, and enlarging the constant  $C$ , we get

$$\begin{aligned} &|\dot{x}_i(t) - \dot{y}_i(t)| \\ &\leq C |u(t)| \left( \sum_{\ell=0}^{\varrho} \frac{t^\ell}{\ell!} \left( \sum_{w_k < w_i - \ell s} |x_k(t) - y_k(t)| \varepsilon^{w_i - w_k - \ell s - 1} + \varepsilon^{(w_i - \ell s)^+} \right) + t^{\varrho+1} \right) \\ &\leq C |u(t)| \left( \sum_{\ell=0}^{\varrho} t^\ell \left( \sum_{w_h < w_i} |x_h(t) - y_h(t)| \varepsilon^{w_i - w_h - 1} + \varepsilon^{(w_i - \ell s)^+} \right) + t^{\varrho+1} \right). \end{aligned}$$

In the last inequality we applied the change of variable  $w_k \mapsto w_h - \ell s$  in each of the sums.

We can integrate the system by induction, since it is in triangular form. For  $w_i = 1$ , since  $(w_i - \ell s)^+ = 0$  for any  $\ell \geq 1$ , the inequality reduces to

$$|\dot{x}_i(t) - \dot{y}_i(t)| \leq C|u(t)| \left( \sum_{\ell=0}^{\varrho} t^\ell \varepsilon^{(w_i - \ell s)^+} + t^{\varrho+1} \right) \leq C|u(t)|(\varepsilon^{w_i} + t).$$

Here we enlarged the constant  $C$ . Thus, integrating the previous inequality, we get  $|x_i(t) - y_i(t)| \leq C\varepsilon(\varepsilon^{w_i} + t) \leq C\varepsilon(\varepsilon + t^{\frac{1}{s}})^{w_i}$ .

Let, then,  $w_i > 1$  and assume that  $|x_h(t) - y_h(t)| \leq C\varepsilon(\varepsilon + t^{\frac{1}{s}})^{w_h}$  for any  $w_h < w_i$ . To complete the proof it suffices to show that  $|\dot{x}_i(t) - \dot{y}_i(t)| \leq C|u(t)|(\varepsilon + t^{\frac{1}{s}})^{w_i}$ , since (25) will follow, as above, by integration. Thus, we have, enlarging again the constant  $C$  and taking  $t$  sufficiently small,

$$\begin{aligned} & |\dot{x}_i(t) - \dot{y}_i(t)| \\ (27) \quad & \leq C|u(t)| \left( \sum_{\ell=0}^{\varrho} t^\ell \left( \sum_{w_h < w_i} \left( \varepsilon + t^{\frac{1}{s}} \right)^{w_h} \varepsilon^{w_i - w_h} + \varepsilon^{(w_i - \ell s)^+} \right) + t^{\varrho+1} \right) \\ & \leq C|u(t)| \left( \sum_{\ell=0}^{\varrho} t^\ell \left( \sum_{w_h < w_i} t^{\frac{w_h}{s}} \varepsilon^{w_i - w_h} + \varepsilon^{(w_i - \ell s)^+} \right) + t^{\varrho+1} \right). \end{aligned}$$

If  $t \leq \varepsilon^s$ , from (27) it is clear that  $|\dot{x}_i(t) - \dot{y}_i(t)| \leq C|u(t)|\varepsilon^{w_i}$ . Here we used the fact that  $\varrho + 1 \geq w_i/s$ . On the other hand, if  $\varepsilon < t^{1/s}$ , it holds

$$|\dot{x}_i(t) - \dot{y}_i(t)| \leq C|u(t)| \left( \sum_{\ell=0}^{\varrho} \left( \sum_{w_h < w_i} t^{\frac{w_h}{s} + \ell + \frac{w_i - w_h}{s}} + t^{\ell + \frac{w_i - \ell s}{s}} \right) + t^{\varrho+1} \right) \leq C|u(t)|t^{\frac{w_i}{s}}.$$

Putting together these two estimates, we get that  $|\dot{x}_i(t) - \dot{y}_i(t)| \leq C|u(t)|(\varepsilon^{w_i} + t^{\frac{w_i}{s}}) \leq C|u(t)|(\varepsilon + t^{\frac{1}{s}})^{w_i}$ , completing the proof of the proposition.  $\square$

*Proof of Theorem 3.19.* We start by claiming that (24) implies (22). In fact, if  $\gamma : q \rightsquigarrow q'$  is the trajectory associated in (17) to a control  $u \in L^1([0, T], \mathbb{R}^m)$ , and  $\hat{\gamma}$  is the trajectory associated with the same control in the homogeneous series approximation (ATD), with  $\hat{\gamma}(0) = q$ , it holds

$$|z_i(q')| \leq |z_i(\hat{\gamma}(T))| + |z_i(\hat{\gamma}(T)) - z_i(\gamma(T))|.$$

Thus, by Proposition 3.20, the claim is proved.

Hence, from now on we assume our system to be in the form (ATD). Let us define, for  $1 \leq j \leq n$  and  $0 \leq \alpha \leq r$ , the vector fields  $\varphi_j^\alpha$  as

$$\varphi_j^\alpha = \sum_{\ell=0}^{\alpha} \frac{t^\ell}{\ell!} F_j^\ell,$$

where  $F_j^\ell$  are defined in (20). We do not explicitly denote the dependence on time, to lighten the notation. Observe that, if  $\alpha = \varrho$ , then, by (21),  $\varphi_j^\alpha = \hat{f}_j^t$ .

We claim that, letting  $x^{(\alpha)}(\cdot)$  be the trajectory associated with a control  $u \in L^1([0, T], \mathbb{R}^m)$  in system (TD) with  $\{\varphi_1^\alpha, \dots, \varphi_m^\alpha\}$  as vector fields, then, for some constant  $C > 0$  and any  $i \in \{1, \dots, n\}$  and  $\alpha \geq 1$ , it holds

$$(28) \quad |x_i^{(\alpha)}(T) - x_i^{(\alpha-1)}(T)| \leq \begin{cases} 0 & \text{if } w_i \leq \alpha s, \\ C\varepsilon(\varepsilon + T^{\frac{1}{s}})^{w_i-1} & \text{if } w_i > \alpha s. \end{cases}$$



In fact, due to the homogeneity of the  $F_j^\ell$ , proceeding as in the proof of Proposition 3.20, we get that for  $w_i \leq \alpha s$  it holds

$$|\dot{x}_i^{(\alpha)}(t) - \dot{x}_i^{(\alpha-1)}(t)| \leq C|u(t)| \sum_{\ell=0}^{\alpha-1} t^\ell \sum_{w_h < w_i} |x_h^{(\alpha)}(t) - x_h^{(\alpha-1)}(t)| \varepsilon^{w_i - w_h - 1}.$$

By induction on  $1 \leq w_i \leq \alpha s$ , this proves the first part of the claim. On the other hand, if  $w_i > \alpha s$ , it holds

$$|\dot{x}_i^{(\alpha)}(t) - \dot{x}_i^{(\alpha-1)}(t)| \leq C|u(t)| \left( \sum_{\ell=0}^{\alpha-1} t^\ell \sum_{w_h < w_i} |x_h^{(\alpha)}(t) - x_h^{(\alpha-1)}(t)| \varepsilon^{w_i - w_h - 1} + t^\alpha \varepsilon^{w_i - \alpha s - 1} \right).$$

Then, again by induction over  $w_i$ , we get that  $|x_i^{(\alpha)}(T) - x_i^{(\alpha-1)}(T)| \leq CT^\alpha \varepsilon^{w_i - \alpha s}$ . Finally, the claim follows considering the cases  $T \leq \varepsilon^s$  and  $T > \varepsilon^s$ .

Due to the fact that  $\varphi_j^0 = \widehat{f}_j$ , by Theorem 2.7 it holds  $|x_i^{(0)}(T)| \leq C\varepsilon^{w_i}$ . Thus, applying (28) and enlarging the constant  $C$ , we get

$$|z_i(q')| = |x_i^{(r)}(T)| \leq \sum_{\ell=1}^r |x_i^{(\ell)}(T) - x_i^{(\ell-1)}(T)| + |x_i^{(0)}(T)| \leq \begin{cases} C\varepsilon^{w_i} & \text{if } w_i \leq s, \\ C\varepsilon(\varepsilon + T^{\frac{1}{s}})^{w_i-1} & \text{if } w_i > s. \end{cases}$$

This proves (23) and (24), completing the proof of the theorem.  $\square$

We end this section by showing that the estimate (23) is sharp, at least in some directions. Indeed, for a system which is series homogeneous at  $q$  in some privileged coordinates  $z$ , and satisfies the hypotheses of Theorem 3.19, it holds that  $z_*((\text{ad}^k f_0)f_j)$  is an homogeneous vector field of weighted degree  $-sk - 1$ . Thus, since  $\varepsilon t^k \leq \varepsilon(\varepsilon + t^{\frac{1}{s}})^{sk}$ , the following proposition shows that (23) is sharp in this direction. The proof is an adaption of an argument from [11].

**Proposition 3.22.** *Let  $\{f_1, \dots, f_m\}$  satisfy the Hörmander condition. Let, moreover  $q \in M$ ,  $i \in \{1, \dots, m\}$  and  $k \geq 0$ . Then, for any coordinate system  $y$  at  $q$ , there exist  $T, \varepsilon_0 > 0$  such that, for any  $\varepsilon < \varepsilon_0$  and  $t < T$  there exists a (TD)-admissible curve  $\gamma : [0, t] \rightarrow M$ , with  $c(\gamma) \leq \varepsilon$ , and such that*

$$y(\gamma(t)) = \varepsilon t^k dy((\text{ad}^k f_0)f_j(q)) + \mathcal{O}(\varepsilon t^{k+1}) \quad \text{as } \varepsilon t \rightarrow 0.$$

*Proof.* Let  $t, \eta > 0$  be fixed, and define  $u \in L^1([0, T], \mathbb{R}^m)$  as  $u_i(\tau) \equiv \eta$ ,  $u_j(\tau) \equiv 0$  for  $j \neq i$ ,  $\tau \in [0, t]$ . Then, fix any  $\Phi \in C^k([0, 1])$  such that  $\Phi^{(i)}(0) = \Phi^{(i)}(1) = 0$ , for  $0 \leq i < k$ . Thus, by integrating by parts and the fact that  $\frac{d}{dt}(e^{-\tau f_0})_* g = (e^{-\tau f_0})_*(\text{ad}(f_0)g)$ , we get

$$\int_0^t \Phi^{(k)}(\tau/t)(e^{-\tau f_0})_* f_i(q) d\tau = t^k \int_0^t \Phi(\tau/t)(e^{-\tau f_0})_* \left( (\text{ad}^k f_0)f_i \right) (q) d\tau,$$

for any  $t$  and  $q$ . This implies that the flows generated by  $\Phi^{(k)}(\tau/t)(e^{-\tau f_0})_* f_i$  and  $t^k \Phi(\tau/t)(e^{-\tau f_0})_* \left( (\text{ad}^k f_0)f_i \right)$  coincide. Using the series expansions of the chronological exponential and  $(e^{-\tau f_0})_*$ , see [5, Section

2.4], there holds, then,

$$\begin{aligned}
\overrightarrow{\text{exp}} \int_0^t \sum_{j=1}^m \Phi^{(k)}(\tau/t) u_j(\tau) (e^{-\tau f_0})_* f_j d\tau &= \overrightarrow{\text{exp}} \int_0^t \eta \Phi^{(k)}(\tau/t) (e^{-\tau f_0})_* f_i d\tau \\
&= \overrightarrow{\text{exp}} \int_0^t \eta t^k \Phi(\tau/t) (e^{-\tau f_0})_* \left( (\text{ad}^k f_0) f_i \right) d\tau \\
&= \overrightarrow{\text{exp}} \int_0^1 \eta t^{k+1} \Phi(s) (e^{-ts f_0})_* \left( (\text{ad}^k f_0) f_i \right) ds \\
&= \overrightarrow{\text{exp}} \int_0^1 \eta t^{k+1} \Phi(s) \left( (\text{ad}^k f_0) f_i + \mathcal{O}(t) \right) ds \\
&= \text{Id} + \eta t^{k+1} (\text{ad}^k f_0) f_i + \mathcal{O}(\eta t^{k+2})
\end{aligned}$$

Finally, considering any coordinate system and letting  $\varepsilon = \eta t$ , this completes the proof.  $\square$

#### 4. CONTROL-AFFINE SYSTEMS

In this section we apply the results of Section 3 to control-affine systems. Let  $\{f_1, \dots, f_m\}$  be a bracket-generating family of vector fields,  $f_0$  be a smooth vector field, called the *drift*, and consider the control-affine system

$$(D) \quad \dot{q} = f_0(q) + \sum_{i=1}^m u_i f_i(q), \quad q \in M, \quad u = (u_1, \dots, u_m) \in \mathbb{R}^m.$$

The assumption on  $\{f_1, \dots, f_m\}$  to be bracket-generating, is called *strong Hörmander condition* for (D).

An absolutely continuous curve  $\gamma : [0, T] \rightarrow M$  is (D)-admissible if  $\dot{\gamma}(t) = f_0(\gamma(t)) + f_{u(t)}(\gamma(t))$  for some control  $u \in L^1([0, T], \mathbb{R}^m)$ . Its cost is defined as

$$c_{f_0}(\gamma) = \min \|u\|_{L^1([0, T], \mathbb{R}^m)},$$

where the minimum is taken over all controls  $u$  such that  $\gamma$  is associated with  $u$ . Then, we define the two value functions we are interested in as

$$\begin{aligned}
\rho_T^{f_0}(q, q') &= \inf \{c(\gamma) : \gamma : [0, T'] \rightarrow M \text{ is (D)-admissible, } \gamma : q \rightsquigarrow q', T' \leq T\}, \\
\rho^{f_0}(q, q') &= \inf \{c(\gamma) : \gamma \text{ (D)-admissible and } \gamma : q \rightsquigarrow q'\}.
\end{aligned}$$

It is clear that  $\rho_T^{f_0}(q, q') \searrow \rho^{f_0}(q, q')$  as  $T \rightarrow +\infty$ , for any  $q, q' \in M$ . Moreover, we observe that,  $\rho_T^{f_0}(q, e^{t f_0} q) = 0$  for any  $0 \leq t \leq T$ . Finally, the reachable sets with respect to these value functions, from any  $q \in M$  and for  $\varepsilon, T > 0$ , are

$$\mathcal{R}_T^{f_0}(q, \varepsilon) = \{q' \in M : \rho_T^{f_0}(q, q') < \varepsilon\}, \quad \mathcal{R}^{f_0}(q, \varepsilon) = \{q' \in M : \rho^{f_0}(q, q') < \varepsilon\}.$$

**4.1. Connection with time-dependent systems.** Applying the variations formula (see [5]), system (D) can be written as a composition of a time-dependent system in the form (17) and of a translation along the drift. Namely, for any  $u \in L^1([0, T], \mathbb{R}^m)$ , it holds

$$(29) \quad \overrightarrow{\text{exp}} \int_0^T \left( f_0 + \sum_{i=1}^m u_i(t) f_i \right) dt = e^{T f_0} \circ \overrightarrow{\text{exp}} \int_0^T \sum_{i=1}^m u_i(t) (e^{-t f_0})_* f_i dt.$$

We call *time-dependent system associated with* (D) the following control system,

$$(30) \quad \dot{q} = \sum_{i=1}^m u_i (e^{-t f_0})_* f_i(q), \quad q \in M, \quad u = (u_1, \dots, u_m) \in \mathbb{R}^m.$$

Observe that, since diffeomorphisms preserve linear independence, the strong Hörmander condition for (D), implies that  $\{(e^{-tf_0})_*f_1, \dots, (e^{-tf_0})_*f_m\}_{t \in [0, +\infty)}$  satisfies the strong Hörmander condition, defined in Definition 3.11.

Exploiting these facts, we can prove the following.

**Proposition 4.23.** *Assume that (D) satisfies the strong Hörmander condition. Then, for any  $T > 0$ , the functions  $\rho_T^{f_0}, \rho^{f_0} : M \times M \rightarrow [0, +\infty)$  are continuous. Moreover, letting  $d_{\text{SR}}$  be the sub-Riemannian distance induced by  $\{f_1, \dots, f_m\}$ , for any  $q, q' \in M$  it holds*

$$\rho_T^{f_0}(q, q') \leq \min_{0 \leq t \leq T} d_{\text{SR}}(e^{tf_0}q, q'), \quad \rho^{f_0}(q, q') \leq \min_{t \geq 0} d_{\text{SR}}(e^{tf_0}q, q').$$

*Proof.* The continuity of the two functions, and the fact that  $\rho_T^{f_0}(q, q'), \rho^{f_0}(q, q') \leq d_{\text{SR}}(q, q')$ , for any  $q, q' \in M$ , follows from the same arguments used in Theorem 3.12, adapting Lemmata 3.13 and 3.14 to the system (D). In particular, one has to consider  $(\mathcal{T}, \mathcal{F}, \xi) \mapsto e^{\mathcal{T}f_0} \circ E_{\mathcal{T}, \mathcal{F}}(\xi)$  instead of  $(\mathcal{T}, \mathcal{F}, \xi) \mapsto E_{\mathcal{T}, \mathcal{F}}(\xi)$ .

To prove the second part of the statement, we let, for any  $t \in [0, T)$ ,

$$\varphi_t(p) = \inf\{c(\gamma) : \gamma : [t, T'] \rightarrow M \text{ is (D)-admissible, } \gamma : p \rightsquigarrow q', T' \leq T\}.$$

It is clear that, as above, it holds  $\varphi_t(p) \leq d_{\text{SR}}(p, q')$ . Moreover, we observe that  $\rho_T^{f_0}(q, e^{tf_0}q) = 0$  for any  $0 \leq t < T$ , and hence that for any such  $t$  it holds

$$\rho_T^{f_0}(q, q') \leq \varphi_t(e^{tf_0}q) \leq d_{\text{SR}}(e^{tf_0}q, q').$$

Taking the minimum for  $0 \leq t < T$ , proves the inequality regarding  $\rho_T^{f_0}$ . To complete the proof it suffices to observe that  $\rho^{f_0}(q, q') \leq \rho_T^{f_0}(q, q')$  for any  $T > 0$ .  $\square$

We point out that in system (D), as in time-dependent systems, the existence of minimizers is not assured. Moreover, this lack of minimizers is possible even if they exist for the associated time-dependent system, as the following example points out.

**Example 4.24.** Consider the following vector fields on  $\mathbb{R}^3$ , with coordinates  $(x, y, z)$ ,

$$f_1(x, y, z) = \partial_x, \quad f_2(x, y, z) = \partial_y + x\partial_z.$$

Since  $[f_1, f_2] = \partial_z$ ,  $\{f_1, f_2\}$  is a bracket-generating family of vector fields. The sub-Riemannian control system associated to  $\{f_1, f_2\}$  on  $\mathbb{R}^3$  corresponds to the Heisenberg group.

Let now  $f_0 = \partial_z$  be the drift. Since  $[f_1, \partial_z] = [f_2, \partial_z] = 0$  it holds that  $(e^{-tf_0})_*f_1 = f_1$  and  $(e^{-tf_0})_*f_2 = f_2$ . Hence, the associated time-dependent system is actually not time-dependent. Thus, by (29), a curve  $\gamma : [0, T] \rightarrow \mathbb{R}^3$  is (SR)-admissible for  $\{f_1, f_2\}$  if and only if  $\tilde{\gamma}(\cdot) = e^{\cdot f_0} \circ \gamma(\cdot)$  is (D)-admissible. As a consequence of this, for any  $q \in \mathbb{R}^3$  and any  $\varepsilon > 0$ ,

$$\mathcal{R}^{f_0}(q, \varepsilon) = \bigcup_{t \geq 0} e^{tf_0} \circ B_{\text{SR}}(q, \varepsilon).$$

As pointed out in Section 2, minimizers for the sub-Riemannian system exist between any pair of points in  $B_{\text{SR}}(q, \varepsilon)$ , if  $\varepsilon$  is sufficiently small. Let us show that, for any point in  $\mathcal{R}^{f_0}(q, \varepsilon)$  with positive  $z$  coordinate, we have an explicit minimizer, while for the others there exists no minimizer. Without loss of generality we can consider  $q = 0$ . Then, since  $e^{t'f_0}(x', y', z') = (x', y', z' + t')$ , every point  $(x, y, z) \in \mathcal{R}^{f_0}(0, \varepsilon)$  with  $z > 0$ , can be reached optimally considering the sub-Riemannian minimizing curve between the origin and  $(x, y, 0)$  rescaled on time  $z$ .

If, instead,  $z \leq 0$ , we cannot construct any sub-Riemannian trajectory from 0 to  $(x, y, z - t)$ ,  $t > 0$ , with cost  $\leq d_{\text{SR}}(0, (x, y, z))$ . This is due to the fact that the minimizing trajectories in Heisenberg group are the lifts of arcs on the plane  $(x, y)$ , spanning area equal to the  $z$  coordinates, and that  $|z - t| = -z + t > |z|$ . Since, by Proposition 4.23,  $\rho^{f_0}(0, (x, y, z)) \leq d_{\text{SR}}(0, (x, y, z))$ , this implies that there exists no minimizer for  $\rho^{f_0}(q, (x, y, z))$ .

**4.2. Estimates on reachable sets.** In this section we apply Theorem 3.19, in order to obtain results in the spirit of Theorem 2.7 and Corollary 2.8. First, we need the following definition.

**Definition 4.25.** The point  $q$  is said to be *regular with respect to the drift  $f_0$* , if  $q' \mapsto \text{ord}_{q'}(f_0)$  is locally constant at  $q$ .

The main result of this section are the following local regularity estimates for  $\rho^{f_0}$ . We cannot expect anything global, since in general the sets  $\mathcal{R}^{f_0}(q, \varepsilon)$  are noncompact.

**Theorem 4.26.** *Assume that (D) satisfies the strong Hörmander condition, and let  $q$  be regular with respect to the drift  $f_0$ . Assume, moreover, that  $z = (z_1, \dots, z_n)$  is a system of privileged coordinates at  $q$  for  $\{f_1, \dots, f_m\}$ , such that  $z_* f_0 = \partial_{z_k}$ , for some  $1 \leq k \leq n$ . Then, there exist  $T_0, \varepsilon_0, C > 0$  such that*

$$\frac{1}{C} \text{dist} \left( z(q'), z(e^{[0, T] f_0} q) \right) \leq \rho_T^{f_0}(q, q') \leq C \text{dist} \left( z(q'), z(e^{[0, T] f_0} q) \right)^{1/r}, \quad q' \in \mathcal{R}_T^{f_0}(q, \varepsilon),$$

where, for any  $x \in \mathbb{R}^n$  and  $A \subset \mathbb{R}^n$ ,  $\text{dist}(x, A) = \inf_{y \in A} |x - y|$  is the Euclidean distance between them and  $r$  is the degree of non-holonomy of  $\{f_1, \dots, f_m\}$  at  $q$ .

Let us define the following sets, for parameters  $\eta > 0$  and  $T > 0$ . We remark that  $\text{Box}(\eta)$  is defined as in (10) and that  $\{\partial_{z_i}\}_{i=1}^n$  is the canonical basis in  $\mathbb{R}^n$ .

$$\begin{aligned} \Xi_T(\eta) &= \bigcup_{0 \leq \xi \leq T} \left( \xi \partial_{z_k} + \text{Box}(\eta) \right), \\ \Pi_T(\eta) &= \text{Box}(\eta) \cup \bigcup_{0 < \xi \leq T} \{z \in \mathbb{R}^n : z_k = \xi, |z_i| \leq \eta^{w_i} + \eta \xi^{\frac{w_i}{s}} \text{ for } w_i \leq s, i \neq k, \\ &\quad \text{and } |z_i| \leq \eta(\eta + \xi^{\frac{1}{s}})^{w_i-1} \text{ for } w_i > s\}, \\ \widehat{\Pi}_T(\eta) &= \text{Box}(\eta) \cup \bigcup_{0 < \xi \leq T} \{z \in \mathbb{R}^n : z_k = \xi, |z_i| \leq \eta^{w_i} \text{ for } w_i \leq s, i \neq k, \\ &\quad \text{and } |z_i| \leq \eta(\eta + \xi^{\frac{1}{s}})^{w_i-1} \text{ for } w_i > s\}. \end{aligned}$$

As in the sub-Riemannian case, Theorem 4.26 is a direct consequence of some estimates on the shape of the accessible sets, contained in the following.

**Theorem 4.27.** *Assume that (D) satisfies the strong Hörmander condition, and let  $q \in M$  be regular with respect to the drift  $f_0$ . Assume, moreover, that  $z = (z_1, \dots, z_n)$  is a system of privileged coordinates at  $q$  for  $\{f_1, \dots, f_m\}$ , such that  $z_* f_0 = \partial_{z_k}$ , for some  $1 \leq k \leq n$ . Then, there exist  $C, \varepsilon_0, T_0 > 0$  such that*

$$(31) \quad \Xi_T \left( \frac{1}{C} \varepsilon \right) \subset \mathcal{R}_T^{f_0}(q, \varepsilon) \subset \Pi_T(C\varepsilon), \quad \text{for } \varepsilon < \varepsilon_0 \text{ and } T < T_0.$$

Here, with abuse of notation, we denoted by  $\mathcal{R}_T^{f_0}(q, \varepsilon)$  the coordinate representation of the reachable set. In particular,

$$\text{Box} \left( \frac{1}{C} \varepsilon \right) \cap \{z_k \leq 0\} \subset \mathcal{R}_T^{f_0}(q, \varepsilon) \cap \{z_k \leq 0\} \subset \text{Box}(C\varepsilon) \cap \{z_k \leq 0\}.$$

Moreover, if the system is nilpotent, it holds

$$(32) \quad \Xi_T \left( \frac{1}{C} \varepsilon \right) \subset \mathcal{R}_T^{f_0}(q, \varepsilon) \subset \widehat{\Pi}_T(C\varepsilon), \quad \text{for } \varepsilon < \varepsilon_0 \text{ and } T < T_0.$$

In order to prove Theorem 4.27, we need the following lemma.

**Lemma 4.28.** *Let  $z = (z_1, \dots, z_n)$  be a system of privileged coordinates at  $q \in M$ . Then there exist  $C, \varepsilon_0, T_0 > 0$  such that, for any  $q' \in \mathcal{R}_T^{f_0}(q, \varepsilon_0)$  for  $\varepsilon < \varepsilon_0$  and  $T < T_0$ , and such that*

- (i) *for any  $t < \varepsilon_0$ ,  $\text{ord}_{q'(t)} f_0 = -s$ , where  $q'(t) = e^{-tf_0}(q')$ ,*
- (ii)  *$dz_k(f_0(z(q'))) \neq 0$ , for some  $k$  with  $w_k = -s$ ,*

*it holds that, if  $u \in L^1([0, T], \mathbb{R}^m)$  is a control steering the system (D) from  $q$  to  $q'$ , with  $\|u\|_1 = \varepsilon$ , then*

$$T \leq C(\varepsilon^s + \max\{z_k(q'), 0\}).$$

*Proof.* For any  $\eta > 0$ , let  $\gamma$  be the trajectory associated with  $u \in L^1([0, T], \mathbb{R}^m)$  in the system (D), and satisfying  $\gamma : q \rightsquigarrow q'$ . Let  $\tilde{\gamma}$  be the trajectory associated with  $u$  and starting from  $q$ , in the time-dependent system (30). Thus  $\gamma(t) = e^{tf_0} \circ \tilde{\gamma}(t)$  and  $\rho(q, \tilde{\gamma}(T)) \leq \varepsilon$ .

Recall that, for any vector field  $g$  and point  $p \in M$ , it holds that  $z_k(e^{Tg}(p)) - z_k(p) = \int_0^T dz_k(g(e^{tg}(p)))$ . Thus, by the mean value theorem, there exists  $\tau \in [0, T]$  such that

$$(33) \quad z_k(q') = z_k(\gamma(T)) = T dz_k(f_0(e^{\tau f_0}(\tilde{\gamma}(T)))) + z_k(\tilde{\gamma}(T)).$$

Since  $e^{\tau f_0}(\tilde{\gamma}(T)) = e^{-(T-\tau)f_0}(q')$ , by hypothesis (ii) and the smoothness of  $f_0$ , there exist  $T_0, C_1 > 0$ , independent of  $\gamma$ , such that  $dz_k(f_0(e^{\tau f_0}(\tilde{\gamma}(T)))) \geq C_1$  for  $T < T_0$ . Hence, by Theorem 3.19 (since  $w_k = s$ ), there exist  $C_2, \bar{\varepsilon} > 0$  such that, if  $\varepsilon < \bar{\varepsilon}$  and  $T < T_0$ ,

$$T \leq \frac{|z_k(\tilde{\gamma}(T))| + \max\{z_k(q'), 0\}}{C_1} \leq \frac{C_2(\varepsilon^s + T\varepsilon) + \max\{z_k(q'), 0\}}{C_1}.$$

Since the constants are independent of  $\gamma$ , taking  $C = C_2/C_1$ ,  $\varepsilon_0 \leq \min\{T_0, \bar{\varepsilon}, (C-1)/C^2\}$  completes the proof.  $\square$

*Proof of Theorem 4.27.* The first inclusion in (31) follows from Theorem 2.7 and Theorem 4.23. In fact, combining them, we have that, for any  $\varepsilon < \varepsilon_0$  and any  $T > 0$ ,

$$\Xi_T \left( \frac{1}{C} \varepsilon \right) \subset \bigcup_{0 \leq t \leq T} B_{\text{SR}}(e^{tf_0} q, \varepsilon) \subset \mathcal{R}_T^{f_0}(q, \varepsilon).$$

To prove the second inclusion, we let  $q' \in \mathcal{R}_T^{f_0}(q, \varepsilon)$ . Fix any  $\eta > 0$  and consider a control  $u \in L^1([0, T], \mathbb{R}^m)$  such that its associated trajectory  $\gamma$ , in the system (D), satisfies  $\gamma : q \rightsquigarrow q'$  and  $c_{f_0}(\gamma) \leq \varepsilon + \eta$ . We distinguish two cases. First we assume that  $z_k(q') \leq 0$ . In this case, by Lemma 4.28 it follows there exists  $C, \varepsilon_0, T_0 > 0$  such that if  $T < T_0$  and  $\varepsilon < \varepsilon_0$ , then  $T \leq C\varepsilon^s$ . Moreover (29) implies that  $e^{-Tf_0}(q') \in \mathcal{R}_T(q, \varepsilon)$ . Then, enlarging the constant  $C$ , Theorem 3.19 yields

$$\begin{aligned} |z_i(q')| &= |z_i(e^{-Tf_0}(q'))| \leq C(\varepsilon^{w_i} + \varepsilon T^{\frac{w_i}{s}}) \leq C\varepsilon^{w_i}, \quad \text{if } w_i \leq s \text{ and } i \neq k, \\ |z_k(q')| &\leq T + |z_k(e^{-Tf_0}(q'))| \leq T + C(\varepsilon + T^{\frac{1}{s}})^s \leq C\varepsilon^s, \\ |z_i(q')| &= |z_i(e^{-Tf_0}(q'))| \leq C\varepsilon(\varepsilon + T^{\frac{1}{s}})^{w_i-1} \leq C\varepsilon^{w_i}, \quad \text{if } w_i > s. \end{aligned}$$

Here, we used the fact that, for any  $p \in M$ , from  $z_* f_0 = \partial_{z_k}$ , it holds  $z_i(p) = z_i(e^{-Tf_0}(p))$  and  $|dz_k(f_0(p))| \equiv 1$ . Thus,  $q' \subset \text{Box}(C\varepsilon) \subset \Pi(C\varepsilon)$ .

On the other hand, if  $z_k(q') > 0$ , Lemma 4.28 yields that  $T \leq C(\varepsilon^s + z_k(q'))$ . Then, applying again Theorem 3.19, we get

$$\begin{aligned} |z_i(q')| &\leq C(\varepsilon^{w_i} + \varepsilon z_k(q')^{\frac{w_i}{s}}), \quad \text{if } w_i \leq s \text{ and } i \neq k, \\ |z_k(q')| &\leq T + C\varepsilon^s, \\ |z_i(q')| &\leq C\varepsilon(\varepsilon + z_k(q')^{\frac{1}{s}})^{w_i-1}, \quad \text{if } w_i > s. \end{aligned}$$

This proves that  $q' \subset \Pi(C\varepsilon)$ , completing the proof of (31).

To prove (32) it suffices to use the same argument as above, applying the result on nilpotent systems in Theorem 3.19.  $\square$

**Remark 4.29.** Theorem 4.27 suggests that the behavior of the system (D), when moving in the direction  $-f_0$ , is essentially sub-Riemannian. However, although this is true locally in time, it is false in general. For example, consider the Euclidean plane endowed with a rotational drift, i.e., such that  $\{e^{tf_0}(q)\}_{t \in (0, +\infty)}$  is diffeomorphic to  $S^1$  for any  $q \neq 0$ . Then,  $\rho^{f_0}(q, e^{-tf_0}(q)) = 0$  for any  $t > 0$  and thus we can move in the direction  $-f_0$  for free.

*Proof of Theorem 4.26.* Since every norm on  $\mathbb{R}^n$  is equivalent,  $\text{dist}(z(q'), [0, T]\partial_{z_k})$  is equivalent to

$$a(q') = \sum_{\substack{1 \leq i \leq n \\ i \neq k}} |z_i(q')| + \min_{t \in [0, T]} |z_k(q') - t|.$$

Thus, to complete the proof it suffices to prove that it holds  $C^{-1}a(q') \leq \rho_T^{f_0}(q, q') \leq Ca(q')^{1/r}$ .

By Theorem 3.19,  $\Xi_T(C^{-1}\varepsilon) \subset \mathcal{R}_T^{f_0}(q, \varepsilon) \subset \Pi_T(C\varepsilon)$  for any  $\varepsilon < \varepsilon_0$ . The first inclusion is equivalent to the fact that, for every  $\varepsilon < \varepsilon_0$  such that  $Ca(q') \leq \varepsilon^r$ , one has  $\rho_T^{f_0}(q, q') \leq \varepsilon$ . From this follows that  $\rho_T^{f_0}(q, q') \leq C^{1/r}a(q')^{1/r}$ . The same reasoning applied to the other inclusion proves that

$$\begin{aligned} |z_i(q')| &\leq C(\rho_T^{f_0}(q, q')^{w_i} + \rho_T^{f_0}(q, q')T^{\frac{w_i}{s}}) \text{ if } w_i \leq s, i \neq k, \\ \min_{t \in [0, T]} |z_k(q') - t| &\leq C\rho_T^{f_0}(q, q')^s, \\ |z_i(q')| &\leq C(\rho_T^{f_0}(q, q')^{w_i} + \rho_T^{f_0}(q, q')T^{\frac{w_i-1}{s}}) \text{ if } w_i > s. \end{aligned}$$

Clearly, this implies that  $a(q') \leq C\rho_T^{f_0}(q, q')$ , for some larger constant, completing the proof of the theorem.  $\square$

## REFERENCES

- [1] A. Agrachev, D. Barilari, and U. Boscain. Introduction to Riemannian and sub-Riemannian geometry (Lecture Notes). [http://people.sissa.it/agrachev/agrachev\\_files/notes.htm](http://people.sissa.it/agrachev/agrachev_files/notes.htm), 2012.
- [2] A. Agrachev, U. Boscain, J.-P. Gauthier, and F. Rossi. The intrinsic hypoelliptic Laplacian and its heat kernel on unimodular Lie groups. *J. Funct. Anal.*, 256(8):2621–2655, 2009.
- [3] A. Agrachev and P. W. Y. Lee. Continuity of optimal control costs and its application to weak KAM theory. *Calc. Var. Partial Differential Equations*, 39(1-2):213–232, 2010.
- [4] A. A. Agrachev, U. Boscain, G. Charlot, R. Ghezzi, and M. Sigalotti. Two-dimensional almost-Riemannian structures with tangency points. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 27(3):793–807, 2010.
- [5] A. A. Agrachev and Y. L. Sachkov. *Control theory from the geometric viewpoint*, volume 87 of *Encyclopaedia of Mathematical Sciences*. Springer-Verlag, Berlin, 2004. Control Theory and Optimization, II.
- [6] Andrei Agrachev, Ugo Boscain, and Mario Sigalotti. A Gauss-Bonnet-like formula on two-dimensional almost-Riemannian manifolds. *Discrete Contin. Dyn. Syst.*, 20(4):801–822, 2008.
- [7] M. Barbero-Liñán and M. Sigalotti. High-order sufficient conditions for configuration tracking of affine connection control systems. *Systems Control Lett.*, 59(8):491–503, 2010.
- [8] A. Bellaïche. The tangent space in sub-Riemannian geometry. In *Sub-Riemannian geometry*, volume 144 of *Progr. Math.*, pages 1–78. Birkhäuser, Basel, 1996.
- [9] U. Boscain and P. Mason. Time minimal trajectories for a spin 1/2 particle in a magnetic field. *J. Math. Phys.*, 47(6):062101, 29, 2006.
- [10] F. Bullo and A. D. Lewis. *Geometric control of mechanical systems*, volume 49 of *Texts in Applied Mathematics*. Springer-Verlag, New York, 2005. Modeling, analysis, and design for simple mechanical control systems.
- [11] J.-M. Coron. *Control and nonlinearity*, volume 136 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2007.
- [12] D. D'Alessandro. *Introduction to quantum control and dynamics*. Chapman & Hall/CRC Applied Mathematics and Nonlinear Science Series. Chapman & Hall/CRC, Boca Raton, FL, 2008.

- [13] L. D. Drager, J. M. Lee, E. Park, and K. Richardson. Smooth distributions are finitely generated. *Ann. Global Anal. Geom.*, 41(3):357–369, 2012.
- [14] G. B. Folland and E. M. Stein. Estimates for the  $\bar{\partial}_b$  complex and analysis on the Heisenberg group. *Comm. Pure Appl. Math.*, 27:429–522, 1974.
- [15] J.-P. Gauthier and V. Zakalyukin. On the codimension one motion planning problem. *J. Dyn. Control Syst.*, 11(1):73–89, 2005.
- [16] J.-P. Gauthier and V. Zakalyukin. On the motion planning problem, complexity, entropy, and nonholonomic interpolation. *J. Dyn. Control Syst.*, 12(3):371–404, 2006.
- [17] H. Hermes. Nilpotent and high-order approximations of vector field systems. *SIAM Rev.*, 33(2):238–264, 1991.
- [18] F. Jean. Complexity of nonholonomic motion planning. *Internat. J. Control*, 74(8):776–782, 2001.
- [19] F. Jean. Entropy and complexity of a path in sub-Riemannian geometry. *ESAIM Control Optim. Calc. Var.*, 9:485–508 (electronic), 2003.
- [20] D. Jerison and A. Sánchez-Calle. Subelliptic, second order differential operators. In *Complex analysis, III (College Park, Md., 1985–86)*, volume 1277 of *Lecture Notes in Math.*, pages 46–77. Springer, Berlin, 1987.
- [21] J. Mitchell. On Carnot-Carathéodory metrics. *J. Differential Geom.*, 21(1):35–45, 1985.
- [22] C. Romero-Meléndez, J. P. Gauthier, and F. Monroy-Pérez. On complexity and motion planning for co-rank one sub-Riemannian metrics. *ESAIM Control Optim. Calc. Var.*, 10(4):634–655, 2004.
- [23] L. P. Rothschild and E. M. Stein. Hypoelliptic differential operators and nilpotent groups. *Acta Math.*, 137(3-4):247–320, 1976.
- [24] H. J. Sussmann. Some properties of vector field systems that are not altered by small perturbations. *J. Differential Equations*, 20(2):292–315, 1976.