

Locally identifying coloring in bounded expansion classes of graphs

Daniel Gonçalves, Aline Parreau, Alexandre Pinlou

► To cite this version:

Daniel Gonçalves, Aline Parreau, Alexandre Pinlou. Locally identifying coloring in bounded expansion classes of graphs. Discrete Applied Mathematics, Elsevier, 2013, 161 (18), pp.2946-2951. 10.1016/j.dam.2013.07.003 . hal-00768472v2

HAL Id: hal-00768472

<https://hal.archives-ouvertes.fr/hal-00768472v2>

Submitted on 10 Jul 2013

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Locally identifying coloring in bounded expansion classes of graphs *

Daniel Gonçalves[†], Aline Parreau[‡], Alexandre Pinlou^{†§}

July 10, 2013

Abstract

A proper vertex coloring of a graph is said to be *locally identifying* if the sets of colors in the closed neighborhood of any two adjacent non-twin vertices are distinct. The lid-chromatic number of a graph is the minimum number of colors used by a locally identifying vertex-coloring. In this paper, we prove that for any graph class of bounded expansion, the lid-chromatic number is bounded. Classes of bounded expansion include minor closed classes of graphs. For these latter classes, we give an alternative proof to show that the lid-chromatic number is bounded. This leads to an explicit upper bound for the lid-chromatic number of planar graphs. This answers in a positive way a question of Esperet *et al.* [L. Esperet, S. Gravier, M. Montassier, P. Ochem and A. Parreau. Locally identifying coloring of graphs. *Electronic Journal of Combinatorics*, **19(2)**, 2012.].

1 Introduction

A vertex-coloring is said to be *locally identifying* if (i) the vertex-coloring is proper (i.e. no adjacent vertices receive the same color), and (ii) for any adjacent vertices u, v , the set of colors assigned to the closed neighborhood of u differs from the set of colors assigned to the closed neighborhood of v whenever these neighborhoods are distinct. The *locally identifying chromatic number* of the graph G (or lid-chromatic number, for short), denoted by $\chi_{lid}(G)$, is the smallest number of colors required in any locally identifying coloring of G .

Locally identifying colorings of graphs have been recently introduced by Esperet et al. [6] and later studied by Foucaud et al. [7]. They are related to identifying codes [8, 9], distinguishing colorings [1, 3, 5] and locating-colorings [4]. For

*This work was partially supported by the ANR grant EGOS 12 JS02 002 01.

[†]LIRMM - Univ. Montpellier 2, CNRS - 161 rue Ada, 34095 Montpellier Cedex 5, France

[‡]LIFL - Univ. Lille 1, INRIA - Parc scientifique de la haute borne, 59650 Villeneuve d'Ascq, France

[§]Second affiliation: Département de Mathématiques et Informatique Appliquées, Université Paul-Valéry, Montpellier 3, Route de Mende, 34199 Montpellier Cedex 5, France.

example, upper bounds on lid-chromatic number have been obtained for bipartite graphs, k -trees, outerplanar graphs and bounded degree graphs. An open question asked by Esperet et al. [6] was to know whether χ_{lid} is bounded for the class of planar graphs. In this paper, we answer positively to this question proving more generally that χ_{lid} is bounded for any class of bounded expansion.

In Section 3, we first give a tight bound of χ_{lid} in term of the tree-depth. Then we use the fact that any class of bounded expansion admits a low tree-depth coloring (that is a k -coloring such that each triplet of colors induces a graph of tree-depth 3, for some constant k) to prove that it has bounded lid-chromatic number.

In Section 4, we focus on minor closed classes of graphs which have bounded expansion and give an alternative bound on the lid-chromatic number, which gives an explicit bound for planar graphs.

The next section is devoted to introduce notation and preliminary results.

2 Notation and preliminary results

Let $G = (V, E)$ be a graph. For any vertex u , we denote by $N_G(u)$ its *neighborhood* in G and by $N_G[u]$ its *closed neighborhood* in G (u together with its adjacent vertices). The notion of neighborhood can be extended to sets as follows: for $X \subseteq V$, $N_G[X] = \{w \in V(G) \mid \exists v \in X, w \in N[v]\}$ and $N_G(X) = N_G[X] \setminus X$. When the considered graph is clearly identified, the subscript is dropped.

The *degree* of vertex u is the size of its neighborhood. The *distance* between two vertices u and v is the number of edges in a shortest path between u and v . For $X \subseteq V$, we denote by $G[X]$ the subgraph of G *induced by* X .

We say that two vertices u and v are *twins* if $N[u] = N[v]$ (although they are often called *true twins* in the literature, we call them *twins* for convenience). In particular, u and v are adjacent vertices. Note that if u and v are adjacent but not twins, there exists a vertex w which is adjacent to exactly one vertex among $\{u, v\}$, i.e. $w \in N[u] \Delta N[v]$ (where Δ is the symmetric difference between sets). We say that w *distinguishes* u and v , or simply w *distinguishes* the edge uv . For a subset $X \subseteq V$, we say that a subset $Y \subseteq V$ *distinguishes* X if for every pair u, v of non-twin vertices of X , there exists a vertex $w \in Y$ that distinguishes the edge uv .

Let $c : V \rightarrow \mathbb{N}$ be a vertex-coloring of G . The coloring c is *proper* if adjacent vertices have distinct colors. We denote by $\chi(G)$ the *chromatic number* of G , i.e. the minimum number of colors in a proper coloring of G . For any $X \subseteq V$, let $c(X)$ be the set of colors that appear on the vertices of X . A *locally identifying coloring* (lid-coloring for short) of G is a proper vertex-coloring c of G such that for any two adjacent vertices u and v that are not twins (i.e. $N[u] \neq N[v]$), we have $c(N[u]) \neq c(N[v])$. A graph G is *k -lid-colorable* if it admits a locally identifying coloring using at most k colors and the minimum number of colors needed for any locally identifying coloring of G is the *locally identifying chromatic number* (lid-chromatic number for short) denoted by $\chi_{lid}(G)$. For a vertex u , we say that u *sees* color a if $a \in c(N[u])$. For two adjacent vertices

u and v , a color that is in the set $c(N[u]) \Delta c(N[v])$ separates u and v , or simply separates the edge uv . The notion of chromatic number (resp. lid-chromatic number) can be extended to a class of graphs \mathcal{C} as follows: $\chi(\mathcal{C}) = \sup\{\chi(G), G \in \mathcal{C}\}$ (resp. $\chi_{lid}(\mathcal{C}) = \sup\{\chi_{lid}(G), G \in \mathcal{C}\}$).

The following theorem is due to Bondy [2]:

Theorem 1 (Bondy's theorem [2]). *Let $\mathcal{A} = \{A_1, \dots, A_n\}$ be a collection of n distinct subsets of a finite set X . There exists a subset X' of X of size at most $n - 1$ such that the sets $A_i \cap X'$ are all distinct.*

Corollary 2. *Let C be a n -clique subgraph of G . There exists a vertex subset $S(C) \subseteq V(G)$ of size at most $n - 1$ that distinguishes all the pair of non-twin vertices of C .*

Proof. Let C be a n -clique subgraph of G induced by the vertex set $V(C) = \{v_1, v_2, \dots, v_n\}$. Let $\mathcal{A} = \{N[v_i] \mid v_i \in V(C)\}$ be a collection of distinct subsets of the finite set $X = \bigcup_{1 \leq i \leq n} N[v_i]$. Note that some v_i 's might be twins in G (i.e. $N[v_i] = N[v_j]$ for some $v_i, v_j \in V(C)$) and therefore $|\mathcal{A}|$ could be smaller than n . By Bondy Theorem, there exists $S(C) \subseteq X$ of size at most $|\mathcal{A}| - 1 \leq n - 1$ such that for any distinct elements A_1, A_2 of \mathcal{A} , we have $A_1 \cap S(C) \neq A_2 \cap S(C)$.

Let us prove that $S(C)$ is a set of vertices that distinguish all the pairs of non-twin vertices of C . For a pair of non-twin vertices v_i, v_j of C , we have $N[v_i] \neq N[v_j]$. By definition of $S(C)$, we have $N[v_i] \cap S(C) \neq N[v_j] \cap S(C)$, then there exists $w \in S(C)$ that belongs to $N[v_i] \Delta N[v_j]$. Therefore, w distinguishes the edge $v_i v_j$. \square

3 Bounded expansion classes of graphs

A rooted tree is a tree with a special vertex, called the *root*. The *height* of a vertex x in a rooted tree is the number of vertices on a path from the root to x (hence, the height of the root is 1). The *height* of a rooted tree T is the maximum height of the vertices of T . If x and y are two vertices of T , x is an *ancestor* of y in T if x belongs to the path between y and the root. The *closure* $\text{clos}(T)$ of a rooted tree T is the graph with vertex set $V(T)$ and edge set $\{xy \mid x \text{ is an ancestor of } y \text{ in } T, x \neq y\}$. The *tree-depth* $\text{td}(G)$ of a connected graph G is the minimum height of a rooted tree T such that G is a subgraph of $\text{clos}(T)$. If G is not connected, the tree-depth of G is the maximum tree-depth of its connected components.

Let p be a fixed integer. A *low tree-depth coloring* of a graph G (relatively to p) is a coloring of the vertices of G such that the union of any $i \leq p$ color classes induces a graph of tree-depth at most i . Let $\chi_p^{\text{td}}(G)$ be the minimum number of colors required in such a coloring. Note that as tree-depth one graphs and tree-depth two graphs are respectively the stables and star forests, χ_1^{td} and χ_2^{td} respectively correspond to the usual chromatic number and the star chromatic number.

In the following of this section, we first give a tight bound on the lid-chromatic number in terms of tree-depth.

Proposition 3. *For any graph G , $\chi_{lid}(G) \leq 2td(G) - 1$ and this is tight.*

Using this bound, we then bound the lid-chromatic number in terms of χ_3^{td} .

Theorem 4. *For any graph G ,*

$$\chi_{lid}(G) \leq 6^{\binom{\chi_3^{td}(G)}{3}}.$$

Classes of graphs of bounded expansion have been introduced by Nešetřil and Ossona de Mendez [10]. These classes contain minor closed classes of graphs and any class of graphs defined by an excluded topological minor. Actually, these classes of graphs are closely related to low tree-depth colorings:

Theorem 5 (Theorem 7.1 [10]). *A class of graphs \mathcal{C} has bounded expansion if and only if $\chi_p^{td}(\mathcal{C})$ is bounded for any p .*

We therefore deduce the following corollary from Theorems 4 and 5:

Corollary 6. *For any class \mathcal{C} of bounded expansion, $\chi_{lid}(\mathcal{C})$ is bounded.*

It is in particular true for a class of bounded tree-width. A consequence is that χ_{lid} is bounded for chordal graphs by a function of the clique number (which is equals to the tree-width plus 1 for a chordal graph). It is conjectured by Esperet *et al.* [6] that $\chi_{lid}(G) \leq 2\omega(G)$ if G is chordal.

We now prove Proposition 3.

Proof of Proposition 3. Let us first prove that the bound is tight. Consider the graph H_n obtained from a complete graph, with vertex set $\{a_1, \dots, a_n\}$, by adding a pendant vertex b_i to every a_i but one, say for $1 \leq i < n$. The tree-depth of this graph is at least n as it contains a n -clique. Indeed, given a rooted tree T , two vertices at the same height are non-adjacent in $clos(T)$, we thus need at least n levels. Actually the tree-depth of this graph is at most n since the tree T rooted at a_1 , and such that a_i has two sons a_{i+1} and b_i , for $1 \leq i < n$, has height n and is such that $clos(T)$ contains H_n as a subgraph.

Let us show that in any lid-coloring of H_n all the vertices must have distinct colors, and thus use $2n - 1 = 2td(H_n) - 1$ colors. Indeed, two vertices a_i must have different colors as the coloring is proper. A vertex b_j cannot use the same color as a vertex a_i , as otherwise the vertex a_j would only see the n colors used in the clique, just as a_n . Similarly if two vertices b_i and b_j would use the same color, the vertices a_i and a_j would see the same set of colors.

Let us now focus on the upper bound. We prove the result for a connected graph and by induction on the tree-depth of G , denoted by k . The result is clear for $k = 1$ (the graph is a single vertex).

Let G be a graph of tree-depth $k > 1$ and let T be a rooted tree of height k such that G is a subgraph of $clos(T)$. If T is a path, the result is clear since there are only k vertices. So assume that T is not a path, and let r be the root

of T . Let s be the smallest height such that there are at least two vertices of height $s + 1$. We name r_i , for $i \in \{1, \dots, s\}$, the unique vertex of height i . Let $R = \{r_1, \dots, r_s\}$. Note that each of the vertices of R is adjacent to all the vertices of $\text{clos}(T)$. Therefore, we can choose the way we label the s vertices in R (i.e. we can choose the height of each of them in T) without changing $\text{clos}(T)$.

Necessarily, $G \setminus R$ has at least two connected components. Let G_1, \dots, G_ℓ be its connected components and thus $\ell \geq 2$. We choose T such that s is minimal. It implies that for each $i \in \{1, \dots, s\}$, r_i has neighbors in all the components G_1, \dots, G_ℓ . Indeed, if it is not the case, by permuting the elements of R (this is possible by the above remark), we can assume without loss of generality that r_s does not have a neighbor in G_ℓ . Therefore, the set of edges $e(r_s, G_\ell) = \{r_s x : x \in V(G_\ell)\}$ of $\text{clos}(T)$ are not used by G . Then let T' be the tree obtained from T by moving the whole component G_ℓ one level up in such a way that the root of the subtree corresponding to G_ℓ is now the son of r_{s-1} (instead of r_s previously). Note that $\text{clos}(T')$ is isomorphic to $\text{clos}(T) \setminus e(r_s, G_\ell)$ and thus G is a subgraph of $\text{clos}(T')$. This new tree T' has two vertices at height s , contradicting the minimality of s .

Any connected component G_j has tree-depth at most $k' = k - s < k$. By induction, for each $j \in \{1, \dots, \ell\}$, there exists a lid-coloring c_j of G_j using colors in $\{1, \dots, 2k' - 1\}$. For each c_j , there is a minimum value s_j such that every vertex r_i sees a color in $\{1, \dots, s_j\}$ in G_j . We choose a $(2k' - 1)$ -lid-coloring c_j of G_j such that s_j is minimized. Note that for each color $a \leq s_j$, there exists $r_i \in R$ such that r_i sees color a in G_j but no other color of $\{1, \dots, s_j\}$. Otherwise, after permuting colors a and s_j , every vertex $r_i \in R$ would see a color in $\{1, \dots, s_j - 1\}$, contradicting the minimality of s_j . Assume without loss of generality that $s_1 \geq s_2 \geq \dots \geq s_\ell$.

We replace in c_1 the colors $1, 2, \dots, s_1$ by $1', 2', \dots, s'_1$. Note that now each vertex r_i sees a color in $\{1', \dots, s'_1\}$ (in G_1) and a color in $\{1, \dots, s_2\}$ (in G_2). Furthermore, the other vertices of G (that is the vertices in G_1, \dots, G_ℓ) do not have this property since $s_1 \geq s_2$. Thus at this step every edge xr_i with x in some G_j is separated.

Now we color each vertex r_i with color i^* . Let $c : V(G) \rightarrow \{1^*, \dots, s^*\} \cup \{1', \dots, s'_1\} \cup \{1, \dots, 2k' - 1\}$ be the current coloring of G .

Note that now every distinguishable edge xy in some G_j is separated. Indeed, either xy was distinguished in G_j and it has been separated by c_j , or xy is distinguished by some r_i and it is separated by the color i^* . Note also that c is a proper coloring.

It remains to deal with the edges $r_i r_j$. For that purpose we will refine some color classes. In the following lemma we show that such refinements do not damage what we have done so far.

Claim. *Consider a graph G and a coloring $\varphi : V(G) \rightarrow \{1, \dots, k\}$. Consider any refinement φ' of φ , obtained from φ by recoloring with color $k + 1$ some vertices colored i , for some i . Any edge xy of G properly colored (resp. separated) by φ is properly colored (resp. separated) by φ' .*

Indeed if $\varphi(x) \neq \varphi(y)$ then $\varphi'(x) \neq \varphi'(y)$, and if $i \in \varphi(N[x]) \Delta \varphi(N[y])$ then

i or $k+1 \in \varphi'(N[x]) \Delta \varphi'(N[y])$.

Let us define a relation \mathcal{R} among vertices in R by $r_i \mathcal{R} r_j$ if and only if $c(N[r_i]) = c(N[r_j])$. Let $R_1, \dots, R_{\bar{s}}$ be the equivalence classes of the relation \mathcal{R} (note that each R_i forms a clique since every r_i has distinct colors). We have $\bar{s} \geq s_1$. Indeed, by definition of s_1 and the coloring c_1 , for each color $a \in \{1', \dots, s'_1\}$, there exists $r_i \in R$ that sees a in G_1 but no other color of $\{1', \dots, s'_1\}$. This vertex r_i belongs to some equivalence class R_j and thus all the vertices of R_j sees color a in G_1 but no other color of $\{1', \dots, s'_1\}$.

By Corollary 2, there is a vertex set $S(R_i)$ of size at most $|R_i| - 1$ which distinguishes all pairs of non-twin vertices in R_i . We give to the vertices of $S(R_i)$ new distinct colors. By the previous claim, this last operation does not damage the coloring, and now all the distinguishable edges are separated.

Since for this last operation we need $s - \bar{s}$ new colors, since we used $2k' - 1$ colors $\{1, \dots, 2k' - 1\}$, s_1 colors $\{1', \dots, s'_1\}$ and s colors $\{1^*, \dots, s^*\}$, the total number of colors is $(s - \bar{s}) + (2k' - 1) + s_1 + s = 2k - 1 + s_1 - \bar{s} \leq 2k - 1$. This concludes the proof of the theorem. \square

We are now ready to prove Theorem 4:

Proof of Theorem 4. Let α be a low tree-depth coloring of G with parameter $p = 3$ and using $\chi_3^{\text{td}}(G)$ colors. Let $A = \{\alpha_1, \alpha_2, \alpha_3\}$ be a triplet of three distinct colors and let H_A be the subgraph of G induced by the vertices colored by a color of A . Since H_A has tree-depth at most 3, by Proposition 3, H_A admits a lid-coloring c_A with five colors (says colors 1 to 5). We extend c_A to the whole graph by giving color 0 to the vertices in $V(G) \setminus V(H_A)$.

Let A_1, A_2, \dots, A_k be the $k = (\chi_3^{\text{td}}(G))$ distinct triplets of colors. We now construct a coloring c of G giving to each vertex x of G the k -uplet

$$(c_{A_1}(x), c_{A_2}(x), \dots, c_{A_k}(x)).$$

The coloring c is using 6^k colors. Clearly it is a proper coloring: each pair of adjacent vertices will be in some common graph H_A and will receive distinct colors in this graph. Let x and y be two adjacent vertices with $N[x] \neq N[y]$. Let w be a vertex adjacent to only one vertex among x and y . Let $A = \{\alpha(x), \alpha(y), \alpha(w)\}$. Vertices x and y are not twins in the graph H_A . Hence $c_A(N[x]) \neq c_A(N[y])$ and therefore, $c(N[x]) \neq c(N[y])$. \square

4 Minor closed classes of graphs

Let G and H be two graphs. H is a *minor* of G if H can be obtained from G with successive edge deletions, vertex deletions and edge contractions. A class \mathcal{C} is *minor closed* if for any graph G of \mathcal{C} , for any minor H of G , we have $H \in \mathcal{C}$. The class \mathcal{C} is *proper* if it is not the class of all graphs. Let H be a graph. A *H -minor free graph* is a graph that does not have H as a minor. We denote by \mathcal{K}_n the K_n -minor-free class of graphs. It is clear that any proper minor closed class of graphs is included in the class \mathcal{K}_n for some n . It is folklore that any proper minor closed class of graphs \mathcal{C} has a bounded chromatic number $\chi(\mathcal{C})$.

The class of graphs of bounded expansion includes all the proper minor closed classes of graphs. Thus, by Corollary 6, proper minor closed classes have bounded lid-chromatic number. In this section, we focus on these latter classes and give an alternative upper bound on the lid-chromatic number. This gives us an explicit upper bound for the lid-chromatic number of planar graphs.

Consider any proper minor closed class of graphs \mathcal{C} . Since \mathcal{C} is proper, there exists n such that \mathcal{C} does not contain K_n , that is $\mathcal{C} \subseteq \mathcal{K}_n$. Let \mathcal{C}^N be the class of graphs defined by $H \in \mathcal{C}^N$ if and only if there exists $G \in \mathcal{C}$ and $v \in G$ such that $H = G[N(v)]$. Note that \mathcal{C}^N is a minor-closed class of graphs. Indeed, given any $H \in \mathcal{C}^N$, let $G \in \mathcal{C}$ and $v \in V(G)$ such that $H = G[N(v)]$. Let H' be any minor of H . Since \mathcal{C} is minor-closed and H is a subgraph of G , there exists a minor G' of G such that $H' = G'[N(v)]$. Therefore, H' belongs to \mathcal{C}^N .

We prove the following result on minor-closed classes of graphs:

Theorem 7. *Let \mathcal{C} be a proper minor closed class of graphs and let $n \geq 3$ be such that $\mathcal{C} \subseteq \mathcal{K}_n$. Then*

$$\chi_{lid}(\mathcal{C}) \leq 4 \cdot \chi_{lid}(\mathcal{C}^N) \cdot \chi(\mathcal{C})^{n-3}$$

The class of trees is exactly the class \mathcal{K}_3 . Esperet et al. [6] proved the following result.

Proposition 8 ([6]). $\chi_{lid}(\mathcal{K}_3) \leq 4$.

It is clear that \mathcal{K}_3^N is the class of stable graphs and therefore, $\chi_{lid}(\mathcal{K}_3^N) = 1$. Note that Theorem 7 implies Proposition 8.

Assume that $\chi_{lid}(\mathcal{K}_{n-1})$ is bounded for some $n \geq 4$. It is clear that $\mathcal{K}_n^N = \mathcal{K}_{n-1}$. Then, by Theorem 7, we have $\chi_{lid}(\mathcal{K}_n) \leq 4 \cdot \chi_{lid}(\mathcal{K}_{n-1}) \cdot \chi(\mathcal{K}_n)^{n-3}$. Since $\chi_{lid}(\mathcal{K}_{n-1})$ and $\chi(\mathcal{K}_n)$ are bounded, $\chi_{lid}(\mathcal{K}_n)$ is bounded.

Esperet et al. [6] also proved the following result.

Proposition 9 ([6]). *If G is an outerplanar graph, $\chi_{lid}(G) \leq 20$.*

We can then deduce from Theorem 7 and Proposition 9 the following corollary:

Corollary 10. *Let \mathcal{P} be the class of planar graphs. Then $\chi_{lid}(\mathcal{P}) \leq 1280$.*

Proof. Any graph $G \in \mathcal{P}$ is $\{K_{3,3}, K_5\}$ -minor free and thus \mathcal{P} is a proper minor closed class of graphs. Moreover, the neighborhood of any vertex of $G \in \mathcal{P}$ is an outerplanar graph. By Proposition 9, we have $\chi_{lid}(\mathcal{P}^N) \leq 20$. Furthermore, the Four-Color-Theorem gives $\chi(\mathcal{P}) = 4$. By Theorem 7, $\chi_{lid}(\mathcal{P}) \leq 4 \times 20 \times 4^2 = 1280$. \square

We finally give the proof of Theorem 7.

Proof of Theorem 7. Let $G \in \mathcal{C}$ and let u be a vertex of minimum degree. For any i , define $V_{u,i}$ as the set of vertices of G at distance exactly i from u and let $G_{u,i} = G[V_{u,i}]$. Let s be the largest distance from a vertex of V to u . In other words, there are $s + 1$ nonempty sets $V_{u,i}$ (note that $V_{u,0} = \{u\}$).

For any i , contracting in G the subgraph $G[V_{u,0} \cup V_{u,1} \cup \dots \cup V_{u,i-1}]$ in a single vertex x gives a graph $G' \in \mathcal{C}$ such that x is exactly adjacent to every vertex of $G_{u,i}$. Therefore, for any i , $G_{u,i} \in \mathcal{C}^N$. Hence, $\chi_{lid}(G_{u,i}) \leq \chi_{lid}(\mathcal{C}^N)$ for any i . Moreover, $\mathcal{C}^N \subseteq \mathcal{K}_{n-1}$. Indeed, suppose that there exists $H \in \mathcal{C}^N$ that admits K_{n-1} as a minor. Therefore there exists $G \in \mathcal{C}$ such that $H = G[N(v)]$ for some $v \in G$. Taking v together with its neighborhood would give K_n as a minor, that contradicts the fact that $\mathcal{C} \subseteq \mathcal{K}_n$. Hence, any $G_{u,i} \in \mathcal{K}_{n-1}$.

We construct a lid-coloring of G using $4 \cdot \chi_{lid}(\mathcal{C}^N) \cdot \chi(\mathcal{C})^{n-3}$ colors. This coloring is constructed with three different colorings of the vertices of G : c_1 which uses 4 colors, c_2 which uses $\chi_{lid}(\mathcal{C}^N)$ colors and c_3 which is itself composed of $n-3$ colorings with $\chi(\mathcal{C})$ colors. The final color $c(v)$ of a vertex v will be the triplet $(c_1(v), c_2(v), c_3(v))$. Hence the coloring c uses at most $4\chi_{lid}(\mathcal{C}^N)\chi(\mathcal{C})^{n-3}$ colors. The coloring c_1 is used to separate the pairs of vertices that lie in distinct sets $V_{u,i}$. The coloring c_2 separates the pairs of vertices that lie in the same set $V_{u,i}$ and are not twins in $G_{u,i}$. Finally, the coloring c_3 separates the pairs of vertices that lie in the same set $V_{u,i}$, that are twins in $G_{u,i}$ but that are not twins in G .

The coloring c_1 is simply defined by $c_1(v) \equiv i \pmod{4}$ if $v \in V_{u,i}$.

To define c_2 , we define for each i , $0 \leq i \leq s$, a lid-coloring c_2^i of $G_{u,i}$ using colors 1 to $\chi_{lid}(\mathcal{C}^N)$. Then c_2 is defined by $c_2(v) = c_2^i(v)$ if $v \in V_{u,i}$.

We now define the coloring c_3 . Let $V_{u,i}^{id}$ be the set of vertices of $V_{u,i}$ that have a twin in $G_{u,i}$:

$$V_{u,i}^{id} = \{v \in V_{u,i} \mid \exists w \in V_{u,i}, N_{G_{u,i}}[v] = N_{G_{u,i}}[w]\}.$$

Let $G_{u,i}^{id} = G_{u,i}[V_{u,i}^{id}]$. Since the relation “be twin” is transitive (i.e. if u and v are twins, and v and w are twins, then u and w are twins), then $G_{u,i}^{id}$ is clearly a union of cliques. In addition, since $G_{u,i} \in \mathcal{K}_{n-1}$, the connected components of $G_{u,i}^{id}$ are cliques of size at most $n-2$.

Let C be a clique of $G_{u,i}^{id}$. By Corollary 2, there exists a subset $S(C) \subseteq V(G)$ of at most $n-3$ vertices that distinguishes all the pairs of non-twin vertices of C . Note that by definition of C , $S(C) \cap V_{u,i} = \emptyset$, and thus $S(C) \subseteq V_{u,i-1} \cup V_{u,i+1}$.

Let $\mathcal{S} = \{(v, C) \mid v \in S(C) \text{ and } C \text{ is a clique in a graph } G_{u,i}^{id}\}$. We partition \mathcal{S} in $s \times (n-3)$ sets S_i^k , $1 \leq i \leq s$, $1 \leq k \leq n-3$, such that:

- if $(v, C) \in S_i^k$ for some k , then $v \in V_{u,i}$;
- if (v, C) and (w, C') are two elements of S_i^k , then $C \neq C'$.

This partition can be done because each set $S(C)$ has size at most $n-3$.

For each $S_i^k = \{(x_1, C_1), (x_2, C_2), \dots, (x_t, C_t)\}$, we define a graph H_i^k as follows. We start from the graph induced by $V_{u,i} \cup V(C_1) \cup V(C_2) \cup \dots \cup V(C_t)$. Then, for each (x_j, C_j) in S_i^k , we contract C_j in a single vertex y_j and finally, we contract the edge $x_j y_j$ on the vertex x_j . Note that $V_{u,i}$ is the vertex set of H_i^k . Note also that $H_i^k \in \mathcal{C}$ since it is obtained from a subgraph of G by successive edge-contractions. Therefore, $\chi(H_i^k) \leq \chi(\mathcal{C})$.

We now define a proper coloring $c_3^{i,k}$ of H_i^k with colors 1 to $\chi(\mathcal{C})$. Let c_3^k be the coloring of vertices of G defined by $c_3^k(v) = c_3^{i,k}(v)$ if $v \in V_{u,i}$. Finally, c_3 is defined by $c_3(v) = (c_3^1(v), \dots, c_3^{n-3}(v))$, and the final color of v is $c(v) = (c_1(v), c_2(v), c_3(v))$.

We now prove that c is a lid-coloring of G . First, c is a proper coloring. Indeed, two adjacent vertices that are not in the same set $V_{u,i}$ lie in consecutive sets $V_{u,i}$ and $V_{u,i+1}$ and thus have different colors in c_1 , and two adjacent vertices in the same set $V_{u,i}$ have different colors in c_2 (which induces a proper coloring on $V_{u,i}$).

Let now x and y be two adjacent vertices with $N[x] \neq N[y]$. We will prove that $c(N[x]) \neq c(N[y])$. We distinguish three cases.

Case 1: $x \in V_{u,i}$ and $y \in V_{u,i+1}$.

If $x = u$, then y has a neighbor v in $V_{u,i+2} = V_{u,2}$. Indeed, u is taken with minimum degree, so y has at least as many neighbors as u and does not have the same neighborhood than u , implying that y has a neighbor in $V_{u,2}$. Then $c_1(v) = 2 \notin c_1(N[u])$ and so $c(N[x]) \neq c(N[y])$.

Otherwise, x has neighbor v in $V_{u,i-1}$ and $c_1(v) \equiv i-1 \pmod{4} \in c_1(N[x])$. On the other hand, all the neighbors of y belong to $V_{u,i} \cup V_{u,i+1} \cup V_{u,i+2}$ and therefore $c_1(N[y]) \subseteq \{i, i+1, i+2 \pmod{4}\}$. Thus, $c(N[x]) \neq c(N[y])$.

Case 2: x and y belong to $V_{u,i}$ and they are not twins in $V_{u,i}$ (i.e. $N_{V_{u,i}}[x] \neq N_{V_{u,i}}[y]$).

By definition of the coloring c_2^i , there exists a color a that separates x and y , i.e. $a \in c_2^i(N_{V_{u,i}}[x]) \Delta c_2^i(N_{V_{u,i}}[y])$. Then we necessarily have $c(N[x]) \neq c(N[y])$.

Case 3: x and y belong to $V_{u,i}$ and they are twins in $V_{u,i}$ (i.e. $N_{V_{u,i}}[x] = N_{V_{u,i}}[y]$).

In this case, vertices x and y are in the set $V_{u,i}^{id}$. Let C be the clique of $G_{u,i}$ containing x and y . Let $v \in S(C)$ that distinguishes x and y ; thus, $v \in V_{u,j}$ for $j = i-1$ or $j = i+1$. Wlog, $v \in N[x]$ but $v \notin N[y]$. Let S_j^k be the part of \mathcal{S} that contains (v, C) . Suppose that there exists a neighbor w of y such that $c(v) = c(w)$. Then w lies in $V_{u,j}$ because of the coloring c_1 . However, in the graph H_j^k , the vertex v is adjacent to all the neighbors of y in $V_{u,j}$, and in particular is adjacent to w ; therefore, $c_3^{j,k}(v) \neq c_3^{j,k}(w)$, a contradiction. Therefore, the vertex y does not have any neighbor that has the same color as v . Hence, $c(v) \notin c(N[y])$, and $c(N[x]) \neq c(N[y])$.

□

References

- [1] P.N. Balister, O.M. Riordan, and R.H. Schelp. Vertex-distinguishing edge-colorings of graphs. *J. Graph Theory*, **42**:95–109, 2003.
- [2] J. A. Bondy. Induced subsets. *J. Combin. Theory Ser. B*, **12**(2):201–202, 1972.
- [3] A.C. Burris and R.H. Schelp. Vertex-distinguishing proper edge-colorings. *J. Graph Theory*, **26**:73–83, 1997.
- [4] G. Chartrand, D. Erwin, M.A. Henning, P.J. Slater and P. Zhang. The locating-chromatic number of a graph. *Bull. Inst. Combin. Appl.*, **36**:89–101, 2002.
- [5] J. Cerný, M. Horňák, and R. Soták. Observability of a graph. *Math. Slovaca*, **46**(1):21–31, 1996.
- [6] L. Esperet, S. Gravier, M. Montassier, P. Ochem and A. Parreau. Locally identifying coloring of graphs. *Electron. J. Combin.*, **19**(2), 2012.
- [7] F. Foucaud, I. Honkala, T. Laihonon, A. Parreau and G. Perarnau. Locally identifying colouring of graphs with given maximum degree. *Discrete Math.*, **312**(10), 2012.
- [8] M. G. Karpovsky, K. Chakrabarty, and L. B. Levitin. On a new class of codes for identifying vertices in graphs. *IEEE Trans. Inform. Theory*, 44:599–611, 1998.
- [9] A. Lobstein. Identifying and locating-dominating codes in graphs, a bibliography. Published electronically at <http://perso.enst.fr/~lobstein/debutBIBidetlocdom.pdf>.
- [10] J. Nešetřil and P. Ossona de Mendez. Grad and classes with bounded expansion I. Decompositions. *European J. Combin.*, **29**:760–776, 2008.