



Homotopes of Symmetric Spaces II. Structure Variety and Classification

Wolfgang Bertram, Pierre Bieliavsky

► To cite this version:

Wolfgang Bertram, Pierre Bieliavsky. Homotopes of Symmetric Spaces II. Structure Variety and Classification. 2012. hal-00535877v2

HAL Id: hal-00535877

<https://hal.archives-ouvertes.fr/hal-00535877v2>

Preprint submitted on 5 Mar 2012

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Homotopes of Symmetric Spaces

II. Structure Variety and Classification

Wolfgang Bertram

IECN Université Nancy I, France

e-mail: bertram@iecn.u-nancy.fr

Pierre Bieliavsky

Université Catholique de Louvain, Belgium.

e-mail: pierre.bieliavsky@gmail.com

Abstract. We classify *homotopes* of classical symmetric spaces (introduced in [Be08] and studied in Part I of this work). Our classification uses the *fibred structure* of homotopes: they are fibred as symmetric spaces, with flat fibers, over a non-degenerate base; the base spaces correspond to *inner ideals* in Jordan pairs. Using that inner ideals in classical Jordan pairs are always complemented in the sense of [LoN94], the classification of homotopes is obtained by combining the classification of inner ideals with the one of isotopes of a given inner ideal.

Subject classification. 17B60, 17B70 17C37, 53C35 32M15

Key words. contraction, complementation, involution, inner ideal, homotope (isotope), Jordan pairs and triple systems, 3-graded Lie algebra, Lie triple system, structural transformation, symmetric space, pseudo-inverse, idempotent

Introduction

Triple systems are trilinear maps $V^3 \rightarrow V$ satisfying certain algebraic properties, giving rise to notions such as *Lie-*, *Jordan-* or *associative triple systems*. Often it is more conceptual to consider a pair of vector spaces (V^+, V^-) and two trilinear maps $T^\pm : V^\pm \times V^\mp \times V^\pm \rightarrow V^\pm$, verifying certain algebraic properties (e.g., *Jordan pairs*, *associative pairs*). Given such an object (V^\pm, T^\pm) , we define a subset of $\text{Hom}(V^+, V^-)$, called the *structure variety* of (V^\pm, T^\pm) :

$$\text{Svar}(V^+) := \{ \alpha : V^+ \rightarrow V^- \mid \alpha \text{ linear, } \forall u, v, w \in V^+ : T^-(\alpha u, v, \alpha w) = \alpha T^+(u, \alpha v, w) \}. \quad (1)$$

Then elements $\alpha \in \text{Svar}(V^+)$ parametrize families of trilinear maps on V^+ , given by

$$T_\alpha : V^+ \times V^+ \times V^+ \rightarrow V^+, \quad T_\alpha(x, y, z) := T^+(x, \alpha(y), z). \quad (2)$$

It is well-known (cf. Lemma 1.6) that, if (V^\pm, T^\pm) is a *Jordan pair*, then T_α is a *Jordan triple system*, and since skew-symmetrizing a Jordan triple system yields a Lie triple system (Lemma 1.3), we conclude that

$$R_\alpha(x, y)z := T_\alpha(x, y, z) - T_\alpha(y, x, z) = T^+(x, \alpha y, z) - T^+(y, \alpha x, z) \quad (3)$$

defines a *Lie triple system*. Thus, given some Jordan pair (V^\pm, T^\pm) , its structure variety parametrizes Lie triple structures on V^+ . Such families of Lie triple systems are called *homotopes* and have been investigated in Part I of this work ([BeBi]). When α is an invertible endomorphism, we speak of *isotopes*; in the examples from Part I, the isotopes belong to semisimple or reductive symmetric spaces, and when α becomes singular, these spaces contract to degenerate homotopes, until, for the most singular value $\alpha = 0$, we get the flat space V^+ . In this work, we prove two kinds of results:

1. A *classification theorem* implying that the lists given in Part I are indeed complete parametrizations of structure varieties of classical symmetric spaces,

2. results on the *structure* of the homotopes: we show that all regular homotopes have a *fibred structure* – they are symmetric spaces fibered over non-degenerate symmetric spaces, having flat fibers, and such that the fibration splits.

Both items are closely related to each other: the key ingredient in both cases is the link with *inner ideals*, and with the notion of *complementation of inner ideals* introduced by Loos and Neher ([LoN94]). More precisely, the image I of the endomorphism $\alpha : V^+ \rightarrow V^-$ is always an *inner ideal in V^-* , and the kernel $\ker \alpha$ is a flat ideal in (V^+, R_α) viewed as a Lie triple system. Thus we get a short exact sequence of Lie triple systems

$$0 \rightarrow \ker \alpha \rightarrow (V^+, R_\alpha) \rightarrow I \rightarrow 0. \quad (4)$$

On the level of symmetric spaces, this corresponds to a fibration over a base belonging to I . In general, I does not carry a Jordan triple structure, nor does this sequence split. We introduce a natural notion of *regularity of homotopies* and prove that both properties hold for regular α . For simple real finite-dimensional Jordan pairs, we show that every α is indeed regular.

The key observation for proving these results is that α is regular if and only if the inner ideal $I = \text{Im} \alpha$ is *complemented*, i.e., there is an inner ideal J in V^+ such that $V^+ = J \oplus \text{Kern}(I)$ and $V^- = I \oplus \text{Kern}(J)$. By results of Loos and Neher ([LoN94]) it is known that, e.g. for simple finite-dimensional real Jordan pairs, every inner ideal admits a complement. This opens the way for classification: one needs, on the one hand, the (known) classification of inner ideals I in V^- , and on the other hand, for given I , in order to describe all α with $I = \text{Im} \alpha$, we have to use the classification of isotopes of I , i.e., of the invertible elements in $\text{Svar}(I)$ (which is also known). Putting these two together, one can finally show that, for classical Jordan pairs, the lists given in Part I ([BeBi], Theorem 4.2) lead to a complete parametrization of structure varieties (with one notable exception: spaces of skew-symmetric matrices $\text{Asym}(n, \mathbb{K})$ contain a “degenerate” class of inner ideals, called “point spaces” by McCrimmon [McC71], and homotopes corresponding to such inner ideals are not given by the construction from Part I).

The organization of this paper is as follows: Chapter 1 contains definitions and basic facts about Jordan pairs and their structure varieties; in Chapter 2 we introduce the notion of regularity and explain the link with inner ideals; Chapter 3 contains classifications and some more results on special cases. While for classification we make quite restrictive assumptions (base field $\mathbb{K} = \mathbb{R}$, finite-dimensionality and simplicity; this framework is most important for the theory of symmetric spaces), most of the results from Chapters 1 and 2 are valid very generally (arbitrary Jordan pairs over general base rings).

Notation. Throughout, \mathbb{K} denotes a commutative base ring in which 2 and 3 are invertible.

1 The structure variety of a Jordan pair

A Γ -graded Lie algebra is a Lie algebra of the form $\mathfrak{g} = \bigoplus_{\gamma \in \Gamma} \mathfrak{g}_\gamma$ with $[\mathfrak{g}_\gamma, \mathfrak{g}_\delta] \subset \mathfrak{g}_{\gamma+\delta}$ where $(\Gamma, +)$ is an abelian group.

1.1 $\mathbb{Z}/(2)$ -graded Lie algebras and Lie triple systems

Recall that $\mathbb{Z}/(2)$ -graded Lie algebras $\mathfrak{g} = \mathfrak{h} \oplus \mathfrak{q}$ (i.e. $[\mathfrak{h}, \mathfrak{h}] \subset \mathfrak{h}$, $[\mathfrak{q}, \mathfrak{q}] \subset \mathfrak{h}$, $[\mathfrak{h}, \mathfrak{q}] \subset \mathfrak{q}$) essentially correspond to *Lie triple systems (LTS)* by considering \mathfrak{q} with the triple Lie bracket $[X, Y, Z] := [[X, Y], Z] =: R(X, Y)Z$. Axiomatically, this triple Lie bracket can be characterized by the three properties

(LT1) $[X, Y, Z] = -[Y, X, Z]$

(LT2) $[X, Y, Z] + [Y, Z, X] + [Z, X, Y] = 0$

(LT3) the endomorphism $D := R(U, V)$ is a derivation of the trilinear product $[X, Y, Z]$.

Every LTS \mathfrak{q} is obtained from a $\mathbb{Z}/(2)$ -graded Lie algebra: we may take for \mathfrak{g} the *standard imbedding* $\mathfrak{q} \oplus [\mathfrak{q}, \mathfrak{q}] \subset \mathfrak{q} \oplus \mathfrak{Der}(\mathfrak{q})$ (see [Lo69]). Then $\sigma := \text{id}_{\mathfrak{h}} \oplus (-\text{id}_{\mathfrak{q}})$ is an automorphism of order 2, and the pair (\mathfrak{g}, σ) is called a *symmetric pair*. See Section 1 of Part I for further remarks on Lie triple systems.

1.2 Jordan pairs, 3-graded Lie algebras and polarized Lie triple systems

A 3-graded Lie algebra is a \mathbb{Z} -graded Lie algebra with $\mathfrak{g}_n = 0$ for $|n| > 1$:

$$\mathfrak{g} = \mathfrak{g}_{-1} \oplus \mathfrak{g}_0 \oplus \mathfrak{g}_1.$$

Note that then the odd part $\mathfrak{q} := \mathfrak{g}_1 \oplus \mathfrak{g}_{-1}$ is a LTS, which, moreover, is *polarized*: the endomorphism $D := \text{id}_{\mathfrak{g}_1} \oplus (-\text{id}_{\mathfrak{g}_{-1}})$ is a polarization of the LTS \mathfrak{q} , that is, $D^2 = 1$, and D is a derivation of the LTS commuting with \mathfrak{h} in the sense that $R(X, Y)D = DR(X, Y)$. Conversely, the standard imbedding of a polarized LTS is a 3-graded Lie algebra. If \mathfrak{g} is a 3-graded Lie algebra, the pair $(V^+, V^-) := (\mathfrak{g}_1, \mathfrak{g}_{-1})$ together with the trilinear maps

$$T^\pm : V^\pm \times V^\mp \times V^\pm \rightarrow V^\pm, \quad (u, v, w) \mapsto T^\pm(u, v, w) := [[u, v], w]$$

is called the *associated (linear) Jordan pair*. It satisfies the identities

$$(J1) \quad T^\pm(u, v, w) = T^\pm(w, v, u)$$

$$(J2) \quad T^\pm(u, v, T^\pm(x, y, z)) = T(T(u, v, x), y, z) - T(x, T^\mp(v, u, y), z) + T(x, y, T(u, v, z))$$

Definition 1.1 A (linear) Jordan pair is a pair of vector spaces (V^+, V^-) with trilinear maps T^\pm satisfying (J1) and (J2). Homomorphisms of Jordan pairs are pairs of linear maps $(f^\pm : V^\pm \rightarrow W^\pm)$ such that $f^\pm(T^\pm(u, v, w)) = T^\pm(f^\pm u, f^\mp v, f^\pm w)$. Obviously, Jordan pairs form a category.

Every Jordan pair is obtained from a 3-graded Lie algebra in the way just described.

Definition 1.2 For any Jordan pair (V^+, V^-) , the opposite pair is (V^-, V^+) . An involution of (V^+, V^-) is an isomorphism $\phi^\pm : V^\pm \rightarrow V^\mp$ onto the opposite pair such that $\phi^- = (\phi^+)^{-1}$.

1.3 Involutions, Jordan triple systems and the Jordan-Lie functor

Assume \mathfrak{g} is a 3-graded Lie algebra and θ an automorphism of \mathfrak{g} of order two and reversing the grading. Then θ restricts to an automorphism of the LTS $\mathfrak{q} = \mathfrak{g}_1 \oplus \mathfrak{g}_{-1}$ anticommuting with the polarization D , called a *para-real form* of (\mathfrak{q}, R, D) , and the restriction $\mathfrak{g}_1 \rightarrow \mathfrak{g}_{-1}$ defines an involution of the Jordan pair (V^+, V^-) . Conversely, every para-real form (every involution) induces an automorphism θ as above. If (\mathfrak{g}, θ) is as above, the space $V := \mathfrak{g}_1$ with trilinear product

$$T(x, y, z) := [[x, \theta(y)], z]$$

is a *Jordan triple system (JTS)*, i.e., it satisfies the identities (J1) and (J2) with superscripts \pm omitted. Every Jordan triple system (V, T) is obtained in this way from an involutive 3-graded Lie algebra (see, e.g., [Be00]): Jordan triple systems are equivalent to Jordan pairs with involution, and the Jordan pair (V^+, V^-) obtained by forgetting the involution is then called the *underlying Jordan pair* of (V, T) . Note that in general there are many grade-reversing involutions, and there is no canonical choice of one of them; therefore it is often conceptually clearer to work with Jordan pairs and not with triple systems (as we will do below).

Lemma 1.3 If (V, T) is a Jordan triple system, then $R := R_T$ with

$$R_T(x, y, z) := T(x, y, z) - T(y, x, z)$$

defines a LTS on V . This defines a functor $T \mapsto R_T$ from the category of JTS to the one of LTS, called the Jordan-Lie functor.

Proof. Direct computation – see [Mey73] or [Be00], Lemma III.2.6. One may note also that the LTS (V, R_T) is isomorphic to the fixed point space under θ of the para-real form of (\mathfrak{q}, R, D) mentioned above. ■

1.4 Definition of the structure variety

Let $V = (V^+, V^-)$ be a Jordan pair. We define the *quadratic operators* for $x \in V^\pm$ by

$$Q^\pm(x) : V^\mp \rightarrow V^\pm, \quad y \mapsto \frac{1}{2}T^\pm(x, y, x).$$

Definition 1.4 *The structure variety (of V^+) is the subset of $\text{Hom}(V^+, V^-)$ defined by Equation (1). An element $\alpha \in \text{Svar}(V^+)$ is called a homotopy of the Jordan pair. Equivalently (by polarization), a linear map α is a homotopy iff for all $u \in V^+$,*

$$Q^-(\alpha u) = \alpha \circ Q^+(u) \circ \alpha.$$

The set of invertible elements in $\text{Svar}(V^+)$ is denoted by $\text{Svar}(V^+)^\times$. For a JTS (V, T) , the structure variety $\text{Svar}(V)$ is defined in the same way, omitting the superscripts \pm . In other words, the structure variety of a JTS is the structure variety of the underlying Jordan pair.

Remarks. (1) If α is an invertible element in $\text{Svar}(V^+)$, then, by change of variables $v' := \alpha v$, we see that (α, α^{-1}) is an involution. Conversely, if (ϕ^+, ϕ^-) is an involution, then $\phi^+ \in \text{Svar}(V^+)^\times$. In this sense $\text{Svar}(V^+)^\times$ corresponds to the space of involutions, and thus $\text{Svar}(V^+)$ can be seen as a sort of completion of this space.

(2) The *fundamental formula* from Jordan theory $Q(Q(x)y) = Q(x)Q(y)Q(x)$ says that all quadratic operators $Q^-(x) : V^+ \rightarrow V^-$, where $x \in V^-$, belong to $\text{Svar}(V^+)$.

(3) Obviously, $\text{Svar}(V^+)$ is stable under multiplication by scalars and contains 0.

(4) There is a similar definition of $\text{Svar}(V^-)$. From a categorial point of view, one may interpret the structure variety as the *space of self-adjoint structural transformations from a Jordan pair to its opposite pair*. Here are the relevant definitions:

Definition 1.5 *A structural transformation between two Jordan pairs (V^+, V^-) , (W^+, W^-) is a pair of linear maps $(g : V^+ \rightarrow W^+, h : W^- \rightarrow V^-)$ such that, for all $u, w \in V^+$ and $v, z \in W^-$:*

$$gT_V^+(u, hv, w) = T_W^+(gu, v, gw), \quad hT_W^-(v, gw, z) = T_V^-(hv, w, hz).$$

Jordan pairs with structural transformations form a category in which composition of morphisms is defined by $(g, h)(g', h') := (g \circ g', h' \circ h)$. In particular, the structural transformations from (V^+, V^-) to itself form a semigroup, called the structure monoid. By definition, the structure group of V is the group of invertible elements of the structure monoid, that is, the group of automorphisms of (V^+, V^-) . A structural transformation (g, h) from (V^+, V^-) to its opposite pair (V^-, V^+) is called self-adjoint if $g = h$.

With these definitions, α belongs to $\text{Svar}(V^+)$ if and only if (α, α) is a structural transformation to the opposite pair. Note also that, if (f^+, f^-) is an isomorphism in the “usual” sense, then $(f^+, (f^-)^{-1})$ is an isomorphism in the new (“structural”) category.

Lemma 1.6 *If $\alpha \in \text{Svar}(V^+)$, then $T_\alpha(x, y, z) := T^+(x, \alpha y, z)$ defines a Jordan triple system on V^+ , called the α -homotope of (V, T) .*

Proof. Straightforward computation – see [Mey73] or [Be00], Lemma III.4.5. ■

Applying the Jordan-Lie functor to T_α , we get by Equation (3) a LTS on V^+ , also called α -homotope of R . Thus we have a map $\alpha \rightarrow R_\alpha$ from the structure variety to the variety of Lie triple products on V^+ .

Lemma 1.7 *Assume $\alpha \in \text{Svar}(V^+)$ and $(g, h) : (W^+, W^-) \rightarrow (V^+, V^-)$ a structural transformation. Then $h \circ \alpha \circ g : W^+ \rightarrow W^-$ belongs to $\text{Svar}(W^+)$, and*

$$g : (W^+, T_{h\alpha g}) \rightarrow (V^+, T_\alpha), \quad v \mapsto gv$$

is a JTS-homomorphism (and hence also a LTS-homomorphism $R_{h\alpha g} \rightarrow R_\alpha$).

Proof. By definition of composition of morphisms, $(h, g)(\alpha, \alpha)(g, h) = (h\alpha g, h\alpha g)$, and this is a structural transformation of W . Hence $h\alpha g : W^+ \rightarrow W^-$ is self-adjoint structural, i.e., it belongs to $\text{Svar}(W^+)$. Since $gT_{h\alpha g}(u, v, w) = gT(u, h\alpha gv, w) = T(gu, \alpha gv, gw) = T_\alpha(gu, gv, gw)$, it follows that g induces a JTS-homomorphism as claimed. ■

Corollary 1.8 *The structure group of the Jordan pair V acts on the structure variety, and if α and α' belong to the same orbit under this action, then the JTS T_α and $T_{\alpha'}$ are isomorphic.*

Proof. This is the special case $V = W$ and (g, h) invertible of the lemma. ■

For a classification of homotopes up to isomorphy it thus suffices to consider structure group orbits in $\text{Svar}(V^+)$. Note that the structure group contains all pairs (rid, rid) with $r \in \mathbb{K}^\times$, and hence it follows that α and $r^2\alpha$ for $r \in \mathbb{K}^\times$ are conjugate under the structure group. In general, if $\mathbb{K} = \mathbb{R}$, α and $-\alpha$ will not be conjugate to each other (they are *c-duals* of each other, cf. remarks in Part I).

Corollary 1.9 *Assume $\alpha \in \text{Svar}(V^+)$ and $\beta \in \text{Svar}(V^-)$. Then $\beta\alpha\beta$ belongs to $\text{Svar}(V^+)$, and $\alpha\beta\alpha$ belongs to $\text{Svar}(V^-)$.*

Proof. This is the special case $(W^+, W^-) = (V^-, V^+)$ and $g = h = \beta$ (resp. $g = h = \alpha$) of the lemma. ■

2 Fibration: Inner ideals, kernels and complementation

The symmetric space M_α corresponding to the LTS R_α has a *fibred structure*: the base corresponds to the image of α and the fiber to the kernel of α . In this chapter we investigate these important features and show that they are also the key for proving classification results. It turns out that classification is made possible by the fact that, in suitably “regular” situations, the above mentioned fibering splits, and that such splittings correspond, in a Jordan theoretic language, to *complementation of inner ideals* in the sense of [LoN94].

2.1 Inner ideals and their kernels

We are going to describe some properties of the image $\text{Im}\alpha \subset V^-$ and the kernel $\ker \alpha \subset V^+$ of a homotopy $\alpha : V^+ \rightarrow V^-$.

Lemma 2.1 *Assume $\alpha \in \text{Svar}(V^+)$. Then $\ker \alpha$ is an ideal in the Jordan triple system (V^+, T_α) , and hence it is also an ideal in the Lie triple system (V^+, R_α) .*

Proof. Let $K := \ker \alpha$; in order to check that $T_\alpha(K, V^+, V^+) + T_\alpha(V^+, K, V^+) + T_\alpha(V^+, V^+, K) \subset K$, it suffices to check that $\alpha T_\alpha(u, v, w) = 0$ whenever one of u, v, w belongs to K . But this follows immediately from the relation $\alpha T_\alpha(u, v, w) = \alpha T(u, \alpha v, w) = T(\alpha u, v, \alpha w)$. ■

The image $\text{Im}\alpha$ is in general not even a “sub-JTS” of V^- (so far there is no JTS-structure on V^-).

Definition 2.2 *Let (V^+, V^-) be a Jordan pair. An inner ideal (in V^-) is a subspace $I \subset V^-$ such that*

$$T^-(I, V^+, I) \subset I.$$

Example 2.3 *Images of structural transformations are inner ideals: assume $(g, h) : (V^+, V^-) \rightarrow (W^+, W^-)$ is a structural transformation, then the $\text{Im}h$ is an inner ideal in W^- since*

$$T^-(hV^-, W^+, hV^-) \subset hT^-(V^-, gW^+, V^-) \subset \text{Im}h.$$

In particular, for $\alpha \in \text{Svar}(V^+)$, the image $\text{Im}\alpha$ is an inner ideal in V^- .

Definition 2.4 *Let I be an inner ideal in V^- . We define, following [LoN94], the kernel of I to be the subspace of V^+ given by*

$$\text{Kern}I := \{x \in V^+ \mid T^-(I, x, I) = 0\} = \bigcap_{y \in I} \ker Q^-(y).$$

Note that the kernel of an inner ideal and the inner ideal itself live in different spaces (V^+ , resp. V^-). We recall also that a Jordan pair (V^+, V^-) is called non-degenerate if $T^\mp(x, V^\pm, x) = 0$ implies $x = 0$.

Example 2.5 *The kernel of an image $\text{Im}g$ of a structural transformation (g, h) is the kernel of h :*

$$\text{Kern}(\text{Im}g) = \ker h.$$

More precisely, if $(g, h) : (V^+, V^-) \rightarrow (W^+, W^-)$ is structural, let $I := \text{Im}g$ and $K := \ker h$; then $T^-(gu, k, gv) = gT^-(u, hk, v) = 0$ for all $k \in K$, whence $\ker h \subset \text{Kern}(\text{Im}g)$. The converse holds if V is non-degenerate ([LoN94], Lemma 1.6 (b)).

2.2 Regularity and pseudo-inverses

An inner ideal I does in general not carry a Jordan triple system structure (we can only say that (V^+, I) is a Jordan pair), and therefore it does not make sense to say that $T_\alpha/\ker \alpha$ and $\text{Im} \alpha$ be isomorphic as JTS (and thus as LTS). However, this will be the case if α is *regular* in the sense to be explained now.

Definition 2.6 *Let (V^+, V^-) be a Jordan pair and $\alpha \in \text{Svar}(V^+)$ (hence $\alpha : V^+ \rightarrow V^-$). We say that α is regular if there is an element $\beta \in \text{Svar}(V^-)$ (so $\beta : V^- \rightarrow V^+$) such that*

$$\alpha \circ \beta \circ \alpha = \alpha.$$

(This implies that α is regular in the Jordan pair $(\text{Hom}(W^+, W^-), \text{Hom}(W^-, W^+))$, see [Lo75].) The element β will then be called a pseudo-inverse of α .

In the situation of the definition, Lemma 1.7 implies that

$$\alpha : (V^+, T_{\alpha\beta\alpha}) = (V^+, T_\alpha) \rightarrow (V^-, T_\beta)$$

is a homomorphism of Jordan triple systems (and hence also a homomorphism of LTS from R_α to R_β).

Example 2.7 *Assume (V, T) is a JTS, hence $V = V^+ = V^-$. If α is a tripotent ($\alpha^3 = \alpha$) element of $\text{Svar}(V)$, then it is regular (take $\beta = \alpha$), and hence $\alpha : T_\alpha \rightarrow T_\alpha$ is a JTS-homomorphism. This holds in particular if $\alpha^2 = \alpha$ or $(-\alpha)^2 = -\alpha$. However, not every regular element is tripotent in this sense.*

If the set of pseudo-inverses of α is not empty, then it is the intersection of $\text{Svar}(V^-)$ with an affine space: the difference $\gamma = \beta - \beta'$ of two pseudo-inverses satisfies the linear equation $\alpha\gamma\alpha = 0$. There is no canonical choice of pseudo-inverse, but once we have fixed a choice, we may “improve” it in a canonical way: by straightforward computation one proves the following

Lemma 2.8 *If β is a pseudo-inverse of α , then the endomorphism $\beta' := \beta\alpha\beta : V^- \rightarrow V^+$ belongs again to $\text{Svar}(V^-)$, is again a pseudo-inverse of α , and it satisfies moreover*

$$\beta' \circ \alpha \circ \beta' = \beta'.$$

Definition 2.9 *Let (W^+, W^-) be a pair of \mathbb{K} -modules and $\alpha : W^+ \rightarrow W^-$, $\beta : W^- \rightarrow W^+$ linear maps. We say that a pair $(\alpha, \beta) \in \text{Hom}(W^+, W^-) \times \text{Hom}(W^-, W^+)$ is idempotent if $\beta \circ \alpha \circ \beta = \beta$ and $\alpha \circ \beta \circ \alpha = \alpha$. (Equivalently, (α, β) is an idempotent in the Jordan pair $(\text{Hom}(W^+, W^-), \text{Hom}(W^-, W^+))$, see [Lo75].)*

Lemma 2.10 *Assume the pair (α, β) is idempotent. Then we have direct sum decompositions*

$$W^+ = \text{Im} \beta \oplus \ker \alpha, \quad W^- = \text{Im} \alpha \oplus \ker \beta, \quad (5)$$

and $\beta|_{\text{Im} \alpha} : \text{Im} \alpha \rightarrow \text{Im} \beta$ is a linear isomorphism with inverse $\alpha|_{\text{Im} \beta} : \text{Im} \beta \rightarrow \text{Im} \alpha$. Conversely, if a pair (α, β) has these properties, then it is idempotent.

Proof. Direct check (cf. [BeL08], Lemma A.8; see also <http://fr.wikipedia.org/wiki/Pseudo-inverse>). ■

Theorem 2.11 *If $\alpha \in \text{Svar}(V^+)$ is regular, then $\text{Im} \alpha$ carries a JTS structure such that*

$$0 \rightarrow \ker \alpha \rightarrow (V^+, T_\alpha) \rightarrow \text{Im} \alpha \rightarrow 0$$

is a split exact sequence of JTS. A similar statement holds for the corresponding LTS.

Proof. Choose β such that (α, β) is idempotent. According to the preceding lemma, α and β induce mutually inverse isomorphisms between $I = \text{Im} \alpha$ and $J := \text{Im} \beta$. Therefore I becomes a JTS with triple product $(u, v, w) \mapsto T^-(u, \beta v, w)$, and $\alpha : (V^+, T_\alpha) \rightarrow I$ becomes a homomorphism which splits via $\beta : I \rightarrow V^+$. ■

By general functoriality of the construction of symmetric spaces associated to Jordan structures ([Be02]), we deduce that the exact sequence from the theorem lifts to the space level: α induces a homomorphism from M_α to the symmetric space B_α belonging to the base $\text{Im}\alpha$; we have a fibration of symmetric spaces

$$F_\alpha \rightarrow M_\alpha \rightarrow B_\alpha,$$

where the fiber F_α is a flat symmetric space, and there exists a splitting $B_\alpha \rightarrow M_\alpha$. The LTS (V^+, R_α) of M_α splits as $V^+ = K \oplus J$ with abelian ideal K and $J := \text{Im}\beta$. Moreover, we have

Lemma 2.12 *In the LTS (V^+, R_α) with ideal $K = \ker \alpha$ we have $R_\alpha(K, K)V^+ = 0$.*

Proof. $T_\alpha(u, v, w) - T_\alpha(v, u, w) = T(u, \alpha v, w) - T(v, \alpha u, w) = 0$ whenever $u, v \in K = \ker \alpha$. ■

However, $R_\alpha(K, J)K$ is in general not zero, and hence the fibration $M_\alpha \rightarrow B_\alpha$ is in general not a *symmetric bundle* in the sense of [BeD09].

2.3 Complementation of inner ideals

Recall from [LoN94] that an inner ideal $J \subset V^+$ is called a *complement* of an inner ideal $I \subset V^-$ if

$$V^- = I \oplus \text{Kern}J, \quad V^+ = J \oplus \text{Kern}I,$$

and that Jordan pair (V^+, V^-) is *complemented* if every inner ideal both in V^+ and V^- admits a complement. Complemented Jordan pairs have been characterized in [LoN94]; we will use their result in the proof of the following theorem, but we will give also a more elementary, independent proof for the positive real case in the next subsection.

Theorem 2.13 *Assume (V^+, V^-) is a non-degenerate Jordan pair that admits an anisotropic involution τ (this means that $T^+(x, \tau(x), x) = 0$ implies $x = 0$; one says also that the JTS (V, T) with $T(x, y, z) = T^+(x, \tau(y), z)$ is anisotropic).*

- (1) *If $\alpha \in \text{Svar}(V^+)$ is regular, then the inner ideal $I := \text{Im}\alpha$ is complemented. A complement is given by $J := \text{Im}\beta \subset V^+$, where $\beta \in \text{Svar}(V^-)$ is such that (α, β) is idempotent.*
- (2) *Assume that (V^+, V^-) is simple and finite-dimensional over a field \mathbb{K} . Then every inner ideal $I \subset V^-$ is complemented, and there exists regular $\alpha \in \text{Svar}(V^+)$ such that $I = \text{Im}\alpha$. Moreover, every element of the structure variety is regular, and after fixing an anisotropic involution, all elements of $\text{Svar}(V^+)$ having same image and kernel as α are naturally parametrized by $\text{Svar}(I)^\times$.*

All assumptions hold, in particular, if (V^+, V^-) is a simple, finite-dimensional Jordan pair over $\mathbb{K} = \mathbb{R}$; we can then choose for τ its Cartan-involution, i.e., the involution coming from a Cartan-involution of the corresponding 3-graded Lie algebra reversing the grading.

Proof. (1) From Lemma 2.10 we get the decomposition

$$V^+ = \text{Im}\beta \oplus \ker \alpha, \quad V^- = \text{Im}\alpha \oplus \ker \beta. \tag{6}$$

Since our assumption implies that (V^+, V^-) is non-degenerate, we have $\ker \alpha = \text{Kern}(\text{Im}\alpha)$ (cf. Example 2.5) and $\ker \beta = \text{Kern}(\text{Im}\beta)$. This shows that $\text{Im}\beta$ is a complement of $\text{Im}\alpha$.

(2) We fix for the moment an anisotropic involution $\tau : V^+ \rightarrow V^-$, giving rise to the JTS (V, T) . By definition, this JTS is then anisotropic, and it is Artinian (i.e., descending chains of inner ideals become stationary) since it is finite-dimensional. Thus [LoN94], Theorem 6.7 implies that I is complemented in the Jordan triple sense, that is, $J := \tau(I) \subset V^+$ is a complement of $I \subset V^-$. Let $\pi^\pm : V^\pm \rightarrow V^\pm$ be the projections onto I , resp. J with kernel $\text{Kern}J$, resp. $\text{Kern}I$; then $\tau \circ \pi^+ = \pi^- \circ \tau$. As shown in [LoN94], Section 3.5, the pair of projections (π^+, π^-) is a structural transformation, and hence also the pair

$$(\tau, \tau)(\pi^+, \pi^-) = (\tau \circ \pi^+, \pi^- \circ \tau) = (\tau \circ \pi^+, \tau \circ \pi^+) =: (\alpha_0, \alpha_0)$$

is structural (mind the definition of composition in the structural category). Similarly, the pair (β_0, β_0) with $\beta_0 := \tau^{-1} \circ \pi^-$ is structural, whence $\alpha_0 \in \text{Svar}(V^+)$, $\beta_0 \in \text{Svar}(V^-)$. Moreover,

$$\alpha_0 \beta_0 \alpha_0 = \tau \circ \pi^+ \circ \tau^{-1} \circ \pi^- \circ \tau \circ \pi^+ = \tau \circ (\pi^+)^3 = \alpha_0$$

and similarly $\beta_0 \alpha_0 \beta_0 = \beta_0$. By definition, the image of α_0 is I and its kernel is $\text{Kern}I$, proving that every inner ideal I is the image of a regular α_0 . Still fixing the anisotropic involution, we have the following lemma describing all elements $\alpha \in \text{Svar}(V^+)$ having same image and kernel as α_0 :

Lemma 2.14 *Let I be a complemented inner ideal in a JTS V and let $\pi : V \rightarrow I$ and $\iota : I \rightarrow V$ be projection and injection corresponding to the decomposition $V = I \oplus \text{Kern}I$. Then $(\pi, \iota) : (V, V) \rightarrow (I, I)$ and $(\iota, \pi) : (I, I) \rightarrow (V, V)$ are structural transformations, and we obtain two maps*

$$\begin{aligned} \phi : \text{Svar}(I) &\rightarrow \text{Svar}(V), & \gamma &\mapsto \alpha := \iota \circ \gamma \circ \pi = \begin{pmatrix} \gamma & 0 \\ 0 & 0 \end{pmatrix} \\ \psi : \text{Svar}(V) &\rightarrow \text{Svar}(I), & \alpha &= \begin{pmatrix} \gamma & \gamma' \\ \gamma'' & \gamma''' \end{pmatrix} \mapsto \gamma = \pi \circ \alpha \circ \iota \end{aligned}$$

(matrices taken w.r.t. the decomposition $V = I \oplus \text{Kern}I$) such that $\psi \circ \phi = \text{id}_{\text{Svar}(I)}$. In particular, ϕ is injective and ψ is surjective, and the image of ϕ is precisely the set of elements $\alpha \in \text{Svar}(V)$ such that $\text{Im}\alpha \subset I$ and $\text{Kern}I \subset \ker \alpha$. In particular, the conditions $\text{Im}\alpha = I$ and $\ker \alpha = \text{Kern}I$ hold if and only if $\alpha = \phi(\gamma)$ with invertible $\gamma \in \text{Svar}(I)$.

Proof. As mentioned above, the pair of projections (π^+, π^-) is structural. Therefore $\pi T(u, \iota v, w) = T(\pi u, v, \pi w)$ for $u, w \in V$, $v \in I$, hence $(\pi, \iota) : (V, V) \rightarrow (I, I)$ is structural. For (ι, π) : let $u, w \in I$, $v = v_I + v_K$ with $v_I \in I$, $v_K \in \text{Kern}(I)$, then

$$\iota T(u, \pi v, w) = T(u, v_I, w) = T(u, v_I + v_K, w) = T(u, v, w) = T(\iota u, v, \iota w).$$

It follows that the pair

$$(\iota, \pi)(\gamma, \gamma)(\pi, \iota) = (\iota \circ \gamma \circ \pi, \iota \circ \gamma \circ \pi) = (\alpha, \alpha) : (V, V) \rightarrow (V, V)$$

is structural, and hence $\alpha \in \text{Svar}(V)$, so ϕ is well-defined. The proof showing that ψ is well-defined is similar. Using the ‘‘matrix description’’, the remaining statements follow immediately. ■

We now finish the proof of part (2) of the theorem: let $\alpha \in \text{Svar}(V)$ be arbitrary and put $I := \text{Im}\alpha$, $J := \tau(I)$. Then J is a complement of I , hence $\text{Im}\alpha = \text{Im}\alpha_0$ and $\ker \alpha = \ker \alpha_0$ with regular α_0 . The lemma shows that $\alpha_0 = \phi(\text{id}_I)$ and $\alpha = \phi(\gamma)$ for some $\gamma \in \text{Svar}(I)^\times$. Then it is easily checked that α is regular with pseudo-inverse $\beta := \phi(\gamma^{-1})$.

In order to prove the final assertion of the theorem, let us show that the Cartan-involution is anisotropic: indeed, the JTS corresponding to this involution is *positive* (cf. next subsection). Let $x \in V$ and fix a Jordan frame e_1, \dots, e_r such that $x = \sum_i r_i e_i$ with $r_i \geq 0$. Then $T(x, x, x) = \sum_i r_i^3$, hence $T(x, x, x) = 0$ implies $r_i = 0$ for all i , hence $x = 0$. ■

The preceding results lead to a method of classification of elements of the structure variety of a finite-dimensional real simple Jordan pair (V^+, V^-) in two steps: Part (2) of the theorem says that the map

$$\text{Svar}(V) \rightarrow \mathcal{X} := \{\text{inner ideals in } V^-\}, \quad \alpha \mapsto I := \text{Im}(\alpha)$$

is surjective with fiber over I being in one-to-one correspondence with $\text{Svar}(I)^\times$. Therefore $\text{Svar}(V)$ can be described by combining the classification of inner ideals I with the one of elements of $\text{Svar}(I)^\times$. Before coming to classification, we give the promised elementary proof for the real case. It relies on the fact that kernels of inner ideals can be expressed in terms of *orthogonal complements* with respect to a suitable bilinear form.

2.4 The trace form

In this section, let (V^+, V^-) be a finite-dimensional Jordan pair over a field \mathbb{K} . The *trace form* is the bilinear form defined by

$$V^+ \times V^- \rightarrow \mathbb{K}, \quad (u, v) \mapsto \langle u, v \rangle := \text{Tr}(T^+(u, v, \cdot))$$

If V is a Jordan triple system, then the same form defines a bilinear form on V , again called *trace form*. It is known and easy to show that, if this form is non-degenerate, then it is symmetric, and that it is *invariant* in the sense that the adjoint of $T(x, y, \cdot)$ is $T(x, y, \cdot)$:

$$\langle T(x, y, u), v \rangle = \langle u, T(y, x, v) \rangle. \quad (7)$$

If $\mathbb{K} = \mathbb{R}$, we say that a JTS (V, T) is *positive*, if the trace form is positive definite, and *negative* or *compact* if it is negative definite. Of course, replacing T by $-T$ makes both notions equivalent.

Theorem 2.15 *Assume (V, T) is a real, positive JTS, and denote by \perp the orthogonal complement with respect to the trace form.*

(1) *Assume I is an inner ideal in V . Then*

$$\text{Kern}I = I^\perp.$$

In particular, the inner ideal I is complemented in the Jordan triple sense: $V = I \oplus \text{Kern}I$.

(2) *Any $\alpha \in \text{Svar}(V)$ is symmetric with respect to the trace form.*

Proof. (1) We will use the following lemma

Lemma 2.16 *Let (V, T) be a positive JTS. Then every sub-JTS $F \subset V$ is again positive, and satisfies in particular $T(F, F, F) = F$.*

Before proving the lemma, let us use it to establish the theorem: for all $u, v, w \in I$ and $k \in \text{Kern}I$,

$$\langle T(u, v, w), k \rangle = \langle w, T(v, u, k) \rangle = \langle w, T(k, u, v) \rangle = \langle T(u, k, w), v \rangle = 0,$$

whence, using the lemma, $\text{Kern}I \subset T(I, I, I)^\perp = I^\perp$. Let us prove that $I^\perp \subset \text{Kern}I$, i.e., $T(I, I^\perp, I) = 0$. By definition of inner ideals, we have $T(I, I^\perp, I) \subset T(I, V, I) \subset I$. On the other hand, using again the lemma,

$$0 = \langle I, I^\perp \rangle = \langle T(I, I, I), I^\perp \rangle = \langle T(I, I^\perp, I), I \rangle,$$

showing that $T(I, I^\perp, I) \subset I^\perp$, whence $T(I, I^\perp, I) \subset (I \cap I^\perp) = 0$, and (1) is proved.

(2) If $\alpha \in \text{Svar}(V)$ is invertible, then, writing $T(u, v) := T(u, v, \cdot)$,

$$T(u, \alpha v) = \alpha \circ T(\alpha u, v) \circ \alpha^{-1},$$

and taking traces we get $\langle u, \alpha v \rangle = \langle \alpha u, v \rangle$. Now consider $\alpha \in \text{Svar}(V)$, not necessarily invertible. Using Lemma 2.14, we write $\alpha = \iota \circ \gamma \circ \pi$, where, according to Part (1), $\pi : V \rightarrow I$ is the orthogonal projection onto $I := \text{Im}(\alpha)$ with kernel $\text{Kern}I = I^\perp$. Therefore the adjoint operator of π is ι . Using this, and that $\gamma \in \text{Svar}(I)$ is invertible, whence $\gamma^* = \gamma$ by the preceding remark, it follows that

$$\alpha^* = (\iota \circ \gamma \circ \pi)^* = \pi^* \circ \gamma^* \circ \iota^* = \iota \circ \gamma \circ \pi = \alpha,$$

proving (2). It remains to prove the lemma. Recall that a real JTS is called *Euclidean* if it is endowed with an invariant Euclidean scalar product: $(T(x, y)u, v) = (u, T(y, x)v)$. By restriction of the trace form (that is, $(u, v) := \langle u, v \rangle_V$), every subsystem F in a positive JTS is Euclidean. Let us show that F is again positive, i.e., $\langle x, x \rangle_F > 0$ for all non-zero $x \in F$. To this end, using the spectral theorem in V , we decompose $x = \sum_{i=1}^k r_i e_i$ with $r_i > 0$ and orthogonal non-zero idempotents e_1, \dots, e_k which can be expressed as certain linear combinations of (odd) powers of x ; in particular they belong again to F since F is a subsystem. It follows that

$$\langle x, x \rangle_F = \sum_i r_i^2 \langle e_i, e_i \rangle_F = \sum_i r_i^2 \langle T(e_i, e_i, e_i), e_i \rangle_F$$

Now, a result due to Backes (see Theorem 1 page 268 of [Backes83])—based on an earlier one of Koecher—asserts that for all u, v in the Euclidean JTS F the inequality

$$\mathrm{Tr} S(S(u, v)u, v) \geq 0 \quad (8)$$

holds with $S(u, v) := \frac{1}{2}(T(u, v) + T(v, u)) \in \mathbf{End}(F)$. Moreover, equality holds in (8) if and only if $S(u, v) = 0$. This implies that $\langle x, x \rangle_F \geq 0$, and that $\langle x, x \rangle_F = 0$ iff $r_i^2 S(e_i, e_i) = 0$ for all i , iff $r_i^2 S(e_i, e_i)e_i = 0$ for all i , iff $r_i^2 = 0$ for all i , iff $x = 0$. Thus F is positive.

Let us show that $T(F, F, F) = F$. Decompose the positive JTS F as $F = U \oplus U^\perp$ with $U = T(F, F, F)$. Then $T(U^\perp, U^\perp, U^\perp) = 0$ (since $0 = (U, U^\perp) = (T(F, F, F), U^\perp) = (F, T(F, F, U^\perp)) \supset (F, T(U^\perp, U^\perp, U^\perp))$), hence U^\perp is a positive JTS with zero product, hence it is zero and $U = F$. ■

For semisimple Jordan *algebras*, statement (2) is essentially already contained in [Ri70], Theorem 3 (with a very different proof). In loc. cit. Corollary 3.1, it is also remarked that this result implies that, if (g, h) is a structural transformation of a real semisimple Jordan algebra, then $h = g^*$ (adjoint operator); the same remark applies in the present context.

Corollary 2.17 *Assume (V^+, V^-) is a finite-dimensional semisimple Jordan pair over \mathbb{R} . Then:*

(1) *For any an inner ideal I in V^- , its kernel $\mathrm{Kern}I \subset V^+$ is the orthogonal complement I^\perp of I with respect to the trace form, and I admits a complement.*

(2) *Any $\alpha \in \mathrm{Svar}(V^+)$ is symmetric with respect to the trace form.*

Proof. It is known that every semi-simple real finite-dimensional Jordan pair admits a positive involution θ (the Cartan-involution, see Theorem 2.13), that is, the JTS $V = V^+$ with $T(x, y, z) := T^+(x, \theta y, z)$ is positive. Applying Part (1) of the theorem to this positive JTS gives Part (1) of the corollary: indeed, both $\mathrm{Kern}I$ and I^\perp are defined independently of the involution; we have seen that under identification of V^+ and V^- under θ the corresponding spaces are equal, therefore $\mathrm{Kern}I = I^\perp$ (and this equality remains true under identification by any involution, be it positive or not). Exactly the same argument shows symmetry of α w.r.t. the trace form. ■

The properties from the corollary concern the trace form of Jordan *pairs*, whereas positivity (and anisotropy) are properties of an involution, hence of triple systems. Positive (more generally, anisotropic) forms permit to choose “simultaneously” complements for *all* inner ideals; for base fields different from \mathbb{R} , such a choice is in general not possible, even if a Jordan pair is complemented.

3 Classification

It goes without saying that in general a classification of the objects introduced in the preceding chapters is out of reach. The assumptions that we will make are fairly restrictive and are mainly imposed by geometric applications: we will look at *simple finite-dimensional Jordan pairs over $\mathbb{K} = \mathbb{R}$* . (From a purely algebraic point of view, it would certainly be interesting to classify objects over more general base fields and by relaxing finite-dimensionality to chain conditions.) Moreover, in order to keep this work in reasonable bounds, we restrict attention to *classical* Jordan pairs. In this chapter we present the (known) classifications of such Jordan pairs and triple systems and of their inner ideals, which then lead to a complete description of their structure varieties.

3.1 Classification of Jordan pairs

I. Simple complex Jordan pairs (cf. [Be00], Table XII.2.1)

label	(V^+, V^-)	T^\pm	3-graded Lie algebra
1.	$(M(p, q; \mathbb{C}), M(q, p; \mathbb{C}))$	$T^\pm(X, Y, Z) = XYZ + ZYX$	$\mathfrak{sl}(p + q; \mathbb{C})$
2.	$(\mathrm{Sym}(n; \mathbb{C}), \mathrm{Sym}(n; \mathbb{C}))$	$T^\pm(X, Y, Z) = XYZ + ZYX$	$\mathfrak{sp}(n, \mathbb{C})$
3.	$(\mathrm{Asym}(n; \mathbb{C}), \mathrm{Asym}(n; \mathbb{C}))$	$T^\pm(X, Y, Z) = XYZ + ZYX$	$\mathfrak{o}(2n, \mathbb{C})$
4.	$(\mathbb{C}^n, \mathbb{C}^n)$	$T^\pm(x, y, z) = \beta(y, z)x + \beta(y, x)z - \beta(x, z)y$	$\mathfrak{o}(\hat{\beta})$
5.	$(\mathrm{Herm}(3, \mathbb{O}_{\mathbb{C}}), \mathrm{Herm}(3, \mathbb{O}_{\mathbb{C}}))$	$T^\pm(X, Y, Z) = XYZ + ZYX$	\mathfrak{e}_7
6.	$(M(1, 2, \mathbb{O}_{\mathbb{C}}), M(1, 2, \mathbb{O}_{\mathbb{C}}))$	$T^\pm(X, Y, Z) = XYZ + ZYX$	\mathfrak{e}_6

Remarks. (1) In type 4, \mathbb{C}^n is equipped with a non-degenerate and symmetric bilinear form β , and $\hat{\beta}$ is the form on \mathbb{C}^{n+2} given by $\hat{\beta}((z_0, z, z_{n+1}), (w_0, w, w_{n+1})) = \beta(z, w) + z_0 w_{n+1} + w_0 z_{n+1}$. The product $T(x, y, z)$ may be realized in the Clifford algebra $Cl(\mathbb{C}^n, \beta)$ as follows: recall the Clifford relation $x \bullet y := \frac{1}{2}(xy + yx) = \beta(x, y)1$ and note that $xyz + zyx = x \bullet (y \bullet z) - y \bullet (x \bullet z) + (x \bullet y) \bullet z$. Using this, one gets for x, y, z belonging to the subspace \mathbb{C}^n of $Cl(\mathbb{C}^n, \beta)$:

$$xyz + zyx = x \bullet (y \bullet z) - y \bullet (x \bullet z) + (x \bullet y) \bullet z = \beta(y, z)x + \beta(y, x)z - \beta(x, z)y.$$

(2) Types 4, 5 and 6 are included here for convenience, but we will mainly consider the matrix cases 1, 2 and 3 in the sequel.

(3) The following lists will be simpler than the ones given in [Be00] in the sense that we will not have to distinguish for Type 1 the cases $p \neq q$ and $p = q$, nor for Type 3 the cases n even and n odd (which had to be done in [Be00] since only *invertible* α were considered there).

II. Simple real Jordan pairs (cf. [Be00], Table XII.2.2). There are two kinds: the complex pairs from above, considered as real, and the following:

label	(V^+, V^-)	T^\pm	3-graded Lie algebra
1.1	$(\text{Herm}(n, \mathbb{C}), \text{Herm}(n, \mathbb{C}))$	$T^\pm(X, Y, Z) = XYZ + ZYX$	$\mathfrak{su}(n, n)$
1.2	$(M(p, q; \mathbb{R}), M(q, p; \mathbb{R}))$	$T^\pm(X, Y, Z) = XYZ + ZYX$	$\mathfrak{sl}(p+q; \mathbb{R})$
1.3	$(M(p, q; \mathbb{H}), M(q, p; \mathbb{H}))$	$T^\pm(X, Y, Z) = XYZ + ZYX$	$\mathfrak{sl}(p+q; \mathbb{H})$
2.1	$(\text{Sym}(n; \mathbb{R}), \text{Sym}(n; \mathbb{R}))$	$T^\pm(X, Y, Z) = XYZ + ZYX$	$\mathfrak{sp}(n, \mathbb{R})$
2.2	$(\text{Herm}(n; \mathbb{H}), \text{Herm}(n; \mathbb{H}))$	$T^\pm(X, Y, Z) = XYZ + ZYX$	$\mathfrak{so}^*(2n)$
3.1	$(\text{Herm}(n; \mathbb{H}), \text{Herm}(n; \mathbb{H}))$	$T^\pm(X, Y, Z) = XYZ + ZYX$	$\mathfrak{sp}(n, n)$
3.2	$(\text{Asym}(n; \mathbb{R}), \text{Asym}(n; \mathbb{R}))$	$T^\pm(X, Y, Z) = XYZ + ZYX$	$\mathfrak{o}(n, n)$
4.	$(\mathbb{R}^{p,q}, \mathbb{R}^{p,q})$ ($p+q = n$)	$T^\pm(x, y, z) = \beta(y, z)x + \beta(y, x)z - \beta(x, z)y$	$\mathfrak{o}(\hat{\beta})$

Remarks. (1) The pair with label $i.j$ is a real form of the complex pair with label i . from the preceding table (with certain restriction of parameters: for instance, 1.1 is a real form in the square case, i.e., $p = q = n$, and quaternionic matrices are real forms for even parameters). For type 4., (p, q) denotes the signature of the form β . Note that $\mathbb{R}^{p,q}$ and $\mathbb{R}^{p',q'}$ are not isomorphic if p is different from p' and q' .

(2) Exceptional Jordan pairs and triple systems have been classified by E. Neher.

3.2 Classification of inner ideals

For the classical Jordan pairs listed above, we describe their inner ideals. This classification is known: for Jordan algebras see [McC71]; for Jordan pairs see [Ne96] or [DFGG08]; for an elementary proof in case of rectangular or symmetric matrices, see [BeL08], Appendix A.

1. Rectangular matrices. $(V^+, V^-) = (M(p, q; \mathbb{K}), M(q, p; \mathbb{K})) = (\text{Hom}(E, F), \text{Hom}(F, E))$ over a field \mathbb{K} : all inner ideals in V^- are of the form

$$I = I_{a,b} := \{f : F \rightarrow E \mid a \subset \ker f, \text{Im} f \subset b\}$$

for linear subspaces $a \subset F, b \subset E$. Thus I can be identified with a matrix space $\text{Hom}(a', b)$ where a' is some vector space complement of a in F . This ideal is *principal* if and only if it is isomorphic to a space of *square* matrices, i.e., if $\dim b = \dim a'$. In the general case, elementary linear algebra shows that the Kernel of $I_{a,b}$ is

$$\text{Kern} I_{a,b} = \{h : E \rightarrow F \mid h(b) \subset a\},$$

and $J := I_{b',a'} \subset \text{Hom}(E, F)$ is a complement of $I_{a,b}$ if a', b' are vector space complements of a , resp. of b .

2. Symmetric and Hermitian matrices. $(V^+, V^-) = (V, V), V = \text{Sym}(n, \mathbb{K})$ or $\text{Herm}(n, \mathbb{F})$. Inner ideals are constructed as above, by taking $E = F = \mathbb{K}^n$ or \mathbb{F}^n and for f self-adjoint maps (w.r.t. standard bilinear or Hermitian forms). Then the condition $\text{Im} f \subset b$ is equivalent to $b^\perp \subset \ker f$, i.e., inner ideals are given as above with $a = b^\perp$:

$$I = I_b := \{f : E \rightarrow E \mid f = f^*, \text{Im} f \subset b\}.$$

The inner ideal I_b is always principal, and it can be identified with the matrix space $\text{Herm}(b)$. The kernel $\text{Kern}I_b$ is characterized by the property $h(b) \subset b^\perp$, and under the identification $V^+ = V^-$, a complement of I is given by I itself.

3. Skew-symmetric matrices. $(V^+, V^-) = (V, V)$, $V = \text{Asym}(n, \mathbb{K})$. There are two different kinds of inner ideals: The first kind is constructed exactly as in the preceding case:

$$I = I_b := \{f : E \rightarrow E \mid f = -f^*, \text{Im}f \subset b\}$$

where $b^\perp \subset \ker f$ follows automatically. Kernel and complement are described as above. The second type (called *point space* in [McC71] and [DFGG08]) is fairly special: fix a non-zero vector $u \in E = \mathbb{K}^n$ and let

$$K_u := \{f : E \rightarrow E \mid f = -f^*, f(u^\perp) \subset \mathbb{K}u\}$$

(where \perp refers to the standard scalar product on \mathbb{K}^n). In matrix realization, taking u as the last base vector e_n and $u^\perp = \text{vect}(e_1, \dots, e_{n-1})$, these are skew matrices having zero upper left $(n-1) \times (n-1)$ -block. We check that K_u is an inner ideal: the key observation is that, for any skew-symmetric $g : E \rightarrow E$, we have $\langle gu, u \rangle = -\langle u, gu \rangle = -\langle gu, u \rangle$, whence $g(\mathbb{K}u) \subset u^\perp$; therefore, for any $f \in K_u$,

$$(fgf)(u^\perp) \subset fg(\mathbb{K}u) \subset f(u^\perp) \subset \mathbb{K}u,$$

whence $fgf \in K_u$. Observe also that $K_u = \text{Kern}I_{u^\perp}$, that is, in this special situation the kernel of I_{u^\perp} is an inner ideal (namely K_u). As above, K_b is complemented, and hence has a natural JTS-structure. Let us describe this structure: realize elements of K_b by matrices as mentioned above; then

$$\begin{pmatrix} 0 & X^t \\ -X & 0 \end{pmatrix} \begin{pmatrix} 0 & Y^t \\ -Y & 0 \end{pmatrix} \begin{pmatrix} 0 & X^t \\ -X & 0 \end{pmatrix} = \begin{pmatrix} 0 & \langle X, Y \rangle X^t \\ -\langle X, Y \rangle X & 0 \end{pmatrix}$$

for row vectors $X, Y \in \mathbb{K}^{n-1}$. This is exactly the JTS-structure of the “projective” JTS $M(1, n-1; \mathbb{K})$. It follows also that fgf is proportional to f for all $f \in K_u$ and $g \in \text{Asym}(n, \mathbb{K})$ (proof: $fgf = \langle f, g \rangle f$ if $g \in K_u$, and $fgf = 0$ if $g \in \text{Kern}(K_u)$). Thus $\mathbb{K}f$ is an inner ideal for all $f \in K_u$ (this property is used to define *point spaces* in [McC71]). Finally, the intersection of a point space K_u with a standard inner ideal I_b is again a point space inner ideal; thus the inner ideals K_u can be characterized as maximal point spaces.

4. Type four. $V^+ = V^- = \mathbb{K}^{p,q}$. The inner ideals are precisely the isotropic subspaces I for the form β . We have $\text{Kern}I = I^\perp$, and an isotropic subspace J is complementary to I as inner ideal iff $\mathbb{K}^n = I \oplus J^\perp = J \oplus I^\perp$. As JTS, the *proper* complementary inner ideal pairs (I, J) are isomorphic to the “projective” JTS $M(1, r; \mathbb{K})$ with $r = \dim I \leq n/2$ ([LoN94], Example 1.14).

Note that in all cases, with the exception of proper inner ideals for type 4 and point spaces for type 3, inner ideals are of the same type as the ambient Jordan pair; in the two exceptional situations, inner ideals are always isomorphic to $M(1, r; \mathbb{K})$. Finally, in low dimensions there are certain isomorphisms between various types, see Subsection 3.8 (in particular, the isomorphism $\text{Asym}(4, \mathbb{R}) \cong \mathbb{R}^{3,3}$ identifies the standard ideals in $\text{Asym}(4, \mathbb{R})$ with the isotropic subspaces of dimension 1 and 3 in $\mathbb{R}^{3,3}$ and the maximal point spaces with the isotropic subspaces of dimension 2).

3.3 Classification of isotopes: invertible elements in structure varieties

The classification of the *invertible* elements in the structure varieties of simple finite dimensional real Jordan pairs is known: it is based on the complete descriptions of their structure groups given by E. Neher, see [Be00], Chapter XII for tables concerning non-exceptional spaces. The lists are fairly long and we will not reproduce them here. Let us just give some comments. The tables given in loc. cit. (especially Table XII.2.6) have the same form as the ones to be given in the next subsection; however, they appear to be longer due to the fact that the description of *invertible* homotopies is less uniform: as a rule, instead of parametrizing by general rectangular matrices A, B as below (e.g., case 1.c), one has to take invertible matrices, which forces to distinguish the cases $p = q$ and $p \neq q$, and instead of parametrizing by general symmetric, Hermitian or skew-symmetric matrices (e.g., case 1.c), one uses invertible such matrices in normal form, involving signatures in the Hermitian case and forcing to distinguish the cases n odd and n even in the skew case. In this sense the tables from loc. cit. form a proper subset of the tables given below.

3.4 Structure varieties of simple classical Jordan pairs

Theorem 3.1 *For each of the classical Jordan pairs (V^+, V^-) of matrix type, the following lists of endomorphisms α give complete parametrizations of the structure variety $\text{Svar}(V^+)$. Moreover, these parametrizations are equivariant with respect to the natural structure group actions; therefore a classification of homotopes T_α up to isomorphism can be deduced by considering structure group orbits in the respective parameter spaces.*

1. Jordan pairs of rectangular matrices

1.1 $V^+ = M(p, q; \mathbb{K})$, $\mathbb{K} = \mathbb{R}, \mathbb{C}$:

label	α	parameter set
1.a	$\alpha(X) = AXA$	$A \in M(q, p; \mathbb{K})$
1.a'	$\alpha(X) = -AXA$	$A \in M(q, p; \mathbb{K})$
1.b	$\alpha(X) = AX^t B$	$A \in \text{Sym}(q, \mathbb{K}), B \in \text{Sym}(p, \mathbb{K})$
1.c	$\alpha(X) = AX^t B$	$A \in \text{Asym}(q, \mathbb{K}), B \in \text{Asym}(p, \mathbb{K})$

1.2, case of antilinear maps: $V^+ = M(p, q; \mathbb{C})$

label	α	parameter set
1.A	$\alpha(X) = A\bar{X}A$	$A \in M(q, p; \mathbb{C})$
1.A'	$\alpha(X) = -A\bar{X}A$	$A \in M(q, p; \mathbb{C})$
1.B	$\alpha(X) = A\bar{X}^t B$	$A \in \text{Herm}(q, \mathbb{C}), B \in \text{Herm}(p, \mathbb{C})$

1.3 $V^+ = M(p, q; \mathbb{H})$:

label	α	parameter set
1.3.a	$\alpha(X) = AXA$	$A \in M(q, p; \mathbb{H})$
1.3.a'	$\alpha(X) = -AXA$	$A \in M(q, p; \mathbb{H})$
1.3.b	$\alpha(X) = A\bar{X}^t B$	$A \in \text{Herm}(q, \mathbb{H}), B \in \text{Herm}(p, \mathbb{H})$
1.3.c	$\alpha(X) = A\tilde{X}^t B$	$A \in \text{Herm}(q, \tilde{\mathbb{H}}), B \in \text{Herm}(p, \tilde{\mathbb{H}})$

2. Jordan pairs of symmetric matrices

2.1, case of linear maps: $V^+ = V^- = \text{Sym}(n, \mathbb{K})$, $\mathbb{K} = \mathbb{R}, \mathbb{C}$:

label	α	parameter set
2.a	$\alpha(X) = AXA$	$A \in \text{Sym}(n, \mathbb{K})$
2.a'	$\alpha(X) = -AXA$	$A \in \text{Sym}(n, \mathbb{K})$
2.b	$\alpha(X) = AXA$	$A \in \text{Asym}(n, \mathbb{K})$
2.b'	$\alpha(X) = -AXA$	$A \in \text{Asym}(n, \mathbb{K})$

2.2, case of antilinear maps: $V^+ = V^- = \text{Sym}(n, \mathbb{C})$

label	α	parameter set
2.A	$\alpha(X) = A\bar{X}A$	$A \in \text{Herm}(n, \mathbb{C})$
2.A'	$\alpha(X) = -A\bar{X}A$	$A \in \text{Herm}(n, \mathbb{C})$

3. Jordan pairs of Skewsymmetric matrices

3.1, case of linear maps: $V^+ = V^- = \text{Asym}(n, \mathbb{K})$, $\mathbb{K} = \mathbb{R}, \mathbb{C}$:

label	α	parameter set
3.a	$\alpha(X) = AXA$	$A \in \text{Asym}(n, \mathbb{K})$
3.a'	$\alpha(X) = -AXA$	$A \in \text{Asym}(n, \mathbb{K})$
3.b	$\alpha(X) = AXA$	$A \in \text{Sym}(n, \mathbb{K})$
3.b'	$\alpha(X) = -AXA$	$A \in \text{Sym}(n, \mathbb{K})$
3.c	$\alpha(X) = u \otimes u^* \circ X \circ A + A \circ X \circ u \otimes u^*$	$A \in \text{Sym}(n, \mathbb{K}), u \in \mathbb{K}^n$
3.c'	$\alpha(X) = -u \otimes u^* \circ X \circ A - A \circ X \circ u \otimes u^*$	$A \in \text{Sym}(n, \mathbb{K}), u \in \mathbb{K}^n$

3.2, case of antilinear maps: $V^+ = V^- = \text{Asym}(n, \mathbb{C})$

label	α	parameter set
3.A	$\alpha(X) = A\bar{X}A$	$A \in i\text{Herm}(n, \mathbb{C})$
3.A'	$\alpha(X) = -A\bar{X}A$	$A \in i\text{Herm}(n, \mathbb{C})$
3.B	$\alpha(X) = u \otimes u^* \circ \bar{X} \circ A + A \circ \bar{X} \circ u \otimes u^*$	$A \in \text{Herm}(n, \mathbb{C}), u \in \mathbb{C}^n$
3.B'	$\alpha(X) = -u \otimes u^* \circ \bar{X} \circ A - A \circ \bar{X} \circ u \otimes u^*$	$A \in \text{Herm}(n, \mathbb{C}), u \in \mathbb{C}^n$

4. Jordan pairs of Hermitian matrices

4.1 $V^+ = V^- = \text{Herm}(n, \mathbb{C})$:

label	α	parameter set
4.1.a	$\alpha(X) = AXA$	$A \in \text{Herm}(n, \mathbb{C})$
4.1.a'	$\alpha(X) = -AXA$	$A \in \text{Herm}(n, \mathbb{C})$
4.1.b	$\alpha(X) = A\bar{X}A^t$	$A \in \text{Sym}(n, \mathbb{C})$
4.1.b'	$\alpha(X) = -A\bar{X}A^t$	$A \in \text{Sym}(n, \mathbb{C})$
4.1.c	$\alpha(X) = A\bar{X}A^t$	$A \in \text{Asym}(n, \mathbb{C})$
4.1.c'	$\alpha(X) = -A\bar{X}A^t$	$A \in \text{Asym}(n, \mathbb{C})$

4.2 $V^+ = V^- = \text{Herm}(n, \mathbb{H})$:

label	α	parameter set
4.2.a	$\alpha(X) = AXA$	$A \in \text{Herm}(n, \mathbb{H})$
4.2.a'	$\alpha(X) = -AXA$	$A \in \text{Herm}(n, \mathbb{H})$
4.2.b	$\alpha(X) = A\bar{X}A$	$A \in \text{Herm}(n, \mathbb{H})$
4.2.b'	$\alpha(X) = -A\bar{X}A$	$A \in \text{Herm}(n, \mathbb{H})$

4.3 $V^+ = V^- = \text{Herm}(n, \tilde{\mathbb{H}})$:

label	α	parameter set
4.3.a	$\alpha(X) = AXA$	$A \in \text{Herm}(n, \tilde{\mathbb{H}})$
4.3.a'	$\alpha(X) = -AXA$	$A \in \text{Herm}(n, \tilde{\mathbb{H}})$
4.3.b	$\alpha(X) = A\tilde{X}A$	$A \in \text{Herm}(n, \tilde{\mathbb{H}})$
4.3.b'	$\alpha(X) = -A\tilde{X}A$	$A \in \text{Herm}(n, \tilde{\mathbb{H}})$

Remark 3.2 *The theorem says that the given parametrizations are surjective; we do not claim that they are injective: for instance, for α of type 1.b the pairs of matrices $(A, \lambda B)$ and $(\lambda A, B)$ for $\lambda \in \mathbb{K}$ yield the same α . Moreover, if $p = 1$ (case of “projective” JTS $M(1, q; \mathbb{K})$), Type 1.c becomes trivial (since $\text{Asym}(1, \mathbb{K}) = 0$), and Type 1.a (the map $\alpha(X) = \langle X, A \rangle A$, i.e., $\alpha = A^* \otimes A$) becomes a special case of Type 1.b: all structural maps are given by $\alpha(X) = BX^t$ with symmetric B , that is, in the “projective case”, structural maps are precisely the self-adjoint operators on \mathbb{K}^q .*

Proof of the theorem. We show first that the parametrizations are well-defined, i.e., the endomorphisms α belong indeed to $\text{Svar}(V^+)$. In all cases, this is checked by elementary calculations. For convenience, let us spell out this, e.g., for the case 1.b: let $X, Y, Z \in M(p, q; \mathbb{K})$, and $A \in \text{Sym}(q, \mathbb{K}), B \in \text{Sym}(p, \mathbb{K})$, then

$$\begin{aligned}
\alpha T^+(X, \alpha Y, Z) &= A(XAY^t BZ + ZAY^t BX)^t B \\
&= (AX^t B)Y(AZ^t B) + (AZ^t B)Y(AX^t B) \\
&= T^-(\alpha X, Y, \alpha Z).
\end{aligned}$$

For $A \in \text{Asym}(q, \mathbb{K}), B \in \text{Asym}(p, \mathbb{K})$ the same calculation applies. Equivalently, one could check that for any $A \in M(q, q; \mathbb{K})$ and $B \in M(p, p; \mathbb{K})$ the pair of linear maps

$$g : V^+ \rightarrow V^-, X \mapsto AX^t B, \quad h : V^+ \rightarrow V^-, X \mapsto A^t X^t B^t$$

is a structural transformation from (V^+, V^-) to its opposite pair (V^-, V^+) , and the self-adjointness condition $g = h$ holds if A and B are both symmetric or both skew-symmetric. Similarly, for Jordan pairs of symmetric matrices, it is useful to observe that, for any square matrix A , the map $X \mapsto AXA^t$ is a well-defined

endomorphism of $\text{Sym}(n; \mathbb{K})$ which together with $X \mapsto A^t X A$ forms a structural pair. The self-adjointness condition is hence satisfied if A is symmetric or skew-symmetric. Moreover, if A is any invertible matrix, the same formula defines an element of the structure group. For Hermitian matrices, similar remarks hold with respect to maps of the form $X \mapsto AX\bar{A}^t$. Finally, cases 3.c, 3.c', 3.B, 3.B' have a different behaviour and will be discussed separately below.

Next let us show that the parametrizations are surjective. First of all, we have to show that all inner ideals I show indeed up in the form $I = \text{Im}\alpha$ for suitable α . For the *principal ideals* this is immediate: by definition, they are images of quadratic operators $Q(A) : X \mapsto AXA$, which appear in the list. Since for Jordan algebras of symmetric or Hermitian matrices all inner ideals are principal, this proves our assertion in these cases. For rectangular matrices, the inner ideal $I_{ab} = \{f : F \rightarrow E \mid a \subset \ker f, \text{Im} f \subset b\}$ defined above is the image of the map $\alpha : X \mapsto AX^t B$ where A and B are the symmetric matrices describing orthogonal projection (w.r.t. the standard scalar product on \mathbb{K}^p and \mathbb{K}^q) onto b , resp. onto a^\perp , hence all inner ideals I in $M(p, q; \mathbb{K})$ are of the form $\text{Im}\alpha$ for some α given in the list. Finally, the principal inner ideals in $\text{Asym}(n, \mathbb{K})$ show up in the cases 3.a in the same way as for $\text{Sym}(n, \mathbb{K})$, whereas the point spaces show up in case 3.c which we shall discuss below separately.

Next, for a given inner ideal I we have to show that $\text{Svar}(I)^\times$ parametrizes in the way described in Lemma 2.14 the set of all α with $\text{Im}\alpha = I$ and $\ker \alpha = \ker(I)$. This amounts to compare, for fixed I , the present list with Table XII.2.6 from [Be00]. Assume first $I = I_{ab}$ is a ‘‘rectangular inner ideal’’ in $M(p, q; \mathbb{K})$. Then we may modify the matrices A and B defined above by a signature (case 1.b), or (if their size is even) by standard skew matrices (case 1.c); this corresponds to the invertible homotopies of I from Table XII.2.6, loc. cit., lines 6.1.a, 6.1.b. If, moreover, I is ‘‘square’’ (i.e., principal), then it is also an image of homotopies from case 1.a and 1.a', which correspond to line 1.1.b in loc. cit. For $\mathbb{K} = \mathbb{C}$, complex conjugation comes in as additional invertible homotopy (case 1.A above; Table XII.2.4 line 1.A in loc. cit.). Summing up, all invertible homotopies of I correspond to some α from the above list. Exactly the same pattern applies to all other cases: the invertible homotopies of I correspond in general to modify by signatures, or by standard skew matrices in even dimension, plus adding possibly a complex conjugation. As explained in section 3.3, this corresponds exactly to the classification pattern from [Be00], Chapter XII.

Let us explain now that the remaining cases 3.c, 3.c', 3.B, 3.B' describe homotopies corresponding to point-space inner ideals in $\text{Asym}(n, \mathbb{K})$. Let $\alpha(X) = u \otimes u^* \circ X \circ A + A \circ X \circ u \otimes u^*$.

Claim 1: α is a homotopy. More generally, let $\tilde{\alpha}(X) := AXB + BXA$ for $A, B \in \text{Sym}(n, \mathbb{K})$. In general, $\tilde{\alpha}$ will not be a homotopy, but we may ask under which conditions on A and B this is the case. Computing $\tilde{\alpha}(X)Y\tilde{\alpha}(X)$ and $\tilde{\alpha}(X\tilde{\alpha}(Y)X)$, we find that equality between these two terms holds if and only if

$$AXAYBXB + BXB YAXA = AXBYBXA + BXYAXB.$$

If B is a symmetric *rank-one* operator, i.e., an operator $u \otimes u^* : v \mapsto \langle u, v \rangle u$, then this equality holds for all $X, Y \in \text{Asym}(n, \mathbb{K})$: indeed, for all $Z \in \text{Asym}(n, \mathbb{K})$, the matrix BZB is then skew-symmetric of rank one, hence must be zero, so $BXB = 0$, $B Y B = 0$, $B(XAYAX)B = 0$. This proves Claim 1.

Claim 2. *The image of α is a point space.* Consider first the case $u = e_n$ and A the diagonal matrix having last diagonal coefficient 0 and the others 1. Then $u \otimes u^* = E_{nn}$, so that $A + u \otimes u^* = 1$ is the identity matrix. Using that $E_{nn} X E_{nn} = 0$ for skew-symmetric X , it follows that

$$\alpha(X) = E_{nn} X A + A X E_{nn} = X - AXA = (\text{id} - \gamma)(X)$$

where $\gamma(X) = AXA$ is the projection onto the principal inner ideal $I_{e_n^\perp}$ (case 3.b) with kernel K_{e_n} , hence $\alpha = \text{id} - \gamma$ is the projection onto the maximal point space K_{e_n} . The general proof of claim 2 (not necessarily maximal point spaces) is similar.

Finally, knowing that the structure variety of K_{e_n} is the one of the projective Jordan pair, that is $\text{Sym}(n-1, \mathbb{K})$ (Remark 3.2), augmented by complex conjugation in the complex case, we see that all α having point spaces as images are covered by cases 3.c, 3.c', 3.B, 3.B'. \blacksquare

Corollary 3.3 *Theorem 4.2 of Part I ([BeBi]) contains a complete classification of homotopes of classical symmetric spaces, in the sense that the symmetric spaces described there exhaust the list of symmetric spaces having curvature R_α , with α a homotopy of one of the classical Jordan pairs of type 1,2 and 3; the only*

exception is that in loc. cit. the symmetric spaces corresponding to the point space inner ideals for type 3 do not appear.

Proof. Labels in Theorem 4.2 loc. cit. are chosen such that curvature tensors $[X, Y, Z]_\alpha$ given there correspond exactly to the LTS R_α corresponding to the α -homotope of the Jordan pair (V^+, V^-) . Thus Part I furnishes a method to calculate the standard imbeddings of these LTS, and hence the symmetric spaces in their form G_α/H_α . \blacksquare

It seems likely that the symmetric spaces belonging to the point space inner ideals cannot be constructed via the two-involution construction from Part I, and moreover, that they are related to the type four series in a profound way. A better understanding of this situation is certainly a topic worth further work.

3.5 Homotopes of polarized Jordan triple systems

Example 3.4 *Let us start with a naive (and important) example: consider the direct product of the JTS (\mathbb{K}, T) with itself, where $T(x, y, z) = 2xyz$, whence $Q(x)y = x^2y$. We wish to compute its structure variety $\text{Svar}(\mathbb{K} \times \mathbb{K})$: write $\alpha : \mathbb{K}^2 \rightarrow \mathbb{K}^2$ as a 2×2 -matrix; then $Q(x, y)$ is the diagonal matrix having coefficients x^2, y^2 , so that the condition $\alpha Q(X)\alpha = Q(\alpha X)$ is equivalent to*

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} x^2 & 0 \\ 0 & y^2 \end{pmatrix} \begin{pmatrix} a & b \\ c & d \end{pmatrix} = \begin{pmatrix} (ax + by)^2 & 0 \\ 0 & (cx + dy)^2 \end{pmatrix}$$

Taking for X the two base vectors e_1, e_2 , and assuming that \mathbb{K} is a field, a short computation leads to distinguish two cases, apart from the trivial case $\alpha = 0$:

1. $a \neq 0$; then $b = c = 0$, and α is a diagonal matrix with $d \in \mathbb{K}$ arbitrary,
2. $a = 0$ and $b \neq 0$; then $d = 0$ and $c = b$, hence α is a multiple of the exchange map $(x, y) \mapsto (y, x)$.

In the first case, homotopes are simply direct products of those in \mathbb{K} ; in the second case, homotopes are polarized Jordan triple systems, corresponding to symmetric spaces of the kind $\text{GL}(2, \mathbb{K})/\text{GL}(1, \mathbb{K}) \times \text{GL}(1, \mathbb{K})$ (for $\mathbb{K} = \mathbb{R}$, this is the one-sheeted hyperboloid, see section 3.8).

The general situation is similar as in the example: for any Jordan pair (V^+, V^-) , the direct sum with its opposite Jordan pair $(V^+ \oplus V^-, V^- \oplus V^+)$ is again a Jordan pair, and this pair carries a canonical involution (the exchange map), so that it becomes a Jordan triple system $V = V^+ \oplus V^-$ with

$$T((x, x'), (y, y'), (z, z')) = (T^+(x, y', z), T^-(x', y, z')).$$

Such JTS are called *polarized*; in the framework of 3-graded Lie-algebras, they correspond to the direct sum $\mathfrak{g} \oplus \mathfrak{g}$ with the exchange involution (where the second copy of \mathfrak{g} carries the opposite grading of the first one). Every structural transformation $(f : V^+ \rightarrow V^-, g : V^- \rightarrow V^+)$ from (V^+, V^-) to itself gives rise to a homotopy of the polarized JTS V via

$$\alpha : V^+ \oplus V^- \rightarrow V^+ \oplus V^-, \quad (u, v) \mapsto (g(v), f(u)).$$

Seen this way, the classification of all structural endomorphisms of a Jordan pair is a subproblem of the one of classifying homotopies. Put differently, classifying structural transformations from Jordan pairs to themselves is a subproblem of classifying the self-adjoint ones from Jordan pairs to their opposite pairs. If the Jordan pair in question admits an involution, then both problems are equivalent (since the involution can be used to reduce one to the other). Note that the classification of invertible such transformations (the structure group) is known, again by work of E. Neher. We conjecture that the following table gives indeed a complete description, but we will not go here into details of the proof.

Table of structural endomorphisms (g, h) of a Jordan pair (V^+, V^-) : $(g : V^+ \rightarrow V^+, h : V^- \rightarrow V^-)$.

label	pair of linear maps (g, h)	parameter set
1.a	$g(X) = AX^tB, h(X') = A^t(X')^tB^t$	$A, B \in M(p, q; \mathbb{K})$
1.b	$g(X) = AXB, h(X') = BX'A$	$A \in M(p, p; \mathbb{K}), B \in M(q, q; \mathbb{K})$
2	$g(X) = AXA^t, h(X') = A^tX'A$	$A \in M(n, n; \mathbb{K})$
3.a	$g(X) = AXA^t, h(X') = A^tX'A$	$A \in M(n, n; \mathbb{K})$
3.b	$g(X) = AXu \otimes u^* + u \otimes u^*XA^t,$ $h(X') = A^tX'u \otimes u^* + u \otimes u^*X'A$	$A \in M(n, n; \mathbb{K}), u \in \mathbb{K}^n$

For $\mathbb{K} = \mathbb{C}$ we have, as usual, \mathbb{C} -antilinear families given by the same formulas, with a preceding complex conjugation of X and X' . Tables of structural endomorphisms of spaces of Hermitian matrices are similar as for symmetric ones (type 2), replacing the transposed matrix by the conjugate transposed matrix. The symmetric spaces corresponding to these homotopies are given in Part I, Theorem 4.3.

3.6 Structure variety of an associative pair

We can define the *structure variety of an associative pair*, exactly as in the Jordan case: recall from [Lo75] that an *associative pair* is given by two vector spaces \mathbb{A}^\pm and trilinear maps $\mathbb{A}^\pm \times \mathbb{A}^\mp \times \mathbb{A}^\pm \rightarrow \mathbb{A}^\pm$, $(x, y, z) \mapsto \langle xyz \rangle^\pm$ such that

$$\langle xy \langle zuv \rangle^\pm \rangle^\pm = \langle \langle xyz \rangle^\pm uv \rangle^\pm = \langle x \langle uzy \rangle^\mp v \rangle^\pm.$$

The symmetrized product $T^\pm(x, y, z) := \langle x, y, z \rangle + \langle z, y, z \rangle$ is then a Jordan pair. If we define the *structure variety of the associative pair* as before by (1), then it will be a subset of the structure variety of the Jordan pair T^\pm . However, there is a second definition of structure variety, given by the ‘‘opposite’’ condition

$$\alpha \langle x, \alpha y, z \rangle = \langle \alpha z, y, \alpha x \rangle, \quad (9)$$

which gives another subset of the structure variety of the Jordan pair T^\pm . In our classification, case 1 (rectangular matrices) corresponds to the associative pair $(M(p, q; \mathbb{K}), M(q, p; \mathbb{K}))$ with product $\langle x, y, z \rangle = xyz$. Here, cases 1.a, 1.a', 1.A, 1.A', 1.3a, 1.3.a' correspond to the first choice of definition of the structure variety, and cases 1.b, 1.c, 1.B, 1.3.b, 1.3.c to the second (opposite) choice. It follows that Theorem 3.1 classifies elements of these kinds of structure varieties.

3.7 Jordan pairs of type four (spin factors)

We recall the following result of Rivillis ([Ri67]):

Proposition 3.5 *Let (V, T) be a real simple Jordan pair of Type 4 (and of dimension at least 3) corresponding to the bilinear form β . Then α belongs to $\text{Svar}(V)$ if and only if α is symmetric with respect to β and $\alpha^2 = \lambda \text{id}$ with $\lambda \in \mathbb{K}$.*

Proof. (Cf. [Ri67].) Note that α belongs to $\text{Svar}(V)$ if and only if

$$\beta(\alpha x, y)\alpha z + \beta(\alpha z, y)\alpha x - \beta(\alpha x, \alpha z)y = \beta(x, \alpha y)\alpha z + \beta(z, \alpha y)\alpha x - \beta(x, z)\alpha^2 y$$

Choosing y linearly independent from αz and αx , we see first that $\alpha^2 y$ must be proportional to y and then $\beta(\alpha x, y) = \beta(x, \alpha y)$. This must hold for all x, y , hence α is symmetric. But then we get the condition $\beta(x, \alpha^2 z)y = \beta(x, z)\alpha^2 y$ which implies that α^2 must be a multiple of the identity. Conversely, it is clear that these conditions imply that $\alpha \in \text{Svar}(V)$. \blacksquare

For $\alpha^2 = 1$, we get endomorphisms that are simultaneously symmetric and orthogonal with respect to β , and for $\alpha^2 = -1$, we get endomorphisms that are simultaneously symmetric and ‘‘anti-orthogonal’’ w.r.t. β . The last case can only occur if β has signature (n, n) : then it is the real (or imaginary) part of a complex quadratic form, and α is the complex structure: $\langle ix, iy \rangle = -\langle x, y \rangle$. The case $\alpha^2 = 0$, $\alpha \neq 0$ can appear whenever the form β is neither positive nor negative definite. The symmetric spaces corresponding to α are described in [Ma79]. However, a global algebraic construction of these spaces, comparable to the ones in the other cases mentioned above, is missing. In particular, we have the impression that the spaces corresponding to $\alpha^2 = 0$ have a close relation to symmetric spaces related to point space ideals in $\text{Asym}(n, \mathbb{R})$, see above.

3.8 Low dimensional cases

In low dimension, certain isomorphisms between members of different families of Jordan pairs or triple systems occur; in particular, matrix algebras of low rank are often isomorphic to type four algebras $\mathbb{R}^{p,q}$ (recall notation from the table in subsection 3.1). In the following, we list such isomorphisms (following [Lo75], p. 196 ff and [Be00], Table XII.1.5) along with some comments on their inner ideals and their geometry. Recall that proper inner ideals in $\mathbb{R}^{p,q}$ are exactly the isotropic subspaces; thus their maximal dimension is $\min(p, q)$; in particular, $\mathbb{R}^n = \mathbb{R}^{n,0}$ contains no non-trivial inner ideals. Thus homotopies in this case are either trivial or invertible, and hence every proper contraction of an isotope is trivial (i.e., flat). The other extreme is the case $M(1, n; \mathbb{K})$: in this case, every vector subspace is an inner ideal, and hence there are many non-trivial contractions.

Before describing the symmetric spaces, recall from Part I ([BeBi], Sections 2 and 4) notation and definition of homotopes of classical Lie algebras and Lie groups, as well as the description of symmetric spaces G_α/H_α belonging to a homotopy α ([BeBi], Theorem 4.1). Recall in particular the *A-orthogonal Lie algebra* (for an arbitrary matrix $A \in \text{Sym}(n, \mathbb{K})$)

$$\mathfrak{o}_n(A; \mathbb{K}) := \text{Asym}(n; \mathbb{K}) \text{ with Lie bracket } [X, Y]_A := XAY - YAX.$$

The case $n = 3$ will be particularly important below: we calculate explicitly the Lie bracket with respect to a *diagonal* matrix $A = \text{dia}(a, b, c)$; let us denote this algebra by $\mathfrak{o}_3((a, b, c); \mathbb{K})$:

$$\left[\begin{pmatrix} 0 & x & y \\ -x & 0 & z \\ -y & -z & 0 \end{pmatrix}, \begin{pmatrix} 0 & x' & y' \\ -x' & 0 & z' \\ -y' & -z' & 0 \end{pmatrix} \right]_A = \begin{pmatrix} 0 & c(zy' - yz') & b(zx' - xz') \\ -c(zy' - yz') & 0 & -a(xy' - x'y) \\ -b(zx' - xz') & a(xy' - x'y) & 0 \end{pmatrix}$$

In other words, denoting by e, f, k the natural basis in $\text{Asym}(3; \mathbb{K})$, this Lie algebra is \mathbb{K}^3 with commutator relations

$$[e, f] = ck, \quad [e, k] = af, \quad [f, k] = -be \quad (10)$$

For $A = \text{dia}(1, 1, 1)$, we get the usual orthogonal algebra $\mathfrak{o}_3((1, 1, 1); \mathbb{K}) = \mathfrak{o}(3; \mathbb{K})$; for $A = \text{dia}(-1, -1, 1)$, we get $\mathfrak{o}_3((-1, -1, 1); \mathbb{K}) = \mathfrak{o}(2, 1; \mathbb{K}) \cong \mathfrak{sl}_2(\mathbb{K})$; for $A = \text{dia}(0, 1, -1)$ and $A = \text{dia}(0, 1, 1)$, we get two solvable 3-dimensional algebras, $\mathfrak{o}_3((0, 1, -1); \mathbb{K})$ and $\mathfrak{o}_3((0, 1, 1); \mathbb{K})$, isomorphic to semidirect products $\mathbb{K}^2 \rtimes \mathfrak{o}(2; \mathbb{K})$, resp. $\mathbb{K}^2 \rtimes \mathfrak{o}(1, 1; \mathbb{K})$. Finally, for $A = \text{dia}(0, 0, 1)$, we get the 3-dimensional Heisenberg algebra. Note also that, for any of these algebras, conjugation by the diagonal matrix $I_{2,1}$ is a Lie algebra automorphism.

3.8.1 Dimension 2

There are exactly three non-isomorphic two-dimensional semisimple Jordan pairs:

A. Two-dimensional conformal geometry: $V = \mathbb{C} = \text{Sym}(1, \mathbb{C}) = M(1, 1; \mathbb{C}) \cong \mathbb{R}^2$ (isomorphic also as Jordan algebras). The only inner ideals are 0 and \mathbb{C} , thus homotopies are either trivial or invertible. We have the following list of isotopes:

- $\alpha(X) = \overline{X}$ gives $M_\alpha = D$ (the unit disc),
- $\alpha(X) = -\overline{X}$ gives $M_\alpha = S^2$ (sphere, c -dual of D); both contract directly to flat spaces;
- $\alpha(X) = X$ gives \mathbb{C}^\times which is self c -dual and flat.

B. Real projective plane: $(V^+, V^-) = (M(1, 2; \mathbb{R}), M(2, 1; \mathbb{R}))$. Every linear subspace in V^- is an inner ideal, hence there are inner ideals of dimension one, and there exist contractions in “two steps”. Recall from Remark 3.2 that in the projective case all homotopies can be put in the form $\alpha(X) = BX$ with a symmetric matrix B . Thus classification of homotopes in the projective case amounts to classification of $\text{GL}(2, \mathbb{R})$ -orbits in $\text{Sym}(2, \mathbb{R})$:

- B.1. Isotopes correspond to the three open orbits which we represent by the three matrices $B = 1$, $B = I_{1,1}$, $B = -1$. The corresponding isotopes are

$$M = D, \quad M = H = \text{O}(2, 1)/\text{O}(1) \times \text{O}(1, 1) \text{ (one-sheeted hyperboloid)}, \quad M = \mathbb{RP}^2 = \text{O}(3)/\text{O}(2) \times \text{O}(1).$$

B.2. There are two rank-one orbits represented by diagonal matrices $B = E_{11} = \text{dia}(1, 0)$ and $B = -E_{11} = \text{dia}(-1, 0)$. The corresponding homotopes are

$$C_H = \text{O}_3((1, 0, 1))/\text{O}_2((1, 0)) \times \text{O}_1(1), \quad C_E = \text{O}_3((-1, 0, 1))/\text{O}_2((-1, 0)) \times \text{O}_1(1)$$

where $\text{O}_3((a, b, c))$ is the group having Lie algebra $\mathfrak{o}_3((a, b, c); \mathbb{R})$, see [BeBi]. We call C_E the *elliptic cylinder* and C_H the *hyperbolic cylinder* (in [BDS09] the terms *Poincaré coset*, resp. *Galilei coset* had been used). The spaces C_E , resp. C_H can be characterized as the non-flat and non-semisimple two-dimensional symmetric spaces, c -dual to each other, and topologically the former is homeomorphic to a cylinder, and the latter to a vector space.

Contraction relations are precisely the closure relations among $\text{GL}(2, \mathbb{R})$ -orbits in $\text{Sym}(2, \mathbb{R})$. Point reflection at the origin corresponds to c -duality. Note that all local isomorphism classes of 2-dimensional symmetric spaces appear in this picture. In particular, the 2-dimensional $ax + b$ -group (affine group of the real line) is among these spaces. Indeed, since the homotopy $\alpha(X) = E_{11}X = (1 \ 0)X(1 \ 0)^t$ is both of type 1.a and 1.b, the space C_H has two interpretations: it can be written as a quotient of two groups (see above), but like all spaces of type 1.a it is also a Lie group, seen as symmetric space, namely the group $\text{GL}_{1,2}((1 \ 0); \mathbb{R})$. The Lie algebra of this group, denoted by $\mathfrak{gl}_{1,2}((1 \ 0); \mathbb{R})$, is $\mathbb{R}^2 = M(1, 2; \mathbb{R})$ with

$$[(x, y), (x', y')] = (x, y)(1 \ 0)^t(x', y') - (x', y')(1 \ 0)^t(x, y) = (0, xy' - x'y)$$

which is the well-known “ $ax + b$ -algebra”, given by the bracket $[e_1, e_2] = e_2$ on \mathbb{R}^2 . Thus $\text{GL}_{1,2}((1 \ 0); \mathbb{R})$ is nothing but the $ax + b$ -group, which is naturally fibered over \mathbb{R}^\times with fiber the translation group of \mathbb{R} . It is quite instructive to check by a direct computation that the transvection algebra of the $ax + b$ -group is a 3-dimensional Lie algebra, given by the bracket relations (10) with $c = 1$, $a = 1$, $b = 0$, hence isomorphic to the Lie algebra $\mathfrak{g} = \mathfrak{o}_3((1, 0, 1); \mathbb{R})$, with symmetric decomposition $\mathfrak{g} = \mathfrak{h} \oplus \mathfrak{p}$ with $\mathfrak{h} = \mathbb{R}k$, $\mathfrak{p} = \mathbb{R}e \oplus \mathbb{R}f$. (Note that the transvection algebra of the $ax + b$ -group is strictly smaller than the algebra of left- and right translations which is of dimension four!) In a similar way, the c -dual of C_H also has two different interpretations: it is at the same time of type 1.a’ and 1.b’:

$$C_E = \text{GL}_{1,2}((1 \ 0); \mathbb{C})/\text{GL}_{1,2}((1 \ 0); \mathbb{R}) = \text{O}_3((-1, 0, 1))/\text{O}_2((-1, 0)) \times \text{O}_1(1)$$

C. Polarized Geometry: direct products. $V = \mathbb{R}^{1,1}$ is not a simple Jordan algebra: it is isomorphic to $\mathbb{R} \times \mathbb{R}$. The structure variety of $\mathbb{R} \times \mathbb{R}$ has been computed in Example 3.4. Isotopes are either uninteresting (direct product of flat spaces), or isomorphic to the one-sheeted hyperboloid H . Although there exist proper inner ideals (the two isotropic lines in $\mathbb{R}^{1,1}$), they do not contribute to non-trivial contractions of H : indeed, the computations from Example 3.4 show that homotopies with image one of these two lines lead to direct product situations, hence to flat spaces. Summing up, no interesting contractions of H arise in this case.

3.8.2 Dimension 3

There are exactly three simple Jordan pairs of dimension 3 over \mathbb{R} :

A. Riemannian conformal space: case of the Jordan algebra \mathbb{R}^3 . Like \mathbb{R}^2 , the Jordan algebra \mathbb{R}^3 has no non-trivial inner ideals, hence there are only isotopes contracting directly to flat space. These isotopes are given by symmetric matrices having coefficients ± 1 on the diagonal, leading to spaces going from the compact S^3 to hyperbolic space via its indefinite isotopes.

B. Projective 3-space: $M(1, 3; \mathbb{R}) \cong \text{Asym}(3, \mathbb{R})$ (see [Lo75], p. 197 for this isomorphism). Looking at this JTS as a space of row matrices, the discussion is similar to the one for $M(1, 2; \mathbb{R})$ above: there are inner ideals of all dimensions, and the structure variety is given by all $\alpha(X) = BX$ with $B \in \text{Sym}(3, \mathbb{R})$; contraction relations are closure relations of $\text{GL}(3, \mathbb{R})$ -orbits there: the 4 open orbits, corresponding the invertible matrices $B = 1, I_{2,1}, I_{1,2}, -1$, give the isotopes

$$\mathbb{R}\mathbb{P}^3 = \text{O}(4)/\text{O}(3) \times \text{O}(1), \quad \text{O}(2, 2)/\text{O}(2, 1) \times \text{O}(1), \quad \text{O}(3, 1)/\text{O}(1) \times \text{O}(2, 1), \quad \text{O}(3, 1)/\text{O}(3) \times \text{O}(1).$$

Next we have three different orbits of rank-2 matrices and two different orbits of rank-1-matrices, corresponding to symmetric spaces resembling the spaces C_E and C_H from above. As above, spaces associated

to the two rank-1 orbits have two different interpretations: semi-positive rank-one matrices correspond to homotopies $X \mapsto AXA$ of type 1.a, hence to group cases: for $A = (1\ 0\ 0)$, the same computation as above shows that this group is the affine group of \mathbb{R}^2 ($ax + b$ -group). Its Lie algebra is 3-dimensional with commutator relations $[k, e] = e$, $[k, f] = f$, $[e, f] = 0$, and it is not isomorphic to any of the 3-dimensional Lie algebras $\mathfrak{o}_3((a, b, c); \mathbb{R})$ described above.

Looking now at this JTS as $\text{Asym}(3, \mathbb{R})$, the picture becomes considerably more complicated: first of all, a glance at the list of homotopes of this JTS shows that the only invertible homotopies are of type 3.b or 3.b' with an invertible symmetric matrix A . We may choose $A = \text{dia}(a, b, c)$, so for type 3.b we get precisely the orthogonal groups $\text{O}_3((a, b, c); \mathbb{R})$ introduced above. For type 3.b', we get their c -duals. This implies the four isotopes mentioned above must be isomorphic to $\text{SO}(3)$, $\text{SO}(2, 1)_o$, $\text{SO}(3, \mathbb{C})/\text{SO}(2, 1)$, $\text{SO}(3, \mathbb{C})/\text{SO}(3)$, respectively (indeed, this could be deduced from known isomorphisms of low dimensional Lie groups). Choosing a degenerate diagonal matrix $A = \text{dia}(a, b, c)$, we get three other group cases, but none of them is isomorphic *as a group* to the $ax + b$ -group mentioned above. However, *as symmetric space*, the $ax + b$ -group is isomorphic to $\mathfrak{o}_3((110); \mathbb{R})$: indeed, both the commutator relations $[k, e] = e$, $[k, f] = f$, $[e, f] = 0$ and $[k, e] = f$, $[k, f] = e$, $[e, f] = 0$ lead to the LTS relations $[k, k, f] = f$, $[k, k, e] = e$. In both cases the formula for the homotopy boils down to $\alpha((x, y, z)) = x$. The case of the group $\mathfrak{o}_3((100); \mathbb{R})$ belongs to the homotopy $\alpha(X) = I_{11}XI_{11}$ which is zero since X is skew-symmetric; indeed, the group $\mathfrak{o}_3((100); \mathbb{R})$ is two-step nilpotent, hence its LTS is zero. Note also that all inner ideals in $\text{Asym}(3; \mathbb{R})$ are ‘‘point spaces’’ (since the principal inner ideal $\text{Asym}(2; \mathbb{R})$ is isomorphic to \mathbb{R}), and hence symmetric spaces are fibered over isotopes of \mathbb{RP}^1 or \mathbb{RP}^2 . (For the time being, we have no good description of homotopes corresponding to point space ideals for $\text{Asym}(n, \mathbb{R})$ in general; in particular, we do not know whether for $n > 3$ new group cases show up.)

C. Pseudo-Riemannian conformal space: Jordan algebra $\text{Sym}(2, \mathbb{R}) \cong \mathbb{R}^{2,1}$. This is the ‘‘small Minkowski space’’. Proper inner ideals are the isotropic lines. Thus there are non-trivial contractions. This case already shows the ‘‘generic’’ feature that the structure variety has several irreducible components. The isotopes are: the two conal spaces $\text{GL}(2, \mathbb{R})/\text{O}(2)$, $\text{GL}(2, \mathbb{R})/\text{O}(1, 1)$ (type 2.a) and their c -duals $\text{U}(2)/\text{O}(2)$, $\text{U}(1, 1)/\text{O}(1, 1)$ (type 2.a'), and the group case $\text{Sp}(1; \mathbb{R}) = \text{SL}(2, \mathbb{R})$ (type 2.b) and its c -dual $\text{SL}(2, \mathbb{C})/\text{SL}(2, \mathbb{R})$ (type 2.b'). Spaces of type 2.a and 2.a' have non-trivial contractions, whereas the two spaces of type 2.b, 2.b' (belonging to another irreducible component of $\text{Svar}(\text{Sym}(2, \mathbb{R}))$) do not.

3.8.3 Dimension 4

There are exactly five simple Jordan pairs of dimension four, and there are also some interesting direct products, such as $\mathbb{R}^2 \times \mathbb{R}^2 = \mathbb{C} \times \mathbb{C} = \mathbb{C}^2$ (which hosts the polarized space $\text{SL}(2, \mathbb{C})/\mathbb{C}^*$). For reasons of space, the following discussions of these spaces will be sketchy and far from being exhaustive.

A. Riemannian Conformal geometry: $\mathbb{R}^4 \cong \mathbb{H}$ (as Jordan algebras). Isotopes are ranking from the compact $\mathbb{HP}^1 = S^4$ to hyperbolic space and contain the group case \mathbb{H}^\times ; they all contract directly to 0.

B. Conformal Artinian geometry: $M(2, 2; \mathbb{R}) \cong \mathbb{R}^{2,2}$ (as Jordan algebras): the principal inner ideals are the isotropic lines (principal), the isotropic planes are inner, but not principal. Isotopes are: group type space $\text{GL}(2, \mathbb{R})$ (type 1.a), its c -dual $\text{GL}(2, \mathbb{C})/\text{GL}(2, \mathbb{R})$ (type 1.a'), $\text{O}(4)/\text{O}(2) \times \text{O}(2)$, $\text{O}(2, 2)/\text{O}(2) \times \text{O}(2)$, $\text{O}(3, 1)/\text{O}(2) \times \text{O}(1, 1)$, $\text{O}(2, 2)/\text{O}(1, 1) \times \text{O}(1, 1)$ (type 1.b), $\text{Sp}(2, \mathbb{R})/\text{Sp}(1, \mathbb{R}) \times \text{Sp}(1, \mathbb{R})$ (type 1.c). The latter space has no non-trivial contractions, whereas all other have proper contractions corresponding to 1-dimensional inner ideals; in particular, $\text{GL}(2, \mathbb{R})$ contracts to groups $\text{GL}_2(A; \mathbb{R})$.

C. Conformal Minkowski space: $\text{Herm}(2, \mathbb{C}) = \mathbb{R}^{3,1}$ (as Jordan algebras). Inner ideals are the isotropic lines; all are principal. Isotopes are: conal spaces $\text{GL}(2, \mathbb{C})/\text{U}(2)$, $\text{GL}(2, \mathbb{C})/\text{U}(1, 1)$ (type 1.1.a) and their duals, the group type spaces $\text{U}(2)$ and $\text{U}(1, 1)$ (type 1.1.a'), $\text{O}(2, 2)/\text{O}(2, \mathbb{C})$ (type 1.1.b; self- c dual in this particular case), $\text{Sp}(2, \mathbb{R})/\text{Sp}(1, \mathbb{C})$ (type 1.1.c), $\text{O}(2, 2)/\text{Sp}(1, \mathbb{C})$ (type 1.1.c'); the latter two have no non-trivial contraction, whereas all other spaces have exactly one non-trivial contraction.

D. Projective 4-space: $M(1, 4; \mathbb{R})$. The situation is similar to the case $M(1, 3; \mathbb{R})$ described above. Since there is no special isomorphism in this case, this case is already ‘‘generic’’ for real projective spaces. In particular, the only group cases correspond exactly to the rank-one homotopes (groups $\text{GL}_{1,4}(A; \mathbb{R})$, still isomorphic to an $ax + b$ -group).

E. Complex projective plane: $M(1, 2; \mathbb{C})$. We distinguish \mathbb{C} -linear and \mathbb{C} -antilinear homotopes: this discussion of the \mathbb{C} -linear case follows exactly the same pattern as the one for $M(1, 2; \mathbb{R})$; since \mathbb{C} is algebraically closed, there are less isomorphism classes of symmetric spaces (3 complex spaces, corresponding to rank B being 0, 1 or 2, compared to the 6 real spaces from above).

For \mathbb{C} -antilinear α , let us first remark that (for $p = 1$, $q = n$; here $n = 2$) types 1.A and 1.A' become special instances of type 1.B; in other words, \mathbb{C} -antilinear homotopies are precisely the maps of the form $F(z) = f(\bar{z})$ with Hermitian $f : \mathbb{C}^2 \rightarrow \mathbb{C}^2$. For invertible f , we get according to the various possible signatures, the isotopes

$$U(3)/U(2) \times U(1), \quad U(2, 1)/U(2) \times U(1), \quad U(2, 1)/U(1) \times U(1, 1).$$

If f is of rank one, then the spaces have two interpretations: type 1.A corresponds to f of rank one and non-negative, and type 1.A' to f of rank one and non-positive. For the corresponding symmetric spaces we have for type 1.A:

$$U_3((1, -1, 0)/U_1(1) \times U_2((1, 0)) \cong GL_{2,1}((1, 0); M(2, 2; \mathbb{R}))/GL_{2,1}((1, 0; \mathbb{C}) .$$

3.8.4 Dimension 6

The cases of dimension 6 and 10 are of particular interest, since low dimensional isomorphisms appear there. We just give the list of simple JTS and of special isomorphisms in dimension 6, and leave a more detailed discussion of the corresponding homotopes and their specific features for subsequent work.

A. Riemannian conformal space: \mathbb{R}^6

B. Lorentzian conformal space: $\mathbb{R}^{5,1} \cong \text{Herm}(2, \mathbb{H})$

C. Artinian conformal space: $\text{Asym}(4, \mathbb{R}) = \mathbb{R}^{3,3}$ (see [Lo75], p. 200, for this isomorphism)

D. Pseudo-Riemannian conformal space of signature (4, 2): $\mathbb{R}^{2,4}$

E. Complex projective space: $M(1, 3; \mathbb{C}) \cong \text{Asym}(3, \mathbb{C})$

F. Complex Lagrangian geometry of a symplectic form on \mathbb{C}^4 : $\text{Sym}(2, \mathbb{C}) \cong \mathbb{C}^3$

G. Real Lagrangian geometry of a symplectic form on \mathbb{R}^6 : $\text{Sym}(3, \mathbb{R})$

H. Real Grassmannian of 2-spaces in \mathbb{R}^5 : $M(2, 3; \mathbb{R})$

References

- [Backes83] Backes E., Geometric applications of Euclidean Jordan triple systems. *Manuscripta math.* **42** (1983), 265 – 272.
- [Be00] Bertram, W., *The geometry of Jordan- and Lie structures*, Springer Lecture Notes in Mathematics **1754**, Berlin 2000
- [Be02] Bertram, W., “Generalized projective geometries: general theory and equivalence with Jordan structures”, *Adv. Geom.* **2** (2002), 329–369 (electronic version: preprint 90 at <http://homepage.uibk.ac.at/~70202/jordan/index.html>)
- [Be08] Bertram, W., “Homotopes and conformal deformations of symmetric spaces.” *J. Lie Theory* **18** (2008), 301 – 333 (arxiv: math.RA/0606449)
- [BeBi] Bertram, W. and Bieliavsky, P., “Homotopes of symmetric spaces. I: Construction by algebras with 2 involutions” (Part I of this work, arxiv, november 2010)
- [BeD09] Bertram, W. and M. Didry, “Symmetric bundles and representations of Lie triple systems”, *Journal of Generalized Theory and Applications* **3** (no.4) (2009), 261-284
- [BeL08] Bertram, W. and H. Löwe, “Inner ideals and intrinsic subspaces of linear pair geometries”, *Adv. Geom.* **8** (2008), 53 – 85

- [BDS09] Bieliavsky, P., S. Detournay and Ph. Spindel, “The Deformation Quantization of the Hyperbolic Plane”, *Com. Math. Phys.* **289** (2), (2009), 529 – 559
- [CGR00] Cahen, Michel; Gutt, Simone; Rawnsley, “John Symmetric symplectic spaces with Ricci-type curvature”. Conference Moshé Flato 1999, Vol. II (Dijon), 81–91, *Math. Phys. Stud.*, 22, Kluwer Acad. Publ., Dordrecht, 2000.
- [DFGG08] Draper , A., Fernandez Lopez, E. Garcia , and M. Gomez Lozano, “The inner ideals of the classical Lie algebras, related gradings and Jordan pairs”, preprint (2008) (electronic version: preprint 264 at <http://homepage.uibk.ac.at/~c70202/jordan/index.html>)
- [Lo69] Loos, O., *Symmetric Spaces I*, Benjamin, New York, 1969
- [Lo75] Loos, O., *Jordan Pairs*, Springer LNM, 1975
- [LoN94] Loos, O. and E. Neher, Complementation of Inner Ideals in Jordan Pairs, *J. of Algebra* **166** (1994), 255 – 295
- [Ma73] Makarevic, B.O., Open symmetric orbits of reductive groups in symmetric R -spaces. *Math. USSR Sbornik* **20** (1973), 406 – 418
- [Ma79] Makarevic, B.O., Jordan algebras and orbits in symmetric R -spaces. *Trans. Moscow. Math. Soc.* **39** (1979), 169 – 192
- [McC71] McCrimmon, K., “Inner ideals in quadratic Jordan algebras”, *Trans. Amer. Math. Soc.*, 159 (1971), 445-468.
- [Mey73] Meyberg, K., *Lectures on algebras and triple systems*, Lecture notes, University of Virginia, Charlottesville, Va., 1973.
- [Ne96] Neher, E., Lie algebras graded by 3-graded root systems and Jordan pairs covered by grids, *Amer. J. Math.*, **118** (1996), 439 – 491
- [Ri67] Rivillis, A.A., A class of symmetric spaces with extendable group of motions and a generalization of the Poincaré model. *Dokl. Akad. Nauk. SSSR* **173** (1967), 423 – 426.
- [Ri70] Rivillis, A.A., Homogeneous locally symmetric regions in homogeneous spaces associated with semisimple Jordan algebras. *Mat. USSR Sbornik* **11** (1970), no. 3, 377 – 389