



Jordan-Hölder sequences and self-adjoint (a, b) -modules

Piotr P. Karwasz

► To cite this version:

| Piotr P. Karwasz. Jordan-Hölder sequences and self-adjoint (a, b) -modules. 2011. hal-00583887

HAL Id: hal-00583887

<https://hal.archives-ouvertes.fr/hal-00583887>

Preprint submitted on 7 Apr 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Jordan-Hölder sequences and self-adjoint (a, b) -modules

Piotr P. Karwasz *

April 7, 2011

Abstract

Given the lack of uniqueness of the Jordan-Hölder composition series in the theory of (a, b) -modules we are interested whether the particularities of certain (a, b) -modules can be transmitted to their composition series. This article will focus on the properties of Jordan-Hölder composition series of self-adjoint (a, b) -modules. In particular we will prove that a self-adjoint composition series always exists for self-adjoint (a, b) -modules.

Mathematics Subject Classification (2010): 32S25, 32S40, 32S50

1 Introduction

The Brieskorn module of a germ of holomorphic function $f : \mathbb{C}^{n+1} \rightarrow \mathbb{C}$ with an isolated singularity at the origin introduced by E BRIESKORN in [Bri70] can be formally completed for the operation $b := df \wedge d^{-1}$. The result is called (a, b) -module and can be defined in an abstract way as:

Definition 1.1. *An (a, b) -module is a free $\mathbb{C}[[b]]$ -module E of finite rank over the ring of formal power series in b , endowed with a \mathbb{C} -linear endomorphism ‘ a ’ which satisfies:*

$$ab - ba = b^2$$

We recall some basic classification of this object: a sub- (a, b) -module F of E is a sub- $\mathbb{C}[[b]]$ -module of E stable by ‘ a ’ and the (a, b) -structure passes onto the quotient $\mathbb{C}[[b]]$ -module E/F which satisfies all the properties of an (a, b) -module except that it has possibly a b -torsion. The sub- (a, b) -module F is called **normal** if E/F is free on $\mathbb{C}[[b]]$.

Since the completion of Brieskorn modules generates **regular** (a, b) -modules ([Bar93]), i.e. (a, b) -modules that can be embedded as a sub- (a, b) -module into an (a, b) -module E satisfying $aE \subset bE$, we’ll limit our inquiries to this subclass of objects.

All regular (a, b) -modules of rank 1 are generated by an element e_λ satisfying $ae_\lambda = \lambda be_\lambda$ for a complex number λ . We will refer to them as **elementary** (a, b) -modules of parameter λ and note them with E_λ .

*Institut Élie Cartan Nancy UMR 7502
Nancy-Université, B.P. 70230, 54506 Vandœuvre-lès-Nancy, France
E-mail: piotr.karwasz@iecn.u-nancy.fr

A basic result shows that a regular (a, b) -module E admits Jordan-Hölder composition series

$$0 = F_0 \subsetneq F_1 \subsetneq \cdots \subsetneq F_n = E$$

with the F_i normal in E and for a regular (a, b) -module the quotients F_i/F_{i-1} are **elementary** (a, b) -modules E_λ of parameter λ .

The isomorphism classes of such quotients vary from a Jordan-Hölder composition series to another and a quotient E_λ may appear as $E_{\lambda+j}$ in another sequence, with $j \in \mathbb{Z}$.

At a first approach we studied the behaviour of such sequences under the duality functor (cf. [Bar97]):

Definition 1.2. *Let E be an (a, b) -module and E_0 the elementary (a, b) -module of parameter 0, then we may define upon the $\mathbb{C}[[b]]$ -module $\text{Hom}_{\mathbb{C}[[b]]}(E, E_0)$ an (a, b) -module structure given by:*

$$(a \cdot \varphi)(x) = a\varphi(x) - \varphi(ax)$$

*This module is called the **dual** (a, b) -module and noted E^* .*

However, as it was proven by R. BELGRADE in [Bel01], an (a, b) -module E associated to a Brieskorn module is isomorphic to the **conjugate** of the dual, i.e. the **adjoint** (a, b) -module. Even if both approaches give similar results concerning the symmetry of Jordan-Hölder composition series, we will prefer the study of self-adjoint (a, b) -modules for the greater interest they play in the theory of singularities.

In the context of (a, b) -modules, the conjugate itself is defined in a way borrowed from the complex vector spaces:

Definition 1.3. *Let E be an (a, b) -module, we call **conjugate** (a, b) -module and note it \check{E} , the set E endowed with the (a, b) -structure given by reversing the signs of both 'a' and 'b': $a \cdot_{\check{E}} v = -a \cdot_E v$ and $b \cdot_{\check{E}} v = -b \cdot_E v$.*

*In particular we call **adjoint** the conjugate of the dual (a, b) -module E^* and we call self-adjoint an (a, b) -module E which is isomorphic to \check{E}^* .*

When working with isomorphisms between an (a, b) -module E and its adjoint \check{E}^* , it is often useful to look at it as a $\mathbb{C}[[b]]$ -bilinear map between $E \times \check{E}$ and E_0 . Such a perspective brought us ([Kar]) to give the following definition of (a, b) -bilinear map and (a, b) -hermitian forms:

Definition 1.4. *Let E, F and G be (a, b) -modules and Φ a $\mathbb{C}[[b]]$ -bilinear map between $E \times F$ and G . We say that Φ is an (a, b) -bilinear map (or form if $G = E_0$) if:*

$$a\Phi(v, w) = \Phi(av, w) + \Phi(v, aw)$$

for every $v \in E$ and $w \in F$.

An (a, b) -bilinear map between $E \times \check{E}$ and E_0 is called an (a, b) -hermitian form if moreover it satisfies

$$\Phi(v, w) = S(b)e_0 \Rightarrow \Phi(w, v) = S(-b)e_0$$

with $v, w \in E$ and $S(b) \in \mathbb{C}[[b]]$. (a, b) -anti-hermitian forms satisfy the same equation with $S(-b)$ replaced by $-S(-b)$.

An (a, b) -bilinear map between $E \times F$ and E_0 is **non-degenerate** if it induces an isomorphism between E and F^* .

An (a, b) -module that admits a non-degenerate hermitian form will be called **hermitian**.

The hermitian (a, b) -bilinear forms on an (a, b) -module E induce, as in the case of complex vector spaces, an isomorphism $\Phi : E \rightarrow \check{E}^*$ which is equal to its image $\check{\Phi}^*$ under the adjunction functor.

We should remark that not every self-adjoint (a, b) -module is hermitian and there are even examples of (a, b) -modules that only admit an anti-hermitian non-degenerate form. In the general case every regular self-adjoint (a, b) -module can be decomposed into a hermitian part and an anti-hermitian part (not necessarily in a unique way) ([Kar]).

2 Self-adjoint composition series

In view of Belgrade's result we will work with the adjunction functor. All proofs however should be valid with only minor modifications if we work with the duality functor.

Consider a regular (a, b) -module E of rank $n \in \mathbb{N}$ and a Jordan-Hölder decomposition of itself:

$$0 = F_0 \subsetneq F_1 \subsetneq \dots \subsetneq F_n = E.$$

with $F_j/F_{j-1} \simeq E_{\lambda_j}$, the elementary (a, b) -module of parameter λ_j . We say that the sequence is **self-adjoint** if $\lambda_{n-j+1} = -\lambda_j$ for all $1 \leq j \leq n$ and the (a, b) -module F_{n-j}/F_j is self-adjoint for all $0 \leq j \leq [n/2]$.

We shall prove the following theorem for regular hermitian (a, b) -modules and we will extend it successively to all self-adjoint (a, b) -modules.

Theorem 2.1. *Let E be a regular hermitian (a, b) -module then it has a Jordan-Hölder sequence*

$$0 = F_0 \subsetneq F_1 \subsetneq \dots \subsetneq F_n = E$$

which is self-adjoint.

Before proving the theorem we shall introduce a couple of lemmas.

Lemma 2.2. *Let E be a regular hermitian (a, b) -module and $\Phi : E \rightarrow \check{E}^*$ a hermitian isomorphism.*

If there exists F_1 normal sub- (a, b) -module isomorphic to E_λ such that

$$\Phi(F_1)(F_1) = 0,$$


then there exists a normal sub- (a, b) -module F_{n-1} of rank $n - 1$ such that $E/F_{n-1} \simeq \check{F}_1^$ and F_{n-1}/F_1 is hermitian.*

Proof. Let e_λ be the generator of F_1 and H the hermitian form associated to Φ by $H(x, y) = \Phi(y)(x)$ and consider the annihilator of this form under H :

$$F_{n-1} := \{x \in E \mid H(e_\lambda, x) = 0\}.$$

We remark that the condition $H(e_\lambda, e_\lambda) = 0$ gives us $F_1 \subset F_{n-1}$ and F_{n-1} is normal, because it is the kernel of a morphism.

Figure 1: Modules in symmetric positions with respect to the dotted line are each other's adjoint.



Let us consider the following exact sequence:

$$0 \rightarrow F_1 \rightarrow E \rightarrow E/F_1 \rightarrow 0$$

from which we can pass to the adjoint sequence:

$$0 \rightarrow (\widetilde{E/F_1})^* \rightarrow \check{E}^* \xrightarrow{\pi} \check{F}_1^* \rightarrow 0.$$

Since π is the restriction morphism of forms on E to the sub- (a, b) -module F_1 , the kernel of π , $K := \text{Ker } \pi$ can be described as follows:

$$K = \{\varphi \in \check{E}^* \mid \varphi(F_1) = 0\}.$$

The adjoint sequence being exact, we can identify from now on $(\widetilde{E/F_1})^*$ with K , i.e. sub- (a, b) -module of \check{E}^* whose elements annihilate F_1 .

If we consider the restriction of the map Φ to F_{n-1}

$$\Phi|_{F_{n-1}} : F_{n-1} \rightarrow \check{E}^*$$

and the fact that by definition $\Phi(x)(e_\lambda) = 0$ for all $x \in F_{n-1}$, we obtain that $\Phi(F_{n-1}) \subset (\widetilde{E/F_1})^*$.

On the other side for all $\varphi \in (\widetilde{E/F_1})^*$ the element $y = \Phi^{-1}(\varphi)$ verifies $\Phi(y)(e_\lambda) = 0$, therefore we have also $(\widetilde{E/F_1})^* \subset \Phi(F_{n-1})$. It follows that $\Phi(F_{n-1}) = (\widetilde{E/F_1})^*$ and since Φ is an isomorphism, F_{n-1} is isomorphic to its image by Φ : $(\widetilde{E/F_1})^*$.

Let us look now at the following exact sequence:

$$0 \rightarrow (F_{n-1}/F_1) \rightarrow (E/F_1) \rightarrow (E/F_{n-1}) \rightarrow 0$$

and its adjoint sequence:

$$0 \rightarrow (\widetilde{E/F_{n-1}})^* \xrightarrow{i} (\widetilde{E/F_1})^* \xrightarrow{\pi} (\widetilde{F_{n-1}/F_1})^* \rightarrow 0.$$

π designates the restriction application on the forms of $(\widetilde{E/F_1})^*$. $\text{Ker } \pi$ is thus the forms of $(\widetilde{E/F_1})^*$ that annihilate $(\widetilde{F_{n-1}/F_1})^*$ or with the convention of the previous paragraph, the forms of \check{E}^* that annihilate F_{n-1} and $F_1 \subset F_{n-1}$:

$$\text{Ker } \pi = \{\varphi \in \check{E}^* \text{ s.t. } \varphi(F_{n-1}) = 0\}$$

We note that the hermiticity of Φ gives us

$$\Phi(e_\lambda)(F_{n-1}) = \Phi(\widetilde{F_{n-1}})(e_\lambda) = 0$$

and therefore we have $\Phi(F_1) \subset \text{Ker } \pi$. An easy calculation shows that $\text{Ker } \pi$ is of rank 1. Since $\Phi(F_1)$ is normal, of rank 1 and included into $\text{Ker } \pi$, they must be equal.

We obtain $(\widetilde{E/F_{n-1}})^* \simeq \text{Ker } \pi \simeq F_1$. Now we know that Φ sends F_{n-1} onto $(E/F_1)^*$ and F_1 onto $\text{Ker } \pi$, so starting with the following exact sequence:

$$0 \rightarrow \text{Ker } \pi \hookrightarrow (E/F_1)^* \xrightarrow{\pi} (\widetilde{F_{n-1}/F_1})^* \rightarrow 0$$

we can obtain another by substituting $\text{Ker } \pi$ with F_1 and $(E/F_1)^*$ with F_{n-1} :

$$0 \rightarrow F_1 \rightarrow F_{n-1} \rightarrow (\widetilde{F_{n-1}/F_1})^* \rightarrow 0.$$

or in other terms $(\widetilde{F_{n-1}/F_1})^* \simeq (F_{n-1}/F_1)$. Note that the isomorphism is given by $x \rightarrow \Phi(x)|_{F_{n-1}}$ and is therefore hermitian.

The proof may be summarized by the graph of interwoven exact sequences presented in figure 1. \square

Remark 2.3. If $ae_\lambda = \lambda be_\lambda$ and $2\lambda \notin \mathbb{N}$, then $H(e_\lambda, e_\lambda) = 0$. In fact $H(e_\lambda, e_\lambda) \in E_0$ must satisfy:

$$\begin{aligned} aH(e_\lambda, e_\lambda) &= H(ae_\lambda, e_\lambda) + H(e_\lambda, -ae_\lambda) = \\ &H(\lambda be_\lambda, e_\lambda) + H(e_\lambda, -\lambda be_\lambda) = 2\lambda bH(e_\lambda, e_\lambda) \end{aligned}$$

which has non-trivial solutions in E_0 only if $2\lambda \in \mathbb{N}$. The double inversion of signs in the second factor are due to the hermitian nature of the form.

Lemma 2.4. *If E is a regular hermitian (a, b) -module and there exist $\lambda \in \mathbb{C}$ such that E contains two distinct normal elementary sub- (a, b) -modules F and G of parameters*

$$f = g = \lambda \text{ mod } \mathbb{Z}$$

then there exists $F_1 \subset F_{n-1}$ two normal sub- (a, b) -modules of rank 1 and $n-1$ respectively such that $(\widetilde{E/F_{n-1}})^ \simeq F_1$ and F_{n-1}/F_1 is hermitian.*

Proof. We will denote by H an hermitian form on E . Let e_f and e_g be generators of F and G and suppose without loss of generality that $f - g \geq 0$. We will show that there exists a normal elementary sub- (a, b) -module F_1 of E whose generator $e \in E$ satisfies $H(e, e) = 0$.

By the fundamental property $ab-ba = b^2$ of (a, b) -modules we have $ab^{f-g}e_g = (g+f-g)b \cdot b^{f-g}e_g$. Let's pose $e_1 = b^{f-g}e_g$. Consider now the complex vector space:

$$V := \{\alpha e_f + \beta e_1 \mid \alpha, \beta \in \mathbb{C}\}$$

Note that every $v \in V$ satisfies $av = fbv$. The b -linearity of H and the definition of the action of a gives us:

$$(a - 2fb)H(v, v) = 0$$

which has in E_0 only solutions of the form $\alpha b^{2f}e_0$, $\alpha \in \mathbb{C}$. There exists therefore an application B from $V \times V$ to \mathbb{C} such that:

$$H(v, w) = B(v, w)b^{2f}e_0 \quad \forall v, w \in V$$

The bilinearity and hermitianity of H imply that B is in fact a \mathbb{C} -bilinear symmetric or anti-symmetric form on a 2 dimensional complex vector space, depending whether $2f$ is even or odd. In the anti-symmetric case every vector will be isotropic, in the symmetric case we have:

$$B(e_f + xe_1, e_f + xe_1) = a_0 + a_1x + a_2x^2$$

for some complex numbers a_i . The vector space V has therefore an isotropic vector $e \neq 0$ such that $B(e, e) = 0$, and hence $H(e, e) = 0$.

By eventually dividing e by a certain power of b , operation that does not change the relation $H(e, e) = 0$, we can assume that $e \notin bE$, hence the module F_1 generated by e is normal.

We can now conclude by applying lemma 2.2 □

Lemma 2.5. *Let E be a regular (a, b) -module and:*

$$0 \subsetneq \dots F_{i-1} \subsetneq F_i \subsetneq F_{i+1} \subsetneq \dots E$$

be a Jordan-Hölder composition series with $F_i/F_{i-1} \simeq E_{\lambda_i}$ for all i and suppose there is a j such that $\lambda_{j+1} \neq \lambda_j \pmod{\mathbb{Z}}$.

Then we can find another Jordan-Hölder composition series that differs only in the j -th term F'_j such that $F'_j/F_{j-1} \simeq E_{\lambda'_j}$ and $F_{j+1}/F'_j \simeq E_{\lambda'_{j+1}}$ with $\lambda_j = \lambda'_{j+1} \pmod{\mathbb{Z}}$ and $\lambda_{j+1} = \lambda'_j \pmod{\mathbb{Z}}$, i.e. we can permute the quotients up to an integer shift of the parameters.

Proof. Let consider $G := F_{j+1}/F_{j-1}$ and the canonical projection $\pi : E \rightarrow E/F_{j-1}$. G is a rank two module. Using the classification of regular (a, b) -modules of rank 2 given by D. Barlet in [Bar93] we see that the only two possibilities for G are:

$$G \simeq E_{\lambda_j} \oplus E_{\lambda_{j+1}}$$

in which case we take $F'_j = \pi^{-1}(E_{\lambda_{j+1}})$ or

$$G \simeq E_{\lambda_{j+1}+1, \lambda_j}$$

generated by y and t satisfying:

$$\begin{aligned} ay &= \lambda_j by \\ at &= \lambda_{j+1} bt + y \end{aligned}$$

that has also another set of generators: t and $x := y + (\lambda_{j+1} - \lambda_j + 1)bt$ which satisfy:

$$\begin{aligned} ax &= (\lambda_j + 1)bx \\ at &= (\lambda_j - 1)bt + x. \end{aligned}$$

In this case we take $F'_j = \pi^{-1}(\langle x \rangle)$. \square

Lemma 2.6. *Let λ be either 0 or $1/2$ and E be a regular hermitian (a, b) -module. Suppose that there is a unique normal elementary sub- (a, b) -module of parameter equal to λ modulo \mathbb{Z} and suppose moreover that every Jordan-Hölder sequence contains at least 2 elementary quotients of parameter equal to λ modulo \mathbb{Z} .*

Then there exists $F_1 \subset \widetilde{F_{n-1}}$ two normal sub- (a, b) -modules of rank 1 and $n - 1$ respectively such that $(\widetilde{E/F_{n-1}})^ \simeq F_1$ and F_{n-1}/F_1 is hermitian.*

Proof. Let $F_1 \simeq E_\mu$ be the elementary sub- (a, b) -module of the hypothesis and $\{F_i\}$ a J-H sequence beginning with F_1 and such that E/F_{n-1} is of parameter μ' equal to $\lambda \bmod \mathbb{Z}$. We can find such a sequence by using repeatedly the previous lemma.

Consider the exact sequence:

$$0 \rightarrow F_{n-1} \rightarrow E \rightarrow (E/F_{n-1}) \rightarrow 0$$

and the adjoint sequence:

$$0 \rightarrow (\widetilde{E/F_{n-1}})^* \xrightarrow{i} \check{E}^* \xrightarrow{\pi} \check{F}_{n-1}^* \rightarrow 0.$$

The image of i is a normal elementary sub- (a, b) -module of \check{E}^* of parameter equal to $-\lambda \bmod \mathbb{Z}$ (since $(\widetilde{E/F_{n-1}})^* \simeq E_{-\mu'}$). But $\lambda = -\lambda \bmod \mathbb{Z}$ and $E \simeq \check{E}^*$ so by the uniqueness of F_1 given in the hypothesis $\text{Im} \left((\widetilde{E/F_{n-1}})^* \right) = \Phi(F_1)$, thus $(\widetilde{E/F_{n-1}})^* \simeq F_1$.

By replacing \check{E}^* by E and $(\widetilde{E/F_{n-1}})^*$ by F_1 in the sequence we obtain:

$$0 \rightarrow F_1 \xrightarrow{i} E \rightarrow \check{F}_{n-1}^* \rightarrow 0$$

which is exact and i is the inclusion of sub- (a, b) -modules, so $\check{F}_{n-1}^* \simeq (E/F_1)$ or equivalently $F_{n-1} \simeq (\widetilde{E/F_1})^*$. Note that the first isomorphism is given by Φ^{-1} , while the second by the restriction of Φ .

Consider the following sequence and its adjoint:

$$\begin{aligned} 0 &\rightarrow F_{n-1}/F_1 \rightarrow E/F_1 \rightarrow E/F_{n-1} \rightarrow 0 \\ 0 &\rightarrow (\widetilde{E/F_{n-1}})^* \rightarrow (\widetilde{E/F_1})^* \rightarrow (\widetilde{F_{n-1}/F_1})^* \rightarrow 0 \end{aligned}$$

by replacing $(\widetilde{E/F_{n-1}})^*$ and $(\widetilde{E/F_1})^*$ with F_1 and F_{n-1} we obtain:

$$0 \rightarrow F_1 \xrightarrow{\varphi} F_{n-1} \xrightarrow{\pi} (\widetilde{F_{n-1}/F_1})^* \rightarrow 0$$

for the uniqueness of F_1 , φ can only be (up to multiplication by a complex number) the inclusion $F_1 \subset F_{n-1}$ and hence $(\widetilde{F_{n-1}/F_1})^* \simeq (F_{n-1}/F_1)$. Note that π is the restriction of Φ to F_{n-1} , so the isomorphism is hermitian. \square

We can now prove the theorem.

Proof of theorem 2.1. We will prove the theorem by induction on the rank of the (a, b) -module. For rank 0 and 1 the theorem is obvious.

Suppose we proved the theorem for every rank $< n$ and let's prove it for rank n . Let find $F_1 \subset F_{n-1}$ of rank 1 and $n-1$ such that $(\widetilde{E/F_{n-1}})^* \simeq F_1$ and F_{n-1}/F_1 is hermitian. We can have different cases which are exhaustive:

- (i) We can find G , a normal elementary sub- (a, b) -module of E of parameter λ not equal to 0 or $1/2 \pmod{\mathbb{Z}}$. Then $\Phi(G)(G) = 0$ by remark 2.3 and we can apply lemma 2.2.

We still need to prove the induction step for (a, b) -modules whose only normal elementary sub- (a, b) -modules have parameter $\lambda = 0$ or $\lambda = 1/2$ modulo \mathbb{Z} .

- (ii) For $\lambda = 0$ or $\lambda = 1/2$ there are two distinct normal elementary sub- (a, b) -modules of parameter equal to $\lambda \pmod{\mathbb{Z}}$. We apply lemma 2.4.

The (a, b) -modules that were not included in the previous points have an unique normal elementary sub- (a, b) -module of parameter equal to $1/2$ modulo \mathbb{Z} and an unique normal elementary sub- (a, b) -module with an integer value of the parameter.

- (iii) There is only one normal elementary sub- (a, b) -module of parameter equal to $\lambda \pmod{\mathbb{Z}}$, where $\lambda = 0$ or $1/2$, but at least two quotients of a J-H sequence are of parameter equal to $\lambda \pmod{\mathbb{Z}}$. We apply lemma 2.6.

Only modules of rank at most 2 (one for each possible value of λ) still need to be checked.

- (iv) The rank of E is 2 and one quotient of a J-H sequence is equal to $0 \pmod{\mathbb{Z}}$, the other equal to $1/2 \pmod{\mathbb{Z}}$. By the classification of rank 2 modules this case is impossible. In fact with the notations of [Bar93]:

$$\begin{aligned} (\widetilde{E_\lambda \oplus E_\mu})^* &\simeq E_{-\lambda} \oplus E_{-\mu} \\ \check{E}_{\lambda, \mu}^* &\simeq E_{1-\lambda, 1-\mu} \end{aligned}$$

so if $\lambda = 0 \pmod{\mathbb{Z}}$ and $\mu = 1/2 \pmod{\mathbb{Z}}$ the (a, b) -module is not self-adjoint.

By induction hypothesis F_{n-1}/F_1 has a J-H composition series that verifies the theorem and by taking the inverse image by the canonical morphism $F_{n-1} \rightarrow F_{n-1}/F_1$ and adding 0 and E we find a J-H sequence of E that satisfies the theorem. \square

Since for an anti-hermitian form A we have $A(e, e) = 0$ for every $e \in E$, by using an anti-hermitian version of lemma 2.2 alone and proceeding by induction, we can prove theorem 2.1 in the anti-hermitian case.

We wish now to extend the result to all regular self-adjoint (a, b) -modules. We have proven in [Kar] that every regular (a, b) -module E can be decomposed into a direct sum of hermitian or anti-hermitian (a, b) -modules. We can hence prove the following theorem:

Theorem 2.7. *Let E be a self-adjoint regular (a, b) -module. Then it admits a self-adjoint Jordan-Hölder composition series.*

Proof. Let decompose E into

$$E = \bigoplus_{i=1}^m H_i$$

where m is an integer, while the H_i are either indecomposable self-adjoint or of the form $G \oplus \check{G}^*$, where G is indecomposable non self-adjoint (a, b) -module.

Each term of this sum admits a self-adjoint composition series. In fact if H_i is indecomposable self-adjoint, then it is hermitian or anti-hermitian. We can therefore apply the previous theorem 2.1.

On the other hand if H_i is the sum $G \oplus \check{G}^*$ of a module and its adjoint, we can easily find a self-adjoint Jordan-Hölder composition series. Take in fact any Jordan-Hölder series of G ,

$$0 = G_0 \subsetneq \cdots \subsetneq G_n = G.$$

and consider the adjoint series

$$0 = (\overline{G/G_n})^* \subsetneq (\overline{G/G_{n-1}})^* \subsetneq \cdots (\overline{G/G_0})^* = \check{G}^*.$$

Then the following composition series of $G \oplus \check{G}^*$ is self-adjoint:

$$\begin{aligned} 0 = G_0 \subsetneq G_1 \subsetneq \cdots \subsetneq G = G \oplus (\overline{G/G_n})^* \subsetneq G \oplus (\overline{G/G_{n-1}})^* \subsetneq \\ \cdots \subsetneq G \oplus (\overline{G/G_0})^* = G \oplus \check{G}^*. \end{aligned}$$

We will now prove the theorem on induction on m . The case $m = 1$ was already proven.

Suppose now $m \geq 2$ and let $E' := H_1$ and $F := \sum_{i=2}^m H_i$. We have therefore $E = E' \oplus F$, and E' and F are both self-adjoint. By the remark above we can find a self-adjoint composition series of E' :

$$0 = E'_0 \subsetneq \cdots \subsetneq E'_r = E'$$

while by induction hypothesis, we can find a self-adjoint composition series of F :

$$0 = F_0 \subsetneq \cdots \subsetneq F_s = F.$$

Then the following composition series is self-adjoint:

$$\begin{aligned} 0 = E'_0 \subsetneq E'_1 \subsetneq \cdots \subsetneq E'_{[r/2]} \subsetneq E'_{[r/2]} \oplus F_1 \subsetneq \cdots \subsetneq E'_{[r/2]} \oplus F_{[s/2]} [\cdots] \\ E'_{[(r+1)/2]} \oplus F_{[(s+1)/2]} \subsetneq E'_{[(r+1)/2]} \oplus F_{[(s+1)/2]+1} \subsetneq \cdots \subsetneq E'_{[(r+1)/2]} \oplus F \\ \subsetneq E'_{[(r+1)/2]+1} \oplus F \subsetneq \cdots \subsetneq E' \oplus F, \end{aligned}$$

where depending on the parity of r and s , $[\cdots]$ stands for

- (i) the $=$ sign if r and s are both even.
- (ii) the \subsetneq sign if one is even and the other odd.

(iii) the subsequence

$$\subsetneq E'_{[r/2]} \oplus F_{[(s+1)/2]} \subsetneq$$

This case needs a short verification. If r and s are odd, then the two central quotients of the series are isomorphic to $E'_{[(r+1)/2]}/E'_{[r/2]}$ and $F_{[(s+1)/2]}/F_{[s/2]}$. Since E'_i and F_i are self-adjoint series both quotients are self-adjoint (a, b) -modules of rank 1. They are therefore isomorphic to E_0 .

□

References

- [Bar93] Daniel Barlet. Theory of (a, b) -modules. i. In *Complex analysis and geometry*, Univ. Ser. Math., pages 1–43. Plenum, New York, 1993.
- [Bar97] D. Barlet. Théorie des (a, b) -modules. II. Extensions. In *Complex analysis and geometry (Trento, 1995)*, volume 366 of *Pitman Res. Notes Math. Ser.*, pages 19–59. Longman, Harlow, 1997.
- [Bel01] R. Belgrade. Dualité et spectres des (a, b) -modules. *J. Algebra*, 245(1):193–224, 2001.
- [Bri70] Egbert Brieskorn. Die Monodromie der isolierten Singularitäten von Hyperflächen. *Manuscripta Math.*, 2:103–161, 1970.
- [Kar] Piotr P. Karwasz. Hermitian (a, b) -modules and Saito’s “higher residue pairings”. available at <http://www.iecn.u-nancy.fr/~karwasz/research>.