



A general definition of influence between stochastic processes

Anne Gégout-Petit, Daniel Commenges

► To cite this version:

Anne Gégout-Petit, Daniel Commenges. A general definition of influence between stochastic processes. Lifetime Data Analysis, Springer Verlag, 2010, 16 (1), pp.33–44. 10.1007/s10985-009-9131-7. hal-00386649

HAL Id: hal-00386649

<https://hal.archives-ouvertes.fr/hal-00386649>

Submitted on 22 May 2009

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A general definition of influence between stochastic processes

Anne Gégout-Petit^{1,2} and Daniel Commenges^{2,3}

May 22, 2009

1 IMB, UMR 5251, INRIA Project CQFD, Talence, F33405, France

2 Université Victor Segalen Bordeaux 2, Bordeaux, F33076, France

3 INSERM, U 897, Bordeaux, F33076, France

Summary.

We extend the study of weak local conditional independence (WCLI) based on a measurability condition made by Commenges and Gégout-Petit (2009) to a larger class of processes that we call \mathcal{D}' . We also give a definition related to the same concept based on certain likelihood processes, using the Girsanov theorem. Under certain conditions, the two definitions coincide on \mathcal{D}' . These results may be used in causal models in that we define what may be the largest class of processes in which influences of one component of a stochastic process on another can be described without ambiguity. From WCLI we can construct a concept of strong local conditional independence (SCLI). When WCLI does not hold, there is a direct influence while

when SCLI does not hold there is direct or indirect influence. We investigate whether WCLI and SCLI can be defined via conventional independence conditions and find that this is the case for the latter but not for the former. Finally we recall that causal interpretation does not follow from mere mathematical definitions, but requires working with a good system and with the true probability.

Keywords: Causality; causal influence; directed graphs; dynamical models; likelihood process; stochastic processes.

1 Introduction

The issue of causality has attracted more and more interest from statisticians in recent years. An approach using the modelling of “potential outcome”, often called the counterfactual approach, has been proposed in the context of clinical trials by Rubin (1974) and further studied by Holland (1986) among others. The counterfactual approach has been extended to the study of longitudinal incomplete data in several papers and books (Gill and Robins, 2001; Robins et al., 2004; van der Laan and Robins, 2002). The counterfactual approach however has been criticised (Dawid, 2000; Geneletti, 2007). Another approach directly based on dynamical models has been developed, starting with Granger (1969) and Schweder (1970), and more recently developed using the formalism of stochastic processes, by Aalen (1987), Florens and Fougère (1996), Fosen et al. (2006) and Didelez (2007, 2008).

Recently we have given more development to the dynamical models approach (Commenges and Gégout-Petit, 2009) using the basic idea of the

Doob-Meyer decomposition proposed in Aalen (1987). We have proposed a definition of weak local independence between processes (WCLI) for a certain class of special semi-martingales (called class \mathcal{D}) which involves the compensator of the Doob-Meyer decomposition of the studied semi-martingale. Although it can be used in discrete time, this definition is especially adapted to continuous-time processes for which as we will see in section 4, definitions based on conventional conditional independence may fail. The aim of this paper is to give an even more general definition of WCLI, and conversely of direct influence. What we call direct influence of one component X_j on another component X_k of a multivariate stochastic process \mathbf{X} (noted $X_j \longrightarrow_{\mathbf{X}} X_k$) is that X_k is not WCLI of X_j (we use WCLI both as the name of the condition and as an adjective, that is the "I" may mean "independence" or "independent" according to the context). This concept of influence is a good starting point for defining *causal* influence (see section 5).

In the perspective of extending WCLI to a larger class of processes, we see two ways. The first one is to stay in the class of semi-martingales and try to be more general about the conditions. In particular we could use the triplet of the characteristics of a semi-martingales. For an exact definition of the characteristics of a semi-martingale, see Jacod and Shiryaev (2003). Roughly speaking, the characteristics of a semi-martingale are represented by the triplet (B, C, ν) where ν is the compensator of the jump part of the semi-martingale, B the finite variation part not included in ν , and C is the angle bracket process of the continuous martingale. The second way is to work with the likelihood of the process which is also tightly linked with the characteristics of the semi-martingale. In this paper we explore these two

ways, extending the WCLI definition to a very large class of processes that we call \mathcal{D}' , and showing that another definition of WCLI is possible by the use of likelihood processes. Another issue that we explore is the link between WCLI and analogous definitions based on conventional conditional independence; this angle of attack is closer to Granger (1969) proposal for time series. The scope of the paper is restricted to these mathematical definitions which may be useful for discussing causality issues. In the core of the paper, we address neither the philosophical nor the inferential issues; we discuss some of the philosophical issues in the last section.

In section 2, we recall the definition of WCLI, showing that it can be expressed in terms of the characteristics of the semi-martingales; this leads us to give a generalized definition of WCLI. We also recall the definition of strong local conditional independence (SCLI). In section 3, we propose another point of view based on the likelihood and we show the equivalence of definitions based on the Doob-Meyer decomposition and the property of certain likelihood processes under certain conditions. In section 4, we show that it is possible to define SCLI by conventional conditional independence, but that this approach falls short for WCLI. We conclude in section 5, where we recall the distinction between the mathematical definition of influences and the construction of a causal interpretation.

2 A generalization of WCLI

2.1 Notations and examples

Consider a filtered space $(\Omega, \mathcal{F}, (\mathcal{F}_t), P)$ and a multivariate stochastic process $\mathbf{X} = (\mathbf{X}_t)_{t \geq 0}$; \mathbf{X}_t takes values in \mathfrak{R}^m , and the whole process \mathbf{X} takes values in $D(\mathfrak{R}^m)$, the Skorohod space of all cadlag functions: $\mathfrak{R}_+ \rightarrow \mathfrak{R}^m$. We suppose that all the filtrations satisfy “the usual conditions”. We have $\mathbf{X} = (X_j, j = 1, \dots, m)$ where $X_j = (X_{jt})_{t \geq 0}$. We denote by \mathcal{X}_t the history of \mathbf{X} up to time t , that is \mathcal{X}_t is the σ -field $\sigma(\mathbf{X}_u, 0 \leq u \leq t)$, and by $(\mathcal{X}_t) = (\mathcal{X}_t)_{t \geq 0}$ the families of these histories, that is the filtration generated by \mathbf{X} . Similarly we shall denote by \mathcal{X}_{jt} and (\mathcal{X}_{jt}) the histories and the filtration associated to X_j . Let $\mathcal{F}_t = \mathcal{H} \vee \mathcal{X}_t$; \mathcal{H} may contain information known at $t = 0$, in addition to the initial value of \mathbf{X} . We shall consider the class of special semi-martingales in the filtration (\mathcal{F}_t) . We denote by (B, C, ν) the characteristics of the semi-martingale \mathbf{X} under probability P , by M_j the martingale part of X_j , and by M_j^c the continuous part of this martingale. We denote by (B^k, C^k, ν^k) the characteristics of the semi-martingale X_k under probability P .

Let us recall the definition of WCLI and see on examples how it involves the characteristics of the semi-martingale at hand. In our previous work (Commenges and Gégout-Petit, 2009) we have imposed the two following conditions bearing on the bracket process of the martingale M :

A1 M_j and M_k are orthogonal martingales, for all $j \neq k$;

A2 X_j is either a counting process or is continuous with a deterministic bracket process, for all j .

We call \mathcal{D} the class of all special semi-martingales satisfying **A1** and

A2. The class of special semi-martingales is stable by change of absolutely continuous probability (Jacod and Shiryaev, 2003, page 43) and this is also true for the the class \mathcal{D} .

Definition 1 [Weak conditional local independence (WCLI)] Let \mathbf{X} be in the class \mathcal{D} . X_k is WCLI of X_j in \mathbf{X} on $[r, s]$ if and only if $\Lambda_{kt} - \Lambda_{ks}$ is (\mathcal{F}_{-jt}) -predictable on $[r, s]$, where $\mathcal{F}_{-jt} = \mathcal{H} \vee \mathcal{X}_{-jt}$ and $\mathcal{X}_{-jt} = \bigvee_{l \neq j} \mathcal{X}_{-lt}$.

Generally, we assess WCLI on $[0, \tau]$, where τ is the horizon of interest, and if X_k is WCLI of X_j on $[0, \tau]$, we note $X_j \not\rightarrow_{\mathbf{X}} X_k$; in the opposite case we say that X_j directly influences X_k and we note $X_j \rightarrow_{\mathbf{X}} X_k$. A graph representation can be given, putting a directed edge when there is a direct influence from one node on another. If there is a directed path from X_j to X_k we say that X_j influences X_k and we note $X_j \rightarrow\rightarrow_{\mathbf{X}} X_k$. If X_j influences X_k but not directly influences it, then the influence is indirect. Inversely, if there is not directed path from X_j to X_k , we say that X_j does not influence X_k . We call this property strong local conditional independence (SCLI), saying that X_j is SCLI of X_k and we note $X_j \rightarrow\rightarrow_{\mathbf{X}} X_k$.

Let us see, using three examples, how the conditions **A1**, **A2** and the definition of WCLI can be expressed in terms of the characteristics of the semi-martingale X_k in the filtration (\mathcal{F}_t) .

Example 1: Let us consider a three-dimensional process $\mathbf{X}^3 \in \mathcal{D}$, $\mathbf{X}_t^3 = (X_{1t}, X_{2t}, X_{3t})$ defined by :

$$\begin{cases} X_{1t} &= \int_0^t f_1(X_{1s}, X_{2s}, X_{3s}) ds + M_{1t} \\ X_{2t} &= \int_0^t f_2(X_{1s}, X_{2s}, X_{3s}) ds + M_{2t} \\ X_{3t} &= \int_0^t f_3(X_{2s}, X_{3s}) ds + W_{3t} \end{cases} \quad (1)$$

where (M_1, M_2, W_3) are independent martingales and W_3 is a Brownian motion. For M_{jt} ($j = 1, 2$), we only assume that X_j is in the class \mathcal{D} . Since $M_3 = W_3$ is a Brownian motion, X_3 is a diffusion, so there is no jump. In this case the characteristics of X_3 are $B_t^3 = \int_0^t f_3(X_{1s}, X_{2s}, X_{3s})ds$ (the finite variation process), $C_t^3 = t$ (the bracket process of W_3) and $\nu_t^3 = 0$ (the compensator of the jump part), for all t . From the fact that f_3 does not involve X_1 , we directly see that X_3 is WCLI of X_1 in \mathbf{X}^3 . Mathematically, the ‘‘compensator’’ of the semi-martingale X_3 (called drift for a diffusion process) is equal to $\int_0^t f_3(X_{2s}, X_{3s})ds$ for all t , and is thus (\mathcal{F}_{-1t}) -predictable; this indeed corresponds to Definition 1 of WCLI. So, when X_k is a continuous semi-martingale, the WCLI condition involves the characteristic B^k of the semi-martingale X_k .

Example 2: Let us consider the following three-dimensional process $\mathbf{X}^3 \in \mathcal{D}$, $\mathbf{X}_t^3 = (X_{1t}, X_{2t}, X_{3t})$ defined by :

$$\begin{cases} X_{1t} &= \int_0^t f_1(X_{1s}, X_{2s}, X_{3s})ds + M_{1t} \\ X_{2t} &= \int_0^t f_2(X_{1s}, X_{2s}, X_{3s})ds + M_{2t} \\ X_{3t} &= \int_0^t \beta_3(X_{2s-}, X_{3s-})ds + M_{3t} \end{cases} \quad (2)$$

where (M_1, M_2, M_3) are independent martingales and X_3 is a counting process. We do not assume the form of M_{jt} ($j = 1, 2$). The WCLI relationships between the X_i 's are the same as in (1). X_3 is a counting process and in this case $B_t^3 = C_t^3 = 0$ and $\nu_t^3 = \int_0^t f_3(X_{2s}, X_{3s})ds$. The compensator of the counting process X_3 is \mathcal{X}_{-1t} -measurable which means that X_3 is WCLI of X_1 in \mathbf{X}^3 . So, when X_k is a counting process the WCLI condition involves the characteristic ν^k of the semi-martingale X_k .

Thus, the WCLI condition of Commenges and Gégout-Petit (2009) in-

volves the characteristic B when X_k is continuous and the characteristic ν when X_k is a counting process. In the framework of class \mathcal{D} , condition **A2** implies that these two characteristics are never simultaneously different from zero. In the following, we will consider processes for which B and ν may be both different from zero. We consider a process each component of which may have both a continuous and a jump part; such a process does not belong to \mathcal{D} .

Example 3:

$$\begin{cases} X_{1t} = \int_0^t f_1(X_{1s}, X_{2s}, X_{3s})ds + \int_0^t \sigma_{1t}dW_{1t} + \int_0^t \beta_1(X_{3s-})dN_{1s} \\ X_{2t} = \int_0^t f_2(X_{1s}, X_{2s})ds + \int_0^t \sigma_{2t}dW_{2t} + \int_0^t \beta_2(X_{2s-}, X_{3s-})dN_{2s} \\ X_{3t} = \int_0^t f_3(X_{2s}, X_{3s})ds + \int_0^t \sigma_{3t}dW_{3t} + \int_0^t \beta_3(X_{2s-}, X_{3s-})dN_{3s} \end{cases} \quad (3)$$

where the W_i 's are independent Brownian motions, the N_j 's are independent Poisson Processes with intensity 1 independent of the W_i 's. We suppose that the σ_{jt} 's are deterministic function of t , with $\sigma_{jt} > 0 \forall t$. It is clear that \mathbf{X} does not belong to class \mathcal{D} . However, the three characteristics of the semi-martingale X_3 are $B_t^3 = \int_0^t f_3(X_{2s}, X_{3s})ds$, $C_t^3 = \int_0^t \sigma_{3s}ds$ and $\nu_t^3 = \int_0^t \beta_3(X_{2s-}, X_{3s-})ds$. So, B_t^3 and ν_t^3 are (\mathcal{F}_{-1t}) -predictable: this will be the conditions of our new WCLI available for a larger class of semi-martingales.

2.2 Generalized definition of WCLI

We use the notations of the beginning of the section. We shall assume two conditions on \mathbf{X} :

A1 M_j and M_k are square integrable orthogonal martingales, for all $j \neq k$.

Under assumption **A1**, the jumping parts of the martingales M_j and M_k are orthogonal. Moreover, the characteristic C of \mathbf{X} (the angle bracket

of the continuous part of the martingale) is a diagonal matrix. Indeed by definition of orthogonality of semi-martingales, $C_{ij} = \langle M_i^c, M_j^c \rangle = 0$ for all $1 \leq i, j \leq m$; we note $C^k = C_{kk}$.

A2' C^j is deterministic for all j .

We call \mathcal{D}' the class of all special semi-martingales satisfying **A1** and **A2'**. In fact, **A1** and **A2'** could be merged into a single compact assumption: the characteristic C of \mathbf{X} is a deterministic diagonal matrix. \mathcal{D}' is stable by change of absolutely continuous probability (C does not change with the probability). \mathcal{D}' is a very large class of processes: it includes random measures, marked point processes, diffusions and diffusions with jumps. .

Definition 2 (Weak conditional local independence (WCLI)) *Let \mathbf{X} be in the class \mathcal{D}' . X_k is WCLI of X_j in \mathbf{X} on $[r, s]$ if and only if the characteristics B^k and ν^k are such that $B_{kt} - B_{kr}$ and $\nu_{kt} - \nu_{kr}$ are (\mathcal{F}_{-jt}) -predictable on $[r, s]$. Equivalently we can say that X_k has the same characteristic triplet (B^k, C^k, ν^k) in (\mathcal{F}_t) and in (\mathcal{F}_{-jt}) on the interval $[r, s]$.*

This new definition coincides with that of Commenges and G out-Petit (2009) for the class $\mathcal{D} \subset \mathcal{D}'$.

3 Link with the likelihood

We consider again the three examples above with a particular attention to the likelihood of the process X_3 . In Example 1, we apply Girsanov theorem to change the current probability using the density process (Z_{1t}^{P/P_0}) :

$$Z_{1t}^{P/P_0} = \exp \left(\int_0^t f_3(X_{2s}, X_{3s}) dX_{3s} - \frac{1}{2} \int_0^t (f_3(X_{2s}, X_{3s}))^2 ds \right). \quad (4)$$

Under the assumption $E_P[\exp(\frac{1}{2} \int_0^{+\infty} (f_3(X_{2s}, X_{3s}))^2 ds)] < +\infty$, the process $Z_{1t}^{P/P_0} = \frac{1}{Z_{1t}^{P_0/P}}$ is a P -martingale and the probability P_0 defined by $\frac{dP_0}{dP}|_{\mathcal{F}_t} = Z_{1t}^{P_0/P}$ for all $t \geq 0$ is equivalent to P on each \mathcal{F}_t ; moreover, under P_0 , X_3 is a Brownian motion independent of (M_1, M_2) .

In Example 2 we consider the density process (Z_{2t}^{P/P_0}) :

$$Z_{2t}^{P/P_0} = \prod_{s \leq t} (\beta_3(X_{2s-}, X_{3s-}))^{\Delta X_{3s}} \exp \left(\int_0^t \beta_3(X_{2s-}, X_{3s-}) ds \right). \quad (5)$$

Under technical conditions given in Lépingle et Mémin (1978), it defines a new probability P_0 such that under P_0 , X_3 is a homogeneous Poisson process.

In Example 3, we consider the density process (Z_{3t}^{P/P_0}) :

$$Z_{3t}^{P/P_0} = \prod_{s \leq t} (\beta_3(\dots))^{\Delta X_{3s}} \exp \left(\int_0^t \frac{f_3(\dots)}{\sigma_{3t}} dX_{3s} + \int_0^t (\beta_3(\dots) - \frac{1}{2} f_3^2(\dots)) ds \right), \quad (6)$$

where $\beta_3(\dots)$ stands for $\beta_3(X_{2s-}, X_{3s-})$ and $f_3(\dots)$ for $f_3(X_{2s}, X_{3s})$. Under technical conditions given in Lépingle et Mémin (1978), P_0 is well defined, and X_3 is the sum of a Brownian motion with variance σ_{3t}^2 and a homogeneous Poisson process under P_0 .

In the three cases, we see that the likelihood processes Z_{jt}^{P/P_0} are \mathcal{X}_{-1t} -measurable. That is, the \mathcal{X}_{-1t} -measurability of the characteristics of X_k implies the \mathcal{X}_{-1t} -measurability of the likelihood process. We want to use a measurability condition on the likelihood process for a new definition of WCLI. We could say that " X_k is weakly locally independent of X_j in \mathbf{X} if the likelihood of X_k is $\mathcal{F}_{-jt} = \mathcal{H} \vee \mathcal{X}_{-jt}$ -measurable". However, we must be cautious because the likelihood is a likelihood ratio between two probabilities, and these probabilities give not only the distribution of X_k but that of the whole process \mathbf{X} . So, the reference measure P_0 must meet some assumptions

given in the definition of this new condition.

Definition 3 [*Likelihood-based weak conditional local independence (LWCLI)*]

Let $\mathbf{X} = (X_j, j = 1, \dots, m)$ be in the class \mathcal{D}' .

1. Suppose the existence of a probability P_0 such that (i) $P \ll P_0$, (ii) the characteristics of the semi-martingales X_i 's with $i \neq k$ are the same under P and P_0 and (iii) the P_0 -characteristics (B_0^k, C_0^k, ν_0^k) of the semi-martingale X_k are deterministic. We say that X_k is LWCLI of X_j in \mathbf{X} on $[0, t]$ if and only if the likelihood ratio process $Z_t^{P/P_0} = \mathcal{L}_{\mathcal{F}_t}^{P/P_0}$ is (\mathcal{F}_{-jt}) -measurable on $[0, t]$. We have denoted $\mathcal{F}_{-jt} = \mathcal{H} \vee \mathcal{X}_{-jt}$ and $\mathcal{X}_{-jt} = \bigvee_{l \neq j} \mathcal{X}_{-lt}$.
2. X_k is LWCLI of X_j in \mathbf{X} on $[r, s]$ if and only if the process $\frac{Z_t^{P/P_0}}{Z_r^{P/P_0}}$ is (\mathcal{F}_{-jt}) -predictable for all $t \in [r, s]$ for all the probabilities P_0 as above.

Let us comment the definition and the conditions imposed to the reference probability P_0 in this definition in the following remarks.

Remark 1. In the examples of this section, we have constructed P_0 by a change of probability. In the definition we are in a context of likelihood writing and we suppose the existence of a "good" reference probability.

Remark 2. We want that the likelihood concerns X_k only in a certain sense given by (ii). It was the case in the three examples considered above. (ii) is true for instance if $\langle Z^{P/P_0}, M^i \rangle = 0$ for all $i \neq k$. Suppose for instance that M^k is not orthogonal to M^j for a $j \neq k$ (assumption **A1** not true) then it is certainly not possible to find a probability which verifies (ii).

Remark 3. We do not want that the "relation" between X_k and X_j under P is hidden by the same relation under P_0 . To make such a condition

explicit, the framework of semi-martingales is again very useful. This condition involves the characteristics (B_0^k, C_0^k, ν_0^k) of X_k under P_0 . They must be deterministic. So (iii) is linked to assumption **A2'** because C^k does not change with the probability and remains deterministic whatever the absolute continuous change of probability. We emphasize that if **A2'** fails, that is the bracket C^k is not deterministic under P , we will never find a probability P_0 which verifies (iii). Moreover, in the examples given above, the process X_3 satisfies the property of independent increments under P_0 . In the case of semi-martingales, this property is verified if and only if the triplet (B, C, ν) is deterministic under P_0 . This is exactly the condition (iii) of definition 3.

Remark 4. If **A1** is not satisfied, this means that at least two components of \mathbf{X} have a common part of martingale: they are driven by the same noise but we can not speak of influence of one on the other. Condition **A2'** is different: even if C^k is driven by another component of \mathbf{X} we will never detect it by a measurability condition because the characteristics C^k is always \mathcal{X}_k -measurable.

LWCLI seems to be more general than WCLI. When X_k is a diffusion with jumps (see Jacod and Shiryaev, 2003, Definition III. 2.18), we can take for P_0 the probability under which X_k is the sum of a Brownian motion and a standard Poisson process with parameter $\lambda = 1$. However, except this standard case, the conditions required on P_0 are not easy to characterize. In the good cases, we have an explicit computation of the likelihood ratio process Z_t^{P/P_0} as function of the characteristics of X_k in the probabilities P and P_0 . This result allows us to lay down the following result:

Proposition 1 *Suppose that \mathbf{X} is a m -dimensional diffusion with bounded*

jumps process satisfying the uniqueness in law conditions and which belongs to the class \mathcal{D}' and suppose the existence of a probability P_0 satisfying the assumptions of the definition (3) then WCLI and LWCLI are equivalent.

Proof: the assumptions of Proposition 1 guarantee the explicit computation of Z_t^{P/P_0} as a function of the characteristics of X_k under P and P_0 (Jacod and Shiryaev 2003: Theorem III. 5.19) and the uniqueness of probability P (Jacod and Shiryaev, 2003: Theorems III. 2.32 and III. 2.33) under which \mathbf{X} has the given characteristics. Under these assumptions, the component X_k is of the form:

$$dX_{kt} = f_k(t, \mathbf{X}_t)dt + \sigma_k(t)dW_{kt} + \beta_k(s, \mathbf{X}_{s-}, z)(p(dt, dz) - q(dt, dz)),$$

where $p(dt, dz)$ is a Poisson random measure with intensity and $q(dt, dz) = dt \otimes F(dx)$ (F is a positive σ -additive measure on $(\mathcal{R}, B(\mathcal{R}))$). So, $B^k = \int_0^t f_k(s, \mathbf{X}_s)ds$, $\nu^k = \int_0^t \beta_k(s, \mathbf{X}_{s-}, z)q(dt, dz)$ and $C^k = \int_0^t \sigma_{3s}^2 ds$ are the characteristics of X_k under P . The likelihood ratio being a function of $(B^k, C^k, \nu^k, B_0^k, C_0^k, \nu_0^k)$, it is obvious that WCLI implies LWCLI. Let us prove the reverse: let X_k be LWCLI of X_j in \mathbf{X} . If B_t^k or ν_t^k were not (\mathcal{F}_{-jt}) -measurable, then Z_t^{P/P_0} would no longer be (\mathcal{F}_{-jt}) -measurable: this contradicts LWCLI !

4 WCLI and SCLI via conditional independence of filtrations

Heuristically, we can state the non-influence of X_j on X_k by saying that, on the basis of the information at time t , we do not need to know $X_{ju}, u < t$ to predict X_k at t , or after t . In the previous sections, we have expressed this

intuition in terms of measurability of certain processes (compensator and likelihood process). Granger (1969), working with stationary time series (in discrete time) proposed a criterion based on the variance of the prediction. Eichler and Didelez (2009) gave a clear definition of Granger non-causality in a more general setting, although still for stationary time series, and they expressed it in terms of conditional independence. They distinguish between “strong Granger-non causality” and “contemporaneous independence”. With our notations, strong Granger-non causality can be expressed as:

$$X_{ks} \perp\!\!\!\perp_{\mathcal{F}_{-jt}} \mathcal{X}_{jt}, t = 0, 1, \dots; s = t + 1, t + 2, \dots, t + h, \quad (7)$$

where h is called “horizon”.

In continuous time it is also tempting to define WCLI and SCLI in terms of conditional independence. Didelez (2008) heuristically proposed the following definition for WCLI when \mathbf{X} is a counting process:

$$X_{kt} \perp\!\!\!\perp_{\mathcal{F}_{-jt-}} \mathcal{X}_{jt-}, 0 \leq t \leq \tau. \quad (8)$$

This formula attempts to express non-influence by requiring that X_{kt} is independent of the past of X_j given the past of the other components of \mathbf{X} . However as remarked in Commenges and Gégout-Petit (2009), this condition is void in general when we consider processes in continuous time. Because conditional independence is defined via conditional probability, and in general, events of \mathcal{X}_{kt} have conditional probabilities equal to one or zero given \mathcal{X}_{kt-} , the condition always holds.

We now propose a rigorous definition of non-influence in continuous time based on conventional conditional independence. Moreover, since independence is defined in probability theory in terms of sigma-fields, we can state

this property directly in terms of the sigma-fields $\mathcal{X}_{jt}, j = 1, \dots, m$, without specifying stochastic processes (as argued in Commenges, 2009, a representation of statistical models in terms of sigma-fields or filtrations is more intrinsic than in terms of random variables or stochastic processes). For simplicity we define it on $(0, \tau)$.

Definition 4 *Filtration-based strong conditional local independence (FSCLI)*

Let $(\mathcal{X}_{jt}), j = 1, \dots, m$ be filtrations, $\mathcal{X}_t = \vee_j \mathcal{X}_{jt}$; $\mathcal{F}_t = \mathcal{H} \vee \mathcal{X}_t$ and $\mathcal{X}_{-jt} = \vee_{l \neq j} \mathcal{X}_{-lt}$, $\mathcal{F}_{-jt} = \mathcal{H} \vee \mathcal{X}_{-jt}$. We say that filtration (\mathcal{X}_{kt}) is FSCLI of (\mathcal{X}_{jt}) in \mathcal{F}_t if and only if:

$$\mathcal{X}_{k\tau} \perp\!\!\!\perp_{\mathcal{F}_{-jt}} \mathcal{X}_{jt}, 0 \leq t \leq \tau. \quad (9)$$

Proposition 2 *Suppose that \mathbf{X} is the unique m -dimensional solution of a given stochastic differential equation with bounded jumps process and which belongs to the class \mathcal{D}' , then FSCLI defined on the filtrations generated by the components of \mathbf{X} and SCLI are equivalent.*

Proof. For a given $X_k \in \mathbf{X}$, denote by $An(k) = \{l_{1k}, \dots, l_{nk}\}$ the set of all the indices l such that X_l is an ancestor of X_k . The assumptions imply that $X_{An(k)}$ is also the unique solution a stochastic differential equation with bounded jumps process generated by the Brownian process $\mathbf{W}_{An(k)} = (W_{l_1}, \dots, W_{l_{nk}})$ and the set of orthogonal Poisson measures $\mathbf{P}_{An(k)} = (p_{l_1}, \dots, p_{l_{nk}})$. Moreover for each t , $X_{An(k)t}$ is a functional of $(\mathbf{W}_{An(k)s}, \mathbf{P}_{An(k)s}, s \leq t)$. If $t \leq \tau$, $X_{An(k)\tau}$ is a functional of $X_{An(k)t}$ and of the processes $\mathbf{W}_{An(k)}^{(t, \cdot)}, \mathbf{P}_{An(k)}^{(t, \cdot)}$ defined by $\mathbf{W}_{An(k)}^{(t, s)} = (\mathbf{W}_{An(k)s} - \mathbf{W}_{An(k)t}), \mathbf{P}_{An(k)}^{(s, t)} = (\mathbf{P}_{An(k)s} - \mathbf{P}_{An(k)t}), t \leq s \leq \tau$. By the independent increments property of the Brownian motion and of the Poisson process, if we denote $\sigma_k^{(t, \tau)} = \sigma((\mathbf{W}_{An(k)}^{(t, s)}, \mathbf{P}_{An(k)}^{(t, s)}, t \leq$

$s \leq \tau$), we have $\sigma_k^{(t,\tau)} \perp\!\!\!\perp \mathcal{F}_t$. Suppose that $X_j \rightarrow\!\!\rightarrow \mathbf{X} X_k$, it implies that $X_j \not\rightarrow\!\!\rightarrow \mathbf{X} X_{An(k)}$ and that $X_{An(k)t}$ is \mathcal{F}_{-jt} -measurable. Using the previous remark and the standard properties of conditional expectation (Jacod, Protter exercise 23.7), for $t \leq s \leq \tau$, we have that $E[f(X_{ks})|\mathcal{F}_t] = E[f(X_{ks})|\mathcal{F}_{-jt}] = G(X_{kt})$ with

$$G(x) = E[f(F(x, \mathbf{W}_{An(k)}^{(t,\cdot)}, \mathbf{P}_{An(k)}^{(t,\cdot)}) | X_{kt} = x)].$$

We have proved $SCLI \Rightarrow FSCLI$.

As for the converse, (9) implies that X_k is perfectly defined by a differential equations with jumps which does not involve the component X_j and thus $X_j \rightarrow\!\!\rightarrow \mathbf{X} X_k$.

Remark 5. We could also define an ‘‘horizon’’ $h > 0$ for FSCLI in a way analogous to formula (7).

$$\mathcal{X}_{k,t+h} \perp\!\!\!\perp_{\mathcal{F}_{-jt}} \mathcal{X}_{jt}, \quad 0 \leq t \leq \tau - h. \quad (10)$$

If we make this horizon tend toward zero the FSCLI requirement tends (heuristically) to the WCLI requirement. In continuous time however, considering an infinitely small h would lead to definition (8), which as already mentioned is void. We conclude that WCLI cannot be rigorously defined by conditional independence; we need the measurability-based definition.

Remark 6. Didelez and Eichler (2009) also defined a concept of contemporaneous independence as:

$$X_{ks} \perp\!\!\!\perp_{\mathcal{F}_t} X_{js}, \quad t = 0, 1, \dots; \quad s = t + 1. \quad (11)$$

For the same reason as for WCLI, contemporaneous independence cannot be defined in continuous time via conventional conditional independence,

because $X_{kt} \perp\!\!\!\perp_{\mathcal{F}_{t-}} X_{jt}$ in void in general. However contemporaneous independence in continuous time might be identified with the assumption of orthogonal martingales.

Remark 7. If the time parameter is discrete, then FSLI defined by (9) is identical to strong Granger-non causality for all horizon. Moreover FSLI defined by (10) for horizon $h = 1$, Granger-non causality for horizon $h = 1$ and WCLI are identical.

5 Discussion and conclusion

We have generalized the definition of WCLI to a larger class of processes and we have proposed another definition through likelihood ratio processes. Under certain conditions the two definitions are equivalent. We have also made the link with definitions based on conventional conditional independence: SCLI can be defined this way but in continuous time WCLI cannot. These results may be used for developing causal models. By definition, there are direct influences where WCLI does not hold: if X_k is not WCLI of X_j , then X_j directly influences X_k in \mathbf{X} . It is to be noted that influence is not a simple lack of (even conditional) independence. WCLI is a dynamical concept which differs markedly from conventional independence concepts. Essentially because it is dynamic, it is not symmetric, while conventional independence is. We can have X_k WCLI of X_j and X_j not WCLI of X_k . This provides a rich set of relationships between two components of a stochastic process \mathbf{X} . We have three possibilities for the influence of X_j on X_k : $X_j \longrightarrow_{\mathbf{X}} X_k$ (direct influence), $X_j \not\rightarrow_{\mathbf{X}} X_k$ (no influence), $X_j \rightarrow\rightarrow_{\mathbf{X}} X_k$ and $X_j \not\rightarrow_{\mathbf{X}} X_k$

(indirect influence). There are also three possibilities of influence of X_k on X_j . Thus, there are nine possibilities for describing the relationship between two components of a stochastic process. Of course it would be of great interest to quantify these influences. Interesting work has been done in this direction, in the time-series framework, by Eichler and Didelez (2009).

The multivariate stochastic process framework is a general framework which incorporates a major feature of causal relationship: time. Thus it is a natural framework to formalize causality in statistics. It is important to know which is the most general class of stochastic processes in which we can work for developing such a formalisation. The class \mathcal{D}' seems to be this class.

However this only describes a mathematical framework which is well suited for formalizing causality. This is why in this paper we speak of influence rather than causal influence. A causal interpretation needs an epistemological act to link the mathematical model to a physical reality. In particular, WCLI is dependent on a filtration and a probability. Commenges and Gégout-Petit (2009) emphasized that the choice of the filtration is related to the choice of the physical system and assumed that there is a true probability, P^* , according to which the events of the universe are generated. Causal influences were defined as influences in a good (or perfect) system and under the true probability.

6 References

Aalen O (1987) Dynamic modelling and causality. Scand. Actuarial J., 1987: 177-190.

- Aalen O (1978) Non parametric inference for a family of counting processes. *The Annals of Statistics* 6: 701-726.
- Aalen O, Frigessi A (2007) What can statistics contribute to a causal understanding? *Scand. J. Statist.* 34: 155-168.
- Commenges,D. (2009) Statistical models, conventional, penalized and hierarchical likelihood. *Statistics Surveys*, 3: 1-17.
- Commenges D, Gégout-Petit A (2009) A general dynamical statistical model with causal interpretation. *J. Roy. Statist. Soc. B*, 71: 1-18.
- Dawid AP (2000) Causal inference without counterfactuals. *J. Am. Statist. Ass.* 95: 407-448.
- Didelez V (2007) Graphical models for composable finite Markov processes, *Scand. J. Statist.* 34: 169-185.
- Didelez, V (2008) Graphical models for marked point processes, *J. Roy. Statist. Soc. B* 70: 245-264.
- Eichler M, Didelez V (2009) On Granger causality and the effect of interventions in time series. Research Report 09-01, Statistics Group, University of Bristol, 2009 <http://www.maths.bris.ac.uk/~maxvd/>
- Florens J-P, Fougère D (1996) Noncausality in continuous time. *Econometrica* 64: 1195-1212.
- Fosen J, Ferkingstad E, Borgan O, Aalen, OO (2006) Dynamic path analysis: a new approach to analyzing time-dependent covariates. *Lifetime Data Analysis* 12: 143-167.

- Geneletti S (2007) Identifying direct and indirect effects in a non-counterfactual framework. *J. Roy. Statist. Soc. A* 69: 199-215.
- Gill RD, Robins J (2001) Causal inference for complex longitudinal data: the continuous case. *Ann. Statist.* 29: 1785-1811.
- Granger CWJ (1969) Investigating causal relations by econometric models and cross-spectral methods. *Econometrica* 37: 424-438.
- Holland P. (1986) Statistics and Causal Influence. *J. Am. Statist. Ass.* 81: 945-960.
- Jacod J, Protter P, (2000) *Essential in probability*, Springer-Verlag.
- Jacod J, Shiryaev AN (2003) *Limit Theorems fo Stochastic Processes*, Berlin: Springer-Verlag.
- Lépingle D, Mémin J (1978) Sur l'intégrabilité uniforme des martingales exponentielles. *Z. Wahrsch. Verw. Gebiete* 42: 175–203.
- Robins J, Lok J, Gill R, van der Vaart, A (2004) Estimating the causal effect of a time-varying treatment on time-to-event using structural nested failure time models. *Statistica Neerlandica* 58: 271-295.
- Robin J., Scheines R, Spirtes P, Wasserman L (2003) Uniform consistency in causal inference. *Biometrika* 90: 491-515
- Rubin DB (1974) Estimating Causal Effects of Treatments in Randomized and Nonrandomized Studies. *Journal of Educational Psychology* 66: 688-701.

Schweder T (1970) Composable Markov processes. J. Appl. Probab. 7: 400-410.

van der Laan M, Robins J (2002) Unified methods for censored longitudinal data and causality, Springer, New-York.