



The Membrane Inclusions Curvature Equations

Jean-Charles Faugère, Hering Milena, Phan Jeff

► To cite this version:

Jean-Charles Faugère, Hering Milena, Phan Jeff. The Membrane Inclusions Curvature Equations. [Research Report] lip6.2000.008, LIP6. 2000. hal-02548285

HAL Id: hal-02548285

<https://hal.archives-ouvertes.fr/hal-02548285>

Submitted on 20 Apr 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

The Membrane Inclusions Curvature Equations

Jean-Charles Faugère* Milena Hering† Jeff Phan‡

*LIP6/CNRS Université Paris 6 E-mail: *jcf@calfor.lip6.fr*

†Cambridge University, UK E-mail: *msch2@cam.ac.uk*

‡University of California, Berkeley E-mail: *phan@math.berkeley.edu*

January 31, 2000

Abstract

We examine a system of equations arising in biophysics whose solutions are believed to represent the stable positions of N conical proteins embedded in a cell membrane. Symmetry considerations motivate two equivalent reformulations of the system which allow the complete classification of solutions for small $N < 13$. The occurrence of regular geometric patterns in these solutions suggests considering a simpler system, which leads to the detection of solutions for larger N up to 280. We use the most recent techniques of Gröbner bases computation for solving non linear systems.

1 Introduction

Both the shapes and positions of proteins which are embedded in a cell membrane can influence their biological function. It is the interaction between the proteins which dictates how they become arranged, but little is known about this interaction and its exact cause is uncertain. However, for conical proteins, a likely explanation is the bending of the membrane caused by the proteins. Specifically, an embedded conical protein induces a curvature in the two dimensional membrane which influences the positions of neighboring proteins. There is an energy associated to this curvature and the proteins will tend to arrange themselves so as to minimize this energy. Recent work in [KJG98] shows that any minimum energy arrangement is a zero energy arrangement. Furthermore, if z_i is the position of the i th protein using complex coordinates, it was also shown that the energy at the i th protein is a constant multiple of $|f_i(z_1, \dots, z_N)|^2$ where

$$f_i(z_1, \dots, z_N) = \sum_{\substack{j=1 \\ j \neq i}}^N \frac{1}{(z_i - z_j)^2} = 0.$$

Therefore the N proteins are at equilibrium if and only if (z_1, \dots, z_N) is a solution to the Membrane Inclusions Curvature Equations, or MICE:

$$f_i(z_1, \dots, z_N) = 0, \quad i = 1, \dots, N. \quad (1)$$

For brevity, we refer to the N -th system of equations as S_N .

One possible application of knowing how these proteins arrange themselves is to deduce the form of proteins by examining the shapes they form. In this case, if they arrange themselves according to our solutions it is very likely that they are conical. Determining the shapes of proteins is still an unsolved problem in biology.

Gröbner bases are used to find the solutions of S_N for several N . In section 2, we review the most efficient algorithms for computing Gröbner bases and their implementations. Direct application of these algorithms gives all the solutions of the problem for $N < 7$ and is described in section 3. Because the difficulty of computing Gröbner bases increases rapidly with respect to the complexity of the input equations, it is necessary to reformulate the system before most of the computations will successfully terminate. Two reformulations of S_N into equivalent systems are given in section 4. The first reformulation employs an algorithm for converting the numerators of the S_N equations into symmetric polynomials, which are then expressed in terms of the elementary symmetric functions prior to computing. The second reformulation uses a differential equation describing the minimum polynomial for the coordinates of a solution and gives directly a system already formulated using the elementary symmetric functions. Both reformulations can be used jointly to decrease the computation time. Finally, we consider a much simplified system obtained from S_N by limiting our search to those solutions which have a certain geometric regularity to them; namely, we look for solutions whose coordinates form concentric rings of regular polygons. While this last approach does not detect all solutions, for a given N , it does allow many to be found.

Our main result is a complete classification of the solutions for small values for N :

Theorem 1.1 *There are no solutions for $N \leq 12$ except for $N = 5$ (finite number of solutions) and $N = 8$ (S_8 form a 1 dimensional variety).*

The proof of this theorem is included in sections 3 and 4. For larger values of N we have only a partial result:

Theorem 1.2 *There exist solutions to S_N for $N = 5, 8, 16, 21, 33, 37, 40, 56, 65, 85, 119, 133, 161, 175, 208, 225, 261$ and 280 . Moreover the number of solutions for S_{16} and S_{21} is infinite.*

We explained in section 5 how we find this list of “regular solutions”.

2 Tools for solving polynomial equations

We now review some major algorithms for solving multivariate polynomial systems. The reader is also referred to [Dav93, Bec93, CLO92, CLO98] for a more detailed introduction.

Let $\mathbb{Q}[x_1, \dots, x_n]$ be the polynomial ring with rational coefficients, F a finite list of equations and I the ideal generated by F .

The main tools we use are Gröbner bases [Buc65, Buc70, Buc79, Buc85]. We recall that, in general, when the number of equations equals the number of variables the shape of the Gröbner basis G for a lexicographical ordering is the following:

$$\begin{cases} h_n(x_n) \\ x_{n-1} = h_{n-1}(x_n) \\ \dots \\ x_1 = h_1(x_n) \end{cases}$$

where all the h_i are univariate polynomials. Of course the shape of a lexicographical Gröbner basis is not always so simple but it will always be the case in this paper (except one very easy non zero dimensional system). From this Gröbner basis it is rather easy to compute numerically all the complex roots: we first solve numerically the first equation [DG99], and we find z_1, \dots, z_N a guaranteed approximation of all the complex roots of h_n . Then we substitute these values into the other coordinates.

Even if all the algorithms for computing Gröbner bases do not depend on a specific order it is well known [Fau93] that it is more efficient to compute first a Gröbner basis for a Degree Reverse Lexicographical ordering and then change the ordering with a specific algorithm. In this paper we have used a standard implementation of the Buchberger algorithm and the FGLM algorithm in Singular [Gre99] for easy cases. When the degree of the univariate is big > 500 we have used:

- the F_4 [Fau99] algorithm for computing a DRL groebner basis.
- the F_2 [Fau94] algorithm to change the ordering. For the bigger computations we found that the dimension of $\mathbb{Q}[x_1, \dots, x_n]/I$ is bigger than 10^6 !

These two algorithms are implemented in an experimental software called FGb [Fau].

For generating the input equations we have used the Maple [Cha91] computer algebra system.

3 First experiments

First, we observe that the set of solutions to S_N is invariant under translation and multiplication by complex scalars. These considerations allow us to change coordinates so that $z_N = 0$ and $z_{N-1} = 1$.

Since the f_i in the system S_N are rational functions we need to transform the system into a polynomial system. In order to avoid "parasite" solutions, where $z_i = z_j$ for some $i \neq j$, we introduce a new variable u and let P_i be the numerator of each f_i in S_N . That is to say

$$P_i(z_1, \dots, z_N) = \sum_{j \neq i} \prod_{k \neq i, j} (z_i - z_k)^2 = 0, \quad i = 1, \dots, N \quad (2)$$

$$S'_N \begin{cases} u \prod_{i=1}^N \prod_{j=i+1}^N (z_i - z_j) = 1 \\ P_i(z_1, \dots, z_N) = 0 \quad i = 1, \dots, N \\ z_1 = 0 \\ z_2 = 1 \end{cases}$$

Proposition 3.1 *There is no solution for $N \leq 4$ and $N = 6$. The only solution for S_5 is a regular pentagon.*

Proof For $N \leq 5$ it takes less than 0.1 second to compute a lexicographic Gröbner basis with FGb on a PC Pentium II 300 Mhz. For $N < 5$ the Gröbner basis is $\{1\}$. For $N = 5$ we can factorize the univariate polynomial and find a decomposition into irreducible varieties: $V = V_1 \cup V_2 \cup V_3 \cup V_4 \cup V_5 \cup V_6$ and

$$V_1 = [z_3 - z_5^3 + z_5^2 - z_5, z_4 + z_5^2 - z_5, z_5^4 - z_5^3 + z_5^2 - z_5 + 1]$$

For any polynomial p in x_1, \dots, x_N and any permutation σ , set $\sigma.p = p(x_{\sigma(1)}, \dots, x_{\sigma(N)})$ and $\sigma(V) = \{\sigma(v) : \forall v \in V\}$. It is easy to check that

$$\begin{aligned} (z_4, z_5)V_1 &= V_6 \\ (z_3, z_5)V_1 &= V_3 \\ (z_3, z_4)V_1 &= V_2 \\ (z_3, z_4, z_5)V_1 &= V_5 \\ (z_3, z_5, z_4)V_1 &= V_4 \end{aligned}$$

Now we have

$$z_5^4 - z_5^3 + z_5^2 - z_5 + 1 = \frac{z_5^5 + 1}{z_5 + 1}$$

so that $z_5 = e^{\frac{\alpha i \pi}{5}}$ and we see that the only solution is the regular pentagon.

The case $N = 6$ is a little more difficult: the degree of the polynomial $u \prod_{i=1}^N \prod_{j=i+1}^N (z_i - z_j) = 1$ is $1 + \frac{N(N-1)}{2} = 16$ and so big that it does not help the Gröbner basis computation. In that case we can replace this condition by $uz_3z_4z_5z_6 = 1$ and it takes only 13.6 seconds to find $\{1\}$ with Fgb. \square

In conclusion the straightforward approach solves the problem for small N but leads to several problems:

- intermediate computations contain the same solution several times (action of the symmetric group), so the degree of the intermediate varieties are big.
- it is not easy to remove the parasite solutions $z_i = z_j$.

We have stopped the computation for $N = 7$ after 2000 seconds.

4 Using the symmetry

It is clear from 1 that if $(z_1, \dots, z_N) \in \mathbb{C}^N$ is a solution of S_N then $(z_{i_1}, \dots, z_{i_N})$ is also a solution of S_N for every possible permutation of (i_1, \dots, i_N) of $(1, \dots, n)$. Hence it is enough to compute the polynomial

$$f(X) = (X - z_1) \cdots (X - z_N) = X^N - e_1 X^{N-1} + \cdots + (-1)^N e_N$$

where the $e_i = e_i(z_1, \dots, z_N)$ are the elementary symmetric functions in z_1, \dots, z_N . In this paper we will say that f is solution of S_N . In general solving efficiently a polynomial system with symmetries is an open issue especially when the group is not the whole symmetric group. In our problem the solutions are invariant under the symmetric group but unfortunately f_i is not a symmetric polynomial in (z_1, \dots, z_n) but only in $\{z_j \mid j \neq i\}$. If we exchange the role of z_j and z_k then f_i remain unchanged while f_j becomes f_k and reciprocally.

$$z_j \longleftrightarrow z_k \quad f_i = f_i \text{ for } i \neq j, k \quad f_j \longleftrightarrow f_k$$

4.1 nilCoxeter algebra

Let e_r be the r th elementary symmetric function in N variables. For $\lambda = (\lambda_1, \dots, \lambda_r)$ let

$$m_\lambda = \sum z_{i_1}^{\lambda_1} \cdots z_{i_r}^{\lambda_r} \quad (3)$$

denote the monomial symmetric functions, where the sum ranges over all monomials whose exponent vector is equal to a permutation of λ . Solving S'_N is equivalent to finding a polynomial

$$\begin{cases} f = X^N - e_1 X^{N-1} + e_2 X^{N-2} - \cdots + (-1)^N e_N \\ f \text{ is squarefree.} \end{cases} \quad (4)$$

whose roots are a solution to S_N . For any polynomial p in z_1, \dots, z_N , set

$$\partial_i(p) = \frac{p(z_1, z_2, \dots, z_N) - p(z_i, z_2, \dots, z_{i-1}, z_1, z_{i+1}, \dots, z_N)}{z_1 - z_i}. \quad (5)$$

Let I_1 be the ideal generated by P_1, \dots, P_N , we define by induction

$$I_k = I_{k-1} : \left(\prod_{i_1 < i_2} (z_{i_1} - z_{i_2}) \right) \quad (6)$$

and $I = I_\infty$. Note that $P_i = (1, i).P_1$ for $1 \leq i \leq N$ and P_1 is symmetric in z_2, \dots, z_N .

Theorem 4.1 Define for $1 \leq i_1 < \cdots < i_{k+1} \leq N$

$$P_{i_1, \dots, i_k, i_{k+1}} = \frac{P_{i_1, \dots, i_k} - P_{i_1, \dots, i_{k-1}, i_{k+1}}}{z_{i_k} - z_{i_{k+1}}}$$

so that $P_{i_1, \dots, i_k} \in I_k$ and P_{i_1, \dots, i_k} is symmetric in z_{i_1}, \dots, z_{i_k} and in the complementary set of variables. Hence

$$H_k = \sum_{1 \leq i_1 < \dots < i_k \leq N} P_{i_1, \dots, i_k}$$

is a true symmetric function

The next theorem gives an efficient method for computing the H_i .

Theorem 4.2 For $1 \leq i_1 < \dots < i_k \leq N$

$$P_{i_1, \dots, i_k} = (1, i_1) \cdot (2, i_2) \cdot \dots \cdot (k, i_k) Q_k$$

where $Q_k = P_{1, 2, \dots, k}$ and we have

$$Q_k = \partial_k Q_{k-1}$$

The H_i were first computed in the monomial basis m_λ using code specifically written for this application in C++ in the small computer algebra system Gb; then the polynomials were expressed in the e_i basis using ACE [AS98], SF [J.98] and symmetrica. If we set $z_n = 0$ and $z_{n-1} = 1$ prior to computing the H_i , the reformulated system \tilde{S}_N consists of the polynomials $H_1, \dots, H_N, P_{N-1}, P_N$ in the variables e_1, \dots, e_{N-2} . It turns out that \tilde{S}_N is easier to solve: it takes 2 minutes to compute a Gröbner basis for $N = 10$ with FGb, while the calculation for S_7 was unsuccessfully stopped after 2000 seconds.

4.2 Harm Derksen formulation

Our second reformulation was found by Harm Derksen[Der99], and appeals to the structure of the polynomial f in (5). First, a lemma.

Lemma 4.1 For any $(z_1, \dots, z_N) \in \mathbb{C}^N$,

$$\sum_{j=1}^N \frac{1}{z_j^2} = \frac{e_{N-1}^2 - 2e_N e_{N-2}}{e_N^2}.$$

Proof Since $\sum_{j=1}^N (1/z_j) = e_{N-1}/e_N$ and $\sum_{i>j} (1/z_i z_j) = e_{N-2}/e_N$, we have

$$\begin{aligned} \sum_{j=1}^N \frac{1}{z_j^2} &= \left(\sum_{j=1}^N \frac{1}{z_j} \right)^2 - 2 \sum_{i>j} \frac{1}{z_i z_j} \\ &= \frac{e_{n-1}^2}{e_n^2} - 2 \frac{e_{n-2}}{e_n}. \end{aligned}$$

□

Theorem 4.3 (z_1, \dots, z_N) is a solution to S_N if and only if

$$\begin{cases} f \text{ is squarefree and} \\ 3(f'')^2 - 4f'f''' \text{ is divisible by } f \end{cases}$$

where $f = \prod_{i=1}^N (x - z_i) = x^N - e_1 x^{N-1} + e_2 x^{N-2} - \dots + (-1)^N e_N$.

Proof Let S_r be the r th elementary symmetric polynomial in $x - z_1, \dots, x - z_N$. Note that replacing x by z_i in S_r gives the r th elementary symmetric polynomial in $z_i - z_1, \dots, z_i - z_{i-1}, z_i - z_{i+1}, \dots, z_i - z_N$, which we denote by E_r^i . Furthermore, the k th derivative of f is $f^{(k)} = k! S_{n-k}$ so that $f^{(k)}(z_i) = k! E_{n-k}^i$. Set $h := 3(f'')^2 - 4f'f'''$. Then

$$\begin{aligned} h(z_i) &= 3(2E_{N-2}^i)^2 - 4E_{N-1}^i(3!E_{N-3}^i) \\ &= 12((E_{N-2}^i)^2 - 2E_{N-1}^i E_{N-3}^i). \end{aligned} \quad (7)$$

By Lemma 4.1

$$f_i = \sum_{j \neq i} \frac{1}{(z_i - z_j)^2} = \frac{(E_{N-2}^i)^2 - 2E_{N-1}^i E_{N-3}^i}{(E_{N-1}^i)^2} \quad (8)$$

so that $h(z_i)$ is a constant multiple of the numerator of f_i . Therefore f divides h and the z_i are distinct $\iff h(z_i) = 0$ for all i and the z_i are distinct $\iff f_i(z_1, \dots, z_N) = 0$ for all $i \iff (z_1, \dots, z_n)$ is a solution of S_N . \square


Let r be the remainder of dividing h by f , and let c_j , $1 \leq j \leq \deg(r)$, be the coefficient of x^j in r . Then each c_j is a polynomial in the e_i and Theorem 2.3 implies the system $c_j(e_1, \dots, e_N) = 0$, $1 \leq j \leq \deg(r)$, is equivalent to S_N .

Computations with Singular [Gre99] using the formulation of Theorem 2.3 reveal a one-dimensional family of solution shapes for $N = 8$:

Proposition 4.1 The coordinates of a solution to S_8 are given by the roots of the polynomial

$$t^8 + \frac{28}{5}t^6 a + 14t^4 a^2 + 28t^2 a^3 - t - 7a^4$$

where a can be arbitrary. Setting $a = 0$, the roots form a regular heptagon with a point in the center. Varying a deforms this into irregular hexagons with two points in the interior.



This is a one dimensional family of solution shapes.

Proposition 4.2 *There are no solutions to S_N for $N = 3, 4, 6, 7, 9, 10, 11,$ and 12 .*

Proof For $N = 3, 4$, short (less than one minute by Maple on a Sun Ultra-5) Gröbner bases computations show \tilde{S}_N , hence S_N , has no solutions. For the remaining N , computations using one or both of the above reformulations show there are no solutions of the equivalent systems. For $N > 7$ we use FGb for the computations. When $N > 9$ another difficulty arises in the computation: it is impossible to compute the discriminant of $g = x^{N-2} - e_1x^{N-3} + e_2x^{N-4} - \dots + (-1)^N e_{N-2}$. At the begining we add only the condition $g(0) = e_{N-2} \neq 0$ and $g(1) \neq 0$ and we compute a lexicographical Gröbner basis. In the last we remove the bad solutions. \square

5 Regular solutions

The geometry of the solutions known thus far lead one to ask: What other regular polygons are solution shapes (with or without a point in the center)? What about two regular polygons, or n regular polygons? We use the notation $[n, m, p]$ to denote a solution shape consisting of n regular concentric m -gons and $p = 1$ or 0 as there is or is not a point in the center. Thus a solution $[n, m, p]$ will be a solution for S_{nm+p} . We begin this section by trying to find “by hand” some regular solutions then give a more systematic way to find these solutions.

5.1 One regular m-gon: $[1, m, p]$

Since the solutions are invariant under translation and multiplication by complex numbers, it suffices to examine the m 'th roots of unity.

The main lemma we need is

Lemma 5.1 *Let ω be a primitive m 'th root of unity. Then*

$$\begin{aligned} \sum_{j=1}^m \frac{1}{(\omega^j)^2} &= 0 \\ \sum_{j=1}^{m-1} \frac{1}{(\omega^j - 1)^2} &= -\frac{(m-1)(m-5)}{12} \\ \sum_{j=1}^m \frac{1}{\left(\frac{a}{b}\omega^j - 1\right)^2} &= \frac{mb^m(b^m + a^m(m-1))}{(b^m - a^m)^2}. \end{aligned} \tag{9}$$

Proof From Lemma 4.1 we know

$$\sum_{j=1}^N \frac{1}{z_j^2} = \frac{e_{N-1}^2 - 2e_N e_{N-2}}{e_N^2}$$

where the e_i are the elementary symmetric polynomials in the z_j . The polynomials with roots ω^j ($1 \leq j \leq m$), $\omega^j - 1$ ($1 \leq j \leq m - 1$), and $\frac{a}{b}\omega^j - 1$ ($1 \leq j \leq m$) are, respectively

$$\begin{aligned} P(X) &= X^m - 1 \text{ and} \\ P(X) &= \frac{(X+1)^m - 1}{X} = X^{m-1} + mX^{m-2} + \dots + \binom{m}{3}X^2 + \binom{m}{2}X + m \quad (10) \\ P(X) &= (X+1)^m - \left(\frac{a}{b}\right)^m \end{aligned}$$

respectively. substituting in the corresponding values of e_N, e_{N-1} and e_{N-2} gives the result.

□

We first consider the case $p = 0$: $[1, m, 0]$. Let $z_i = \omega^i$ for all i , where ω is a primitive m 'th root of unity. Then the i 'th equation is

$$f_i = \sum_{j \neq i} \frac{1}{(z_i - z_j)^2} = \sum_{j \neq i} \frac{1}{(\omega^i - \omega^j)^2} = \frac{1}{(\omega^i)^2} \sum_{j \neq i} \frac{1}{(\omega^{j-i} - 1)^2} = \frac{1}{(\omega^i)^2} \sum_{j=1}^{m-1} \frac{1}{(\omega^j - 1)^2}.$$

By lemma 5.1, for all i the i 'th equation is zero if and only if

$$\sum_{j=1}^{m-1} \frac{1}{(\omega^j - 1)^2} = -\frac{(m-1)(m-5)}{12} = 0,$$

i.e., if and only if $m = 5$ or $m = 1$. Thus the regular pentagon is the only solution shape for this case. If $p = 1$, i.e., $[1, m, 1]$, we have $z_i = \omega^i$ for $i = 1, \dots, m$ and $z_N = 0$. Then the N 'th equation

$$f_N = \sum_{j=1}^m \frac{1}{(\omega^j - 0)^2} = 0$$

by lemma 5.1. For $i = 1, \dots, m$, the i 'th equation is

$$\begin{aligned} f_i &= \sum_{j \neq i} \frac{1}{(z_i - z_j)^2} = \sum_{j \neq i} \frac{1}{(\omega^i - \omega^j)^2} + \frac{1}{(\omega^i)^2} = \frac{1}{(\omega^i)^2} \left(\sum_{j \neq i} \frac{1}{(\omega^{j-i} - 1)^2} + 1 \right) \\ &= \frac{1}{(\omega^i)^2} \left(\sum_{j=1}^{m-1} \frac{1}{(\omega^j - 1)^2} + 1 \right). \end{aligned} \quad (11)$$

So for all $i = 1, \dots, m$ the i 'th equation is zero if and only if

$$\sum_{j=1}^{m-1} \frac{1}{(\omega^j - 1)^2} = -\frac{(m-1)(m-5)}{12} = -1,$$

i.e., $m = 7$ or $m = -1$. Therefore the regular heptagon with a point in the center is the only solution shape in this case.

5.2 Two regular m -gons $[2, m, x]$

Again we may fix one m -gon, P_1 , to be the m 'th roots of unity. We introduce a new complex variable, x , to describe the second m -gon, $P_2 = xP_1$, where multiplication of a polygon P with x means x times each vertex of the polygon.

Proposition 5.1 *There are no solution shapes of the form $[2, m, 0]$ or $[2, m, 1]$.*

Proof We include in square brackets facts for the case $[2, m, 1]$. Let

$$z_i = \begin{cases} \omega^i & \text{if } i = 1, \dots, m; \\ x\omega^i & \text{if } i = m + 1, \dots, 2m; \\ [0 & \text{if } i = 2m + 1]. \end{cases} \quad (12)$$

Dividing by $(\frac{1}{\omega^i})^2$ in the i 'th equation when $i = 1, \dots, m$, or $(\frac{1}{x\omega^i})^2$ when $i = m + 1, \dots, 2m$, we get two equations in one unknown:

$$-\frac{(m-1)(m-5)}{12}[+1] + \frac{m(1+x^m(m-1))}{(1-x^m)^2} = 0 \quad (13)$$

$$-\frac{(m-1)(m-5)}{12}[+1] + \frac{mx^m(x^m+m-1)}{(x^m-1)^2} = 0. \quad (14)$$

where we have used the third part of lemma 5.1. Subtracting one equation from the other gives

$$1 - x^{2m} = 0, \quad (15)$$

so the solution set would have to consist of $2m$ th roots of unities. But we have already seen that in the single polygon case the only solutions are $m = 5$ and $m = 7$, neither of which is divisible by two. Therefore no shapes of the form $[2, m, 0]$ or $[2, m, 1]$ can be a solution. \square

5.3 The Generalization

Using the differential equation of theorem 4.3 we can find some more conditions not only for the case of regular polygons but for any set of roots to a polynomial $N(X)$. For the case of regular polygons this raises the chances of succesful computations since we can add the new equations to our old systems.

Definition 5.1 *Let N, M, P be univariate polynomials of degree n, m, p . We use the notation $[N, M, P]$ to denote the set of solutions of S_{nm+p} with the shape $P(X)N(M(X))$. In the particular case $P(X) = X^p$, $M(X) = X^m$ we use the simplified notation $[N, m, p]$.*

Theorem 5.1 Let $N(x) = \sum_{i=0}^n a_i X^i$ be a squarefree polynomial of degree n such that $a_0 \neq 0$. Then $[N, m, p]$ (with $m > 1$) is a solution of S_{nm+p} if and only if $p \leq 1$ and $N(X)$ divides

$$\sum_{\substack{i=0 \\ j=0}}^n i j a_i a_j (m i - 1) (3 m j + 5 - 4 m i) X^{i+j} \text{ if } p = 0$$

and $N(X)$ divides

$$\sum_{\substack{i=0 \\ j=1}}^n a_i a_j j m (j m + 1) (i m + 1) (3 i m - 4 j m + 4) X^{i+j} \text{ if } p = 1.$$

Proof Let $f(X) = X^p N(X^m)$. We know from theorem 4.3 that f is a solution of S_{nm+p} if and only if f is squarefree and $U(X) = 3(f'')^2 - 4f'f'''$ is divisible by $f(X)$. The first condition is true as soon as $p \leq 1$ since 0 is not a root of $N(X)$.

Considering the case $p = 0$, we find:

$$U(X) = 3 \left(\sum_{i=1}^n i m (m i - 1) a_i X^{m i - 2} \right)^2 - 4 \left(\sum_{i=1}^n i m a_i X^{m i - 1} \right) \left(\sum_{i=i_3}^n i m (m i - 1) (m i - 2) a_i X^{i m - 3} \right)$$

where $i_3 = 2$ if $m = 2$ and $i_3 = 1$ else. Since X and $f(X)$ are relative prime, f divides U iff f divides $X^4 U = V$ with

$$V(X) = 3 \left(\sum_{i=1}^n i m (m i - 1) a_i X^{m i} \right)^2 - 4 \left(\sum_{i=1}^n i m a_i X^{m i} \right) \left(\sum_{i=i_3}^n i m (m i - 1) (m i - 2) a_i X^{i m} \right)$$

hence $V = W(X^m)$ is divisible by $N(X^m)$ iff $W(X)$ is divisible by $N(X)$. We can rewrite the sum:

$$W(X) = m^2 \sum_{\substack{i=1 \\ j=1}}^n i j a_i a_j (m i - 1) (3 m j + 5 - 4 m i) X^{i+j}.$$

We consider now the case $p = 1$ and find:

$$U(X) = 3 \left(\sum_{i=1}^n a_i (i m + 1) (i m) X^{i m - 1} \right)^2 - 4 \left(\sum_{i=0}^n a_i (i m + 1) X^{i m} \right) \left(\sum_{i=1}^n a_i (i m + 1) (i m) (i m - 1) X^{i m - 2} \right)$$

must be divisible by X and $N(X^m)$ so that $m > 2$ and $V_1(X) = X^2 U(X)$ should be divisible by $N(X^m)$

$$V_1(X) = 3 \left(\sum_{i=1}^n a_i (i m + 1) (i m) X^{i m} \right)^2 - 4 \left(\sum_{i=0}^n a_i (i m + 1) X^{i m} \right) \left(\sum_{i=1}^n a_i (i m + 1) (i m) (i m - 1) X^{i m} \right)$$

this equivalent to divisibility of

$$W_1(X) = 3 \left(\sum_{i=1}^n a_i (im + 1)(im) X^i \right)^2 - 4 \left(\sum_{i=0}^n a_i (im + 1) X^i \right) \left(\sum_{i=1}^n a_i (im + 1)(im)(im - 1) X^i \right)$$

$$= \sum_{\substack{i=0 \\ j=1}}^n a_i a_j j m (j m + 1) (im + 1) (3im - 4jm + 4) X^{i+j}$$

□

Remark 5.1 We can always suppose that $N(X) = X^n + X^{n-1} + \sum_{i=0}^{n-2} a_i X^i$

Remark 5.2 In the following we give an explicit value to n and p and we consider m as a variable.

Corollary 5.1 There are no solutions of the form $[N, 2, 1]$.

Proof From the proof of Theorem 5.1, $f(X) = XN(X)$ does not divide $U(X)$ because X does not divide $U(X)$. □

Corollary 5.2 For $\deg(N) = 1$, $[N, m, 0]$ is a solution iff $(m-1)(m-5) = 0$ and $N(X) = 1 + X$

Proof We apply the theorem 5.1 to $N = 1 + X$ and we find $W(X) = -X^2(m-1)(m-5)$. □

Corollary 5.3 For $\deg(N) = 1$, $[N, m, 1]$ is a solution iff $m = 7$ and $N(X) = 1 + X$

Proof We apply the theorem 5.1 to $N = 1 + X$ and we find

$$W_1(X) = X(-4 + 4m^2) + X^2(m^3 - 2m^2 - 7m - 4)$$

and the remainder of W_1 divided by N should be zero:

$$-m(-7 + m)(m + 1)x$$

□

Corollary 5.4 $\deg(N) = 2$, $[N, m, 0]$ there is no solution.

Proof We apply the theorem 5.1 to $N = a_0 + X + X^2$ and we find

$$W(X) = - (4 (2m - 1) (2m - 5) X^2 + 4 (m - 1) (4m - 5) X + (m - 1) (m - 5)) X^2$$

and the remainder of W divided by N should be zero:

$$- (-5 - m^2 + 18m - 60a_0m + 16a_0m^2 + 20a_0) X$$

$$- (-m^2 + 18m - 5 + 16a_0m^2 - 48a_0m + 20a_0) a_0 = 0$$

we can compute a lexico Gröbner of the coefficients:

$$[20a_0 - m^2 + 18m - 5, m(m^2 - 18m + 5)]$$

and the number of solutions is 0.

5.4 Summary of the regular solutions

An extended version of this paper including a complete list of solutions, pictures and all the polynomials can be found at <http://calfor.lip6.fr/~jcf/MICE/mice.ps.gz>. We summarize all the results:

Theorem 5.2 *For fixed values of n and p we give all the possible values of m and for each m all the solutions $[n, m, p]$. The results are summarized in the following table.*

Shape	Values of m	Values of N
$[1, m, 0]$	$m = 5$	$N = 5$
$[1, m, 1]$	$m = 7$	$N = 8$
$[2, m, 0]$	\emptyset	
$[2, m, 1]$	\emptyset	
$[3, m, 0]$	$m = 7, m = 11$	$N = 21, N = 33$
$[3, m, 1]$	$m = 5, m = 13$	$N = 16, N = 40$
$[4, m, 0]$	$m = 2, m = 4$	$N = 8, N = 16$
$[4, m, 1]$	$m = 5$	$N = 21$
$[5, m, 0]$	$m = 13, m = 17$	$N = 65, N = 85$
$[5, m, 1]$	$m = 11, m = 19$	$N = 56, N = 96$
$[6, m, 0]$	\emptyset	
$[6, m, 1]$	\emptyset	
$[7, m, 0]$	$m = 19, m = 23$	$N = 133, N = 161$
$[7, m, 1]$	$m = 1, m = 17, m = 25$	$N = 8, N = 119, N = 175$
$[8, m, 0]$	$m = 5, m = 7$	$N = 40, N = 56$
$[8, m, 1]$	$m = 4, m = 8$	$N = 37, N = 65$
$[9, m, 0]$	$m = 25, m = 29$	$N = 225, N = 261$
$[9, m, 1]$	$m = 23, m = 31$	$N = 208, N = 280$

Corollary 5.5 *Using this information we could find the following solution families:*

$$f(x) = -32\lambda x^5 + \lambda + x^{16} + x^{11} + \frac{11}{8}x^6 - \frac{11}{128}x$$

$$f(x) = -\lambda x + 25\lambda x^8 + x^{21} + x^{14} - \frac{13}{10}x^7 + \frac{13}{400}$$

for $N = 16$ and $N = 21$.

Conjecture 5.1 *For n odd, there will be solutions for $[n, m, 0]$ with $m = 3n - 2$ and $m = 3n + 2$ and for $[n, m, 1]$ with $m = 3n - 4, m = 3n + 4$.*

□

Solutions [9,31,1]




Figure 1: one regular solution for $N = 280$.


6 Extended version of the paper.

6.1 Equations

$$[3, 7, 0]$$

$$N(X) = X^3 + X^2 - \frac{13}{10}X + \frac{13}{400}$$


Solutions [3,7,0]



The following polynomial is also a solution for all λ :

$f(X) = -\lambda x + 25 \lambda x^8 + x^{21} + x^{14} - \frac{13}{10} x^7 + \frac{13}{400}$
 for instance for $\lambda = 400$:

MICE



In fact the solution when “ λ is big” are from one part:

$$f_1(X) = -\lambda x + 25 \lambda x^8 = \lambda (25x^7 - 1) x$$

that is to say 0 and $25^{-\frac{1}{7}} e^{\frac{2k\pi}{7}}$ for $k = 0, \dots, 6$. The second set of solution is


$$f_2(X) = 25 x^{13} + 26 x^6 + 625 \lambda \approx 25 x^{13} + 625 \lambda$$

and the solutions are $(-25\lambda)^{\frac{1}{13}} e^{\frac{2k\pi}{13}}$ for $k = 0, \dots, 12$

[3, 11, 0]

$$N(X) = X^3 + X^2 - \frac{17}{70} X + \frac{17}{7840}$$


Solutions [3, 11, 0]



[3, 5, 1]

$$N(X) = X^3 + X^2 + \frac{11}{8}X - \frac{11}{128}$$

Solutions [3,5,1]




$$f(x) = -32\lambda x^5 + \lambda + x^{16} + x^{11} + \frac{11}{8}x^6 - \frac{11}{128}x$$

[3, 13, 1]

$$N(X) = X^3 + X^2 - \frac{19}{112}X + \frac{19}{17920}$$


Solutions [3,13,1]



[4, 2, 0]

$$N(X) = X^4 + X^3 + \frac{25}{56}X^2 + \frac{125}{784}X - \frac{625}{87808}$$


Solutions [4,2,0]



[4, 4, 0]

$$N(X) = X^4 + X^3 + \frac{49}{88} X^2 - \frac{343}{1936} X - \frac{2401}{1703680}$$


Solutions [4,4,0]



[5, 13, 0]

$$N(X) = X^5 + X^4 - \frac{26246}{10773} X^3 + \frac{264143}{6786990} X^2 + \frac{1331}{12216582} X + \frac{14641}{1169859892320}$$


Solutions [5,13,0]



[5, 17, 0]

$$N(X) = X^5 + X^4 - \frac{505466}{1071455} X^3 + \frac{517478}{129646055} X^2 + \frac{191139}{49913731175} X + \frac{828269}{4861846985100875}$$


Solutions [5,17,0]



[5, 11, 1]

$$N(X) = X^5 + X^4 + \frac{1355}{544} X^3 - \frac{1885}{30464} X^2 - \frac{345}{974848} X - \frac{115}{1856110592}$$


Solutions [5,11,1]



[5, 19, 1]

$$N(X) = X^5 + X^4 - \frac{623}{1872} X^3 + \frac{35231}{16061760} X^2 + \frac{10633}{7709644800} X + \frac{10633}{264595009536000}$$


Solutions [5,19,1]



[7, 19, 0]

$$N(X) = X^7 + X^6 - \frac{250573}{70304} X^5 + \frac{15738440127}{380204032000} X^4 + \frac{1616936130527}{9895190136832000} X^3 + \frac{118110645017}{8232798193844224000} X^2 - \frac{5415437771}{44522972632309563392000} X + \frac{92062442107}{1854007817165581914944307200000}$$


Solutions [7,19,0]



[7, 23, 0]

$$N(X) = X^7 + X^6 - \frac{13439}{19250} X^5 + \frac{115793631}{22586795000} X^4 + \frac{4543346861}{549084986450000} X^3 + \frac{5295509}{13727124661250000} X^2 - \frac{843657}{628908216355168750000} X + \frac{5343161}{22571767448273548505000000000}$$


Solutions [7,23,0]



[7, 17, 1]

$$N(X) = X^7 + X^6 + \frac{673177}{186340} X^5 - \frac{232819511}{4148673760} X^4 - \frac{13989941649}{35139266747200} X^3 - \frac{562316989}{13528617697672000} X^2 + \frac{317057}{654785096567324800} X - \frac{317057}{1275586846622805442880000}$$


Solutions [7,17,1]



[7, 25, 1]

$$N(X) = X^7 + X^6 - \frac{1203647}{2434432} X^5 + \frac{1513386395}{504725917696} X^4 + \frac{3365775485}{997338413367296} X^3 + \frac{564207339}{4914883701074034688} X^2 - \frac{111082029}{427319648506180871913472} X + \frac{1240334593196852549369068519424}{37027343}$$


Solutions [7,25,1]



[8, 5, 0]

$$X^8 + X^7 + \frac{189}{832} X^6 - \frac{1519}{13312} X^5 + \frac{36015}{5537792} X^4 - \frac{215061}{143982592} X^3 - \frac{45619}{9214885888} X^2 + \frac{319333}{958348132352} X + \frac{319333}{1594691292233728}$$


Solutions [8,5,0]



[8, 7, 0]

$$N(X) = X^8 + X^7 + \frac{28}{85} X^6 - \frac{14}{187} X^5 + \frac{7}{9350} X^4 - \frac{1932}{6755375} X^3 + \frac{161}{148618250} X^2 + \frac{23}{6316275625} X + \frac{94491483350000}{23}$$


Solutions [8,7,0]



[8, 4, 1]

$$N(X) = X^8 + X^7 + \frac{13}{176} X^6 - \frac{187083}{862400} X^5 + \frac{17064099}{303564800} X^4 - \frac{40299571}{6678425600} X^3 + \frac{246167259}{747983667200} X^2 - \frac{1066724789}{164556406784000} X - \frac{13867422257}{810933972631552000}$$


Solutions [8,4,1]



[8, 8, 1]

$$N(X) = X^8 + X^7 + \frac{1207}{3344} X^6 - \frac{851683}{9563840} X^5 - \frac{12729583}{8140740608} X^4 - \frac{87613529}{309348143104} X^3 + \frac{3403397229}{3538942757109760} X^2 + \frac{410338673}{268959649540341760} X + \frac{6975757441}{233844277696354739814400}$$


Solutions [8,8,1]



[9, 25, 0]

$$N(X) = X^9 + X^8 - \frac{268466}{57239} X^7 + \frac{377495813029}{8840907899824} X^6 + \frac{413656879904907}{2119368964469508752} X^5 + \frac{12304828054251749}{996713791562436331960576} X^4 - \frac{2016962935969613}{13792525447640993961670450688} X^3 + \frac{57217346608047}{1049280165954736256628041206540288} X^2 + \frac{2902085007}{2130338531221244525599674518192939008} X + \frac{22249318387}{12640627189356212335708507331037005997751533568}$$


Solutions [9,25,0]



[9, 29, 0]

$$N(X) = X^9 + X^8 - \frac{3171229688}{3433022619} X^7 + \frac{12964501623920}{2205535082508693} X^6 + \frac{13450720325129528144}{1110030908637142840678785} X^5 + \frac{68635259350872716128}{140189133544510680777845794005} X^4 - \frac{563378090146829056}{196573203056112876586695372353811} X^3 + \frac{6736550307020061440}{2835847336220800} X^2 + \frac{22686778689766400}{3262885539298427619562253503112758259154206613} X + \frac{11261775713673813363299935122968402618823}{658915285859971494652443714988122706538556183025198791}$$


Solutions [9,29,0]



[9, 31, 1]

$$N(X) = X^9 + X^8 - \frac{87004837}{132721650} X^7 + \frac{988698212699}{276694777878750} X^6 + \frac{7649466909707441}{1433863961228011500000} X^5 + \frac{778364618387927033}{4619802143279560952137500000} X^4 - \frac{31235560755437909}{45292540212712815574756050000000} X^3 + \frac{886646005540711}{8584068683814396371805640376250000000} X^2 + \frac{44262730811}{5591907599741949636490531445100000000000} X - \frac{1018042808653}{315310387339618467525703271464968060067500000000000}$$


Solutions [9,31,1]



[9, 23, 1]

$$N(X) = X^9 + X^8 + \frac{446725777}{94325000} X^7 - \frac{14305770645413}{267147265000000} X^6 - \frac{13328514939091617283}{3226738240302500000000} X^5 - \frac{247787301154250111}{907593465045085000000000} X^4 + \frac{1342628342958548400883}{34243501436151057050000000000000} X^3 - \frac{784362891765830249}{50937208386274697361875000000000000} X^2 - \frac{193968889534934047554020000000000000000}{852798576538163125190767927480000000000000000000000} X - \frac{55014634972391767}{852798576538163125190767927480000000000000000000000}$$

Solutions [9,23,1]



6.2 Another proof of Derksens formulation

Let $f = \prod_{j=1}^n (x - z_j)$. Define

$$f_j = \prod_{i \neq j} (x - z_i), j = 1, \dots, n.$$

Then

$$\frac{f'_j}{f_j} = \sum_{i \neq j} \frac{1}{(x - z_j)^2}. \quad (16)$$

Derivating this and plugging in z_j we get

$$\frac{f''_j f_j - f_j'^2}{f_j^2}(z_j) = - \sum_{i \neq j} \frac{1}{(x - z_i)^2}(z_j) = - \sum_{i \neq j} \frac{1}{(z_j - z_i)^2}. \quad (17)$$

Therefore for all j , $(f''_j f_j - f_j'^2)(z_j) = 0$ if and only if $\sum_{i \neq j} \frac{1}{(z_j - z_i)^2}$, i.e. if the MICE equations are satisfied and z_j is a root of f . On the other hand we have

$$\begin{aligned} f &= f_j(x - z_j), \\ f' &= f'_j(x - z_j) + f_j, \\ f'' &= f''_j(x - z_j) + 2f'_j, \\ f''' &= f'''_j(x - z_j) + 3f''_j, \end{aligned} \quad (18)$$

and therefore

$$\begin{aligned} (3(f'')^2 - 4f'f''')(z_j) &= 12((f'_j)^2 - f_j f''_j)(z_j) \\ &= 0. \end{aligned} \quad (19)$$

6.3 Three regular m-gons

Proposition 6.1 *The only solution shapes $[3, m, 0]$ are for $m = 7$ or $m = 11$. The only solution shapes $[3, m, 1]$ are for $m = 5$ or $m = 13$.*

Proof Proceeding as before, we fix one m -gon to be the m 'th roots of unity and use two variables x and y to describe the remaining two. Again using formula in lemma 5.1, S_{3m+p} is reduced to a system of three equations in two unknowns. The case $p = 1$ is included in square brackets:

$$-\frac{(m-1)(m-5)}{12}[+1] + \frac{m(1+x^m(m-1))}{(1-x^m)^2} + \frac{m(1+y^m(m-1))}{(1-y^m)^2} = 0 \quad (20)$$

$$-\frac{(m-1)(m-5)}{12}[+1] + \frac{mx^m(x^m+m-1)}{(x^m-1)^2} + \frac{mx^m(x^m+my^m-y^m)}{(x^m-y^m)^2} = 0 \quad (21)$$

$$-\frac{(m-1)(m-5)}{12}[+1] + \frac{my^m(y^m+m-1)}{(y^m-1)^2} + \frac{my^m(y^m+mx^m-x^m)}{(y^m-x^m)^2} = 0 \quad (22)$$

Remark 6.1 *In general, this method reduces S_{nm+p} to a system of n equations in $n - 1$ unknowns.*

We can also make the variable changes $X = x^m$ and $Y = y^m$ to eliminate m from the exponents, which allows m to be treated as a variable in the calculations. A lexicographic Gröbner basis for $[3, m, 0]$ with $X > Y > m$ was obtained after 536 seconds by computing a degree reverse-lexicographic Gröbner basis then using FGLM. This was done using Singular[Gre99]. The single polynomial involving m alone is $9m^{10} - 990m^9 + 41657m^8 - 833256m^7 + 8052130m^6 - 35019540m^5 + 56505450m^4 - 33100200m^3 + 8675125m^2 - 1050750m + 48125$ which factors as $(m - 7)(m - 11)(3m - 5)(3m - 1)(m^2 - 30m + 5)^3$. We are only interested in the integer partial solutions $m = 7$ and $m = 11$, both of which lift to solutions. Similarly, the same calculation took 657 seconds for the case $[3, m, 1]$ and the relevant polynomial is $9m^{10} - 990m^9 + 41213m^8 - 796104m^7 + 6974386m^6 - 22637556m^5 + 9529122m^4 + 27373752m^3 + 12827269m^2 + 2364642m + 156065$. Its factorization is $(m - 5)(m - 13)(3m - 7)(3m + 1)(m^2 - 30m - 7)^3$ and both integer partial solutions lift. \square

$$U(X) = 12 \left(\sum_{i=1}^n a_i X^{mi-1} im \right)^2 + 3 \left(\sum_{i=0}^n a_i X^{mi-1} im (mi-1) \right)^2 - 12 \sum_{i=0}^n a_i X^{mi} \sum_{i=0}^n a_i X^{mi-2} im (mi-1) - 4 \sum_{i=0}^n a_i X^{mi} \sum_{i=0}^n a_i X^{mi-2} im (mi-1) (mi-2) - 4 \sum_{i=0}^n a_i X^{mi-1} im \sum_{i=0}^n a_i X^{mi-1} im (mi-1) (mi-2)$$

must be divisible by X and $N(X^m)$ so that $m > 2$ and $V_1(X) = X^2 U(X)$ should be divisible by $N(X^m)$

$$V_1(X) = 12 \left(\sum_{i=1}^n a_i X^{mi} im \right)^2 + 3 \left(\sum_{i=0}^n a_i X^{mi} im (mi-1) \right)^2 - 12 \sum_{i=0}^n a_i X^{mi} \sum_{i=0}^n a_i X^{mi} im (mi-1) - 4 \sum_{i=0}^n a_i X^{mi} \sum_{i=0}^n a_i X^{mi} im (mi-1) (mi-2) - 4 \sum_{i=0}^n a_i X^{mi} im \sum_{i=0}^n a_i X^{mi} im (mi-1) (mi-2)$$

this equivalent to divisibility of

$$W_1(X) = 12 \left(\sum_{i=1}^n a_i X^i im \right)^2 + 3 \left(\sum_{i=0}^n a_i X^i im (mi-1) \right)^2 - 12 \sum_{i=0}^n a_i X^i \sum_{i=0}^n a_i X^i im (mi-1) - 4 \sum_{i=0}^n a_i X^i \sum_{i=0}^n a_i X^i im (mi-1) (mi-2) - 4 \sum_{i=0}^n a_i X^i im \sum_{i=0}^n a_i X^i im (mi-1) (mi-2)$$

by $N(X)$.

6.4 Different Polygons

We tried using 3 different polygons for the case $N = 16$ and there were no solutions, neither with a point nor without a point. We tried the partitions 1+2+2+11, 1+2+3+10, 1+2+4+9, 1+2+5+8, 1+2+6+7, 1+3+3+9, 1+3+4+8, 1+3+5+7, 1+3+6+6, 1+4+4+7, 1+4+5+6, 1+5+5+5 (there is the solution we know), 2+2+12, 2+3+11, 2+4+10, 2+5+9, 2+6+8, 2+7+7, 3+3+10, 3+4+9, 3+5+8, 3+6+7, 4+4+8, 4+5+7, 4+6+6, 5+5+6.

7 Conclusion

We have a new application of computer algebra in biological physics. We were able to solve the system completely up to $N = 12$ using the symmetry and the most recent techniques for the Gröbner bases computation. Starting with solution shapes of regular polygons we found solution

families for $N = 8, 16, 21$ as well as single solutions for N up to 280 for which we have reason to assume that they are part of solution families as well.

From the biophysical point of view, solutions for N about 1000 are needed since there are thousands of proteins in a cell membrane [Kim99]. But even small numbers of proteins can give some interesting insights. We have extended the results in the original paper[KJG98] from $N = 5$ to 12.

This work is a particular instance of the more general problem of finding a global minimum of an energy function and in particular we want to point out similar work related to the classification of the stable solutions of the n body problem.

Acknowledgments

We are deeply indebted to Prof. Bernd Sturmfels for bringing this problem to our attention. Two of the authors (M.H. and J.P.) gratefully acknowledge several discussions with K.S. Kim as well as with Eric Antokoletz and Frank Calegari.

References

- [AS98] Lascoux A. and Veigneau S. Algebraic combinatorics environment. Technical report, Institut Gaspard Monge, Universite de Marne-la-Vallée, 1998. Version 3.0.
- [Bec93] Becker T. and Weispfenning V. *Groebner Bases, a Computational Approach to Commutative Algebra*. Graduate Texts in Mathematics. Springer-Verlag, 1993.
- [Buc65] Buchberger B. *Ein Algorithmus zum Auffinden der Basiselemente des Restklassenringes nach einem nulldimensionalen Polynomideal*. PhD thesis, Innsbruck, 1965.
- [Buc70] Buchberger B. An Algorithmical Criterion for the Solvability of Algebraic Systems. *Aequationes Mathematicae*, 4(3):374–383, 1970. (German).
- [Buc79] Buchberger B. A Criterion for Detecting Unnecessary Reductions in the Construction of Gröbner Basis. In *Proc. EUROSAM 79*, volume 72 of *Lect. Notes in Comp. Sci.*, pages 3–21. Springer Verlag, 1979.
- [Buc85] Buchberger B. Gröbner Bases : an Algorithmic Method in Polynomial Ideal Theory. In Reidel, editor, *Recent trends in multidimensional system theory*. Bose, 1985.
- [Cha91] Char B. and Geddes K. and Gonnet G. and Leong B. and Monagan M. and Watt S. *Maple V Library Reference Manual*. Springer-Verlag, 1991. Third Printing, 1993.
- [CLO92] D. Cox, J. Little, and D. O’Shea. *Ideals, Varieties and Algorithms*. Springer Verlag, New York, 1992.
- [CLO98] D. Cox, J. Little, and D. O’Shea. *Using Algebraic Geometry*. Springer Verlag, New York, 1998.

- [Dav93] Davenport J.H. and Siret Y. and Tournier E. *Calcul Formel*. Masson, 1993. 2^e édition révisée.
- [Der99] H. Derksen. Private communication. Seminar Berkeley, 1999.
- [DG99] Bini D. and Fiorentino G. Mpsolve. Technical report, University of Pisa, 1999.
- [Fau] Faugère J.C. *FGb available online*. available on the WEB <http://calfor.lip6.fr/~jcf/FGb.html>.
- [Fau93] Faugère, J.C., Gianni, P., Lazard, D. and Mora T. Efficient Computation of Zero-Dimensional Gröbner Basis by Change of Ordering. *Journal of Symbolic Computation*, 16(4):329–344, October 1993.
- [Fau94] Faugère J.C. *Résolution des systèmes d'équations algébriques*. PhD thesis, Université Paris 6, Feb. 1994.
- [Fau99] Faugère J.C. A new efficient algorithm for computing Gröbner bases (F4). *Journal of Pure and Applied Algebra*, 139(1–3):61–88, June 1999.
- [Gre99] Greuel G.-M. and Pfister G. and Schoenemann H. *SINGULAR 1.2.3*, Feb 1999. <http://www.mathematik.uni-kl.de/~zca/Singular/Welcome.html>.
- [J.98] Stembridge J. The symmetric functions package. Technical report, Maple share library, 1998.
- [Kim99] K.S. Kim. Private communication. Cambridge, 1999.
- [KJG98] K.S. Kim, Neu J., and Oster G. Curvature-mediated interactions between membrane proteins. *Biophysical Journal*, 75:2274–2291, 1998.