



The distribution of integers with at least two divisors in a short interval

Kevin Ford, Gérald Tenenbaum

► To cite this version:

Kevin Ford, Gérald Tenenbaum. The distribution of integers with at least two divisors in a short interval. 2006. hal-00091331

HAL Id: hal-00091331

<https://hal.archives-ouvertes.fr/hal-00091331>

Preprint submitted on 5 Sep 2006

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

The distribution of integers with at least two divisors in a short interval

Kevin Ford* Gérald Tenenbaum

Version 7 : 12/8/2006, 1h49

1 Introduction

Whereas, in usual cases, sieving by a set of primes may be fairly well controlled, through Buchstab's identity, sieving by a set of integers is a much more complicated task. However, some fairly precise results are known in the case where the set of integers is an interval. We refer to the recent work [1] of the first author for specific statements and references.

Define

$$\begin{aligned}\tau(n; y, z) &:= |\{d|n : y < d \leq z\}|, \\ H(x, y, z) &:= |\{n \leq x : \tau(n; y, z) \geq 1\}|, \\ H_r(x, y, z) &:= |\{n \leq x : \tau(n; y, z) = r\}|, \\ H_2^*(x, y, z) &:= |\{n \leq x : \tau(n; y, z) \geq 2\}| = \sum_{r \geq 2} H_r(x, y, z).\end{aligned}$$

Thus, the numbers $H_r(x, y, z)$ ($r \geq 1$) describe the local laws of the function $\tau(n; y, z)$. When y and z are close, it is expected that, if an integer has at least a divisor in $(y, z]$, then it usually has exactly one, in other words

$$H(x, y, z) \sim H_1(x, y, z). \tag{1.1}$$

In this paper, we address the problem of determining the exact range of validity of such behavior. In other words, we search a necessary and sufficient condition so that $H_2^*(x, y, z) = o(H(x, y, z))$ as x and y tend to infinity.

As shown in [5], for given y , the threshold for the behavior of the function $H(x, y, z)$ lies near the critical value

$$z = z_0(y) := y \exp\{(\log y)^{1-\log 4}\} \approx y + y/(\log y)^{\log 4 - 1}.$$

We concentrate on the case $z_0(y) \leq z \leq 2y$. Define

$$\begin{aligned}z &= e^\eta y, \quad \eta = (\log y)^{-\beta}, \quad \beta = \log 4 - 1 - \Xi/\sqrt{\log_2 y}, \quad \lambda = \frac{1 + \beta}{\log 2}, \\ Q(w) &= \int_1^w \log t \, dt = w \log w - w + 1.\end{aligned}$$

*First author supported by National Science Foundation grant DMS-0301083.

Here \log_k denotes the k th iterate of the logarithm.

With the above notation, we have

$$\log(z/y) = \frac{e^{\Xi\sqrt{\log_2 y}}}{(\log y)^{\log 4 - 1}}, \quad \log\{z/z_0(y)\} = \frac{e^{\Xi\sqrt{\log_2 y}} - 1}{(\log y)^{\log 4 - 1}},$$

so

$$\begin{aligned} 0 \leq \Xi &\leq (\log 4 - 1)\sqrt{\log_2 y} + \frac{\log_2 2}{\sqrt{\log_2 y}}, \\ \frac{|\log_2 2|}{\log_2 y} &\leq \beta \leq \log 4 - 1, \\ \frac{1}{\log 2} + \frac{\kappa}{\log_2 y} &\leq \lambda \leq 2, \end{aligned}$$

with $\kappa := |\log_2 2|/\log 2$.

From Theorem 1 of [1], we know that, uniformly in $10 \leq y \leq \sqrt{x}$, $z_0(y) \leq z \leq 2y$,

$$H(x, y, z) \asymp \frac{\beta x}{(\Xi + 1)(\log y)^{Q(\lambda)}}. \quad (1.2)$$

By Theorems 5 and 6 of [1], for any $c > 0$ and uniformly in $y_0(r) \leq y \leq x^{1/2-c}$, $z_0(y) \leq z \leq 2y$, we have

$$\begin{aligned} \frac{H_1(x, y, z)}{H(x, y, z)} &\asymp_c 1, \\ \frac{\Xi + 1}{\sqrt{\log_2 y}} &\ll_{r,c} \frac{H_r(x, y, z)}{H(x, y, z)} \leq 1 \quad (r \geq 2). \end{aligned} \quad (1.3)$$

When $0 \leq \Xi \leq o(\sqrt{\log_2 y})$ and $r \geq 2$, the upper and lower bounds above for $H_r(x, y, z)$ have different orders. We show in this paper that the lower bound represents the correct order of magnitude.

Theorem 1. *Uniformly in $10 \leq y \leq \sqrt{x}$, $z_0(y) \leq z \leq 2y$, we have*

$$\frac{H_2^*(x, y, z)}{H(x, y, z)} \ll \frac{\Xi + 1}{\sqrt{\log_2 y}}.$$

Corollary 2. *Let $r \geq 2$ and $c > 0$. Uniformly in $y_0(r, c) \leq y \leq x^{1/2-c}$, $z_0(y) \leq z \leq 2y$, we have*

$$\frac{H_r(x, y, z)}{H(x, y, z)} \asymp_{r,c} \frac{\Xi + 1}{\sqrt{\log_2 y}}.$$

Corollary 3. *Uniformly in $10 \leq y \leq \sqrt{x}$, $y < z \leq y + y(\log y)^{1-\log 4+o(1)}$, we have*

$$H_1(x, y, z) \sim H(x, y, z).$$

Since we know from (1.3) that $H_2^*(x, y, z) \gg H(x, y, z)$ when $\beta \leq \log 4 - 1 - \varepsilon$ for any $\varepsilon > 0$ we have therefore completely answered the question raised at the beginning of this introduction concerning the exact validity range for the asymptotic formula (1.1). This may be viewed as a complement a theorem of Hall (see [2], ch. 7; following a note mentioned by Hall in private correspondence, we slightly modify the statement) according to which

$$H(x, y, z) \sim F(-\Xi) \sum_{r \geq 1} r H_r(x, y, z) = F(-\Xi) \sum_{n \leq x} \tau(n; y, z) \quad (1.4)$$

in the range $\Xi = o(\log_2 y)^{1/6}$, $x > \exp\{\log z \log_2 z\}$ with

$$F(\xi) := \frac{1}{\sqrt{\pi}} \int_{-\infty}^{\xi/\log^4} e^{-u^2} du.$$

It is likely that (1.4) still holds in the range $(\log_2 y)^{1/6} \ll \Xi \leq o(\sqrt{\log_2 y})$.

2 Auxiliary estimates

In the sequel, unless otherwise indicated, constants implied by O - and \ll - symbols are independent of any parameter.

Let m be a positive integer. We denote by $P^+(m)$ and $P^-(m)$ the smallest prime factor and largest prime factor of m , respectively, with the convention that $P^-(1) = \infty$, $P^+(1) = 1$. We write $\omega(m)$ for the number of distinct prime factors of m and $\Omega(m)$ the number of prime power divisors of m . We further define

$$\omega(m; t, u) = \sum_{\substack{p^\nu \parallel m \\ t < p \leq u}} 1, \quad \Omega(m; t, u) = \sum_{\substack{p^\nu \parallel m \\ t < p \leq u}} \nu, \quad \bar{\Omega}(m; t) = \Omega(m; 2, t), \quad \bar{\Omega}(m) = \Omega(m; 2, m).$$

Also, we let $\mathcal{P}(u, v)$ denote the set of integers all of whose prime factors are in $(u, v]$ and write $\mathcal{P}^*(u, v)$ for the set of squarefree members of $\mathcal{P}(u, v)$. By convention, $1 \in \mathcal{P}^*(u, v)$.

Lemma 2.1. *There is an absolute constant $C > 0$ so that for $\frac{3}{2} \leq u < v$, $v \geq e^4$, $0 \leq \alpha \leq 1/\log v$, we have*

$$\sum_{\substack{m \in \mathcal{P}(u, v) \\ \omega(m) = k}} \frac{1}{m^{1-\alpha}} \leq \frac{(\log_2 v - \log_2 u + C)^k}{k!}.$$

Proof. For a prime $p \leq v$, we have $p^\alpha \leq 1 + 2\alpha \log p$, thus the sum in question is

$$\leq \frac{1}{k!} \left(\sum_{u < p \leq v} \frac{1}{p^{1-\alpha}} + \frac{1}{p^{2-2\alpha}} + \dots \right)^k \leq \frac{\{\log_2 v - \log_2 u + O(1)\}^k}{k!}.$$

□

Lemma 2.2. *Uniformly for $u \geq 10$, $0 \leq k \leq 2.9 \log_2 u$, and $0 \leq \alpha \leq 1/(100 \log u)$, we have*

$$\sum_{\substack{P^+(m) \leq u \\ \bar{\Omega}(m) = k}} \frac{1}{m^{1-\alpha}} \ll \frac{(\log_2 u)^k}{k!}.$$

Proof. We follow the proof of Theorem 08 of [3]. Let w be a complex number with $|w| \leq \frac{29}{10}$. If p is prime and $3 \leq p \leq u$, then $|w/p^{1-\alpha}| \leq \frac{99}{100}$ and $p^\alpha \leq 1 + 2\alpha \log p$. Thus,

$$S(w) := \sum_{P^+(m) \leq u} \frac{w^{\overline{\Omega}(m)}}{m^{1-\alpha}} = \left(1 - \frac{1}{2^{1-\alpha}}\right)^{-1} \prod_{3 \leq p \leq u} \left(1 - \frac{w}{p^{1-\alpha}}\right)^{-1} \ll e^{(\Re w) \log_2 u}.$$

Put $r := k/\log_2 u$. By Cauchy's formula and Stirling's formula,

$$\begin{aligned} \sum_{\substack{P^+(m) \leq u \\ \overline{\Omega}(m) = k}} \frac{1}{m^{1-\alpha}} &= \frac{1}{2\pi r^k} \int_{-\pi}^{\pi} e^{-ik\vartheta} S(re^{i\vartheta}) d\vartheta \\ &\ll \frac{(\log_2 u)^k}{k^k} \int_{-\pi}^{\pi} e^{k \cos \vartheta} d\vartheta \ll \frac{(\log_2 u)^k}{k!}. \end{aligned}$$

□

Lemma 2.3. *Suppose z is large, $0 \leq a + b \leq \frac{5}{2} \log_2 z$ and*

$$\exp\{(\log x)^{9/10}\} \leq w \leq z \leq x, \quad xz^{-1/(10 \log_2 z)} \leq Y \leq x.$$

The number of integers n with $x - Y < n \leq x$, $\overline{\Omega}(n; w) = a$ and $\omega(n; w, z) = \Omega(n; w, z) = b$, is

$$\ll \frac{Y}{\log z} \frac{\{\log_2 w\}^a (b+1) \{\log_2 z - \log_2 w + C\}^b}{a! b!},$$

where C is a positive absolute constant.

Proof. There are $\ll x^{9/10}$ integers with $n \leq x^{9/10}$ or $2^j | n$ with $2^j \geq x^{1/10}$. For other n , write $n = rst$, where $P^+(r) \leq w$, $s \in \mathcal{P}^*(w, z)$ and $P^-(t) > z$. Here $\overline{\Omega}(r) = a$ and $\omega(s) = b$. We have either $t = 1$ or $t > z$. In the latter case $x/rs > z$, whence $Y/rs > \sqrt{z}$. We may therefore apply a standard sieve estimate to bound, for given r and s , the number of t by

$$\ll \frac{Y}{rs \log z}.$$

By Lemmas 2.1 and 2.2,

$$\sum_{r,s} \frac{1}{rs} \ll \frac{(\log_2 w)^a (\log_2 z - \log_2 w + C)^b}{a! b!}.$$

If $t = 1$, then we may assume $a + b \geq 1$. Set $p = P^+(n)$. If $b \geq 1$, then $p|s$ and we put $r_1 := r$ and $s_1 := s/p$. Otherwise, let $r_1 := r/p$ and $s_1 := s = 1$. Let $A := \overline{\Omega}(r_1)$ and $B := \omega(s_1)$, so that $A + B = a + b - 1$ in all circumstances. We have

$$p \geq x^{1/2\overline{\Omega}(n)} \geq x^{1/5 \log_2 z} \geq (x/Y)^2.$$

Define the non-negative integer h by $z^{e^{-h-1}} < p \leq z^{e^{-h}}$. By the Brun-Titchmarsh theorem, we see that, for each given r_1 and s_1 , the number of p is $\ll Ye^h/(r_1 s_1 \log z)$. Set $\alpha := 0$ if $h = 0$ and $\alpha := e^h/(100 \log z)$ otherwise. For $h \geq 1$, we have $r_1 s_1 > x^{3/4} z^{-1/e} > \sqrt{z}$. Therefore, for $h \geq 0$,

$$\frac{1}{r_1 s_1} \leq \frac{z^{-\alpha/2}}{(r_1 s_1)^{1-\alpha}} \ll \frac{e^{-e^h/200}}{(r_1 s_1)^{1-\alpha}}.$$

Now, Lemmas 2.1 and 2.2 imply that

$$\begin{aligned} \sum_{r_1, s_1} \frac{1}{(r_1 s_1)^{1-\alpha}} &\ll \frac{(\log_2 w)^A (\log_2 z - \log_2 w + C)^B}{A! B!} \\ &\ll (b+1) \frac{(\log_2 w)^a (\log_2 z - \log_2 w + C)^b}{a! b!}, \end{aligned}$$

where we used the fact that $a \ll \log_2 w$. Summing over all h , we derive that the number of those integers $n > x^{9/10}$ satisfying the conditions of the statement is

$$\ll \frac{Y}{\log z} (b+1) \frac{(\log_2 w)^a (\log_2 z - \log_2 w + C)^b}{a! b!}.$$

Since $a! b! \leq (3 \log_2 z)^{3 \log_2 z}$, this last expression is $> x^{9/10}$. This completes the proof. \square

Our final lemma is a special case of a theorem of Shiu (Theorem 03 of [3]).

Lemma 2.4. *Let f be a multiplicative function such that $0 \leq f(n) \leq 1$ for all n . Then, for all x, Y with $1 < \sqrt{x} \leq Y \leq x$, we have*

$$\sum_{x-Y < n \leq x} f(n) \ll \frac{Y}{\log x} \exp \left\{ \sum_{p \leq x} \frac{f(p)}{p} \right\}.$$

3 Decomposition and outline of the proof

Throughout, ε will denote a very small positive constant. Since Theorem 1 holds trivially for $\beta \leq \log 4 - 1 - \varepsilon$, we henceforth assume that

$$\log 4 - 1 - \varepsilon \leq \beta \leq \log 4 - 1. \quad (3.1)$$

Let

$$K := \lfloor \lambda \log_2 z \rfloor,$$

so that $(2 - \frac{3}{2}\varepsilon) \log_2 z \leq K \leq 2 \log_2 z$. In light of (1.2), Theorem 1 reduces to

$$H_2^*(x, y, z) \ll \frac{x}{(\log y)^{Q(\lambda)} \sqrt{\log_2 y}}. \quad (3.2)$$

At this stage, we notice for further reference that, by Stirling's formula, for $k \leq K$ we have

$$\frac{\eta(2 \log_2 z)^k}{k! (\log z)^2} \leq \frac{\eta(2 \log_2 z)^K}{K! (\log z)^2} \asymp \frac{1}{(\log y)^{Q(\lambda)} \sqrt{\log_2 y}}. \quad (3.3)$$

Let \mathcal{H} denote the set of integers $n \leq x$ with $\tau(n; y, z) \geq 2$. We count separately the integers $n \in \mathcal{H}$ lying in 6 classes. In these definitions, we write $k = \overline{\Omega}(n; z) = K - b$ and for brevity we put $z_h = z^{e^{-h}}$. Let

$$K_0 := (2 - 3\varepsilon) \log_2 z$$

and define

$$\begin{aligned}\mathcal{N}_0 &:= \{n \in \mathcal{H} : n \leq x/\log z \text{ or } \exists d > \log z : d^2 | n\}, \\ \mathcal{N}_1 &:= \{n \in \mathcal{H} \setminus \mathcal{N}_0 : k \notin (K_0, K]\}, \\ \mathcal{N}_2 &:= \bigcup_{1 \leq h \leq 5\epsilon \log_2 z} \mathcal{N}_{2,h}, \\ \text{with } \mathcal{N}_{2,h} &:= \left\{n \in \mathcal{H} \setminus (\mathcal{N}_0 \cup \mathcal{N}_1) : \bar{\Omega}(n; z_h, z) \leq \frac{19}{10}h - \frac{1}{100}b\right\}.\end{aligned}$$

For integers $n \in \mathcal{N}_2$, we will only use the fact that $\tau(n; y, z) \geq 1$. Integers in other classes do not have too many small prime factors and it is sufficient to count pairs of divisors d_1, d_2 of n in $(y, z]$. For each such pair, write $v = (d_1, d_2)$, $d_1 = vf_1$, $d_2 = vf_2$, $n = f_1 f_2 v u$ and assume $f_1 < f_2$. Let

$$F_1 = \bar{\Omega}(f_1), \quad F_2 = \bar{\Omega}(f_2), \quad V = \bar{\Omega}(v), \quad U = \bar{\Omega}(u, z), \quad (3.4)$$

and

$$Z := \exp\{(\log z)^{1-4\epsilon}\}. \quad (3.5)$$

For further reference, we note that if $n \notin \mathcal{N}_0$ and $h \leq 5\epsilon \log_2 z$, then

$$\bar{\Omega}(n; z_h, z) = \omega(n; z_h, z).$$

Now we define $\mathcal{H}^* := \mathcal{H} \setminus (\mathcal{N}_0 \cup \mathcal{N}_1 \cup \mathcal{N}_2)$ and

$$\begin{aligned}\mathcal{N}_3 &:= \{n \in \mathcal{H}^* : \min(u, f_2) \leq Z\}, \\ \mathcal{N}_4 &:= \{n \in \mathcal{H}^* : \min(u, f_2) > z^{1/10}\}, \\ \mathcal{N}_5 &:= \{n \in \mathcal{H}^* : Z < \min(u, f_2) \leq z^{1/10}\}.\end{aligned}$$

In the above decomposition, the main parts are \mathcal{N}_2 and \mathcal{N}_5 . We expect \mathcal{N}_2 to be small since, conditionally on $\bar{\Omega}(n; z) = k$, the normal value of $\bar{\Omega}(n; z_h, z)$ is $hk/\log_2 z > \frac{19}{10}h$. It is more difficult to see that \mathcal{N}_5 is small too. This follows from the fact that we count integers in this set according to their number of factorizations in the form $n = uvf_1 f_2$ with $y < vf_1 < vf_2 \leq z$. Suppose for instance that $f_1, f_2 \leq z_j$. For $\bar{\Omega}(n; z) = k$ and $\bar{\Omega}(n; z_j, z) = G$, then, ignoring the localization of vf_1 and vf_2 in $(y, z]$, there are $4^{k-G}2^G = 4^k 2^{-G}$ such factorizations. Thus, larger G means fewer factorizations. On probabilistic grounds, larger G should also mean fewer factorizations with the localization of vf_1 and vf_2 .

We now briefly consider the cases of \mathcal{N}_0 and \mathcal{N}_1 .

Trivially,

$$|\mathcal{N}_0| \leq \frac{x}{\log z} + \sum_{d > \log z} \frac{x}{d^2} \ll \frac{x}{\log z} \ll \frac{x}{(\log y)^{Q(\lambda)} \sqrt{\log_2 y}}, \quad (3.6)$$

since $Q(\lambda) \leq Q(2) = \log 4 - 1$ in the range under consideration.

By the argument on pages 40–41 of [3],

$$\sum_{\substack{n \leq x \\ \bar{\Omega}(n; z) > K}} 1 \ll \frac{x}{(\log y)^{Q(\lambda)} \sqrt{\log_2 y}}.$$

Setting $t := 1 - \frac{3}{2}\varepsilon$, Lemma 2.4 gives

$$\begin{aligned} \sum_{\substack{n \leq x \\ \tau(n; y, z) \geq 1 \\ \bar{\Omega}(n; z) \leq K_0}} 1 &\leq t^{-(2-3\varepsilon)\log_2 z} \sum_{\substack{dm \leq x \\ y < d \leq z}} t^{\bar{\Omega}(d) + \bar{\Omega}(m; z)} \ll x(\log z)^{2t-2-\beta-(2-3\varepsilon)\log t} \\ &\ll x(\log y)^{-\beta-2\varepsilon^2} \ll x(\log y)^{-Q(\lambda)-\varepsilon^2/2}. \end{aligned}$$

Therefore,

$$|\mathcal{N}_1| \ll \frac{x}{(\log y)^{Q(\lambda)} \sqrt{\log_2 y}}. \quad (3.7)$$

In the next four sections, we show that

$$|\mathcal{N}_j| \ll \frac{x}{(\log y)^{Q(\lambda)} \sqrt{\log_2 y}} \quad (2 \leq j \leq 5). \quad (3.8)$$

Together with (3.6) and (3.7), this will complete the proof of Theorem 1.

4 Estimation of $|\mathcal{N}_2|$

We plainly have $|\mathcal{N}_2| \leq \sum_h |\mathcal{N}_{2,h}|$. For $1 \leq h \leq 5\varepsilon \log_2 z$, the numbers $n \in \mathcal{N}_{2,h}$ satisfy

$$\begin{cases} x/\log z < n \leq x, \\ k := \bar{\Omega}(n; z) = K - b, \quad 0 \leq b \leq 3\varepsilon \log_2 z, \\ \bar{\Omega}(n; z_h, z) \leq \frac{19}{10}h - \frac{1}{100}b, \end{cases}$$

We note at the outset that $\mathcal{N}_{2,h}$ is empty unless $h \geq b/190$.

Write $n = du$ with $y < d \leq z$ and $u \leq x/y$. Let

$$\bar{\Omega}(d; z_h) = D_1, \quad \Omega(d; z_h, z) = D_2, \quad \bar{\Omega}(u; z_h) = U_1, \quad \Omega(u; z_h, z) = U_2,$$

so that $D_1 + D_2 \geq 1$, $D_2 + U_2 \leq \frac{19}{10}h - \frac{1}{100}b$ and $D_1 + D_2 + U_1 + U_2 = k$.

Fix $k = K - b$, h , D_1 , D_2 , U_1 and U_2 . By Lemma 2.3 (with $w = z_h$, $a = U_1$, $b = U_2$), the number of u is

$$\ll \frac{x}{y \log z} \frac{(\log_2 z - h)^{U_1}}{U_1!} (U_2 + 1) \frac{(h + C)^{U_2}}{U_2!}.$$

A second application of Lemma 2.3 yields that the number of d is

$$\ll \frac{\eta y}{\log z} \frac{(\log_2 z - h)^{D_1}}{D_1!} (D_2 + 1) \frac{(h + C)^{D_2}}{D_2!}.$$

Since $D_2 + U_2 < 2h$, we have $(h + C)^{U_2 + D_2} \leq e^{2C} h^{U_2 + D_2}$. Summing over D_1, D_2, U_1, U_2 with $G = D_2 + U_2$ fixed and using the binomial theorem, we find that the number of n in question is

$$\ll \frac{\eta x}{(\log z)^2} (\log_2 z - h)^{k-G} h^G (G + 1)^2 \sum_{\substack{U_1 + D_1 = k - G \\ D_2 + U_2 = G}} \frac{1}{U_1! D_1! D_2! U_2!} \ll \frac{\eta x 2^k}{(\log z)^2} A(h, G),$$

where

$$A(h, G) = (G + 1)^2 \frac{(\log_2 z - h)^{k-G} h^G}{(k - G)! G!}.$$

Since $G + 1 \leq G_h := \lfloor \frac{19}{10} h \rfloor$, we have

$$\frac{A(h, G + 1)}{A(h, G)} \geq \frac{h(k - G)}{(G + 1)(\log_2 z - h)} \geq \frac{k - 10\varepsilon \log_2 z}{1.9(1 - 5\varepsilon) \log_2 z} > \frac{21}{20}$$

if ε is small enough. Next,

$$\begin{aligned} A(h, G_h) &\leq (G_h + 1)^2 \frac{(\log_2 z - h)^{k-G_h} (hk)^{G_h}}{k! (G_h/e)^{G_h}} \\ &\ll (h + 1)^2 \frac{(\log_2 z)^k}{k!} \left(\frac{20}{19}e\right)^{19h/10} e^{-h(k-G_h)/\log_2 z} \\ &\ll \frac{(\log_2 z)^k}{k!} e^{-h/500}, \end{aligned}$$

since $(k - G_h)/\log_2 z > 2 - 13\varepsilon$ and $\frac{19}{10} \log(\frac{20}{19}e) < 2 - 1/400$. Thus,

$$\sum_{b/190 \leq h \leq 5\varepsilon \log_2 z} \sum_{0 \leq G \leq G_h} A(h, G) \ll \sum_{b/190 \leq h \leq 5\varepsilon \log_2 z} A(h, G_h) \ll \frac{(\log_2 z)^k}{k!} e^{-b/95000}$$

and so

$$\sum_{\substack{n \in \mathcal{N}_2 \\ \Omega(n; z) = k}} 1 \ll \frac{\eta x (2 \log_2 z)^k}{(\log z)^2 k!} e^{-(K-k)/95000} \ll \frac{x e^{-(K-k)/95000}}{(\log y)^{Q(\lambda)} \sqrt{\log_2 y}},$$

by (3.3). Summing over the range $K_0 \leq k \leq K$ furnishes the required estimate (3.8) for $j = 2$.

5 Estimation of $|\mathcal{N}_3|$

All integers $n = f_1 f_2 u v$ counted in \mathcal{N}_3 verify

$$\begin{cases} x/\log z < n \leq x, \\ \bar{\Omega}(n; z) \leq K, \\ y < v f_1 < v f_2 \leq z, \quad \min(u, f_2) \leq Z, \end{cases}$$

where Z is defined in (3.5). This is all we shall use in bounding $|\mathcal{N}_3|$.

Let $\mathcal{N}_{3,1}$ be the subset corresponding to the condition $f_2 \leq Z$ and let $\mathcal{N}_{3,2}$ comprise those $n \in \mathcal{N}_3$ such that $u \leq Z$.

If $f_2 \leq Z$, then $v > z^{1/2}$ and $u > x/\{v Z^2 \log z\} > x^{1/3}$. For $\frac{1}{2} \leq t \leq 1$ we have

$$\begin{aligned} |\mathcal{N}_{3,1}| &\leq \sum_{f_1, f_2, v, u} t^{\bar{\Omega}(f_1 f_2 u v; z) - K} \\ &= t^{-K} \sum_{f_1 \leq Z} t^{\bar{\Omega}(f_1)} \sum_{f_1 < f_2 \leq e^\eta f_1} t^{\bar{\Omega}(f_2)} \sum_{y/f_1 < v \leq z/f_1} t^{\bar{\Omega}(v)} \sum_{u \leq x/f_1 f_2 v} t^{\bar{\Omega}(u; z)}. \end{aligned}$$

Apply Lemma 2.4 to the three innermost sums. The u -sum is

$$\ll \frac{x}{f_1 f_2 v} (\log z)^{t-1} \leq \frac{x}{f_1 y} (\log z)^{t-1}.$$

and the v -sum is

$$\ll \frac{\eta y}{f_1} (\log z)^{t-1}.$$

The f_2 -sum is $\ll \eta f_1 (\log f_1)^{t-1}$ if $f_1 > \eta^{-3}$ and otherwise is $\ll \eta f_1$ trivially (note that $\eta f_1 \gg 1$ follows from the fact that $(f_1 + 1)/f_1 \leq f_2/f_1 \leq e^\eta$). Next

$$\begin{aligned} \sum_{f_1 \leq \eta^{-3}} \frac{1}{f_1} + \sum_{2 \leq f_1 \leq Z} \frac{t^{\bar{\Omega}(f_1)}}{f_1} (\log f_1)^{t-1} &\ll \log_2 z + (\log_2 z) \max_{j \leq \log_2 Z} e^{j(t-1)} \sum_{f_1 \leq \exp\{e^j\}} \frac{t^{\bar{\Omega}(f_1)}}{f_1} \\ &\ll (\log_2 z) (\log Z)^{2t-1}. \end{aligned}$$

Thus,

$$|\mathcal{N}_{3,1}| \ll x (\log_2 x) (\log x)^E$$

with $E = -2\beta - \lambda \log t + 2t - 2 + (2t - 1)(1 - 4\varepsilon)$. We select optimally $t := \frac{1}{4}\lambda/(1 - 2\varepsilon)$, and check that $t \geq \frac{1}{2}$ since $\lambda \geq 2 - \varepsilon/\log 2$. Then

$$\begin{aligned} E &= -Q(\lambda) + \lambda \log(1 - 2\varepsilon) + 4\varepsilon \leq -Q(\lambda) + (2 - \varepsilon/\log 2)(-2\varepsilon - 2\varepsilon^2) + 4\varepsilon \\ &< -Q(\lambda) - \varepsilon^2. \end{aligned}$$

Next, we consider the case when $u \leq Z$. We observe that this implies

$$\frac{1}{4}vz^2 \leq vx \leq vn \log z = u f_1 v f_2 v \log z \leq Z z^2 \log z$$

hence $v \leq 4Z \log z \leq Z^2$, and therefore

$$\min(f_1, f_2) > z^{1/2}.$$

Also, $z > x^{1/3}$ since $x/\log z < n = uv f_1 f_2 \leq Z z^2$. Thus, for $\frac{1}{2} \leq t \leq 1$, we have

$$\begin{aligned} |\mathcal{N}_{3,2}| &\leq \sum_{f_1, f_2, v, u} t^{\bar{\Omega}(f_1 f_2 uv; z) - K} \\ &= t^{-K} \sum_{v \leq Z^2} t^{\bar{\Omega}(v)} \sum_{u \leq xv/y^2} t^{\bar{\Omega}(u)} \sum_{y/v < f_1 \leq z/v} t^{\bar{\Omega}(f_1)} \sum_{y/v < f_2 \leq z/v} t^{\bar{\Omega}(f_2)}. \end{aligned}$$

The sums upon f_1 and f_2 are each

$$\ll \frac{\eta y}{v} (\log z)^{t-1}$$

and the u -sum is

$$\ll \frac{xv}{y^2} (\log 2xv/y^2)^{t-1} \leq \frac{xv}{y^2} (\log 2v)^{t-1}.$$

Thus, selecting the same value $t := \frac{1}{4}\lambda/(1 - 2\varepsilon)$, we obtain

$$\begin{aligned} |N_{3,2}| &\ll t^{-K} x \eta^2 (\log z)^{2t-1} \sum_{v \leq Z^2} \frac{t^{\bar{\Omega}(v)} (\log 2v)^{t-1}}{v} \\ &\ll x (\log_2 z) (\log z)^E \leq x (\log_2 z) (\log z)^{-Q(\lambda) - \varepsilon^2}. \end{aligned}$$

This completes the proof of (3.8) with $j = 3$.

6 Estimation of $|\mathcal{N}_4|$

We now consider those integers $n = f_1 f_2 u v$ such that

$$\begin{cases} x/\log z < n \leq x, \\ k := \bar{\Omega}(n; z) = K - b, \quad 0 \leq b \leq 3\varepsilon \log_2 z, \\ y < v f_1 < v f_2 \leq z, \quad \min(u, f_2) > z^{1/10}. \end{cases}$$

With the notation (3.4), fix k, F_1, F_2, U and V . Here u, f_1 and f_2 are all $> \frac{1}{2}z^{1/10}$. By Lemma 2.3 (with $w = z$), for each triple f_1, f_2, v the number of u is

$$\ll \frac{x}{f_1 f_2 v \log z} \frac{(\log_2 z)^U}{U!}.$$

Using Lemma 2.3 two more times, we obtain, for each v ,

$$\sum_{y/v < f_1 \leq z/v} \frac{1}{f_1} \sum_{y/v < f_2 \leq z/v} \frac{1}{f_2} \ll \frac{\eta^2}{(\log z)^2} \frac{(\log_2 z)^{F_1+F_2}}{F_1! F_2!}.$$

Now, Lemma 2.2 gives

$$\sum_v \frac{1}{v} \ll \frac{(\log_2 z)^V}{V!}.$$

Gathering these estimates and using (3.3) yields

$$\begin{aligned} |\mathcal{N}_4| &\ll \frac{x\eta^2}{(\log z)^3} \sum_{(2-3\varepsilon)\log_2 z \leq k \leq K} \sum_{F_1+F_2+U+V=k} \frac{(\log_2 z)^k}{F_1! F_2! U! V!} \\ &= \frac{x\eta^2}{(\log z)^3} \sum_{(2-3\varepsilon)\log_2 z \leq k \leq K} \frac{(2\log_2 z)^k}{k!} 2^k \\ &\ll \frac{x}{(\log y)^{Q(\lambda)} \sqrt{\log_2 y} \log z} \frac{2^K \eta}{\sqrt{\log_2 y} \log z} \ll \frac{x}{(\log y)^{Q(\lambda)} \sqrt{\log_2 y}}. \end{aligned}$$

Thus (3.8) holds for $j = 4$.

7 Estimation of $|\mathcal{N}_5|$

It is plainly sufficient to bound the number of those $n = f_1 f_2 u v$ satisfying the following conditions

$$\begin{cases} x/\log z < n \leq x, \\ k := \bar{\Omega}(n; z) = K - b, \quad 0 \leq b \leq 3\varepsilon \log_2 z, \\ \bar{\Omega}(n; z_h, z) > \frac{19}{10}h - \frac{1}{100}b \quad (1 \leq h \leq 5\varepsilon \log_2 z) \\ y < v f_1 < v f_2 \leq z, \quad Z < \min(u, f_2) \leq z^{1/10}. \end{cases}$$

Define j by $z_{j+2} < \min(u, f_2) \leq z_{j+1}$. We have $1 \leq j \leq 5\varepsilon \log_2 z$. Let $\mathcal{N}_{5,1}$ be the set of those n satisfying the above conditions with $u \leq z_{j+1}$ and let $\mathcal{N}_{5,2}$ be the complementary set, for which $f_2 \leq z_{j+1}$.

If $u \leq z_{j+1}$, then $v \leq (z^2 u \log z)/x \leq 4u \log z \leq z_j$ and $f_2 > f_1 > z^{1/2}$. Recall notation (3.4) and write

$$F_{11} := \overline{\Omega}(f_1; z_j), \quad F_{12} := \Omega(f_1; z_j, z), \quad F_{21} := \overline{\Omega}(f_2; z_j), \quad F_{22} := \Omega(f_2; z_j, z),$$

so that the initial condition upon $\overline{\Omega}(n; z_h, z)$ with $h = j$ may be rewritten as

$$F_{12} + F_{22} \geq G_j := \max(0, \lfloor \frac{19}{10}j - b/100 \rfloor).$$

We count those n in a dyadic interval $(X, 2X]$, where $x/(2 \log z) \leq X \leq x$. Fix k, j, X, U, V, F_{rs} and apply Lemma 2.3 to sums over u, f_1, f_2 . The number of n is question is

$$\begin{aligned} &\leq \sum_{v \leq z_j} \sum_{vX/z^2 \leq u \leq 2vX/y^2} \sum_{y/v < f_1 \leq z/v} \sum_{y/v < f_2 \leq z/v} 1 \\ &\ll \frac{\eta^2 X e^j}{(\log z)^3} \frac{(\log_2 z - j)^{U+F_{11}+F_{21}}}{U!F_{11}!F_{21}!} (F_{12} + 1)(F_{22} + 1) \frac{(j + C)^{F_{12}+F_{22}}}{F_{12}!F_{22}!} \sum_{v \leq z_j} \frac{1}{v}. \end{aligned}$$

Bounding the v -sum by Lemma 2.2, and summing over X, U, V, F_{rs} with $F_{12} + F_{22} = G$ yields

$$|\mathcal{N}_{5,1}| \ll \frac{\eta^2 x}{(\log z)^3} \sum_{(2-3\varepsilon) \log_2 z \leq k \leq K} 4^k \sum_{1 \leq j \leq 5\varepsilon \log_2 z} \sum_{G_j \leq G \leq k} M(j, G),$$

where

$$M(j, G) := e^j (G + 1)^2 \frac{(\log_2 z - j)^{k-G} (j + C)^G}{2^G (k - G)! G!}.$$

Let $j_b = \lfloor \frac{1}{2}b + 100C + 100 \rfloor$. If $j \leq j_b$, then $j + C \leq \frac{99}{100}(j + C_b)$ with $C_b := 3C + 2 + \frac{b}{100}$ and, introducing $R := \max_{G \geq 0} \{(G + 1)^2 (\frac{99}{100})^G\}$, we have

$$\begin{aligned} \sum_{1 \leq j \leq j_b} \sum_{G_j \leq G \leq k} M(j, G) &\leq R \sum_{1 \leq j \leq j_b} e^j \sum_{0 \leq G \leq k} \frac{(\log_2 z - j)^{k-G} (j + C_b)^G}{2^G G! (k - G)!} \\ &\ll \frac{1}{k!} \sum_{1 \leq j \leq j_b} e^j (\log_2 z - \frac{1}{2}j + \frac{1}{2}C_b)^k \\ &\ll \frac{(\log_2 z)^k}{k!} \sum_{1 \leq j \leq j_b} e^{j+(b/200-j/2)k/\log_2 z} \\ &\ll \frac{(\log_2 z)^k}{k!} e^{b/100+2\varepsilon j_b} \ll \frac{(\log_2 z)^k}{k!} e^{b/50}. \end{aligned}$$

When $j > j_b$, then

$$G_j \geq \frac{9}{5}(j + C) + \frac{1}{10}(j_b + C + 1) - \frac{1}{100}b - 1 \geq \frac{9}{5}(j + C) + 9 \geq 189.$$

Thus, for $G \geq G_j$ we have

$$\frac{M(j, G+1)}{M(j, G)} = \left(\frac{G+2}{G+1} \right)^2 \frac{j+C}{2(G+1)} \frac{k-G}{\log_2 z - j} \leq \frac{4}{7}.$$

Therefore,

$$\begin{aligned} \sum_{G_j \leq G \leq k} M(j, G) &\ll M(j, G_j) \ll \frac{j^2 e^j (\log_2 z - j)^{k-G_j} (jk)^{G_j}}{k! 2^{G_j} G_j!} \\ &\leq \frac{j^2 e^j (\log_2 z)^k}{k!} e^{-j(k-G_j)/\log_2 z} \left(\frac{e j k}{2 G_j \log_2 z} \right)^{G_j} \ll \frac{(\log_2 z)^k}{k!} e^{-j/5}, \end{aligned}$$

since $k - G_j \geq (2 - 10\varepsilon) \log_2 z$, $e j k / (2 G_j \log_2 z) \leq \frac{5}{9} e$, and $-1 + \frac{19}{10} \log(\frac{5}{9} e) < -\frac{1}{5}$. We conclude that

$$\sum_{1 \leq j \leq 5\varepsilon \log_2 z} \sum_{G_j \leq G \leq k} M(j, G) \ll \frac{(\log_2 z)^k}{k!} e^{b/50} \quad (7.1)$$

and hence, by (3.3),

$$|\mathcal{N}_{5,1}| \ll \frac{\eta^2 x}{(\log z)^3} \sum_{k \leq K} \frac{(2 \log_2 z)^k}{k!} 2^{K-b/2} \ll \frac{\eta^2 2^K x}{(\log z)^3} \frac{(2 \log_2 z)^K}{K!} \ll \frac{x}{(\log y)^{Q(\lambda)} \sqrt{\log_2 y}}.$$

Now assume $f_2 \leq z_{j+1}$. Then $\min(u, v) > \sqrt{z}$. Fix F_1, F_2 and

$$\bar{\Omega}(v; z_j) = V_1, \quad \Omega(v; z_j, z) = V_2, \quad \bar{\Omega}(u; z_j) = U_1, \quad \Omega(u; z_j, z) = U_2.$$

By Lemma 2.3, given f_1, f_2 and v , the number of u is

$$\ll \frac{x}{f_1 f_2 v \log z} \frac{(\log_2 z - j)^{U_1} (U_2 + 1) (j + C)^{U_2}}{U_1! U_2!}.$$

Applying Lemma 2.3 again, for each f_1 we have

$$\sum_{\substack{f_1 < f_2 \leq e^\eta f_1 \\ y/f_1 < v \leq z/f_1}} \frac{1}{f_2 v} \ll \frac{\eta^2 e^j}{(\log z)^2} \frac{(V_2 + 1) (\log_2 z - j)^{V_1 + F_2} (j + C)^{V_2}}{V_1! V_2! F_2!}.$$

By Lemma 2.2,

$$\sum_{f_1 \leq z_j} \frac{1}{f_1} \ll \frac{(\log_2 z - j)^{F_1}}{F_1!}.$$

Combine these estimates, and sum over $F_1, F_2, U_1, U_2, V_1, V_2$ with $V_2 + U_2 = G$. As in the estimation of $|\mathcal{N}_{5,1}|$, sum over k, j, G using (3.3) and (7.1). We obtain

$$\begin{aligned} |\mathcal{N}_{5,2}| &\ll \frac{\eta^2 x}{(\log z)^3} \sum_{(2-3\varepsilon) \log_2 z \leq k \leq K} 4^k \sum_{1 \leq j \leq 5\varepsilon \log_2 z} \sum_{G_j < G \leq k} M(j, G) \\ &\ll \frac{x}{(\log y)^{Q(\lambda)} \sqrt{\log_2 y}}. \end{aligned}$$

References

- [1] K. Ford, The distribution of integers with a divisor in a given interval, (2004).
- [2] R.R Hall, *Sets of multiples*, Cambridge Tracts in Mathematics 118, Cambridge University Press, Cambridge 1996.
- [3] R. R. Hall and G. Tenenbaum, *Divisors*, Cambridge Tracts in Mathematics, vol. 90, Cambridge University Press, Cambridge, 1988.
- [4] R. R. Hall and G. Tenenbaum, The set of multiples of a short interval, in : D.V. Chudnovsky *et al.* (eds), *Number Theory* (New York Seminar 1989–90), 119–128 (Springer Verlag 1991).
- [5] G. Tenenbaum, Sur la probabilité qu'un entier possède un diviseur dans un intervalle donné, *Compositio Math.* **51** (1984), 243–263.