

# On the uncapacitated plant location problem I: valid inequalities and facets

D. Chinhyung Cho, E.L. Johnson, M.W. Padberg, M.R. Rao

► **To cite this version:**

D. Chinhyung Cho, E.L. Johnson, M.W. Padberg, M.R. Rao. On the uncapacitated plant location problem I: valid inequalities and facets. RR-0086, INRIA. 1981. inria-00076475

**HAL Id: inria-00076475**

**<https://hal.inria.fr/inria-00076475>**

Submitted on 24 May 2006

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# IRIA

CENTRE DE ROCQUENCOURT

Institut National  
de Recherche  
en Informatique  
et en Automatique

Domaine de Voluceau  
Rocquencourt  
B.P.105  
78153 Le Chesnay Cedex  
France  
Tél: 954 90 20

## Rapports de Recherche

N° 86

### ON THE UNCAPACITATED PLANT LOCATION PROBLEM I: VALID INEQUALITIES AND FACETS

D. Chinyung CHO  
Ellis L. JOHNSON  
Manfred W. PADBERG  
M. R. RAO

Juillet 1981

ON THE UNCAPACITATED PLANT LOCATION PROBLEM I :

VALID INEQUALITIES AND FACETS

by

D. CHINHYUNG CHO\*)

Ellis L. JOHNSON\*\*)

Manfred W. PADBERG\*\*\*)

M.R. RAO\*\*\*\*)

- 
- \* ) GBA, New York University, New York, N.Y., U.S.A.
  - \*\* ) IBM Research, Yorktown Heights, N.Y., U.S.A. and Universität Bonn, Bonn, R.F.A.
  - \*\*\* ) GBA, New York University, New York, N.Y., U.S.A. and INRIA, Rocquencourt, France.
  - \*\*\*\* ) Indian Institute of Management, Bangalore, India.

Résumé :

Nous considérons un problème standard de la recherche opérationnelle concernant l'implantation des installations appelé "uncapacitated plant location problem" (PLP) dans la littérature anglaise. Nous voyons ce problème ici comme un problème défini sur un graphe où il faut trouver un ensemble de sommets indépendants. Pour le problème PLP, nous montrons plusieurs inégalités valides dont quelques unes définissent des facettes du polytope avec des sommets zéro ou un associé à ce problème. Nous déduisons des conditions nécessaires et suffisantes pour les facettes dites triviales et, en plus, des conditions nécessaires pour des facettes générales. Pour le cas particulier de ce problème à trois installations et à trois destinations ou plus, nous donnons une description complète du polytope associé à des inégalités linéaires.

Abstract :

The uncapacitated plant location problem is considered as a node-packing problem. For this problem, several valid inequalities and facets are discussed. Necessary and sufficient conditions for trivial facets along with necessary conditions for non-trivial facets are derived. In addition, all of the facets for the case of 3 plants and 3 or more destinations are identified.

## 1. Introduction.

The plant location problem is the problem of finding the optimal location of plants so as to minimize a cost function while satisfying given demands. The problem is called the uncapacitated (or simple) plant location problem if each plant under consideration has the ability to satisfy the total demand. The plant location problem is a standard problem of the OR/MS literature and is of interest not only for its own merits but also for its close similarities to other classes of real-world problems. (See [10] for a detailed bibliography.)

Weber [24], who may be credited with the formulation of the problem, and other researchers attempted to solve the problem by way of locating the centeroids. This approach often resulted in locating the plants at impossible locations; for example, at the top of a mountain or in the middle of a lake. This impossibility of the optimal location was later resolved by constraining the problem to allow only a finite number of feasible locations.

In fact, since 1954 when Hirsch and Dantzig [17] introduced a linear programming method for solving the fixed charge problem, most of the studies have been carried out along a linear programming approach using 0-1 decision variables. The majority of the researchers attempted to solve this mixed integer programming problem by using the branch and bound technique, see e.g. Efrøymson and Ray [9] and Spielberg [25]. However, the computational results have not been satisfactory, especially when faced with problems of a large size. This can be expected because the problem is NP-complete, i.e. it belongs to the class of hard combinatorial optimization problems which to date cannot be solved by polynomially bounded algorithms [12].

Recently, several researchers have begun to study facets of the uncapacitated plant location problem. (See Guignard [16] and Cornuejols and Thizy [6].) This approach deserves attention since facets are "strongest cutting planes". One can thus reasonably expect to improve computational results substantially

even if one can identify only a subset of these facets. This point has been well illustrated recently with related work on the travelling salesman problem by Grötschel and Padberg [14, 15], Crowder and Padberg [7] and Padberg and Hong [23].

Our objective, then, is to investigate some of these facet inequalities that describe the integer polyhedron of the uncapacitated plant location problem.

In section 2, we formulate the problem as the node-packing problem (PLP) and introduce some notation that will be used through-out the study. Once formulated in this way, it follows that the constraint matrix of (PLP) is totally unimodular if the number of plants,  $p$ , is less than 3 or the number of destinations,  $d$ , is less than 3. This implies that the linear programming relaxation of (PLP) will give an integer solution and thus it is of no interest to us here. However, if  $p \geq 3$  and  $d \geq 3$ , then linear programming solution will not always be integer and thus our concern is how we can eliminate these non-integer solutions.

Section 3 deals with a particular class of valid inequalities which remove some fractional extreme points of the polyhedron associated with the linear programming relaxation of (PLP). It is demonstrated that all of the valid inequalities studied to date are members of this class.

In section 4, we show that the non-negativity constraints and clique constraints are all facets and prove several necessary and sufficient conditions that characterize facets.

The results of section 4 and the necessary conditions for non-trivial facets of section 5 enable us to give a description of all facets of (PLP) when  $p = 3$  and  $d \geq 3$ . In fact, theorem 6.1 states that all of these facets correspond to odd-cycles of length 9 in the associated intersection graph.

In a sequel to this paper [4], we discuss several lifting procedures and

present necessary and sufficient conditions for non-trivial facets with 0-1 coefficients. We also derive a family of facets and describe all facets for the case of several plants ( $p \geq 3$ ) and three destinations.

## 2. Problem formulation and notation.

The uncapacitated plant location problem arises from the following situation.

Suppose that, at  $p$  possible locations, plants can be built necessitating a fixed investment  $f_i$  at location  $i$  ( $i = 1, 2, \dots, p$ ) for producing a homogeneous good demanded from destination  $j$  ( $j = 1, 2, \dots, d$ ). Assuming that the capacity of each plant is sufficiently large to satisfy the total demand, one has to decide at which locations to build the plants so as to minimize the total cost of satisfying the destination's demands.

Denoting by  $x_{ij}$  the fraction of destination  $j$ 's demand filled by a plant  $i$  and letting  $y_i' = 1$  if a plant is located in location  $i$  ( $y_i' = 0$  if not), the problem can be formulated as the following mixed integer programming problem.

$$\text{Min. } \sum_j \sum_i c_{ij} x_{ij} + \sum_i f_i y_i'$$

subject to

$$\sum_i x_{ij} = 1 \quad \text{for } j = 1, 2, \dots, d$$

$$0 \leq x_{ij} \leq y_i' \quad \text{for } i = 1, 2, \dots, p; j = 1, 2, \dots, d$$

$$y_i' = 0 \text{ or } 1 \quad \text{for } i = 1, 2, \dots, p$$

Here,  $c_{ij}$  are the transportation costs involved in satisfying the total demand of destination  $j$  from plant  $i$  and, moreover, transportation costs are assumed to be linear.

This formulation can be transformed into a set-packing problem by interchanging the variables  $y_i'$  with  $1 - y_i'$  and by converting the set of  $d$  equality constraints



into less-than-or-equal-to constraints. The latter can be accomplished by introducing an artificial variable for each equation having a "large enough" cost (e.g.  $\theta_j > \sum_i c_{ij}$ ) and subsequent elimination of these variables. More precisely, the interchange of variables  $y_i$  with  $1-y_i$  transforms the problem into the form

$$\begin{aligned} & \text{Min. } \sum_j \sum_i c_{ij} x_{ij} - \sum_i f_i y_i + \sum_i f_i \\ & \text{subject to} \\ & \sum_i x_{ij} = 1 \quad \text{for } j = 1, 2, \dots, d \\ & x_{ij} + y_i \leq 1 \quad \text{for } i = 1, 2, \dots, p; j = 1, 2, \dots, d \\ & y_i = 0 \text{ or } 1 \quad \text{for } i = 1, 2, \dots, p \\ & x_{ij} \geq 0 \quad \text{for } i = 1, 2, \dots, p; j = 1, 2, \dots, d. \end{aligned}$$

(PLPE)

Applying the second transformation, (PLPE) can be brought to the following equivalent form:

$$\begin{aligned} & \text{Max. } \sum_j \sum_i (\theta_j - c_{ij}) x_{ij} + \sum_i f_i y_i - \sum_i f_i - \sum_j \theta_j \\ & \text{subject to} \\ & \sum_i x_{ij} \leq 1 \quad \text{for } j = 1, 2, \dots, d \quad (2.1) \\ & x_{ij} + y_i \leq 1 \quad \text{for } i = 1, 2, \dots, p; j = 1, 2, \dots, d \quad (2.2) \\ & y_i = 0 \text{ or } 1 \quad \text{for } i = 1, 2, \dots, p \\ & x_{ij} \geq 0 \quad \text{for } i = 1, 2, \dots, p; j = 1, 2, \dots, d. \end{aligned}$$

(PLP)

This formulation is the required set-packing version of the plant location problem.

Remark 2.1. (PLPE) differs from (PLP) by the fact that the former must satisfy the constraints (2.1) with equalities.

Remark 2.2. In this formulation,  $y_i = 0$  means that a plant is located at  $i$  and  $y_i = 1$  means the opposite case. Also, even though  $x_{ij}$  is allowed to take a fractional value, there exists an optimal solution with  $x_{ij} = 0$  or  $1$  if all variables  $y_i$  are fixed at  $0$  or  $1$ .

Let  $A$  be the matrix whose elements are the coefficients of the constraints (2.1) and (2.2) of (PLP). Note that  $A$  is a  $(pd+p) \times (pd+p)$  matrix whose elements are either  $0$  or  $1$ . The intersection graph  $G = (N,E)$  is constructed by assigning a node for each column  $a^i$  of  $A$  and an undirected edge for each pair of nodes whose corresponding columns are non-orthogonal, i.e.  $|N| = pd+p$  and there is an edge between the node  $i$  and the node  $j$  if and only if  $a^i \cdot a^j \neq 0$ . It follows that (PLP) is equivalent to the node-packing problem (NP) in this intersection graph in the sense that they both have the same solution set (including the set of optimal solutions) if for each node  $i$  the node weight is defined to be the cost associated with column  $i$ . (see [8,13, 19,21].)

For convenience, we define some notation.

1)  $P = \{1,2,\dots,p\}$  with  $|P| = p$  and  $D = \{1,2,\dots,d\}$  with  $|D| = d$ .

2)  $L^{pd} = \{(x,y) \in R^{p(d+1)} \mid \sum_i x_{ij} \leq 1 \forall j \in D; x_{ij} + y_i \leq 1, x_{ij} \geq 0, y_i \geq 0 \forall i \in P, j \in D\}$

This is the polyhedron associated with the linear programming relaxation of (PLP) with  $p$  plants and  $d$  destinations. Note that  $\dim(L^{pd}) = p(d+1)$ .

3)  $LE^{pd} = \{(x,y) \in R^{p(d+1)} \mid \sum_i x_{ij} = 1 \forall j \in D; x_{ij} + y_i \leq 1, x_{ij} \geq 0, y_i \geq 0 \forall i \in P, j \in D\}$

This is the polyhedron associated with the linear programming relaxation of (PLPE) with  $p$  plants and  $d$  destinations.

4)  $L_I^{pd} = \text{conv}\{L^{pd} \cap Z^{p(d+1)}\}$  where conv means the convex hull and  $Z^{p(d+1)}$  is the set of all 0-1 vectors of dimension  $p(d+1)$ . This is the integer polyhedron associated with (PLP), i.e. the convexified solution space of (PLP).

$$5) Q^{pd} = \{(\pi, \mu) \in \mathbb{R}^{p(d+1)} \mid \sum_j \pi_{\sigma(j)j} + \sum_{i \in S} \mu_i \leq 1 \forall S \subseteq P; \pi_{ij} \geq 0, \mu_i \geq 0 \forall i \in P, j \in D\}$$

where  $\sigma$  is an onto function from  $D$  into  $\bar{S}$  ( $= P-S$ ). This is the anti-blocker [1] of  $L_I^{pd}$  and thus, if  $\pi x + \mu y \leq 1$  is a facet of  $L_I^{pd}$ , then  $(\pi, \mu)$  is a vertex of  $Q^{pd}$ .

$$6) F^{pd} = \{(\pi, \mu) \in \mathbb{R}^{p(d+1)} \mid \pi x + \mu y \leq \pi_0 \text{ is a facet of } L_I^{pd} \text{ for some } \pi_0\}$$

This is the set consisting of all the possible facets.

$$7) F_c^{pd} = \{(\pi, \mu) \in \mathbb{R}^{p(d+1)} \mid \pi x + \mu y \leq \pi_0 \text{ where } \pi_0 = 0 \text{ or } 1 \text{ and } \pi x + \mu y \text{ is equal to } \sum_i x_{ij} \text{ or } x_{ij} + y_i \text{ or } -x_{ij} \text{ or } -y_i \text{ for some } i \in P, j \in D\}$$

This is the set consisting of all clique constraints and non-negativity constraints.

8)  $H_{\pi, \mu} = L_I^{pd} \cap \{(x, y) \in \mathbb{R}^{p(d+1)} \mid \pi x + \mu y = 1\}$ , where  $\pi x + \mu y \leq 1$  is a valid inequality for  $L_I^{pd}$ . This is the set consisting of all the feasible points that lie on the plane  $\pi x + \mu y = 1$ .

9)  $I_j = \{i \in P \mid \pi_{ij} > 0\}$  for  $j = 1, 2, \dots, d$  for a given  $(\pi, \mu)$  and, likewise,

$$J_i = \{j \in D \mid \pi_{ij} > 0\} \text{ for } i = 1, 2, \dots, p.$$

10) Let  $(\pi, \mu) \in Q^{pd}$ . Then, we partition  $\pi$  as  $\pi = (\pi^1, \pi^2, \dots, \pi^d)$  where

$$\pi^j = (\pi_{1j}, \pi_{2j}, \dots, \pi_{pj}) \text{ for } j = 1, 2, \dots, d \text{ and accordingly for } x \text{ as}$$

$$x = (x^1, x^2, \dots, x^d).$$

- 11) We let  $G = (N, E)$  be the intersection graph associated with (PLP). Recall that each node in  $N$  corresponds to a column of  $A$  whose elements are the coefficients of the constraints (2.1) and (2.2). Hence, we denote the nodes accordingly by their corresponding variables of the constraints (2.1) and (2.2). In other words,  $N$  consists of all the nodes  $y_i$ ,  $i = 1, 2, \dots, p$  and  $x_{ij}$ ,  $i = 1, 2, \dots, p$ ;  $j = 1, 2, \dots, d$  while  $E$  contains all the edges  $(x_{ij}, y_i)$ ,  $i = 1, 2, \dots, p$ ;  $j = 1, 2, \dots, d$  and  $(x_{\mu j}, x_{\nu j})$ ,  $\mu \neq \nu$ ;  $j = 1, 2, \dots, d$ .
- 12) Let  $G^S = (N^S, E^S)$  be a subgraph of  $G$  induced by  $N^S$  where  $N^S \subseteq N$ , i.e.  $G^S$  has the node set  $N^S$  and all edges of  $G$  whose both end points are in  $N^S$ . Let  $I^S = \{i \in P \mid y_i \in N^S\}$  and  $J^S = \{j \in D \mid x_{ij} \in N^S \text{ for some } i \in P\}$ . Also, let  $S$  be an  $|I^S| \times |J^S|$  matrix whose elements are such that  $s_{ij} = 1$  if  $(y_i, x_{ij}) \in E^S$  and  $s_{ij} = 0$  if  $(y_i, x_{ij}) \notin E^S$ . Note that  $s_{ij} = 1$  implies that a plant  $i$  can cover a destination  $j$ . We say that  $S$  is an adjacency matrix if there is no zero column, i.e. every destination in  $J^S$  can be covered by some plant  $i \in I^S$ , and no zero row, i.e. each plant  $i \in I^S$  can cover at least one destination  $j \in J^S$ . Vice versa, any  $|I^S| \times |J^S|$  zero-one matrix  $S = (s_{ij})$  with  $I^S \subseteq P$  and  $J^S \subseteq D$  having no zero column and no zero row is an adjacency matrix and there exists a corresponding subgraph  $G^S = (N^S, E^S)$  of  $G$ , where  $N^S = \{x_{ij} \in N \mid i \in I^S, j \in J^S \text{ and } s_{ij} \neq 0\} \cup \{y_i \in N \mid i \in I^S\}$  and  $E^S = \{(x_{\mu j}, x_{\nu j}) \in E \mid \mu \neq \nu, \mu \in I^S, \nu \in I^S, j \in J^S \text{ and } s_{\mu j}, s_{\nu j} \neq 0\} \cup \{(y_i, x_{ij}) \in E \mid i \in I^S, j \in J^S \text{ and } s_{ij} \neq 0\}$ .
- 13) For any subgraph  $G^S$  of  $G$ , we let  $\alpha(G^S)$  be the independence number of  $G^S$ , i.e. maximum node packing for the intersection graph  $G^S$ , and we let  $\beta(G^S)$  be the covering number of  $G^S$ , i.e. the minimum number of plants  $i \in I^S$  necessary to

cover all destinations  $j \in J^S$ . If no ambiguity arises, we simply write  $\alpha$  or  $\beta$ . We call  $S$  to be a maximal adjacency matrix if changing the zero element to one decreases  $\beta(G^S)$  by one or increases  $\alpha(G^S)$  by one. We call an edge of  $E^S$  to be critical if its removal increases  $\alpha(G^S)$ .

14) For an adjacency matrix  $S$ , let  $L_I^S = L_I^{pd} \cap \{(x,y) \in \mathbb{R}^{P(d+1)} \mid x_{ij} = 0 \text{ for } s_{ij} = 0, \text{ and } y_i = 0, x_{ij} = 0 \text{ for } i \notin I^S, j \notin J^S\}$ .

15) An adjacency matrix  $S$  is called a pd-adjacency matrix if i)  $G^S$  is connected, ii) there exist at least one zero element in each column, and iii)  $|I^S| \geq 3$  and  $|J^S| \geq 3$ . The corresponding subgraph  $G^S$  is called a pd-subgraph.

16)  $\lceil a \rceil$  is the smallest integer greater than or equal to  $a$ .

### 3. Valid inequalities.

An inequality  $ax + by \leq a_0$  is called a valid inequality with respect to  $L_I^{pd}$  if and only if  $L_I^{pd} \subseteq \{(x,y) \in \mathbb{R}^{p(d+1)} \mid ax + by \leq a_0\}$ . The valid inequalities (or cuts) are very useful in removing the fractional extreme vertices that result from the linear relaxation of the integer programming problems. A facet may be viewed as the 'strongest' valid inequality in a sense that it is a valid inequality that intersects the integer polyhedron in a face of the highest possible dimension. However, it should be noted that intersecting  $L_I^{pd}$  by a valid inequality may introduce new fractional extreme points that in turn have to be removed again.

By noting the correspondence between a pd-subgraph  $G^S = (N^S, E^S)$  and its  $|I^S| \times |J^S|$  adjacency matrix  $S$ , we get the following theorem.

Theorem 3.1. For any pd-subgraph  $G^S = (N^S, E^S)$ , the inequality

$$\sum_{i \in I^S} \sum_{j \in J^S} s_{ij} x_{ij} + \sum_{i \in I^S} y_i \leq |I^S| + |J^S| - k \quad (3.1)$$

is a valid inequality of  $L_I^{pd}$  if and only if  $k$  is an integer satisfying  $k \leq \beta(G^S)$ .

Furthermore,  $\alpha(G^S) = |I^S| + |J^S| - \beta(G^S)$ .

Proof:

( $\Rightarrow$ ) Suppose (3.1) is a valid inequality and suppose  $k > \beta(G^S) = \beta$ . Since  $\beta$  plants are sufficient to cover all destinations  $j \in J^S$ , the left hand side of (3.1) can become  $|J^S| + |I^S| - \beta$  which is greater than the right hand side of (3.1). Hence, we have a contradiction.

( $\Leftarrow$ ) Consider any feasible solution such that  $\eta$  plants from  $I^S$  are open.

Case 1.  $\eta \geq \beta$ . Then,  $y_i = 1$  for exactly  $|I^S| - \eta$  plants. But,  
 $|I^S| - \eta \leq |I^S| - \beta \leq |I^S| - k$ . Thus, even if all  $|J^S|$  destinations are covered, the  
left hand side of (3.1) will be less than or equal to  $|I^S| + |J^S| - k$ .

Case 2.  $\eta < \beta$ . Since  $\beta$  plants are required to cover all  $|J^S|$  destinations,  
 $\eta$  plants can cover at most  $|J^S| - (\beta - \eta)$  destinations. Therefore, the left hand  
side of (3.1) can be at most  $|J^S| - (\beta - \eta) + |I^S| - \eta$  which is less than or equal to  
 $|I^S| + |J^S| - k$ .

Now, we prove that  $\alpha(G^S) = |I^S| + |J^S| - \beta$ . First, we note that  $\alpha(G^S) \leq |I^S| + |J^S| - \beta$   
because (3.1) holds when  $k = \beta$ . Without loss of generality, assume that the  
 $\beta$  plants that can cover all destinations are numbered 1 to  $\beta$ , i.e.  $y_i = 0$  for  
 $i = 1, 2, \dots, \beta$  and  $y_i = 1$  for the remaining plants in  $I^S$ . Since the  $\beta$  plants  
can cover all destinations, for each  $j \in J^S$  we can set  $x_{ij} = 1$  where  $i$  is such  
that  $(x_{ij}, y_i)$  is an edge of  $G^S$  and  $1 \leq i \leq \beta$ . Hence,  $\alpha(G^S) \geq |I^S| + |J^S| - \beta$ .  
Therefore,  $\alpha(G^S) = |I^S| + |J^S| - \beta$ . △

Remark 3.1. If (3.1) is a facet, then  $k = \beta(G^S)$  and the right hand side becomes  
 $\alpha(G^S)$ .

A valid inequality is useful only when it can remove some fractional extreme  
points. Cornuejols, Fisher and Nemhauser (C.F.N) showed the following characterization  
of an extreme point for the LP relaxation of (PLPE). (See [5].)

Theorem 3.2. For a given non-integer solution  $(x^f, y^f)$  contained in  $LE^{pd}$ , let  
 $P_f = \{i \in P \mid 0 < v_i^f < 1\}$ ,  $D_f = \{j \in D \mid x_{ij}^f \text{ non-integer for some } i, \text{ and } x_{ij}^f = 0 \text{ or}$   
 $1 - v_i^f \text{ for all } i \in P_f\}$  and  $F$  be the  $|P_f| \times |D_f|$  matrix whose elements are  
 $f_{ij} = \begin{cases} 1 & \text{if } x_{ij}^f > 0 \\ 0 & \text{if } x_{ij}^f = 0 \end{cases}$ . Then, the non-integer solution  $(x^f, y^f)$  is an extreme

point of  $LE^{pd}$  if and only if

$$1) \underline{y_i^f = 1 - \max_j x_{ij}^f \text{ for all } i \in P_f,}$$

$$2) \underline{\text{for each } i \in D, \text{ there exists at most one } i(j) \in P \text{ with } 0 < x_{i(j)j}^f < 1 - y_{i(j)}^f,}$$

$$3) \underline{\text{rank of } F \text{ equals } |P_f|}.$$

Proof: See [5, theorem 6].

If we further define  $P_0 = \{i \in P \mid y_i^f = 0\}$ ,  $P_1 = \{i \in P \mid y_i^f = 1\}$ ,  
 $D_1 = \{j \in D \mid x_{ij}^f \text{ integer for all } i \in P\}$ ,  $D_2 = \{j \in D \mid 0 < x_{i(j)j}^f < 1 - y_{i(j)}^f \text{ for precisely}$   
 one  $i(j) \in P\}$ , then it is easy to see that  $\{P_0, P_1, P_f\}$  and  $\{D_1, D_2, D_f\}$  partition  
 $P$  and  $D$ , respectively. In fact, any fractional extreme point must have the  
 following  $x_{ij}$ 's.

		destinations		
		$D_1$	$D_f$	$D_2$
plants	$P_1$	0	0	0
	$P_f$	0	0 or ** fractionals	0 or fractionals
	$P_0$	0 or *1	0	0 or fractionals

\*: Each column has exactly one 1.

\*\* : There are no zero-rows and each column has at least two fractional numbers. This  $|P_f| \times |D_f|$  matrix corresponds to  $F$ .

If the rank of  $F$  equals  $|P_f|$ ,  $F$  contains a  $|P_f| \times |D_f'|$  non-singular submatrix  $B$  where  $|P_f| = |D_f'|$  and  $D_f' \subseteq D_f$ , and the fractional parts of the extreme point  $(x^f, y^f)$  can be fully identified by solving  $z^t B = e^t$  where  $z^t$  and  $e^t$  are the transpose of  $z_i$ 's and one's, respectively. This is so because an extreme point of  $LE^{pd}$  must satisfy  $\sum_i x_{ij} = 1$ ,  $x_{ij} + y_i \leq 1$  and the three conditions above in theorem 3.2. In other words, we let



$$y_i^f = 1 - z_i \quad \text{for all } i \in P_f$$

$$x_{ij}^f = \begin{cases} z_i & \text{if } f_{ij} = 1 \\ 0 & \text{if } f_{ij} = 0 \end{cases} \quad \forall i \in P_f, j \in D_f \text{ or } \forall i \in P_f - i(j), j \in D_2$$

$$x_{i(j)j}^f = 1 - \sum_{i \in P_f - i(j)} x_{ij}^f \quad \forall j \in D_2$$

Note that B is an adjacency matrix.

Vice versa, it can be seen that there corresponds an extreme point of  $LE^{pd}$  for any non-singular  $k \times k$  adjacency matrix B if  $k \leq \min\{p, d\}$  and  $z^t B = e^t$  has a non-negative solution. Furthermore, an extreme point of  $LE^{pd}$  is also an extreme point of  $L^{pd}$ . Hence, we can derive a valid inequality of  $L_I^{pd}$  that removes some extreme points of  $L^{pd}$  as follows.

Theorem 3.3. Let B be an  $|I^B| \times |J^B|$  non-singular adjacency matrix where  $|I^B| = |J^B|$  and for which  $z^t B = e^t$  has non-negative  $z_i$ 's. If  $\theta(G^B) > \sum_{i \in I^B} z_i$ ,

then the inequality

$$\sum_{i \in I^B} \sum_{j \in J^B} b_{ij} x_{ij} + \sum_{i \in I^B} y_i \leq |I^B| + |J^B| - \theta(G^B) \quad (3.2)$$

is a valid inequality of  $L_I^{pd}$  and removes some fractional extreme points of  $L^{pd}$ .

Proof:

From theorem 3.1, (3.2) is a valid inequality of  $L_I^{pd}$ . Furthermore, from theorem 3.2, there exists a fractional point satisfying

$$\sum_{i \in I^B} \sum_{j \in J^B} b_{ij} x_{ij} + \sum_{i \in I^B} y_i = |J^B| + |I^B| - \sum_{i \in I^B} z_i$$

Therefore, the valid inequality (3.2) removes some fractional extreme points

because  $\theta(G^B) > \sum_{i \in I^B} z_i$ .

△

We now consider some special cases of (3.2). Let  $C$  be a  $k \times k$  cyclic matrix whose rows are 0-1 vectors in which  $t$  consecutive ones are successively moved one position to the right. Furthermore, assume  $k$  and  $t$  are relatively prime with  $t < k$ . Then,  $C$  is a non-singular adjacency matrix and

$\beta(G^C) = \left\lceil \frac{k}{t} \right\rceil > \sum_{i \in I^C} z_i = \frac{k}{t}$ . Hence, the valid inequality (3.2) becomes

$$\sum_{i \in I^C} \sum_{j \in J^C} c_{ij} x_{ij} + \sum_{i \in I^C} y_i \leq 2k - \left\lceil \frac{k}{t} \right\rceil \quad (3.3)$$

In particular, if there are  $k-1$  one's per row, i.e.  $t = k-1$ , then  $\beta(G^C) = 2$  and the valid inequality (3.2) becomes

$$\sum_{i \in I^C} \sum_{j \in J^C} c_{ij} x_{ij} + \sum_{i \in I^C} y_i \leq 2k - 2 \quad (3.4)$$

On the other hand, if  $k$  is odd and  $t = 2$ , then  $C$  is non-singular with  $\beta(G^C) = \left\lceil \frac{k}{2} \right\rceil$  and the valid inequality (3.2) becomes

$$\sum_{i \in I^C} \sum_{j \in J^C} c_{ij} x_{ij} + \sum_{i \in I^C} y_i \leq 2k - \left\lceil \frac{k}{2} \right\rceil \quad (3.5)$$

We note that the inequality (3.3) is exactly the same as the one obtained in [5], (3.4) is the same as the one obtained in [16] and (3.5) is the odd-cycle inequality obtained in [21]. We shall show, in a sequel to this paper [4], that one can lift the inequality (3.3) and (3.5) to be a facet of  $L_I^{pd}$  whereas in [16] it has been shown that (3.4) defines a facet of  $L_I^{pd}$ . (See also [6].) Note that (3.3) remains a valid inequality for  $L_I^{pd}$  if the assumption that  $k$  and  $t$  are relatively prime is dropped. This is an immediate consequence of Theorem 3.1. In the sequel [4] we show how these valid inequalities can be extended to be facets of the polyhedron  $L_I^{pd}$ .

#### 4. Trivial facets.

An inequality  $\pi t \leq \pi_0$  is called a facet of a polyhedron  $P$  if i)  $\pi t \leq \pi_0$  for all  $t \in P$ , and ii) there exists  $n$  affinely independent points  $t^i \in P$  such that  $\pi t^i = \pi_0$  for  $i = 1, 2, \dots, n$  where  $\dim(P) = n$ . The first property implies that  $\pi t \leq \pi_0$  is a valid inequality for  $P$  and the second property implies that there are  $n$  linear independent vectors on the face of the polyhedron if  $\pi_0 \neq 0$ .

From this definition of a facet, we immediately get the following theorem about the non-negativity constraints. (See Balas and Padberg [1] for a similar argument.)

Theorem 4.1. Let  $\pi x + \mu y \leq \pi_0$  be a facet of  $L_I^{pd}$ . Then,  $\pi_0 = 0$  if and only if

$$\pi x + \mu y = -x_{-1j} \text{ or } \pi x + \mu y = -y_{-1i}.$$

Proof:

For notational convenience, we let  $t_{(i-1)d+j} = x_{ij}$  for  $i = 1, 2, \dots, p$ ;  $j = 1, 2, \dots, d$  and  $t_{pd+i} = y_i$  for  $i = 1, 2, \dots, p$ . Then, the above statement can be written as follows: Let  $\pi t \leq \pi_0$  be a facet of  $L_I^{pd}$ . Then  $\pi_0 = 0$  if and only if  $\pi t = -t_{-1i}$ .

( $\Rightarrow$ ) Suppose  $\pi_0 = 0$ . Then, since any unit vector is feasible, we must have  $\pi_i \leq 0$  for all  $i = 1, 2, \dots, pd+p$ . Let  $N^- = \{i \mid \pi_i < 0\}$ . Then, there are  $pd+p - |N^-|$  linearly independent vectors satisfying  $\pi t = 0$ . However, for  $\pi t \leq 0$  to be a facet, we must have  $pd+p-1$  linearly independent vectors satisfying  $\pi t = 0$ . Therefore,  $|N^-| = 1$ .

( $\Leftarrow$ ) Suppose  $\pi_0 \neq 0$ , but  $\pi t = -t_{-1i}$ . Clearly,  $\pi_0 < 0$  is not possible because 0 vector

is a feasible solution. If  $\pi_0 > 0$ , then there exists no feasible vector satisfying  $\pi t = \pi_0$  contradicting the fact that  $\pi t \leq \pi_0$  is a facet. Therefore,  $\pi_0 = 0$ .  $\Delta$

Remark 4.1. If  $\pi_0 \neq 0$ , we can assume without loss of generality that  $\pi_i \geq 0$  for

$i = 1, 2, \dots, pd+p$  and  $\pi_0 = 1$ . This can be seen as follows. Suppose  $N^- \neq \emptyset$ .

Let  $i \in N^-$ . Then, since  $\pi_0 \neq 0$ , there must exist  $\bar{t} \in L_I^{pd}$  such that  $\bar{t}_i = 1$

and  $\bar{\pi} = \pi_0$ . (Otherwise, if  $\bar{t}_i = 0$  for all solutions such that  $\bar{\pi} = \pi_0$ , then all

such solutions also satisfy the facet  $-t_i \leq 0$  with equality. Consequently,

$\pi t \leq \pi_0$  can not be a facet.) But, if we define  $t^*$  to be  $t_j^* = \bar{t}_j$  for  $j \neq i$  and

$t_i^* = 0$  for  $i \in N^-$ , then from the non-negativity of  $A$ , we have  $At^* \leq A\bar{t} \leq e$ , i.e.

$t^*$  feasible, and  $\pi t^* = \bar{\pi} + |\pi_i| > \pi_0$  contradicting the fact that  $\pi t \leq \pi_0$  is a

facet. Consequently,  $N^- = \emptyset$  and we must have  $\pi_i \geq 0$  for  $i = 1, 2, \dots, pd+p$ .

Furthermore,  $\pi_0 > 0$ , which enables us to divide both sides of inequality by  $\pi_0$ .

We, now, give necessary and sufficient conditions for the facets to be clique constraints. Clique constraints have been shown to be facets by Padberg [21] and we will use this fact in the proof of the next theorem repeatedly.

Theorem 4.2. Let  $\pi x + \mu y \leq 1$  be a facet of  $L_I^{pd}$ . Then, the following statements

are true.

1)  $\pi x + \mu y = x_{i,j} + y_i$  for some  $i \in P, j \in D$  if and only if  $\sum_1 \mu_i = 1$ .

2)  $\pi x + \mu y = x_{i,j} + y_i$  if and only if there exists  $j$  such that  $\pi^j \neq 0$  and  
exactly one  $\pi_{i,j} > 0$ .

3)  $\pi x + \mu y = \sum_i x_{i,j}$  for some  $j \in D$  if and only if  $\pi^j > 0$ .

Proof:

1)( $\Rightarrow$ ) This is obvious because we have  $\mu_i = 1$ , and  $\mu_k = 0$  for  $k \neq i$ .

(\*) If  $(\pi, \mu) \in F^{pd}$ , then  $\sum_j \pi_{\sigma(j)j} + \sum_{i \in S} \mu_i \leq 1$  must hold for all  $S \subseteq P$  where

$\sigma$  is an onto function from  $D$  into  $\bar{S}$ . Take  $\bar{S} = \{i\}$ . Then, we must have

$\sum_i \pi_{it} + \sum_{\substack{k \\ k \neq i}} \mu_k \leq 1$  because it is feasible for a plant  $i$  to supply all destinations.

This is equivalent to  $\sum_t \pi_{it} \leq \mu_i$  because of the condition  $\sum_k \mu_k = 1$ . Suppose  $\pi_{ij} > 0$ ,

but  $\pi x + \mu y \neq x_{1j} + y_i$ . Then, all solutions  $(x, y) \in H_{\pi, \mu}$  must have either  $y_i = 1$  or  $y_i = 0$  and  $x_{1j} = 1$ . This is so because, if there exist a solution  $(x^*, y^*) \in H_{\pi, \mu}$

such that  $y_i^* = 0$  and  $x_{1j}^* = 0$ , then we can let  $(\bar{x}, \bar{y})$  be same with  $(x^*, y^*)$  except

$\bar{y}_i = 1$  and  $\bar{x}_{1t} = 0$  for  $t \in J_1$  so that  $(\bar{x}, \bar{y})$  is feasible but

$\pi \bar{x} + \mu \bar{y} = \pi x^* + \mu y^* + \mu_i - \sum_{t \in J_1} \pi_{1t} + \pi_{1j} > 1$  leading to the contradiction.

2)(=) Obvious.

(\*) Without loss of generality, suppose  $\pi^j \neq 0$  and  $\pi_{1j} > 0$ ,  $\pi_{kj} = 0$  for  $k \geq 2$ .

Furthermore, suppose  $\pi x + \mu y \neq x_{1j} + y_1$ . Now, there must exist an integer vector

$(x, y) \in H_{\pi, \mu}$  such that  $x_{1j} + y_1 < 1$ , which implies  $x_{1j} = 0 = y_1$ . This is so

because  $x_{1j} + y_1 \leq 1$  is a facet and  $(x, y)$  are integers. Since  $\pi_{1j}$  is the only

positive component (i.e.  $\pi_{qj} = 0$  if  $q \neq 1$ ), if we define a new vector  $(\bar{x}, \bar{y})$  to be

$\bar{x}_{1j} = 1$ ,  $\bar{x}_{qj} = 0$  if  $x_{qj} = 1$ , and all other components being same with  $(x, y)$ , then

$(\bar{x}, \bar{y}) \in L_1^{pd}$  and  $\pi \bar{x} + \mu \bar{y} = \pi x + \mu y + \pi_{1j} > 1$ . This is a contradiction to the fact

$\pi x + \mu y \leq 1$  is a facet.

3)(=) This is obvious because  $\pi^j = (1, 1, \dots, 1)$ .

(\*) Suppose  $\pi^j > 0$  but  $\pi x + \mu y \neq \sum_i x_{ij}$ . Then, we must have an integer vector

$(x, y) \in H_{\pi, \mu}$  such that  $\sum_i x_{ij} < 1$  which implies  $x^j = 0$ . This is so because

$\sum_i x_{ij} \leq 1$  defines a facet and  $(x, y)$  are integers. Consequently,  $y_i = 1$  holds

for all  $i = 1, 2, \dots, p$ . Otherwise, if  $y_k = 0$  for some  $k$ , then we can set  $x_{kj} = 1$  while maintaining feasibility for  $L_I^{pd}$  but, then,  $\pi x + \mu y + \pi_{kj} > 1$  which contradicts the assumption that  $\pi x + \mu y \leq 1$  is a facet. Hence,  $\sum_l \mu_l = 1$  holds and by 1) we can not have  $\pi_j^i > 0$ . This is a contradiction.

We have thus shown that all inequalities of the set  $F_c^{pd}$  defined under point 7) of section 2 define facets of  $L_I^{pd}$ . We will call these facets trivial facets of  $L_I^{pd}$ .

## 5. Non-trivial facets.

In this section, we will discuss some necessary conditions for non-trivial facets. By remark 4.1, we can restrict ourselves to non-negative  $(\pi, \mu)$  when we consider nontrivial facets. Furthermore, the right hand side of the related inequality,  $\pi_0$ , is understood to equal 1 in the following discussion.

Theorem 5.1. Let  $(\pi, \mu) \in F^{pd} - F_c^{pd}$ . Then, the following statements are true.

1) If  $\pi_{ij} > 0$  for some  $j$ , then  $\mu_i \geq \pi_{ij}$  for all  $i$ .

2) If  $\mu_i > 0$ , then there exist  $j \neq k$  such that  $\pi_{ij} > 0$  and  $\pi_{ik} > 0$ , i.e.  $|J_i| \geq 2$ .

3) If  $I_j \neq \emptyset$ , then  $|I_j| \geq 2$  and there exist  $i \neq k$  such that

$$\max\{\pi_{qj} \mid q \in I_j\} = \pi_{ij} = \pi_{kj}.$$

4) If  $I_j \neq \emptyset$  and  $I_k \neq \emptyset$  for  $j \neq k$ , then  $I_j \neq I_k$ , i.e.  $I_j$  and  $I_k$  must be distinct.

Proof:

1) Since  $(\pi, \mu)$  is not a clique constraint, there exists an integer vector  $(x, y) \in H_{\pi, \mu}$  such that  $x^j = 0$  for some  $j \in D$ . Otherwise,  $\sum_i x_{ij} = 1$  for all  $j = 1, 2, \dots, d$  which would be clique constraints. Consequently, without loss of generality, we can let  $y_i = 1$  for all  $i \in I_j$ . Because, if not, then  $y_k = 0$  for some  $k \in I_j$  and so by defining a new vector  $(\bar{x}, \bar{y})$  to be  $\bar{x}_{kj} = 1$  for some  $\pi_{kj} > 0$  others being same with  $(x, y)$  we get  $(\bar{x}, \bar{y}) \in L_I^{pd}$  and  $\pi\bar{x} + \mu\bar{y} = \pi x + \mu y + \pi_{kj} > 1$  contradicting the fact that  $\pi x + \mu y \leq 1$  is a facet.

Suppose, now, that  $\mu_q < \pi_{qj}$  for some  $q \in I_j$ . Then, by defining a new vector  $(x^*, y^*)$  to be  $x_{qj}^* = 1$ ,  $y_q^* = 0$  others being same with  $(x, y)$ , we get  $(x^*, y^*) \in L_I^{pd}$  and  $\pi x^* + \mu y^* = \pi x + \mu y - \mu_q + \pi_{qj} > 1$ . This is a contradiction.

2) Since  $(\pi, \mu)$  is not a clique constraint, there exists an integer vector

$(x, y) \in H_{\pi, \mu}$  such that  $x_{ij} = 0$  and  $y_i = 0$  for  $i, j$  such that  $\pi_{ij} > 0$  and  $\mu_i > 0$ .

Otherwise, for such  $i$  and  $j$ ,  $x_{ij} + y_i = 1$ , which would be a clique constraint.

Suppose, now, that there exists no other  $k$  such that  $\pi_{ik} > 0$ , i.e.  $\pi_{ik} = 0$  for

all  $k \neq j$ . Then, by defining a new vector  $(\bar{x}, \bar{y})$  to be  $\bar{x}_{ij} = 0$  for all  $j \in D$ ,

$\bar{y}_i = 1$ , others being same with  $(x, y)$ , we get  $(\bar{x}, \bar{y}) \in L_I^{pd}$  and

$\pi \bar{x} + \mu \bar{y} = \pi x + \mu y + \mu_i > 1$ . This is a contradiction.

3) Since  $(\pi, \mu)$  is not a clique constraint, from 2) of theorem 4.2,  $|I_j| \geq 2$ .

Suppose, now, that the statement is wrong. In other words, assume  $\pi_{ij} > \pi_{qj}$

for all  $q$  in  $P$  different from  $i$ . Since  $\pi x + \mu y \leq 1$  is not a clique constraint,

there exists an integer vector  $(x, y) \in H_{\pi, \mu}$  such that  $x_{ij} = 0$  and  $y_i = 0$  for

some  $i \in I_j$ ,  $j \in D$ . Otherwise, we will have  $x_{ij} + y_i = 1$  which would be a clique

constraint. Now, if we define a new vector  $(\bar{x}, \bar{y})$  such that  $\bar{x}_{qj} = 0$  for  $q \neq i$ ,

$\bar{x}_{ij} = 1$  and all other components equal to the components of  $(x, y)$ , then we get

$(\bar{x}, \bar{y}) \in L_I^{pd}$  and  $\pi \bar{x} + \mu \bar{y} = \pi x + \mu y + \pi_{ij} - \pi_{qj} > 1$ . This is a contradiction.

4) Suppose  $I_j = I_k \neq \emptyset$ . First, we note that if  $(x, y) \in H_{\pi, \mu}$  and  $x_{ij} = 0$  for

all  $i \in I_j$ , then we must have  $y_i = 1$  for all  $i \in I_j$ . This is so because, if there

exists  $k \in I_j$  such that  $y_k = 0$ , then by defining a new vector  $(\bar{x}, \bar{y})$  to be  $\bar{x}_{kj} = 1$ ,

and  $\bar{x}_{qj} = 0$  if  $x_{qj} = 1$  (note:  $q \notin I_j$ ), and all others being the same as  $(x, y)$ , we

get  $(\bar{x}, \bar{y}) \in L_I^{pd}$  and  $\pi \bar{x} + \mu \bar{y} = \pi x + \mu y + \pi_{kj} > 1$ , which leads to a contradiction.

Thus, if  $x_{ij} = 0$  for all  $i \in I_j$ , then  $x_{ik} = 0$  for all  $i \in I_k$  because  $y_i = 1$  for all

$i \in I_j$  and  $I_j = I_k$ . In other words, if  $x_{ij} = 1$  for some  $i \in I_j$ , then there must



exist  $i' \in I_k$  (possibly,  $i = i'$ ) such that  $x_{i',j} = 1$ . Therefore, we have

$$\sum_{i \in I_j} x_{ij} = \sum_{i \in I_k} x_{ik} \text{ for } (x,y) \in H_{\pi,\mu} \text{ implying that we can not have } p(d+1) \text{ linearly}$$

independent vectors satisfying  $\pi x + \mu y = 1$ . This is a contradiction.  $\Delta$

Remark 5.1. Theorem 4.2 and theorem 5.1 together imply that, if  $(\pi, \mu) \in F^{pd} - F_t^{pd}$ ,

then either  $\pi^j = 0$  or if  $\pi^j \neq 0$  then at least two components  $\pi_{ij}$  and  $\pi_{qj}$ , say, of  $\pi^j$  are positive. Furthermore, for at least one  $j \in D$  we have  $\pi^j \neq 0$ . If  $\pi^j \neq 0$ , then by part 2) of theorem 5.1 it follows that there exists a  $k \neq j$  such that  $I_k \neq \emptyset$ ,  $\mu_i > 0$  where  $i \in I_k$  and  $i \in I_j$ . Also, by part 4) of theorem 5.1, we have  $I_k \neq I_j$ . Finally, for each  $j$  such that  $\pi^j \neq 0$  there exists an  $i$  such that  $\mu_i > 0$  and  $\pi_{ij} = 0$ .

Remark 5.2. Theorem 4.2 and theorem 5.1 together imply that, if  $(\pi, \mu) \in F^{pd} - F_t^{pd}$ ,

then there must exist at least 3 plants  $i$  such that  $\mu_i > 0$  and at least 3 destinations  $j$  such that  $\pi^j \neq 0$ . This is so because, if  $\pi^j \neq 0$ , then we must have  $|I_j| \geq 2$  and at least one plant  $i$  such that  $\pi_{ij} = 0$ . Thus, we must have at least 3 plants  $i$  such that  $\mu_i > 0$ . Furthermore, if  $i \in I_j$ , then there must exist a destination  $q$  different from  $j$  such that  $\pi_{iq} > 0$  in order to satisfy  $|J_i| \geq 2$ . Now, if  $\pi^t = 0$  for  $t \neq j, q$ , then, for each  $i$  such that  $\mu_i > 0$ , we have  $\pi_{ij} > 0$  if and only if  $\pi_{iq} > 0$  in order to satisfy  $|J_i| \geq 2$ . This is a contradiction to the fact that  $I_j \neq I_q$ . Thus, there must exist at least 3 destinations  $j$  such that  $\pi^j \neq 0$ .

6. Facets for the case of 3 plants and several destinations ( $d \geq 3$ ).

The results of the previous section enable us now to describe all facets for the case of three plants and several destinations.

Theorem 6.1. Let  $(\pi, \mu) \in F_c^{pd} - F_c^{pd}$  where  $p = 3$  and  $d \geq 3$ . Then  $\pi x + \mu y \leq 1$  is a facet if and only if  $\mu_i = \frac{1}{4}$  for all  $i \in P = \{1, 2, 3\}$  and  $\pi_{i_j} = \frac{1}{4}$  for  $i \in I_j$ ,  $j \in J$  where  $J$  is any subset of  $D$  consisting of exactly 3 elements and  $I_j$ 's are distinct subsets of  $I$  with  $|I_j| = 2$ .

Proof:

( $\Leftarrow$ ) Without loss of generality, assume  $J = \{1, 2, 3\}$ ,  $I_1 = \{1, 2\}$ ,  $I_2 = \{2, 3\}$  and  $I_3 = \{1, 3\}$ . Then, we have to show  $\sum_{j \in J} \sum_{i \in I_j} \frac{1}{4} x_{ij} + \sum_{i \in P} \frac{1}{4} y_i \leq 1$  is a facet for

$L_1^{pd}$ . This follows from Guignard [16].

( $\Rightarrow$ ) By remark 5.1, we have  $I_j = \emptyset$  or  $|I_j| = 2$  for all  $j = 1, 2, \dots, d$  and  $|I_j| = 2$  for at least one  $j$ . Furthermore, it follows from remark 5.1 that  $\mu_i > 0$  for  $i = 1, 2, 3$ .

From 2) of theorem 5.1, we know that  $|J_i| \geq 2$  for all  $i = 1, 2, 3$ . Suppose  $|J_1| = 3$ , say,  $J_1 = \{1, 2, 3\}$ . Then, in order to satisfy  $|I_1| = 2$ , we must have either  $I_1 = \{1, 2\}$  or  $I_1 = \{1, 3\}$ . If  $I_1 = \{1, 2\}$ , then  $I_2$  must be  $\{1, 3\}$  in order to satisfy  $I_1 \neq I_2$ . But, this would force  $I_3$  to be equal to either  $I_1$  or  $I_2$  contradicting the fact that  $\pi x + \mu y \leq 1$  is a facet. Similar result holds if we start with  $I_1 = \{1, 3\}$ . Also, it can be easily seen that if  $|J_i| > 3$ , then we would get a similar contradiction. Therefore,  $|J_i| = 2$ .

Now, since  $|J_i| = 2$  for all  $i = 1, 2, 3$  and  $I_j$ 's are distinct with  $|I_j| = 2$  for all  $j = 1, 2, \dots, |J|$ , it is obvious that  $|J| = 3$  where  $J = \{j \in D \mid \pi^j \neq 0\}$ .

Note that  $I_j$ 's must be distinct subsets of  $I$  with  $|I_j| = 2$ , and thus

$I_j$  for  $j = 1, 2, 3$  is one element of  $\{\{1, 2\}, \{1, 3\}, \{2, 3\}\}$ .

Next, we show that all of the  $\pi_{ij}$ 's and  $\mu_i$ 's have to be same. Without loss of generality, assume  $I_1 = \{1, 2\}$ ,  $I_2 = \{2, 3\}$  and  $I_3 = \{1, 3\}$ . We note that there exists a solution vector  $(x, y) \in H_{\pi, \mu}$  such that  $x_{11} = 0$  and  $y_1 = 0$ . This, in turn, implies that  $x_{13} = 1$ . (Otherwise, if  $x_{13} = 0$ , then letting  $y_1 = 1$  would contradict the fact that  $\pi x + \mu y \leq 1$  is a facet.) From theorem 5.1, we know that  $\mu_1 \geq \pi_{13}$ .

Suppose  $\mu_1 > \pi_{13}$ . Then, letting  $y_1 = 1$  and  $x_{13} = 0$  would contradict the fact that  $\pi x + \mu y \leq 1$  is a facet. Therefore,  $\mu_1 = \pi_{13}$ . Again, from theorem 5.1,

$\pi_{13} = \pi_{33}$ . Repeating this argument, one can show that all the coefficients are

same and so dividing by  $\pi_{11}$ ,  $\pi x + \mu y \leq 1$  becomes  $\sum_{j \in J} \sum_{i \in I_j} x_{ij} + \sum_{i \in P} y_i \leq \frac{1}{\pi_{11}}$ .

However, the left hand side of the above inequality corresponds to a chordless odd cycle of length 9 and thus the right hand side is equal to 4. Therefore,

$$\pi_{11} = \frac{1}{4}.$$

## References

1. E. Balas and M.W. Padberg, Set Partitioning: A Survey, SIAM Review 18 (1976), 710-760.
2. E. Balas and E. Zemel, Critical Cutsets of Graphs and Canonical Facets of Set-Packing Polytopes, Math. of Operations Research 2 (1977), 15-20.
3. V. Chvatal, On Certain Polytopes Associated with Graphs, J. Comb. Theory B 18 (1975), 138-154.
4. D. Cho, M.W. Padberg and M.R. Rao, On the Uncapacitated Plant Location Problem II: Facets and Lifting Theorems, GBA Working Paper, New York University, New York, December 1980.
5. G. Cornuejols, M.L. Fisher and G.L. Nemhauser, On the Uncapacitated Location Problem, Annals of Discrete Math. 1 (1977), 163-177.
6. G. Cornuejols and J.M. Thizy, Some Facets of the Simple Plant Location Polytope, Working Paper, Carnegie-Mellon University.
7. H.P. Crowder and M.W. Padberg, Solving Large-Scale Symmetric Travelling Salesman Problems to Optimality, Management Science 26 (1980), 495-509.
8. J. Edmonds, Covers and Packings in a Family of Sets, Bull. Am. Math. Soc. 68 (1962), 494-499.
9. M.A. Efronson and T.L. Ray, A Branch-Bound Algorithm for Plant Location, Operations Research 14 (1966), 361-368.
10. R.L. Francis and J.M. Goldstein, Location Theory: A Selected Bibliography, Operations Research 22 (1974), 400-410.
11. D.R. Fulkerson, Blocking and Anti-blocking Pairs of Polyhedra, Math. Programming 1 (1971), 168-194.
12. M. Garey and D. Johnson, Computer and Intractability. A Guide to the Theory of NP-Completeness (W.H. Freedman Co., San Francisco 1979)
13. R. Garfinkel and G.L. Nemhauser, A survey of Integer Programming Emphasizing Computation and Relations among Models, in T.C. Hu and S.M. Robinson, eds; Mathematical Programming (Academic Press, 1973).

14. M. Grötschel and M.W. Padberg, On the symmetric Travelling Salesman Problem I Inequalities, Math. Programming 16 (1979), 265-280.
15. M. Grötschel and M.W. Padberg, On the Symmetric Travelling Salesman Problem II: Lifting Theorems and Facets, Math. Programming 16 (1979), 281-302.
16. M. Guignard, Fractional Vertices, Cuts and Facets of the Simple Plant Location Problem, Mathematical Programming Study 12 (1980), 150-162.
17. W.M. Hirsch and G.B. Dantzig, The Fixed Charge Problem, Naval Research Logistics Quarterly 15 (1968), 413-424.
18. E.L. Johnson and M.W. Padberg, On the Simple Plant Location Problem, Unpublished notes (Nov. 1972).
19. R.M. Karp, Reducibility among Combinatorial Problems, in R.E. Miller et.al., eds., Complexity of Computer Computations (Plenum Press, New York, 1972).
20. G.L. Nemhauser and L.E. Trotter, Properties of Vertex Packing and Independence System Polyhedra, Math. Programming 6 (1974), 48-61.
21. M.W. Padberg, On the Facial Structure of Set Packing Polyhedra, Math. Programming 5 (1973), 199-215.
22. M.W. Padberg, On the Complexity of Set Packing Polyhedra, Annals of Discrete Math. 1 (1977), 421-434.
23. M.W. Padberg and S. Hong, On the Symmetric Travelling Salesman problem: A Computational Study, Mathematical Programming Study 12 (1980) 78-107.
24. A. Weber, Über den Standort der Industrien, Tübingen 1909. Translated as Alfred Weber's Theory of the Location of Industries, by C.J. Friedrich University of Chicago Press, 1929.
25. K. Spielberg, Algorithms for the Simple Plant Location Problem with Some Side Conditions, Operations Research 17 (1) (1969), 85-111.

