



C1-curved finite elements with numerical integration for thin plate and thin shell problems: part 1: construction and interpolation properties of curved C1 finite elements

Michel Bernadou

► To cite this version:

Michel Bernadou. C1-curved finite elements with numerical integration for thin plate and thin shell problems: part 1: construction and interpolation properties of curved C1 finite elements. [Research Report] RR-1626, INRIA. 1992. inria-00074935

HAL Id: inria-00074935

<https://hal.inria.fr/inria-00074935>

Submitted on 24 May 2006

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

INRIA

UNITÉ DE RECHERCHE
INRIA-ROCQUENCOURT

Institut National
de Recherche
en Informatique
et en Automatique

Domaine de Voluceau
Rocquencourt
B.P.105
78153 Le Chesnay Cedex
France
Tél.: (1) 39 63 55 11

Rapports de Recherche

1992



25^{ème}
anniversaire

N° 1626

Programme 6

*Calcul Scientifique, Modélisation et
Logiciel numérique par Ordinateur*

C¹-CURVED FINITE ELEMENTS WITH NUMERICAL INTEGRATION FOR THIN PLATE AND THIN SHELL PROBLEMS

Part 1 :

Construction and interpolation properties
of curved C¹ finite elements

Michel BERNADOU

Février 1992



★ R R - 1 6 2 6 ★

C^1 -CURVED FINITE ELEMENTS WITH NUMERICAL INTEGRATION FOR THIN PLATE AND THIN SHELL PROBLEMS (*)(**)

Part 1 : Construction and interpolation properties
of curved C^1 finite elements

Michel BERNADOU
INRIA, Domaine de Voluceau,
Rocquencourt, 78153 Le Chesnay Cedex, France

Abstract

The purpose of this paper is to construct curved finite elements of class C^1 , compatible with the elements of ARGYRIS-FRIED-SCHARPF and BELL. We start by the approximation of the curved boundary and, next, we construct curved finite elements of class C^1 . We conclude by proving that the corresponding asymptotic interpolation error has the same order than for the associate straight elements.

In the second part of this work, we will study the approximation of thin plate and thin shell problems by such curved C^1 finite elements. The effectiveness of these methods will be illustrated in a subsequent work (en collaboration avec J.M. BOISSERIE).

(*) To appear in Comput. Meth. Appl. Mech. Engng.

(**) This work is part of

- the Project "Numerical Solution of Partial Differential Equations for Engineering Problems" of the Programm "INRIA-NSF"

- the Project "Junctions in Elastic Multi-Structures" of the Program "S.C.I.E.N.C.E." of the Commission of the European Communities (contract n° SC1 * 0473 - C(EDB))

ELEMENTS FINIS COURBES DE CLASSE C^1 AVEC INTEGRATION NUMERIQUE POUR DES PROBLEMES DE PLAQUES ET DE COQUES MINCES

Partie 1 : Construction et propriétés d'interpolation des éléments finis courbes de classe C^1

Résumé

L'objet de cette première partie est de construire des éléments finis courbes de classe C^1 , compatibles avec les triangles d'ARGYRIS et de BELL. Nous commençons par l'approximation de frontières courbes puis nous construisons des éléments finis courbes de classe C^1 . Nous concluons par l'étude de l'erreur d'interpolation correspondante et nous montrons que pour la même régularité de la solution, elle est du même ordre que dans le cas des éléments finis droits associés.

Dans la seconde partie de ce travail, à l'aide de ces éléments finis courbes de classe C^1 , nous étudierons l'approximation de problèmes de plaques et de coques minces. Dans un travail ultérieur (en collaboration avec J.M. BOISSERIE), nous illustrerons l'efficacité de ces méthodes par quelques exemples numériques.

1 INTRODUCTION

Many structural problems are set on curved boundary plane domains and are modeled by fourth order partial differential equations or by systems of fourth order partial differential equations. In this way, let us mention :

*) many thin plate problems whose middle surfaces are curved boundary plane domains. It is well known that the approximation of such problems by straight finite elements can produce some lack of convergence as reported by [1] and bibliography of this work.

***) many thin shell problems : in [2] we have thoroughly shown that a high accuracy approximation method can be easily developed when a general thin shell problem is formulated on a plane reference domain. This is generally obtained through the use of an exact or an approximated (for instance, by B-spline methods) mapping of the middle surface of the shell. In general, such plane reference domains have curved boundaries.

***) many structural problems which include junctions between different thin shells : for example let us mention junctions between circular cylinders, or pipes, which are used in oil platform constructions. Each shell element can be associated to a plane reference domain and, due to the complexity of surface intersections, the part of the boundary which is associate to such intersections is generally curved. Corresponding open problems are mentioned in [3].

Then, the development of approximations of high degree of accuracy by finite element methods requires the use of \mathcal{C}^1 -curved elements. Thus, in this paper, we develop and analyze such methods. The whole paper comprises two parts :

i) the construction of curved finite elements of class \mathcal{C}^1 and the study of their interpolation properties ;

ii) the use of these curved \mathcal{C}^1 -finite elements in order to approximate the solutions of plate or shell problems posed over plane reference domains with curved boundary. This second part [4] will include the study of convergence and the obtention of asymptotic error estimates ;

The detailed description of how to implement such curved \mathcal{C}^1 -elements and some numerical results obtained from benchmark plate problems will be reported in an additional paper [5].

In this first part, we shall be only concerned with the construction of curved finite elements which have a connection of class \mathcal{C}^1 with some classical \mathcal{C}^1 -elements like ARGYRIS triangle (see [6]) or BELL triangle (see [7]). In the sequel, we shall say that such elements are \mathcal{C}^1 -compatible with ARGYRIS or BELL triangles.

In the second part, we will generalize the results of [8] relative to the approximation of solutions of thin shell problems posed on polygonal domains to the case of curved boun-

dary domains. In this way, we will analyze the simultaneous use of curved C^1 -elements, considered here, and of numerical integration techniques.

In addition to the introduction, this paper comprises three paragraphs. In paragraph 2, we consider the approximation of the domain Ω . We first define an exact triangulation of the domain Ω . Next, to every triangle having a curved side located on the boundary, we associate an approximate triangle by interpolating this curved side. Thus, we get an approximate domain Ω_h whose constitutive triangles are in bijective correspondence with a reference triangle \hat{K} through applications F_K , affine or not. Similar constructions were first analyzed by [9,10,11] and next some modifications were considered in [12 to 17]. Let us also mention [18] who obtains optimal error estimates for the finite element solution of second order elliptic problems by using isoparametric simplicial elements, [19] who considers the case of the interpolation on curved domains of functions which are not smooth and [20] who analyzes special exact curved finite elements useful for solving contact problems of second order in domain with piecewise circular boundaries..

Paragraph 3 is devoted to the definition of curved finite elements C^1 -compatible with ARGYRIS or BELL triangles. Each curved finite element is in correspondence with a reference element on \hat{K} which is obtained by constraining a suitable polynomial space \hat{P} . In our knowledge, the results concerning the first curved element, C^1 -compatible with ARGYRIS triangle, are new. For the second curved element, C^1 -compatible with BELL triangle, we obtain, in an independent way, results comparable with some of [17]. In this direction we also mention [21,22]. In addition, other curved C^1 -elements were considered by different Authors : when the plane curved domain Ω is the image of a plane polygonal domain ω through a parametric representation $\vec{\psi}$, [23] introduces a curved version of the Argyris element - named TUBAC 6 - which is the image of the associated straight element defined on straight triangles of the polygonal domain ω through the mapping $\vec{\psi}$. This construction is really fruitful when such a mapping $\vec{\psi}$ can be explicitated like for circular or elliptical domain for instance. In addition [24,25] have considered curved C^1 -elements for which the finite dimensional space is polynomial over the current triangle. This is convenient for the approximation of constant coefficient operator as in some plate problems. Nevertheless, for corresponding finite element methods, it seems difficult to analyze the effect of numerical integration. A significant improvement of the method, suggested in [24], is given in [26].

In paragraph 4, we give estimates of the asymptotic interpolation errors for every family of these curved C^1 -elements. It is worth to note that these asymptotic error estimates are of the same order than the one obtained for the associate straight elements.

Throughout this paper, we use notations of [27,28]. In particular, let $W^{k,p}(\Omega)$ be the Sobolev space of real-valued functions which, together with all their partial distributional derivatives of order k or less, belong to $L^p(\Omega)$. We set $H^k(\Omega) = W^{k,2}(\Omega)$. On $W^{k,p}(\Omega)$ we shall use the norms and semi-norms defined by

$$\|u\|_{k,p,\Omega} = \left(\sum_{|\alpha| \leq k} \|D^\alpha u\|_{L^p(\Omega)}^p \right)^{1/p}, \quad |u|_{k,p,\Omega} = \left(\sum_{|\alpha|=k} \|D^\alpha u\|_{L^p(\Omega)}^p \right)^{1/p}$$

for $1 \leq p \leq \infty$, with the standard modification for $p = +\infty$ (see for example [29,30,31]).

Contents

1 - INTRODUCTION

2 - APPROXIMATION OF THE DOMAIN Ω

2.1. Exact triangulation of the domain Ω

2.2. Construction of an approximate triangulation of the domain Ω

2.3. Some properties of the application F_K

2.4. Examples

3 - DEFINITION OF CURVED C^1 FINITE ELEMENTS

3.1. Basic principles

3.2. Definition of curved finite elements C^1 -compatible with ARGYRIS or BELL triangles when $F_K \in (P_5)^2$

3.3. Definition of curved finite elements C^1 -compatible with ARGYRIS or BELL triangles when $F_K \in (P_3)^2$

3.4. Definition of curved finite elements C^0 -compatible with the Hermite triangle of type (3) when $F_K \in (P_n)^2, n = 3$ or 5

4 - ESTIMATES OF THE INTERPOLATION ERRORS

4.1. Estimates of the interpolation error for curved finite elements C^1 -compatible with ARGYRIS or BELL triangles when $F_K \in (P_5)^2$

4.2. Estimates of the interpolation error for curved finite elements C^1 -compatible with ARGYRIS or BELL triangles when $F_K \in (P_3)^2$

4.3. Estimates of the interpolation error for curved finite elements C^0 -compatible with Hermite triangle of type (3) ($F_K \in (P_n)^2, n = 3$ or $n = 5$)

5 - CONCLUDING REMARKS

References

2 APPROXIMATION OF THE DOMAIN Ω

Let there be given the plane \mathbf{R}^2 referred to an orthonormal system $(0, \vec{e}_1, \vec{e}_2)$, the corresponding system of coordinates (x_1, x_2) and a bounded domain Ω in \mathbf{R}^2 . We assume that the boundary Γ can be subdivided into a finite number of arcs, each of them admitting the following parametric representation

$$(2.1) \quad x_1 = \chi_1(s), \quad x_2 = \chi_2(s), \quad s_m \leq s \leq s_M.$$

The functions $\chi_1(s), \chi_2(s)$ belong to C^{q+1} , q sufficiently large, and they are such that $[(\chi_1)']^2 + [(\chi_2)']^2$ is different from 0 on the interval $[s_m, s_M]$.

2.1 Exact triangulation of the domain Ω

Let us subdivide the set $\bar{\Omega}$ into a finite number of triangles K with straight or curvilinear sides. More precisely, given two distinct triangles of this triangulation \mathcal{T} , we assume that either they are disjoint or they have a common vertex or they have a common side. Moreover, we assume that every "interior" triangle K (i.e. a triangle having at most one vertex on Γ) has only straight sides and that every "boundary" triangle K_c has at most one curved side located over an arc of type (2.1) upon the boundary Γ . In the following, we shall denote \tilde{K} the (rectilinear) triangle having the same vertices than the curvilinear triangle K_c .

From now on, we consider regular families of triangulations \mathcal{T} of the domain Ω , i.e.,

(i) There exists a constant σ such that

$$(2.2) \quad \forall \mathcal{T}, \quad \forall K (\text{ or } \tilde{K}) \in \mathcal{T}, \quad \frac{h_K}{\rho_K} \leq \sigma,$$

where $h_K = \text{diam}(K)$ and $\rho_K = \sup\{\text{diam}(C); C \text{ is a disk contained in } K\}$ (in case of curved triangles K_c , we replace K_c by \tilde{K}).

(ii) The parameter

$$(2.3) \quad h = \max_{K \in \mathcal{T}} h_K \text{ approaches zero.}$$

Remark 2.1 : For h sufficiently small, it was proved in [10, Theorem 1] and [16, Theorem 1] that every triangle K_c is the image of a reference triangle \tilde{K} through a diffeomorphism of order q . ■

2.2 Construction of an approximate triangulation of the domain Ω

In order to construct finite elements of class C^1 , it is convenient to approach the curvilinear side of every triangle K_c by an arc parameterized with the help of polynomial functions. This amounts to associate to every curved triangle K_c an approximate curved triangle K (see Figure 2.1). Thus, the initial domain Ω is replaced by an approximate domain Ω_h whose corresponding triangulation is denoted by \mathcal{T}_h . Every triangulation \mathcal{T}_h is the union

(i) of a triangulation \mathcal{T}_h^1 constituted by straight sides triangles K in correspondence with a fixed reference triangle \tilde{K} through an affine mapping F_K ;

(ii) of a triangulation \mathcal{T}_h^2 constituted by triangles K having two straight sides and a third curved side approximating an arc of the boundary Γ . These triangles K are the

images of the reference triangle \hat{K} through a nonlinear mapping F_K .




Figure 2.1 : curved triangles, exact K_c and approximate K

In both cases, we indicate how to construct an application $F_K : \hat{K} \rightarrow K$, for all $K \in \mathcal{T}_h$. Firstly, if the triangle $K \in \mathcal{T}_h^1$, then the application F_K is affine : there exists an invertible matrix B_K and a vector b_K of \mathbb{R}^2 such that

$$(2.4) \quad F_K : (\hat{x}_1, \hat{x}_2) \in \hat{K} \rightarrow F_K(\hat{x}_1, \hat{x}_2) = B_K \begin{bmatrix} \hat{x}_1 \\ \hat{x}_2 \end{bmatrix} + b_K \in K.$$

For clarity, using notations of Figure 2.2 :

$$(2.5) \quad F_K(\hat{x}_1, \hat{x}_2) = \begin{cases} x_1 = x_{13} + (x_{11} - x_{13})\hat{x}_1 + (x_{12} - x_{13})\hat{x}_2, \\ x_2 = x_{23} + (x_{21} - x_{23})\hat{x}_1 + (x_{22} - x_{23})\hat{x}_2, \end{cases}$$

where $x_{\alpha i}$, $\alpha = 1, 2$ denote the coordinates of the vertices a_i , $i = 1, 2, 3$ of the triangle K .

Secondly, when the triangle $K \in \mathcal{T}_h^2$, the application $F_K : \hat{K} \rightarrow K$ is in general non-linear and can be conveniently defined in two steps :

Step 1 : Interpolation of the curved side $a_1 a_2$ of the triangle K_c :

Consider the new parametrization of the arc $a_1 a_2$ by using variable \hat{x}_2

$$(2.6) \quad \begin{cases} x_1 = \psi_1(\hat{x}_2), \quad x_2 = \psi_2(\hat{x}_2), \quad 0 \leq \hat{x}_2 \leq 1 \\ \text{where} \\ \psi_\alpha(\hat{x}_2) = \chi_\alpha(\underline{s} + (\bar{s} - \underline{s})\hat{x}_2), \quad \alpha = 1, 2. \end{cases}$$




Figure 2.2 : Case of a triangle $K \in \mathcal{T}_h^1$

Then, the approximate arc γ_h is defined by the parametric equations

$$x_1 = \psi_{1h}(\hat{x}_2), \quad x_2 = \psi_{2h}(\hat{x}_2), \quad 0 \leq \hat{x}_2 \leq 1,$$

where the functions ψ_{1h} and ψ_{2h} satisfy the following hypothesis :

Hypothesis 2.1 : The functions $\psi_{\alpha h}$ are interpolation polynomials, of Lagrange or Hermite type, with degree $n \geq 1$, of the functions ψ_α , $\alpha = 1, 2$, over the interval $[0, 1]$ and such that

$$(2.7) \quad \begin{cases} \psi_{\alpha h}(0) = \psi_\alpha(0), \quad \psi_{\alpha h}(1) = \psi_\alpha(1), \quad \alpha = 1, 2; \\ |\psi_\alpha - \psi_{\alpha h}|_{p, \infty} \leq ch_K^{n+1-p} |\psi_\alpha|_{n+1, \infty}, \quad p = 0, \dots, n+1, \end{cases}$$

where c is a constant independent of h_K . ■

From Hypothesis 2.1, we immediately derive

$$(2.8) \quad \psi_{\alpha h}(\hat{x}_2) = x_{\alpha 1} + (x_{\alpha 2} - x_{\alpha 1})\hat{x}_2 + \begin{cases} 0 & \text{if } n = 1, \\ \hat{x}_2(1 - \hat{x}_2)P_{n-2, \alpha}(\hat{x}_2) & \text{if } n \geq 2, \end{cases}$$

where $P_{n-2, \alpha}$, $\alpha = 1, 2$, denotes polynomials of degree $n - 2$ with respect to \hat{x}_2 , completely determined by the choice of the interpolation method. Some examples are given in section 2.4.

Step 2 : Definition of the application $F_K : \hat{K} \longrightarrow K$:

To any point \hat{M} of the reference triangle \hat{K} , the application F_K associates the point M_h as follows :

$$(2.9) \quad \overrightarrow{OM}_h = F_{K1}(\hat{x}_1, \hat{x}_2)\vec{e}_1 + F_{K2}(\hat{x}_1, \hat{x}_2)\vec{e}_2$$

where the functions $F_{K\alpha}$, $\alpha = 1, 2$ are defined by the relations

$$(2.10) \quad \left\{ \begin{array}{l} F_{K\alpha}(\hat{x}_1, \hat{x}_2) = x_{\alpha 3} + (x_{\alpha 1} - x_{\alpha 3})\hat{x}_1 + (x_{\alpha 2} - x_{\alpha 3})\hat{x}_2 \\ + \begin{cases} 0 \text{ if } n = 1, \\ \frac{1}{2} \hat{x}_1 \hat{x}_2 [P_{n-2;\alpha}(1 - \hat{x}_1) + P_{n-2;\alpha}(\hat{x}_2)] \text{ if } n \geq 2 \end{cases} \\ \text{(where polynomials } P_{n-2;\alpha} \text{ are defined by (2.8)).} \end{array} \right.$$

Let us note that these functions $F_{K\alpha}$ are constructed so that $F_{K\alpha}(1 - \hat{x}_2, \hat{x}_2) \equiv \psi_{\alpha h}(\hat{x}_2)$, i.e., the image of the side $\hat{a}_1\hat{a}_2$ of the reference triangle \hat{K} through the application F_K , is the approximate arc γ_h . Moreover, if we denote by \tilde{M} the point of barycentric coordinates $(\hat{x}_1, \hat{x}_2, 1 - \hat{x}_1 - \hat{x}_2)$ in the (straight) triangle $\tilde{K} = (a_1, a_2, a_3)$, relations (2.10) can be rewritten in the vectorial form

$$(2.11) \quad \overrightarrow{OM}_h = \overrightarrow{O\tilde{M}} + \frac{1}{2}(\overrightarrow{\tilde{M}M}_h^1 + \overrightarrow{\tilde{M}M}_h^2).$$




Figure 2.3 : Construction of $\tilde{M} \rightarrow M_h$ through $\overrightarrow{\tilde{M}M}_h = \frac{1}{2}(\overrightarrow{\tilde{M}M}_h^1 + \overrightarrow{\tilde{M}M}_h^2)$ (for clarity, the constructions of $\tilde{M} \rightarrow M_h^1$ and $\tilde{M} \rightarrow M_h^2$ are displayed separately)

Corresponding geometrical interpretation is illustrated in Figure 2.3 and obtained from relations

$$(2.12) \quad \left\{ \begin{array}{l} \overrightarrow{\tilde{M}M}_h^1 = \frac{\hat{x}_2}{1 - \hat{x}_1} \overrightarrow{\tilde{P}^1 P_h^1} \text{ if } \hat{x}_1 \neq 1, \quad \tilde{M} \equiv M_h^1 \equiv a_1 \text{ if } \hat{x}_1 = 1, \\ \overrightarrow{\tilde{M}M}_h^2 = \frac{\hat{x}_1}{1 - \hat{x}_2} \overrightarrow{\tilde{P}^2 P_h^2} \text{ if } \hat{x}_2 \neq 1, \quad \tilde{M} \equiv M_h^2 \equiv a_2 \text{ if } \hat{x}_2 = 1, \end{array} \right.$$

since $\overrightarrow{\tilde{P}^1 P_h^1} = \sum_{\alpha=1}^2 \hat{x}_1(1 - \hat{x}_1) P_{n-2;\alpha}(1 - \hat{x}_1) \vec{e}_\alpha$, $\overrightarrow{\tilde{P}^2 P_h^2} = \sum_{\alpha=1}^2 \hat{x}_2(1 - \hat{x}_2) P_{n-2;\alpha}(\hat{x}_2) \vec{e}_\alpha$.

2.3 Some properties of the application F_K

In the following theorem, we prove directly some properties of the application F_K defined by relation (2.10). Some of these results have been obtained in [10,11] by using properties of the diffeomorphism $\hat{K} \rightarrow K_c$ mentioned in Remark 2.1.

We will use the following notations :

$$(2.13) \quad J_{F_K}(\hat{x}) = \text{Jacobian of } F_K \text{ at point } \hat{x} = (\hat{x}_1, \hat{x}_2),$$

$$(2.14) \quad J_{F_K^{-1}}(\hat{x}) = \text{Jacobian of } F_K^{-1} \text{ at point } x = (x_1, x_2),$$

$$(2.15) \quad \|F_K\|_{\ell, \infty, \hat{K}} = \sup_{\hat{x} \in \hat{K}} \|D^\ell F_K(\hat{x})\|,$$

$$(2.16) \quad \|F_K^{-1}\|_{\ell, \infty, K} = \sup_{x \in K} \|D^\ell F_K^{-1}(x)\|.$$

Theorem 2.1 : Let \mathcal{T} be a regular family of triangulations of the domain Ω , i.e., satisfying conditions (2.2) and (2.3). To any curved triangle K_c of the "exact" triangulation of the domain Ω , we associate the approximate curved triangle K obtained through the interpolation of the curved side located on the boundary Γ . We assume that this interpolation satisfies Hypothesis 2.1, and that the boundary Γ of the bounded domain $\Omega \subset \mathbb{R}^2$ is piecewise of class C^{q+1} , $q \geq n$ ($n = \text{degree of the components } F_{K_\alpha}$).

Then, for h_K sufficiently small, we have the following properties that are independent of the degree n of the components F_{K_α} :

(i) The application $F_K : \hat{K} \rightarrow \bar{K}$, defined by relations (2.9) (2.10) is a C^∞ -diffeomorphism from \hat{K} onto \bar{K} ;

(ii) The application F_K and its inverse $F_K^{-1} : \bar{K} \rightarrow \hat{K}$, satisfy following estimates

$$(2.17) \quad \|F_K\|_{\ell, \infty, \hat{K}} \leq ch_K^\ell, \quad \ell = 0, 1, \dots$$

$$(2.18) \quad \|F_K^{-1}\|_{\ell, \infty, K} \leq ch_K^{-1}, \quad \ell = 1, 2, \dots$$

(iii) The Jacobians $J_{F_K}(\hat{x})$ and $J_{F_K^{-1}}(x)$ satisfy the following estimates

$$(2.19) \quad c_1 h_K^2 \leq |J_{F_K}|_{0, \infty, \hat{K}} \leq c_2 h_K^2 ; |J_{F_K}|_{\ell, \infty, \hat{K}} \leq ch_K^{2+\ell}, \quad \ell = 0, \dots, n$$

$$(2.20) \quad \frac{c_1}{h_K^2} \leq |J_{F_K^{-1}}|_{0, \infty, K} \leq \frac{c_2}{h_K^2},$$

where, in the inequalities (2.17) to (2.20), letters c, c_1, c_2 denote strictly positive constants which are not necessarily the same from an inequality to the other.

Proof : When $n = 1$ (F_K affine) these results are obvious. Henceforth, we assume $n \geq 2$. Then, the proof takes seven steps which can be summarized as follows (more details can be found in [32]) :

Step 1 : F_K verifies estimates (2.17) :

From hypothesis 2.1, we have

$$(2.21) \quad |\psi_\alpha - \psi_{\alpha h}|_{p,\infty} \leq ch_K^{n+1-p} |\psi_\alpha|_{n+1,\infty}, \quad p = 0, \dots, n+1.$$

Taking into account the hypothesis $[(\chi_1)']^2 + [(\chi_2)']^2 \neq 0$ on $[s_m, s_M]$ and the definition (2.6), we obtain $|\psi_\alpha|_{p,\infty} = 0(h_K^p), p = 0, \dots, n+1$, so that

$$(2.22) \quad |\psi_{\alpha h}|_{p,\infty} \leq ch_K^p, p = 0, 1, \dots \text{ (note that } |\psi_{\alpha h}|_{p,\infty} = 0 \text{ when } p \geq n+1).$$

Substituting the estimate (2.22) into definition (2.8), we derive

$$(2.23) \quad \sup_{\hat{x}_2 \in [0,1]} |[P_{n-2;\alpha}(\hat{x}_2)]^{(m)}| \leq ch_K^{m+2}, m = 0, 1, \dots$$

and hence, with (2.10),

$$(2.24) \quad \sup_{(\hat{x}_1, \hat{x}_2) \in \hat{K}} \left| \frac{\partial^{i_1+i_2} F_{K\alpha}}{(\partial \hat{x}_1)^{i_1} (\partial \hat{x}_2)^{i_2}}(\hat{x}_1, \hat{x}_2) \right| \leq ch_K^{i_1+i_2}, \quad 0 \leq i_1 + i_2, \quad \alpha = 1, 2.$$

Thus, we obtain

$$\|D^\ell F_K(\hat{x})\| \leq c \max_{|\beta|=\ell} |\partial^\beta F_K(\hat{x})| \leq ch_K^\ell$$

so that the definition (2.15) involves estimates (2.17).

Step 2 : F_K verifies estimates (2.19)

Expressions (2.10) and estimates (2.23) imply

$$(2.25) \quad |J_{F_K}(\hat{x})| = |\overline{a_3 a_1} \times \overline{a_3 a_2}| + 0(h_K^3).$$

Then, the hypotheses (2.2) and (2.3) give the two first estimates (2.19). The last inequalities (2.19) are direct consequences of estimates (2.24).

Step 3 : For h_K sufficiently small, F_K is injective :

We use a result from [33, Corollary 2] expressing that if \mathcal{O} is an open set of \mathbb{R}^n , if M is a compact of \mathcal{O} with a connected boundary ∂M , if $g : \mathcal{O} \rightarrow \mathbb{R}^n$ is of class C^1 such that $J_g(x) > 0$ for all $x \in M$, and if $g|_{\partial M}$ is injective, then $g|_M$ is injective.

Here we take $M = \bar{\hat{K}}$, $g = F_K$ which can be extended to an open set \mathcal{O} of \mathbb{R}^n containing \hat{K} without any difficulty. Since estimates (2.19) imply $J_{F_K}(\hat{x}) > 0$, $\forall \hat{x} \in \bar{\hat{K}}$, it remains to prove that $F_K|_{\partial\hat{K}}$ is injective.




Figure 2.4 : Basic situations for which F_K is not injective

Firstly, the definition (2.10) implies immediately that the restriction of F_K to the sides $\hat{a}_3\hat{a}_1$ and $\hat{a}_3\hat{a}_2$ is injective. Now, it remains to prove that situations as illustrated in Figure 2.4 cannot occur. Situation (i) is avoided if the restriction of F_K to the side $\hat{a}_1\hat{a}_2$ is injective. Since $F_{K_\alpha}(1 - \hat{x}_2, \hat{x}_2) \equiv \psi_{\alpha h}(\hat{x}_2)$, we only need to prove that the system $\psi_{\alpha h}(X) = \psi_{\alpha h}(Y)$, $\alpha = 1, 2$, $X, Y \in [0, 1]$ has the unique solution $X = Y$. This property is easy to prove by using estimates (2.23) and by assuming h_K sufficiently small. In the same way we prove that situations (ii) and (iii) cannot occur for h_K sufficiently small. For instance, situation (ii) leads to the system

$$\begin{cases} x_{\alpha 3} - x_{\alpha 1} + (x_{\alpha 2} - x_{\alpha 3})X - (x_{\alpha 2} - x_{\alpha 1})Y - Y(1 - Y)P_{n-2;\alpha}(Y) = 0 \\ 0 \leq X, Y \leq 1 \end{cases}$$

which admits the unique solution $X = Y = 1$ (i.e., point a_2) when h_K is sufficiently small.

Step 4 : F_K is an homeomorphism from $\bar{\hat{K}}$ onto \bar{K} :

Since F_K is continuous and injective, F_K is an homeomorphism from $\bar{\hat{K}}$ onto $F_K(\bar{\hat{K}})$. By construction, the image of the boundary $\partial\hat{K}$ is the boundary ∂K . Since an homeomorphism applies the interior of a Jordan curve onto the interior of its image (see [34, §§110 to 113]), the application F_K is an homeomorphism from $\bar{\hat{K}}$ onto \bar{K} .

Step 5 : F_K is a C^∞ -diffeomorphism from $\bar{\hat{K}}$ onto \bar{K} :

Application F_K is C^∞ and its Jacobian $J_{F_K}(\hat{x})$ is different from 0 over $\bar{\hat{K}}$. It follows that the inverse application of F_K is also C^∞ (see [35]).

Step 6 : Application F_K^{-1} verifies estimates (2.18) :

We give a proof by induction. Starting from identity $F_K \circ F_K^{-1} = I$ and using estimates (2.17) and (2.24), we obtain

$$\|F_K^{-1}\|_{1,\infty,K} = \sup_{x \in K} \|DF_K^{-1}(x)\| \leq c \sup_{x \in K} \max_{\alpha=1,2} \left| \frac{\partial F_K^{-1}}{\partial x_\alpha}(x) \right| \leq \frac{c}{h_K},$$

i.e., the estimate (2.18) for $\ell = 1$.

Next, let us assume that estimate (2.18) is true for $\ell = 2, \dots, m-1$ and let us show that (2.18) is true for $\ell = m$. For $m \geq 2$, we have (see [35, section 7.5]) :

$$(2.26) \quad \left\{ \begin{array}{l} m! \sum_{\ell=1}^m \sum_{j \in J(\ell,m)} \frac{1}{\ell!} D^\ell F_K^{-1}(x_1, x_2) \left[\frac{1}{j_1!} D^{j_1} F_K(\hat{x}_1, \hat{x}_2)(\xi_1, \dots, \xi_{j_1}), \dots, \right. \\ \left. \frac{1}{j_\ell!} D^{j_\ell} F_K(\hat{x}_1, \hat{x}_2)(\xi_{m-j_\ell+1}, \dots, \xi_m) \right] = 0, \end{array} \right.$$

where

$$J(\ell, m) = \{j = (j_1, \dots, j_\ell) \in \mathbf{N}^\ell ; 1 \leq j_1, \dots, j_\ell \leq m, j_1 + j_2 + \dots + j_\ell = m\}.$$

Since the application F_K is a C^∞ -diffeomorphism from \bar{K} onto \bar{K} , the application $DF_K(\hat{x}_1, \hat{x}_2) : \bar{K} \rightarrow \bar{K}$ is bijective. Let $\eta_i = DF_K(\hat{x}_1, \hat{x}_2)\xi_i, i = 1, \dots, m$ and, conversely, $\xi_i = DF_K^{-1}(x_1, x_2)\eta_i$. Relation (2.26) can be written as

$$(2.27) \quad \left\{ \begin{array}{l} D^m F_K^{-1}(x)(\eta_1, \dots, \eta_m) = -m! \sum_{\ell=1}^{m-1} \sum_{j \in J(\ell,m)} \frac{1}{\ell!} D^\ell F_K^{-1}(x) \\ \left\{ \frac{1}{j_1!} D^{j_1} F_K(\hat{x})[DF_K^{-1}(x)\eta_1, \dots, DF_K^{-1}(x)\eta_{j_1}], \dots, \right. \\ \left. \frac{1}{j_\ell!} D^{j_\ell} F_K(\hat{x})[DF_K^{-1}(x)\eta_{m-j_\ell+1}, \dots, DF_K^{-1}(x)\eta_m] \right\} \end{array} \right.$$

Then, estimates (2.17) and the hypothesis of the proof by induction give $\|D^m F_K^{-1}(x)\| \leq \frac{c}{h_K}$, hence $\|F_K^{-1}\|_{m,\infty,K} \leq ch_K^{-1}$ and thus, estimates (2.18) are proved.

Step 7 : F_K^{-1} satisfies estimates (2.20)

It suffices to combine equality $J_{F_K}(\hat{x})J_{F_K^{-1}}(x) = 1$, with the estimates (2.19). ■

2.4 Examples

In this section we give three examples of functions F_K which seem to be practically the most attractive. According to equations (2.8) and (2.10) it suffices to define the approximate arc γ_h .

Example 2.1 : Construction of γ_h by using polynomials of order 2

The degrees of freedom of the Lagrange-type interpolation are given by

$$(2.28) \quad \begin{cases} x_{\alpha 1} = \psi_{\alpha}(0) = \chi_{\alpha}(\underline{s}), & x_{\alpha 2} = \psi_{\alpha}(1) = \chi_{\alpha}(\bar{s}), \\ x_{\alpha 0} = \psi_{\alpha}\left(\frac{1}{2}\right) = \chi_{\alpha}\left(\frac{\underline{s} + \bar{s}}{2}\right), & \alpha = 1, 2 \end{cases}$$

so that relations (2.8) and (2.10) lead to :

$$(2.29) \quad \psi_{\alpha h}(\hat{x}_2) = x_{\alpha 1} + (x_{\alpha 2} - x_{\alpha 1})\hat{x}_2 + 4\hat{x}_2(1 - \hat{x}_2) \left[x_{\alpha 0} - \frac{x_{\alpha 1} + x_{\alpha 2}}{2} \right]$$

and then

$$(2.30) \quad \begin{cases} F_{K\alpha}(\hat{x}_1, \hat{x}_2) = x_{\alpha 3} + (x_{\alpha 1} - x_{\alpha 3})\hat{x}_1 \\ + (x_{\alpha 2} - x_{\alpha 3})\hat{x}_2 + 4 \left[x_{\alpha 0} - \frac{x_{\alpha 1} + x_{\alpha 2}}{2} \right] \hat{x}_1 \hat{x}_2. \end{cases}$$

Example 2.2 : Construction of γ_h by using polynomials of order 3

The degrees of freedom of the Hermite-type interpolation are given by

$$(2.31) \quad \begin{cases} x_{\alpha 1} = \psi_{\alpha}(0) = \chi_{\alpha}(\underline{s}), & x_{\alpha 2} = \psi_{\alpha}(1) = \chi_{\alpha}(\bar{s}), \\ \psi'_{\alpha}(0) = (\bar{s} - \underline{s})\chi'_{\alpha}(\underline{s}), & \psi'_{\alpha}(1) = (\bar{s} - \underline{s})\chi'_{\alpha}(\bar{s}), & \alpha = 1, 2 \end{cases}$$

so that

$$(2.32) \quad \begin{cases} \psi_{\alpha h}(\hat{x}_2) = x_{\alpha 1} + (x_{\alpha 2} - x_{\alpha 1})\hat{x}_2 \\ + \hat{x}_2(1 - \hat{x}_2) \{ [2(x_{\alpha 2} - x_{\alpha 1}) - (\bar{s} - \underline{s})[\chi'_{\alpha}(\underline{s}) + \chi'_{\alpha}(\bar{s})]] \hat{x}_2 \\ + x_{\alpha 1} - x_{\alpha 2} + (\bar{s} - \underline{s})\chi'_{\alpha}(\underline{s}) \} \end{cases}$$

and then

$$(2.33) \quad \begin{cases} F_{K\alpha}(\hat{x}_1, \hat{x}_2) = x_{\alpha 3} + (x_{\alpha 1} - x_{\alpha 3})\hat{x}_1 + (x_{\alpha 2} - x_{\alpha 3})\hat{x}_2 \\ + \frac{1}{2}\hat{x}_1\hat{x}_2 \{ [2(x_{\alpha 2} - x_{\alpha 1}) - (\bar{s} - \underline{s})[\chi'_{\alpha}(\underline{s}) + \chi'_{\alpha}(\bar{s})]](\hat{x}_2 - \hat{x}_1) \\ + (\bar{s} - \underline{s})[\chi'_{\alpha}(\underline{s}) - \chi'_{\alpha}(\bar{s})] \} \end{cases}$$

Example 2.3 : Construction of γ_h by using polynomials of order 5

Now, the degrees of freedom of the Hermite-type interpolation are given by

$$(2.34) \quad \begin{cases} x_{\alpha 1} = \psi_{\alpha}(0) = \chi_{\alpha}(\underline{s}), & x_{\alpha 2} = \psi_{\alpha}(1) = \chi_{\alpha}(\bar{s}), \\ \psi_{\alpha}^{(\ell)}(0) = (\bar{s} - \underline{s})^{\ell} \chi_{\alpha}^{(\ell)}(\underline{s}), & \psi_{\alpha}^{(\ell)}(1) = (\bar{s} - \underline{s})^{\ell} \chi_{\alpha}^{(\ell)}(\bar{s}), & \ell = 1, 2; \alpha = 1, 2 \end{cases}$$

so that

$$(2.35) \quad \begin{cases} \psi_{\alpha h}(\hat{x}_2) = x_{\alpha 1} + (x_{\alpha 2} - x_{\alpha 1})\hat{x}_2 \\ \quad + \hat{x}_2(1 - \hat{x}_2)[\beta_{\alpha 3}(\hat{x}_2)^3 + \beta_{\alpha 2}(\hat{x}_2)^2 + \beta_{\alpha 1}\hat{x}_2 + \beta_{\alpha 0}] \end{cases}$$

where the coefficients $\beta_{\alpha \ell}$, $\ell = 0, 1, 2, 3$ are given by

$$(2.36) \quad \begin{cases} \beta_{\alpha 0} = x_{\alpha 1} - x_{\alpha 2} + (\bar{s} - \underline{s})\chi'_{\alpha}(\underline{s}) \\ \beta_{\alpha 1} = x_{\alpha 1} - x_{\alpha 2} + (\bar{s} - \underline{s})\chi'_{\alpha}(\underline{s}) + \frac{(\bar{s} - \underline{s})^2}{2}\chi''_{\alpha}(\underline{s}) \\ \beta_{\alpha 2} = 9(x_{\alpha 2} - x_{\alpha 1}) - (\bar{s} - \underline{s})[5\chi'_{\alpha}(\underline{s}) + 4\chi'_{\alpha}(\bar{s})] - \frac{(\bar{s} - \underline{s})^2}{2}[2\chi''_{\alpha}(\underline{s}) - \chi''_{\alpha}(\bar{s})] \\ \beta_{\alpha 3} = 6(x_{\alpha 1} - x_{\alpha 2}) + 3(\bar{s} - \underline{s})[\chi'_{\alpha}(\underline{s}) + \chi'_{\alpha}(\bar{s})] + \frac{(\bar{s} - \underline{s})^2}{2}[\chi''_{\alpha}(\underline{s}) - \chi''_{\alpha}(\bar{s})] \end{cases}$$

Hence, from relations (2.10)

$$(2.37) \quad \begin{cases} F_{K\alpha}(\hat{x}_1, \hat{x}_2) = x_{\alpha 3} + (x_{\alpha 1} - x_{\alpha 3})\hat{x}_1 + (x_{\alpha 2} - x_{\alpha 3})\hat{x}_2 \\ \quad + \frac{1}{2}\hat{x}_1\hat{x}_2[\beta_{\alpha 3}(\hat{x}_2)^3 + \beta_{\alpha 2}(\hat{x}_2)^2 + \beta_{\alpha 1}\hat{x}_2 + \beta_{\alpha 0}] \\ \quad + \tilde{\beta}_{\alpha 3}(\hat{x}_1)^3 + \tilde{\beta}_{\alpha 2}(\hat{x}_1)^2 + \tilde{\beta}_{\alpha 1}\hat{x}_1 + \tilde{\beta}_{\alpha 0} \end{cases}$$

where the coefficients $\tilde{\beta}_{\alpha \ell}$, $\ell = 0, 1, 2, 3$ are given by

$$(2.38) \quad \begin{cases} \tilde{\beta}_{\alpha 0} = x_{\alpha 2} - x_{\alpha 1} - (\bar{s} - \underline{s})\chi'_{\alpha}(\bar{s}) \\ \tilde{\beta}_{\alpha 1} = x_{\alpha 2} - x_{\alpha 1} - (\bar{s} - \underline{s})\chi'_{\alpha}(\bar{s}) + \frac{(\bar{s} - \underline{s})^2}{2}\chi''_{\alpha}(\bar{s}) \\ \tilde{\beta}_{\alpha 2} = 9(x_{\alpha 1} - x_{\alpha 2}) + (\bar{s} - \underline{s})[5\chi'_{\alpha}(\bar{s}) + 4\chi'_{\alpha}(\underline{s})] - \frac{(\bar{s} - \underline{s})^2}{2}[2\chi''_{\alpha}(\bar{s}) - \chi''_{\alpha}(\underline{s})] \\ \tilde{\beta}_{\alpha 3} = 6(x_{\alpha 2} - x_{\alpha 1}) - 3(\bar{s} - \underline{s})[\chi'_{\alpha}(\bar{s}) + \chi'_{\alpha}(\underline{s})] + \frac{(\bar{s} - \underline{s})^2}{2}[\chi''_{\alpha}(\bar{s}) - \chi''_{\alpha}(\underline{s})] \end{cases}$$

3 DEFINITION OF CURVED \mathcal{C}^1 FINITE ELEMENTS


In this paragraph, we define curved finite elements which have a connection of class \mathcal{C}^1 (we will say \mathcal{C}^1 -compatible) with ARGYRIS and BELL triangles. We state the basic principles in section 3.1. In section 3.2, we detail this definition for a connection between a curved finite element and an ARGYRIS triangle in case of an approximate boundary parameterized by polynomials of degree five. Following a remark of [17], we indicate in


section 3.3 the simplifications that it is possible to realize when the boundary conditions are of homogeneous Dirichlet type : the approximate boundary γ_h can be parameterized by polynomials of degree three only. In both cases, we mention the modifications which permit to obtain a curved element which is C^1 -compatible with the BELL triangle. Finally, in section 3.4, we define curved elements which have a connection of class C^0 with an Hermite element of degree 3 ; these elements can be used for instance to approximate tangential components of the displacement for thin shell problems.

3.1 Basic principles

Figures 3.1 and 3.2 recall the definitions of the ARGYRIS and BELL triangles as well as their interpolation properties. By P_K and Σ_K we note respectively the functional space and the set of degrees of freedom of the finite element while π_K means the associate interpolation operator.

Among the principles we need to observe when constructing such curved finite elements, we shall consider the "essential" conditions and the "desirable" conditions.





"Essential" conditions :

The connections between the curved elements, associated to curved triangles K of Figure 2.1, and the adjacent (straight or curved) finite elements are realized through the straight sides $a_3 a_1$ and $a_3 a_2$, as mentioned in Figure 3.3.

First, consider the connection of a curved finite element associated to the curved triangle K with the ARGYRIS (or BELL) triangle associated to the triangle K_D of Figure 3.3. Such a connection will be of class C^1 if, and only if, the traces of functions $p \in P_K$ and of their normal derivatives $\partial p / \partial n_{31}$ along the side $a_3 a_1$ coincide with their homologues of the adjacent finite element. In order to satisfy these conditions, it is sufficient to meet the following requirements :

$$(3.1) \left\{ \begin{array}{l} \text{(i) the degrees of freedom of the curved finite element relative to the sides} \\ a_3 a_\alpha, \alpha = 1, 2, \text{ are identical to that of the adjacent finite elements.} \end{array} \right.$$

$$(3.2) \left\{ \begin{array}{l} \text{(ii) the traces } p|_{[a_3, a_\alpha]} \left(\text{resp. } \frac{\partial p}{\partial n_{3\alpha}} \Big|_{[a_3, a_\alpha]} \right), \alpha = 1, 2, \text{ of functions} \\ p \in P_K \text{ associated to the curved triangle } K \text{ are one-variable} \\ \text{polynomials of degree 5 (resp. 4 for ARGYRIS triangle and 3} \\ \text{for BELL triangle), entirely determined by the degrees of} \\ \text{freedom relative to the sides } a_3 a_\alpha, \alpha = 1, 2. \end{array} \right.$$

Likewise, it is worth to note that these conditions insure a C^1 connection between two adjacent curved finite elements.




Figure 3.3 : Adjacent triangles to a curved triangle K

"Desirable" conditions :

The application F_K , studied in paragraph 2, associates the curved triangle K to the reference triangle \hat{K} . We use the same application F_K to associate to any function v defined over the triangle K , a function \hat{v} defined over the triangle \hat{K} , i.e.,

$$(3.3) \quad v = \hat{v} \circ F_K^{-1}, \quad \hat{v} = v \circ F_K.$$

Then, it is "desirable" that the following condition is satisfied :

$$(3.4) \quad \left\{ \begin{array}{l} \text{(iii) To any function } p \in P_K, \text{ defined over the curved triangle } K, \\ \text{the correspondance (3.3) associates a polynomial function } \hat{p} = p \circ F_K. \end{array} \right.$$

This condition (3.4) is convenient for the study of the approximation error and to take into account the numerical integration and the boundary conditions.

On the other hand, this condition leads to the definition of reference finite elements which are most complicated than those associated to corresponding straight finite elements. Indeed, let \hat{a} be any point of the side $\hat{a}_3\hat{a}_1$ of the triangle \hat{K} (see Figure 3.4) and set $a = F_K(\hat{a})$. Then, relation $\hat{p} = p \circ F_K$ involves

$$(3.5) \quad \frac{\partial \hat{p}}{\partial \hat{x}_2}(\hat{a}) = \left\langle \frac{\partial F_K}{\partial \hat{x}_2}(\hat{a}), t_{31} \right\rangle \frac{\partial p}{\partial t_{31}}(a) + \left\langle \frac{\partial F_K}{\partial \hat{x}_2}(\hat{a}), n_{31} \right\rangle \frac{\partial p}{\partial n_{31}}(a),$$

and a similar relation for any point $\hat{a} \in \hat{a}_3\hat{a}_2$:

$$(3.6) \quad \frac{\partial \hat{p}}{\partial \hat{x}_1}(\hat{a}) = \left\langle \frac{\partial F_K}{\partial \hat{x}_1}(\hat{a}), t_{32} \right\rangle \frac{\partial p}{\partial t_{32}}(a) + \left\langle \frac{\partial F_K}{\partial \hat{x}_1}(\hat{a}), n_{32} \right\rangle \frac{\partial p}{\partial n_{32}}(a),$$

where $\langle \cdot, \cdot \rangle$ denotes the usual scalar product of \mathbb{R}^2 .




Figure 3.4 : Local reference systems along the straight sides of the curved triangle K ($t_{3\alpha}, n_{3\alpha}$ are the unit tangential and normal vectors to the side a_3a_α).

By consideration of the hypotheses (3.2), we verify that

$$(3.7) \quad \left\{ \begin{array}{l} \text{the derivatives } \frac{\partial p}{\partial t_{3\alpha}}(a), \text{ for } a \in a_3a_\alpha, \alpha = 1, 2, \text{ are polynomials} \\ \text{of degree 4 with respect to } \hat{x}_\alpha \text{ (note that } F_K \text{ is affine along } \hat{a}_3\hat{a}_\alpha \text{)}; \end{array} \right.$$

$$(3.8) \quad \left\{ \begin{array}{l} \text{the derivatives } \frac{\partial p}{\partial n_{3\alpha}}(a), \text{ for } a \in a_3a_\alpha, \alpha = 1, 2, \text{ are polynomials} \\ \text{of degree 4 (ARGYRIS) or 3 (BELL) with respect to } \hat{x}_\alpha; \end{array} \right.$$

Thus, relations (2.10) (3.5) (3.6) (3.7) and (3.8) involve that, for $\hat{a} \in \hat{a}_3\hat{a}_1$, $\frac{\partial \hat{p}}{\partial \hat{x}_2}(\hat{a})$ is a polynomial of degree $n + 3$ with respect to \hat{x}_1 , while for $\hat{a} \in \hat{a}_3\hat{a}_2$, $\frac{\partial \hat{p}}{\partial \hat{x}_1}(\hat{a})$ is a polynomial of degree $n + 3$ with respect to \hat{x}_2 . Let us denote

$$\hat{P}_K = \{ \hat{p} : \hat{K} \rightarrow \mathbb{R}; \hat{p} = p \circ F_K, p \in P_K \}$$

the space of functions defined over the reference triangle \hat{K} from the space P_K and through the application F_K . Then, in order to satisfy condition (3.4), above remarks show that we need to satisfy the inclusion

$$(3.9) \quad \hat{P}_K \subset P_{n+4}.$$

By using these considerations, we show in section 3.2 that it is possible to define curved finite elements compatible with ARGYRIS or BELL triangles in case of $n = 5$, i.e.,

$F_K \in (P_5)^2$. In section 3.3, we examine the more simple case $n = 3$ which is sufficient for homogeneous Dirichlet boundary conditions.

3.2 Definition of curved finite elements \mathcal{C}^1 -compatible with Argyris or Bell triangles when $F_K \in (P_5)^2$

Subsequently, we detail the definition of a curved finite element \mathcal{C}^1 -compatible with ARGYRIS triangle. Similarly a curved finite element \mathcal{C}^1 -compatible with BELL triangle can be constructed ; the modifications will be mentioned by "resp. BELL :...". Since all the curved elements into consideration satisfy conditions (3.1) (3.2) and (3.4), the \mathcal{C}^1 -compatibility between two adjacent curved finite elements is automatically satisfied.

For our definition, we associate to the reference triangle \hat{K} a basic finite element $(\hat{K}, \hat{P}, \hat{\Sigma})$ described in Figure 3.5. The choice of $\hat{P} = P_9$ takes into account inclusion (3.9). We say that this finite element is "basic" by opposition to the notion of reference finite element (see the theory of affine finite elements). To get the reference finite element associated to the curved finite element into consideration we will have to impose 24 (resp. BELL : 27) constraints to the set \hat{P} .

Considering Figure 3.5, it remains to prove :

Theorem 3.1 : The triple $(\hat{K}, \hat{P}, \hat{\Sigma})$ of Figure 3.5 defines a finite element

Proof : Since $\dim \hat{P} = \text{card}(\hat{\Sigma})$, we need to prove that 0 is the unique element of \hat{P} whose corresponding degrees of freedom $\hat{\Sigma}$ are zero. First we remark that such a function \hat{p} is identically zero when restricted to the sides of \hat{K} so that $\hat{p}(\hat{x}_1, \hat{x}_2) = [\hat{x}_1 \hat{x}_2 (1 - \hat{x}_1 - \hat{x}_2)]^2 \hat{q}(\hat{x}_1, \hat{x}_2)$ where $\hat{q} \in P_3$. Next, we have $\hat{q}(\hat{e}_i) = 0, i = 1, \dots, 10$ so that $\hat{q} \equiv 0$. ■


Construction of the interpolate function $v \longrightarrow \pi_K v$:

By using this basic finite element we are going to associate to any regular function v (for instance $v \in C^2(K)$) defined over the curved triangle K its interpolate $\pi_K v$. It takes three steps :

Step 1 : Definition of the set Σ_K of degrees of freedom of the curved element

In all section 3.2, application F_K is that of Example 2.3, i.e., (2.37). Let us set

$$(3.10) \quad \left\{ \begin{array}{l} a_i = F_K(\hat{a}_i), \quad b_i = F_K(\hat{b}_i), \quad i = 1, 2, 3 ; \quad d_i = F_K(\hat{d}_i), \quad i = 1, \dots, 12 ; \\ e_i = F_K(\hat{e}_i), \quad i = 1, \dots, 10 ; \\ t_{3\alpha}, n_{3\alpha}, \alpha = 1, 2, \text{ unit vectors defined according to Figure 3.4} \end{array} \right.$$



\hat{K} = unit right-angled triangle ;

$\hat{P} = P_9$; $\dim \hat{P} = 55$;

$$\hat{\Sigma}(\hat{w}) = \left\{ \begin{aligned} & \hat{w}(\hat{a}_i), \frac{\partial \hat{w}}{\partial \hat{x}_1}(\hat{a}_i), \frac{\partial \hat{w}}{\partial \hat{x}_2}(\hat{a}_i), \frac{\partial^2 \hat{w}}{\partial \hat{x}_1^2}(\hat{a}_i), \frac{\partial^2 \hat{w}}{\partial \hat{x}_1 \partial \hat{x}_2}(\hat{a}_i), \frac{\partial^2 \hat{w}}{\partial \hat{x}_2^2}(\hat{a}_i), \quad i = 1, 2, 3 ; \\ & -\frac{\partial \hat{w}}{\partial \hat{x}_1}(\hat{b}_1) ; -\frac{\partial \hat{w}}{\partial \hat{x}_2}(\hat{b}_2) ; \frac{\sqrt{2}}{2} \left(\frac{\partial \hat{w}}{\partial \hat{x}_1} + \frac{\partial \hat{w}}{\partial \hat{x}_2} \right) (\hat{b}_3) ; \hat{w}(\hat{d}_i), \quad i = 1, \dots, 12 ; \\ & -\frac{\partial \hat{w}}{\partial \hat{x}_1}(\hat{d}_i), \quad i = 1, \dots, 4 ; -\frac{\partial \hat{w}}{\partial \hat{x}_2}(\hat{d}_i), \quad i = 5, \dots, 8 ; \\ & \frac{\sqrt{2}}{2} \left(\frac{\partial \hat{w}}{\partial \hat{x}_1} + \frac{\partial \hat{w}}{\partial \hat{x}_2} \right) (\hat{d}_i), \quad i = 9, \dots, 12 ; \hat{w}(\hat{e}_i), \quad i = 1, \dots, 10 \end{aligned} \right\}$$

Figure 3.5 : "Basic" P_9 -Hermite finite element

Then, the set $\Sigma_K(v)$ of values of degrees of freedom of v is given by (see Fig. 3.6) :

$$(3.11) \quad \left\{ \begin{array}{l} \Sigma_K(v) = \{(D^\alpha v(a_i), |\alpha| = 0, 1, 2), i = 1, 2, 3; Dv(b_2)n_{31}; \\ Dv(b_1)n_{32}; Dv(b_3)DF_K(\hat{b}_3)(\hat{a}_3 - \hat{b}_3); v(e_i), i = 1, \dots, 10\}; \end{array} \right.$$

$$(3.12) \quad \left\{ \begin{array}{l} \text{(resp. BELL :} \\ \Sigma_K(v) = \{(D^\alpha v(a_i), |\alpha| = 0, 1, 2), i = 1, 2, 3; v(e_i), i = 1, \dots, 10\}. \end{array} \right.$$




Figure 3.6 : Sets Σ_K for C^1 -compatible curved finite element ($F_K \in (P_5)^2$)

Note that in addition to the usual degrees of freedom of the corresponding classical elements, we find ten additional degrees of freedom inside of the triangle K . Also, note that the sets Σ_K given by (3.11) (3.12) satisfy condition (3.1).

Step 2 : Definition of the set $\hat{\Delta}_K(v)$ from $\Sigma_K(v)$

Consider the following partition of the set $\hat{\Sigma} : \hat{\Sigma} = \hat{\Sigma}_1 \cup \hat{\Sigma}_2 \cup \hat{\Sigma}_3$ where

$$(3.13) \quad \left\{ \begin{array}{l} \hat{\Sigma}_1 = \{(D^\alpha \hat{p}(a_i), |\alpha| = 0, 1, 2), i = 1, 2, 3; \hat{p}(\hat{e}_i), i = 1, \dots, 10\}, \\ \hat{\Sigma}_2 = \{\hat{p}(\hat{d}_i), i = 1, \dots, 8; -\frac{\partial \hat{p}}{\partial \hat{x}_1}(\hat{b}_1); -\frac{\partial \hat{p}}{\partial \hat{x}_1}(\hat{d}_i), i = 1, \dots, 4; \\ -\frac{\partial \hat{p}}{\partial \hat{x}_2}(\hat{b}_2); -\frac{\partial \hat{p}}{\partial \hat{x}_2}(\hat{d}_i), i = 5, \dots, 8\}; \\ \hat{\Sigma}_3 = \{\hat{p}(\hat{d}_i), i = 9, \dots, 12; \frac{\sqrt{2}}{2} \left(\frac{\partial \hat{p}}{\partial \hat{x}_1} + \frac{\partial \hat{p}}{\partial \hat{x}_2} \right) (\hat{b}_3); \\ \frac{\sqrt{2}}{2} \left(\frac{\partial \hat{p}}{\partial \hat{x}_1} + \frac{\partial \hat{p}}{\partial \hat{x}_2} \right) (\hat{d}_i), i = 9, \dots, 12\} \end{array} \right.$$

To the set of values $\Sigma_K(v)$, we are going to associate the sets of values $\hat{\Delta}_i(v)$ that we need to attribute to $\hat{\Sigma}_i, i = 1, 2, 3$ to obtain a suitable function $\hat{w} \in \hat{P}$. Let us note $\hat{\Delta}_K(v) = \hat{\Delta}_{K1}(v) \cup \hat{\Delta}_{K2}(v) \cup \hat{\Delta}_{K3}(v)$. By means of the application F_K we define the function

$$(3.14) \quad \hat{v} = v \circ F_K.$$

Then, from the set $\Sigma_K(v)$ we immediately derive

$$(3.15) \quad \hat{\Delta}_{K1}(v) = \{(D^\alpha \hat{v}(\hat{a}_i), |\alpha| = 0, 1, 2), i = 1, 2, 3; \hat{v}(\hat{e}_i), i = 1, \dots, 10\}.$$

Now, consider the set $\hat{\Delta}_{K2}(v)$. We first examine the case of the degrees of freedom of $\hat{\Sigma}_2$ located on the side $\hat{a}_3\hat{a}_1$. In order to obtain an interpolate function $\pi_K v$ which satisfies conditions (3.2) we need that :

- on the one hand, its trace $\pi_K v|_{[a_3, a_1]}$ coincides with the one-variable P_5 -Hermite polynomial defined by the data of the following degrees of freedom

$$(3.16) \quad \left\{ \begin{array}{l} \{v(a_1), v(a_3), Dv(a_1)(a_3 - a_1), Dv(a_3)(a_1 - a_3), \\ D^2v(a_1)(a_3 - a_1)^2, D^2v(a_3)(a_1 - a_3)^2\}. \end{array} \right.$$

We parameterize the side a_3a_1 by using \hat{x}_1 , i.e.,

$$(3.17) \quad x_1 = x_{13} + (x_{11} - x_{13})\hat{x}_1, \quad x_2 = x_{23} + (x_{21} - x_{23})\hat{x}_1$$

and we note

$$(3.18) \quad \hat{f}_1 \text{ (which will coincide with } (\pi_K v) \circ F_K|_{[\hat{a}_3, \hat{a}_1]})$$

the so-defined Hermite polynomial ;

- on the other hand, its normal derivative $\frac{\partial \pi_K v}{\partial n_{31}}$ has a trace $\frac{\partial \pi_K v}{\partial n_{31}}|_{[a_3, a_1]}$ which coincides with the one-variable P_4 -Hermite polynomial (resp. BELL : P_3), defined by the data of the following degrees of freedom

$$(3.19) \quad \{Dv(a_i)n_{31}, D^2v(a_i)(n_{31}, t_{31}), i = 1, 3; Dv(b_2)n_{31}\},$$

$$(3.20) \quad (\text{resp. BELL : } \{Dv(a_i)n_{31}, D^2v(a_i)(n_{31}, t_{31}), i = 1, 3\})$$

Similarly to (3.17) and (3.18), we note

$$(3.21) \quad \hat{g}_1 \text{ (which will coincide with } \left(\frac{\partial \pi_K v}{\partial n_{31}}\right) \circ F_K|_{[\hat{a}_3, \hat{a}_1]}),$$

$$(3.22) \quad (\text{resp. BELL : } \hat{h}_1 \text{ (which will coincide with } \left(\frac{\partial \pi_K v}{\partial n_{31}}\right) \circ F_K|_{[\hat{a}_3, \hat{a}_1]}),$$

the so-defined Hermite polynomial. Next, we define Hermite polynomials over the second straight side, i.e.,

$$(3.23) \quad x_1 = x_{13} + (x_{12} - x_{13})\hat{x}_2, \quad x_2 = x_{23} + (x_{22} - x_{23})\hat{x}_2,$$

$$(3.24) \quad \hat{f}_2 \text{ (which will coincide with } (\pi_K v) \circ F_K|_{[\hat{a}_3, \hat{a}_2]}),$$

$$(3.25) \quad \hat{g}_2 \text{ (which will coincide with } \left(\frac{\partial \pi_K v}{\partial n_{32}} \right) \circ F_K|_{[\hat{a}_3, \hat{a}_2]}),$$

$$(3.26) \quad \text{(resp. BELL : } \hat{h}_2 \text{ (which will coincide with } \left(\frac{\partial \pi_K v}{\partial n_{32}} \right) \circ F_K|_{[\hat{a}_3, \hat{a}_2]})).$$

At this stage, note that the polynomial functions $\hat{f}_\alpha, \hat{g}_\alpha, \hat{h}_\alpha, \alpha = 1, 2$, are only dependent of the values of the degrees of freedom of the function v , given in the set $\Sigma_K(v)$, and relative to the sides $a_3 a_1$ and $a_3 a_2$. Then, the set of values $\hat{\Delta}_{K_2}(v)$ associated to the set of degrees of freedom $\hat{\Sigma}_2$ (see (3.13)) is given by :

$$(3.27) \quad \left\{ \begin{array}{l} \hat{\Delta}_{K_2}(v) = \{ \hat{f}_2(\hat{d}_i), i = 1, \dots, 4 ; \hat{f}_1(\hat{d}_i), i = 5, \dots, 8 ; \\ - \frac{1}{|a_3 a_2|} \left\{ \left\langle \frac{\partial F_K}{\partial \hat{x}_1}(\hat{b}_1), t_{32} \right\rangle \frac{d\hat{f}_2}{d\hat{x}_2}(\hat{b}_1) + |a_3 a_2| \left\langle \frac{\partial F_K}{\partial \hat{x}_1}(\hat{b}_1), n_{32} \right\rangle \hat{g}_2(\hat{b}_1) \right\} ; \\ - \frac{1}{|a_3 a_2|} \left\{ \left\langle \frac{\partial F_K}{\partial \hat{x}_1}(\hat{d}_i), t_{32} \right\rangle \frac{d\hat{f}_2}{d\hat{x}_2}(\hat{d}_i) + |a_3 a_2| \left\langle \frac{\partial F_K}{\partial \hat{x}_1}(\hat{d}_i), n_{32} \right\rangle \hat{g}_2(\hat{d}_i) \right\}, \\ \quad \quad \quad i = 1, \dots, 4 ; \\ - \frac{1}{|a_3 a_1|} \left\{ \left\langle \frac{\partial F_K}{\partial \hat{x}_2}(\hat{b}_2), t_{31} \right\rangle \frac{d\hat{f}_1}{d\hat{x}_1}(\hat{b}_2) + |a_3 a_1| \left\langle \frac{\partial F_K}{\partial \hat{x}_2}(\hat{b}_2), n_{31} \right\rangle \hat{g}_1(\hat{b}_2) \right\} ; \\ - \frac{1}{|a_3 a_1|} \left\{ \left\langle \frac{\partial F_K}{\partial \hat{x}_2}(\hat{d}_i), t_{31} \right\rangle \frac{d\hat{f}_1}{d\hat{x}_1}(\hat{d}_i) + |a_3 a_1| \left\langle \frac{\partial F_K}{\partial \hat{x}_2}(\hat{d}_i), n_{31} \right\rangle \hat{g}_1(\hat{d}_i) \right\}, \\ \quad \quad \quad i = 5, \dots, 8 \} \\ \text{(resp. BELL : replace } \hat{g}_\alpha \text{ by } \hat{h}_\alpha, \alpha = 1, 2). \end{array} \right.$$

To find the expression of the ten last elements of $\hat{\Delta}_{K_2}(v)$ we have used relations (3.5) and (3.6), by observing that from conditions (3.18) and (3.24), we obtain :

$$(3.28) \quad \left\{ \begin{array}{l} \frac{d\hat{f}_2}{d\hat{x}_2} \text{ will coincide with } |a_3 a_2| \left(\frac{\partial \pi_K v}{\partial t_{32}} \right) \circ F_K \Big|_{[\hat{a}_3, \hat{a}_2]} \\ \frac{d\hat{f}_1}{d\hat{x}_1} \text{ will coincide with } |a_3 a_1| \left(\frac{\partial \pi_K v}{\partial t_{31}} \right) \circ F_K \Big|_{[\hat{a}_3, \hat{a}_1]} \end{array} \right.$$

In a third set $\hat{\Delta}_{K3}(v)$ we give suitable values for the nine degrees of freedom of $\hat{\Sigma}_3$. First, let us observe that the correspondence $\hat{v} = v \circ F_K$ involves

$$(3.29) \quad D\hat{v}(\hat{b}_3)(\hat{a}_3 - \hat{b}_3) = Dv(b_3)DF_K(\hat{b}_3)(\hat{a}_3 - \hat{b}_3).$$

The second member is known since it appears among the degrees of freedom of Σ_K (see (3.11)). Moreover, expression (3.15) contains the values of $D^\alpha \hat{v}(\hat{a}_i)$, $|\alpha| = 0, 1, 2$. The side $\hat{a}_1\hat{a}_2$ of triangle \hat{K} is parameterized by

$$(3.30) \quad \hat{x}_1 = \hat{x}_1, \quad \hat{x}_2 = 1 - \hat{x}_1.$$

Then, we note $\hat{f}_3(\hat{x}_1)$ the P_5 -Hermite polynomial defined by the data of the degrees of freedom

$$(3.31) \quad \left\{ \begin{array}{l} \{\hat{v}(\hat{a}_1), \hat{v}(\hat{a}_2), D\hat{v}(\hat{a}_1)(\hat{a}_2 - \hat{a}_1), D\hat{v}(\hat{a}_2)(\hat{a}_1 - \hat{a}_2), \\ D^2\hat{v}(\hat{a}_1)(\hat{a}_2 - \hat{a}_1)^2, D^2\hat{v}(\hat{a}_2)(\hat{a}_1 - \hat{a}_2)^2\} \end{array} \right.$$

and by $\hat{g}_3(\hat{x}_1)$ [resp. BELL : $\hat{h}_3(\hat{x}_1)$] the P_4 -Hermite polynomial (resp. BELL : P_3), defined by the data of the degrees of freedom

$$(3.32) \quad \left\{ \begin{array}{l} \{D\hat{v}(\hat{a}_\alpha)(\hat{a}_3 - \hat{b}_3), \alpha = 1, 2; D^2\hat{v}(\hat{a}_1)(\hat{a}_3 - \hat{b}_3, \hat{a}_2 - \hat{a}_1); \\ D^2\hat{v}(\hat{a}_2)(\hat{a}_3 - \hat{b}_3, \hat{a}_1 - \hat{a}_2); D\hat{v}(\hat{b}_3)(\hat{a}_3 - \hat{b}_3)\}; \end{array} \right.$$

$$(3.33) \quad \left\{ \begin{array}{l} [\text{resp. BELL : } \{D\hat{v}(\hat{a}_\alpha)(\hat{a}_3 - \hat{b}_3), \alpha = 1, 2; D^2\hat{v}(\hat{a}_1)(\hat{a}_3 - \hat{b}_3, \hat{a}_2 - \hat{a}_1); \\ D^2\hat{v}(\hat{a}_2)(\hat{a}_3 - \hat{b}_3, \hat{a}_1 - \hat{a}_2)\}]. \end{array} \right.$$

Then, we set

$$(3.34) \quad \left\{ \begin{array}{l} \hat{\Delta}_{K3}(v) = \{\hat{f}_3(\hat{d}_i), i = 9, \dots, 12; -\sqrt{2} \hat{g}_3(\hat{b}_3); -\sqrt{2} \hat{g}_3(\hat{d}_i), i = 9, \dots, 12\}; \\ [\text{resp. BELL : replace } \hat{g}_3 \text{ by } \hat{h}_3]. \end{array} \right.$$

Here again, note that the set of values $\hat{\Delta}_{K3}(v)$ is only dependent of the values $\Sigma_K(v)$ of the degrees of freedom of the function v .

Thus, the set $\hat{\Delta}_K(v) = \{\hat{\Delta}_{K1}(v), \hat{\Delta}_{K2}(v), \hat{\Delta}_{K3}(v)\}$ (see (3.15) (3.27) and (3.34)) specifies a value to every degree of freedom of $\hat{\Sigma}$ (see Fig. 3.5).

Step 3 : Definition of function $\pi_K v$ from the set $\hat{\Delta}_K(v)$

Let \hat{w} be the function of \hat{P} which takes the set of values $\hat{\Delta}_K(v)$ over the set of degrees of freedom $\hat{\Sigma}$ (see Fig. 3.5). Then, the function w is obtain through the mapping F_K^{-1} , i.e.,

$$(3.35) \quad w = \hat{w} \circ F_K^{-1}.$$

In Theorem 3.2, we prove that w interpolates the function v and verifies properties (3.1) (3.2) and (3.4), so that we are allowed to pose $w = \pi_K v$. ■

Remark 3.1 : To the function \hat{v} defined by (3.14), the basic finite element $(\hat{K}, \hat{\Sigma}, \hat{P})$ (see Fig. 3.5) associates an interpolate function $\hat{\pi}\hat{v}$ which is generally different of \hat{w} . The difference $\hat{w} - \hat{\pi}\hat{v}$ is studied in paragraph 4. ■

Now, we check that the function w verifies the desirable interpolation properties.

Theorem 3.2 : The function w , defined by (3.35), is determined in an unique way by the data of the set $\Sigma_K(v)$ of the values of the degrees of freedom of the function v , i.e.,

$$(3.36) \quad \left\{ \begin{array}{l} \Sigma_K(v) = \{(D^\alpha v(a_i), |\alpha| = 0, 1, 2), i = 1, 2, 3 ; Dv(b_2)n_{31} ; Dv(b_1)n_{32} \\ Dv(b_3)DF_K(\hat{b}_3)(\hat{a}_3 - \hat{b}_3) ; v(e_i), i = 1, \dots, 10\} \\ \\ (\text{resp. BELL : } \Sigma_K(v) = \{(D^\alpha v(a_i), |\alpha| = 0, 1, 2), i = 1, 2, 3 ; \\ v(e_i), i = 1, \dots, 10\}). \end{array} \right.$$

Moreover the function w verifies the relations :

$$(3.37) \quad D^\alpha w(a_i) = D^\alpha v(a_i), \quad |\alpha| = 0, 1, 2 ; i = 1, 2, 3 ;$$

$$(3.38) \quad Dw(b_2)n_{31} = Dv(b_2)n_{31},$$

$$(3.39) \quad Dw(b_1)n_{32} = Dv(b_1)n_{32},$$

$$(3.40) \quad Dw(b_3)DF_K(\hat{b}_3)(\hat{a}_3 - \hat{b}_3) = Dv(b_3)DF_K(\hat{b}_3)(\hat{a}_3 - \hat{b}_3),$$

$$(3.41) \quad w(e_i) = v(e_i), \quad i = 1, \dots, 10,$$

(resp. BELL : w verifies relations (3.37) and (3.41)) and w satisfies conditions (3.1) (3.2) and (3.4) so that we have $w = \pi_K v$.

Proof : The functions \hat{w} and w are only dependent on the set of values $\Sigma_K(v)$. By construction, the function \hat{w} verifies (see (3.15))

$$\left\{ \begin{array}{l} D^\alpha \hat{w}(\hat{a}_i) = D^\alpha \hat{v}(\hat{a}_i), \quad |\alpha| = 0, 1, 2, \quad i = 1, 2, 3 \\ \\ \hat{w}(\hat{e}_i) = \hat{v}(\hat{e}_i), \quad i = 1, \dots, 10, \end{array} \right.$$

so that, by using (3.14) (3.35), we get (3.37) and (3.41).

In the same way, by construction of $\hat{\Delta}_{K3}(v)$, we get

$$Dw(b_3)DF_K(\hat{b}_3)(\hat{a}_3 - \hat{b}_3) = D\hat{w}(\hat{b}_3)(\hat{a}_3 - \hat{b}_3) = \hat{g}_3 \left(\frac{1}{2} \right) = D\hat{v}(\hat{b}_3)(\hat{a}_3 - \hat{b}_3) = Dv(b_3)DF_K(\hat{b}_3)(\hat{a}_3 - \hat{b}_3),$$

and hence we obtain relation (3.40).

To prove (3.38) (3.39), we show that

$$(3.42) \quad \left[\frac{\partial w}{\partial n_{3\alpha}} \right] \circ F_K|_{[\hat{a}_3, \hat{a}_\alpha]} = \hat{g}_\alpha, \quad \alpha = 1, 2.$$

Then, relations (3.38) (3.39) are direct consequences of definitions (3.19) (3.21) and (3.25) for interpolating functions \hat{g}_α which verify

$$\hat{g}_1(\hat{b}_2) = Dv(b_2)n_{31}, \quad \hat{g}_2(\hat{b}_1) = Dv(b_1)n_{32}.$$

Therefore let us prove (3.42). First, by construction

$$(3.43) \quad \hat{w}|_{[\hat{a}_3, \hat{a}_\alpha]} = \hat{f}_\alpha.$$

Since application F_K is affine along the sides $\hat{a}_3\hat{a}_\alpha, \alpha = 1, 2$, the traces $w|_{[a_3, a_\alpha]}$ are one-dimensional P_5 -polynomials entirely determined by the degrees of freedom relative to the sides $a_3a_\alpha (\alpha = 1, 2)$. Thus, the function $w|_{[a_3, a_\alpha]}$ realizes the first part of condition (3.2).

Now, we prove relation (3.42) for $\alpha = 1$. Relations (3.5) and $\hat{w} = w \circ F_K$ involve for any $\hat{a} \in [\hat{a}_3, \hat{a}_1]$:

$$(3.44) \quad D\hat{w}(\hat{a})\vec{e}_2 = \left\langle \frac{\partial F_K}{\partial \hat{x}_2}(\hat{a}), t_{31} \right\rangle \frac{\partial w}{\partial t_{31}}(a) + \left\langle \frac{\partial F_K}{\partial \hat{x}_2}(\hat{a}), n_{31} \right\rangle \frac{\partial w}{\partial n_{31}}(a).$$

The functions $\frac{d\hat{f}_1}{d\hat{x}_1}, \hat{g}_1, \frac{\partial F_K}{\partial \hat{x}_2}|_{[\hat{a}_3, \hat{a}_1]}$ are P_4 -polynomials in \hat{x}_1 . Then, the expression

$$\frac{1}{|a_3a_1|} \left\langle \frac{\partial F_K}{\partial \hat{x}_2}(\hat{a}), t_{31} \right\rangle \frac{d\hat{f}_1}{d\hat{x}_1}(\hat{a}) + \left\langle \frac{\partial F_K}{\partial \hat{x}_2}(\hat{a}), n_{31} \right\rangle \hat{g}_1(\hat{a}),$$

is a P_8 -polynomial in \hat{x}_1 . This polynomial is identical to $D\hat{w}(\hat{a})\vec{e}_2$, which is a P_8 -polynomial in \hat{x}_1 , since both polynomials take the same values over the set

$$\left\{ \hat{p}(\hat{a}_3); \hat{p}(\hat{a}_1); \hat{p}(\hat{b}_2); \hat{p}(\hat{d}_i), i = 5, \dots, 8; \frac{\partial \hat{p}}{\partial \hat{x}_1}(\hat{a}_3), \frac{\partial \hat{p}}{\partial \hat{x}_1}(\hat{a}_1) \right\}.$$

In particular, we use the equalities

$$\frac{d\hat{f}_1}{d\hat{x}_1}(\hat{a}_i) = |a_3a_1|Dv(a_i)t_{31}, \quad \hat{g}_1(\hat{a}_i) = Dv(a_i)n_{31}, \quad i = 1, 3,$$

$$\frac{d^2 \hat{f}_1}{(d\hat{x}_1)^2}(\hat{a}_i) = |a_3 a_1|^2 D^2 v(a_i)(t_{31}, t_{31}), \quad \frac{d\hat{g}_1}{d\hat{x}_2}(\hat{a}_i) = |a_3 a_1| D^2 v(a_i)(n_{31}, t_{31}), \quad i = 1, 3,$$

which are themselves direct consequences of the definitions of \hat{f}_1 and \hat{g}_1 . Thus for any $\hat{a} \in [\hat{a}_3, \hat{a}_1]$

$$(3.45) \quad D\hat{w}(\hat{a})\vec{e}_2 = \frac{1}{|a_3 a_1|} \left\langle \frac{\partial F_K}{\partial \hat{x}_2}(\hat{a}), t_{31} \right\rangle \frac{d\hat{f}_1}{d\hat{x}_1}(\hat{a}) + \left\langle \frac{\partial F_K}{\partial \hat{x}_2}(\hat{a}), n_{31} \right\rangle \hat{g}_1(\hat{a}).$$

But, relations (3.43) imply

$$\frac{\partial w}{\partial t_{31}}(a) = \frac{1}{|a_3 a_1|} \frac{d\hat{f}_1}{d\hat{x}_1}(\hat{a}), \quad a = F_K(\hat{a}), \quad \forall \hat{a} \in \hat{a}_3 \hat{a}_1.$$

Then, relations (3.44) and (3.45) involve relation (3.42) for $\alpha = 1$. Indeed, for h_K sufficiently small, we have

$$\left| \left\langle \frac{\partial F_K}{\partial \hat{x}_2}(\hat{a}), n_{31} \right\rangle \right| = \frac{1}{|a_3 a_1|} | \overrightarrow{a_3 a_1} \times \overrightarrow{a_3 a_2} | + 0(h_K^2) \neq 0.$$

Similarly, we could show that relation (3.42) is true for $\alpha = 2$. Since application F_K is affine along the sides $\hat{a}_3 \hat{a}_\alpha, \alpha = 1, 2$, the traces $\frac{\partial w}{\partial n_{3\alpha}}|_{[a_3, a_\alpha]}$ are one-variable P_4 (resp. BELL : P_3) polynomials entirely determined by the degrees of freedom relative to the sides $a_3 a_\alpha, \alpha = 1, 2$. Thus, the function $\frac{\partial w}{\partial n_{3\alpha}}|_{[a_3, a_\alpha]}$ realizes the second part of condition (3.2).

By construction, \hat{w} is a polynomial function. Hence, condition (3.4) is verified by $w = \hat{w} \circ F_K^{-1}$. Finally, the definition of Σ_K (see (3.11) (resp. BELL : (3.12))) insures condition (3.1). \blacksquare

Thus, to the set $\Sigma_K(v)$ of values of degrees of freedom of a function v , the previous construction associates a suitable interpolate function w that we denote $\pi_K v$ in the sequel. It remains to study the interpolation error : this is the object of paragraph 4.

3.3 Definition of curved finite elements \mathcal{C}^1 -compatible with ARGYRIS or BELL triangles when $F_K \in (P_3)^2$

The construction of such elements follows the same lines than in the case $F_K \in (P_5)^2$. Thus we just indicate the main changes.


Now, the basic finite element is described in Figure 3.7 (see also [36]) and by similarity with Theorem 3.1, we get

Theorem 3.3 : In Figure 3.7, the triple $(\hat{K}, \hat{P}, \hat{\Sigma})$ defines a finite element. ■

Construction of the interpolate function $v \rightarrow \pi_K v$

By using the basic finite element, we associate to any regular function v its interpolate $\pi_K v$. First, with notations of Fig. 3.4 and 3.8 we define the set $\Sigma_K(v)$ of values of degrees of freedom of the curved element. Next, from the 24 (resp. BELL : 21) - elements of the set $\Sigma_K(v)$, we associate a set $\Delta_K(v)$ with 36-elements from which we define a suitable interpolating function $\hat{w} \in \hat{P}$:

$$(3.46) \quad \left\{ \begin{array}{l} \Sigma_K(v) = \{(D^\alpha v(a_i), |\alpha| = 0, 1, 2), i = 1, 2, 3 ; Dv(b_2)n_{31} ; \\ Dv(b_1)n_{32} ; Dv(b_3)DF_K(\hat{b}_3)(\hat{a}_3 - \hat{b}_3) ; v(e_i), i = 1, 2, 3\} ; \end{array} \right.$$



\hat{K} = unit right-angled triangle

$\hat{P} = P_7$; $\dim \hat{P} = 36$;

$$\hat{\Sigma}(\hat{w}) = \left\{ \begin{array}{l} \hat{w}(\hat{a}_i), \frac{\partial \hat{w}}{\partial \hat{x}_1}(\hat{a}_i), \frac{\partial \hat{w}}{\partial \hat{x}_2}(\hat{a}_i), \frac{\partial^2 \hat{w}}{\partial \hat{x}_1^2}(\hat{a}_i), \frac{\partial^2 \hat{w}}{\partial \hat{x}_1 \partial \hat{x}_2}(\hat{a}_i), \frac{\partial^2 \hat{w}}{\partial \hat{x}_2^2}(\hat{a}_i), i = 1, 2, 3 ; \\ -\frac{\partial \hat{w}}{\partial \hat{x}_1}(\hat{b}_1) ; -\frac{\partial \hat{w}}{\partial \hat{x}_2}(\hat{b}_2) ; \frac{\sqrt{2}}{2} \left(\frac{\partial \hat{w}}{\partial \hat{x}_1} + \frac{\partial \hat{w}}{\partial \hat{x}_2} \right) (\hat{b}_3) ; \hat{w}(\hat{d}_i), i = 1, \dots, 6 ; \hat{w}(\hat{e}_i), i = 1, 2, 3 ; \\ -\frac{\partial \hat{w}}{\partial \hat{x}_1}(\hat{d}_i), i = 1, 2 ; -\frac{\partial \hat{w}}{\partial \hat{x}_2}(\hat{d}_i), i = 3, 4 ; \frac{\sqrt{2}}{2} \left(\frac{\partial \hat{w}}{\partial \hat{x}_1} + \frac{\partial \hat{w}}{\partial \hat{x}_2} \right) (\hat{d}_i), i = 5, 6 \end{array} \right\}$$

Figure 3.7 : "Basic" P_7 -Hermite finite-element

$$(3.47) \quad \left\{ \begin{array}{l} \text{(resp. BELL :} \\ \Sigma_K(v) = \{(D^\alpha v(a_i), |\alpha| = 0, 1, 2), i = 1, 2, 3 ; v(e_i), i = 1, 2, 3\}. \end{array} \right.$$

Finally we set

$$(3.48) \quad w = \hat{w} \circ F_K^{-1}$$

and by similarity with Theorem 3.2, we prove in the next theorem that w satisfies the desirable properties.

Theorem 3.4 : The function w , defined in (3.48), is determined in a unique way by the data of the set $\Sigma_K(v)$ of the values of the degrees of freedom of the function v , i.e.,

$$(3.49) \quad \left\{ \begin{array}{l} \left\{ \begin{array}{l} \Sigma_K(v) = \{(D^\alpha v(a_i), |\alpha| = 0, 1, 2), i = 1, 2, 3 ; Dv(b_2)n_{31} ; \\ Dv(b_1)n_{32} ; Dv(b_3)DF_K(\hat{b}_3)(\hat{a}_3 - \hat{b}_3) ; v(e_i), i = 1, 2, 3\} ; \\ \text{(resp. BELL :} \\ \Sigma_K(v) = \{(D^\alpha v(a_i), |\alpha| = 0, 1, 2), i = 1, 2, 3 ; v(e_i), i = 1, 2, 3\} \end{array} \right. \end{array} \right.$$




Figure 3.8 : Sets Σ_K for \mathcal{C}^1 -compatible curved finite elements ($F_K \in (P_3)^2$).

Moreover the function w verifies the relations

$$(3.50) \quad D^\alpha w(a_i) = D^\alpha v(a_i), \quad |\alpha| = 0, 1, 2 ; i = 1, 2, 3 ;$$

$$(3.51) \quad Dw(b_2)n_{31} = Dv(b_2)n_{31},$$

$$(3.52) \quad Dw(b_1)n_{32} = Dv(b_1)n_{32},$$

$$(3.53) \quad Dw(b_3)DF_K(\hat{b}_3)(\hat{a}_3 - \hat{b}_3) = Dv(b_3)DF_K(\hat{b}_3)(\hat{a}_3 - \hat{b}_3)$$

$$(3.54) \quad w(e_i) = v(e_i), \quad i = 1, 2, 3,$$

(resp. *BELL* : w verifies relations (3.50) and (3.54)) and w satisfies conditions (3.1) (3.2) and (3.4)) so that we have $w = \pi_K v$. ■

3.4 Definition of curved finite elements \mathcal{C}^0 -compatible with the Hermite triangle of type (3) when $F_K \in (P_n)^2, n = 3$ or 5

These elements will be used in the approximation of tangential components of the displacement for thin shell problems. They are defined by

$$(3.55) \quad \left\{ \begin{array}{l} K = F_K(\hat{K}), \quad F_K \in (P_n)^2, \quad n = 3 \text{ (see Example 2.2)} \\ \quad \text{or } n = 5 \text{ (see Example 2.3)} ; \\ P_K = \{p : K \longrightarrow \mathbb{R} ; p = \hat{p} \circ F_K^{-1} ; \hat{p} \in P_3\}, \\ \Sigma_K = \{D^\alpha p(a_i), \quad |\alpha| \leq 1, \quad i = 1, 2, 3 ; p(F_K(\hat{a}_0)), \\ \quad \hat{a}_0 = \text{barycenter of } \hat{K}\} \end{array} \right.$$

Thus, to any regular function v (for instance $v \in \mathcal{C}^1(K)$) we associate the interpolating function $\pi_K v$ defined by

$$(3.56) \quad \pi_K v = \hat{\pi} \hat{v} \circ F_K^{-1}, \quad \hat{v} = v \circ F_K,$$

where $\hat{\pi} \hat{v}$ is the P_3 -Hermite interpolating function of \hat{v} over the reference triangle \hat{K} .

Remark 3.2 : When $F_K \in (P_3)^2$, the curved finite element defined in (3.56) is similar, but not the same, than the isoparametric Hermite triangle of type (3) studied by [37, Example 6]. Indeed, our definition of application F_K is different. ■

4 ESTIMATES OF THE INTERPOLATION ERRORS

In this paragraph, we give estimates of the interpolation error for the curved finite elements of class \mathcal{C}^1 constructed in the previous paragraph. It is worth to note that

(i) these estimates have the same order of those obtained for the associate straight finite elements ;

(ii) these estimates use the interpolate triangle K instead of the "original" triangle K_c (see Figure 2.1). The "geometrical" approximation resulting of the substitution of triangle K_c by triangle K will be analyzed in Part 2.

4.1 Estimates of the interpolation error for curved finite elements \mathcal{C}^1 -compatible with ARGYRIS or BELL triangle when $F_K \in (P_5)^2$

We are going to extend to the associate curved elements the interpolation error estimates obtained for ARGYRIS (resp. BELL) triangle in [27, chapter 6].

Theorem 4.1 : *There exists a constant c , independent of h_K , such that for all curved finite elements \mathcal{C}^1 -compatible with ARGYRIS (resp. BELL) triangles we have :*

$$(4.1) \quad \begin{cases} |v - \pi_K v|_{m,K} \leq ch_K^{k+1-m} \|v\|_{k+1,K}, \forall v \in H^{k+1}(K), \\ k = 3, \dots, 5; 0 \leq m \leq k+1, \end{cases}$$

$$(4.2) \quad \begin{cases} (\text{resp. BELL : } |v - \pi_K v|_{m,K} \leq ch_K^{k+1-m} \|v\|_{k+1,K}, \forall v \in H^{k+1}(K), \\ k = 3, 4; 0 \leq m \leq k+1), \end{cases}$$

where $\pi_K v$ is the interpolate function of v defined in section 3.2.

Proof : We prove estimates (4.1) (resp. BELL : (4.2)) in the most significant cases, i.e., $k = 5$ (resp. BELL : $k = 4$). The other cases can be obtained similarly. Let \hat{v} and $\widehat{\pi_K v}$ corresponding functions over the reference triangle \hat{K} , i.e.

$$(4.3) \quad \hat{v} = v \circ F_K; \widehat{\pi_K v} = \pi_K v \circ F_K.$$

The assumption $v \in H^6(K)$ involves $\hat{v} \in H^6(\hat{K})$. Moreover, note that function $\widehat{\pi_K v}$ (resp. $\pi_K v$) is denoted \hat{w} (resp. w) in section 3.2.

According to [37, (3.10)], there exists a constant c , independent of h_K such that

$$(4.4) \quad \begin{cases} \forall v \in W^{k,p}(K), |v|_{k,p,K} \\ \leq c |J_{F_K}|_{0,\infty,\hat{K}}^{1/p} \sum_{j=1}^k |\hat{v}|_{j,p,\hat{K}} \sum_{i \in I(j,k)} \left(|F_K^{-1}|_{1,\infty,K}^{i_1} |F_K^{-1}|_{2,\infty,K}^{i_2} \dots |F_K^{-1}|_{k,\infty,K}^{i_k} \right) \end{cases}$$

where $|F_K^{-1}|_{m,\infty,K}$, $1 \leq m \leq k$, is defined by relation (2.16) and where

$$(4.5) \quad \begin{cases} I(j, k) = \{i = (i_1, \dots, i_k) \in N^k; i_1 + i_2 + \dots + i_k = j, \\ i_1 + 2i_2 + \dots + ki_k = k\}, 1 \leq j \leq k. \end{cases}$$

Thus, $\forall v \in H^m(K)$, we have for $1 \leq m \leq 6$ (resp. BELL : $1 \leq m \leq 5$)

$$(4.6) \quad \begin{cases} |v - \pi_K v|_{m,K} \leq \\ c |J_{F_K}|_{0,\infty,\hat{K}}^{1/2} \sum_{j=1}^m |\hat{v} - \widehat{\pi_K v}|_{j,\hat{K}} \sum_{i \in I(j,m)} (|F_K^{-1}|_{1,\infty,K}^{i_1} \cdots |F_K^{-1}|_{m,\infty,K}^{i_m}) \end{cases}$$

We prove below the existence of a constant c such that for any $v \in H^6(K)$ (resp. BELL : $v \in H^5(K)$) :

$$(4.7) \quad |\hat{v} - \widehat{\pi_K v}|_{j,\hat{K}} \leq ch_K^5 \|v\|_{6,K}, \quad 0 \leq j \leq 6,$$

$$(4.8) \quad (\text{resp. BELL : } |\hat{v} - \widehat{\pi_K v}|_{j,\hat{K}} \leq ch_K^4 \|v\|_{5,K}, \quad 0 \leq j \leq 5),$$

where $v = \hat{v} \circ F_K^{-1}$.

Moreover, from Theorem 2.1,

$$(4.9) \quad \begin{cases} |F_K|_{\ell,\infty,\hat{K}} \leq ch_K^\ell, \quad \ell = 0, 1, \dots; \quad |F_K^{-1}|_{\ell,\infty,K} \leq ch_K^{-\ell}, \quad \ell = 1, 2, \dots \\ |J_{F_K}|_{0,\infty,\hat{K}} \leq ch_K^2; \quad |J_{F_K^{-1}}|_{0,\infty,K} \leq ch_K^{-2}. \end{cases}$$

Then, by combining estimates (4.7) (resp. BELL : (4.8)) and (4.9) with inequality (4.6), we obtain the result :

$$(4.10) \quad \begin{cases} |v - \pi_K v|_{m,K} \leq ch_K^{6-m} \|v\|_{6,K}, \quad \forall v \in H^6(K), \quad 1 \leq m \leq 6, \\ (\text{resp. BELL : } |v - \pi_K v|_{m,K} \leq ch_K^{5-m} \|v\|_{5,K}, \quad \forall v \in H^5(K), \quad 1 \leq m \leq 5). \end{cases}$$

Note that estimates (4.10) are still valid for $m = 0$. It suffices to replace inequality (4.6) by the following

$$(4.11) \quad |v - \pi_K v|_{0,K} \leq |J_{F_K}|_{0,\infty,\hat{K}}^{1/2} |\hat{v} - \widehat{\pi_K v}|_{0,\hat{K}}.$$

To complete the proof, it remains to obtain estimates (4.7) (resp. BELL : (4.8)).

Let $\hat{\pi}$ be the P_9 -interpolation operator associated to the basic finite element of Figure 3.5. Then, for any $0 \leq j \leq 6$ (resp. BELL : $0 \leq j \leq 5$), we have

$$(4.12) \quad |\hat{v} - \widehat{\pi_K v}|_{j,\hat{K}} \leq |\hat{v} - \hat{\pi} \hat{v}|_{j,\hat{K}} + |\hat{\pi} \hat{v} - \widehat{\pi_K v}|_{j,\hat{K}}.$$

In particular, polynomials of degree 5 (resp. BELL : 4) are invariant by $\hat{\pi}$; hence

$$(4.13) \quad |\hat{v} - \hat{\pi} \hat{v}|_{j,\hat{K}} \leq c |\hat{v}|_{6,\hat{K}}, \quad 0 \leq j \leq 6,$$

$$(4.14) \quad (\text{resp. BELL : } |\hat{v} - \hat{\pi}\hat{v}|_{j,\hat{K}} \leq c|\hat{v}|_{5,\hat{K}}, \quad 0 \leq j \leq 5).$$

Similarly to relation (4.4), we obtain

$$(4.15) \quad \left\{ \begin{array}{l} |\hat{v}|_{k,p,\hat{K}} \leq c|J_{F_K^{-1}}|_{0,\infty,K}^{1/p} \sum_{j=1}^k |v|_{j,p,K} \sum_{i \in I(j,k)} \left(|F_K|_{1,\infty,\hat{K}}^{i_1} |F_K|_{2,\infty,\hat{K}}^{i_2} \cdots |F_K|_{k,\infty,\hat{K}}^{i_k} \right), \\ \forall \hat{v} \in W^{k,p}(\hat{K}). \end{array} \right.$$

From inequalities (4.9) and (4.15), we have

$$(4.16) \quad |\hat{v}|_{6,\hat{K}} \leq ch_K^5 \|v\|_{6,K}, \quad \forall \hat{v} \in H^6(\hat{K}),$$

$$(4.17) \quad (\text{resp. BELL : } |\hat{v}|_{5,\hat{K}} \leq ch_K^4 \|v\|_{5,K}), \quad \forall \hat{v} \in H^5(\hat{K}),$$

so that, with (4.13) (4.16) (resp. BELL : (4.14) (4.17)) :

$$(4.18) \quad |\hat{v} - \hat{\pi}\hat{v}|_{j,\hat{K}} \leq ch_K^5 \|v\|_{6,K}, \quad \forall \hat{v} \in H^6(\hat{K}), \quad 0 \leq j \leq 6,$$

$$(4.19) \quad (\text{resp. BELL : } |\hat{v} - \hat{\pi}\hat{v}|_{j,\hat{K}} \leq ch_K^4 \|v\|_{5,K}, \quad \forall \hat{v} \in H^5(\hat{K}), \quad 0 \leq j \leq 5).$$

Subsequently, we show (in 3 steps) that $\forall \hat{v} \in H^6(\hat{K})$ (resp. BELL : $\forall \hat{v} \in H^5(\hat{K})$), we have

$$(4.20) \quad |\hat{\pi}\hat{v} - \widehat{\pi_K v}|_{j,\hat{K}} \leq ch_K^5 \|v\|_{6,K}, \quad 0 \leq j \leq 6,$$

$$(4.21) \quad (\text{resp. BELL : } |\hat{\pi}\hat{v} - \widehat{\pi_K v}|_{j,\hat{K}} \leq ch_K^4 \|v\|_{5,K}, \quad 0 \leq j \leq 5).$$

Step 1 : Method of estimation :

The difference

$$(4.22) \quad \hat{\delta}(\hat{x}) = \hat{\pi}\hat{v}(\hat{x}) - \widehat{\pi_K v}(\hat{x})$$

is a polynomial of degree 9, hence $\hat{\pi}\hat{\delta} = \hat{\delta}$. Let $\widehat{DL}_i, \hat{p}_i, 1 \leq i \leq 55$, be the degrees of freedom and the corresponding basis polynomials of the basic element of Fig. 3.5, arranged according to (3.13). For any $\hat{x} \in \hat{K}$, we have

$$(4.23) \quad \hat{\pi}\hat{\delta}(\hat{x}) = \hat{\delta}(\hat{x}) = \sum_{i=1}^{55} \widehat{DL}_i(\hat{\delta}) \hat{p}_i(\hat{x}).$$

The sets of values of degrees of freedom used to construct $\widehat{\pi_K v} = \hat{w}$ are given by $\hat{\Delta}_{K1}(v) \cup \hat{\Delta}_{K2}(v) \cup \hat{\Delta}_{K3}(v)$ (see (3.15) (3.27) and (3.34)). Comparing with (3.13), we observe that $\sum_1(\hat{v}) \equiv \hat{\Delta}_{K1}(v)$ so that

$$(4.24) \quad \widehat{DL}_i(\hat{\delta}) = 0, \quad i = 1, \dots, 28.$$

Step 2 : Estimates of $\widehat{DL}_i(\hat{\delta}), i = 29, \dots, 46 :$

Next, consider $\widehat{\Sigma}_2(\hat{v})$ and $\widehat{\Delta}_{K_2}(v)$. We have from (3.13) and (3.27) :

$$(4.25) \quad \widehat{DL}_{28+i}(\hat{\delta}) = \hat{v}(\hat{d}_i) - \hat{f}_2(\hat{d}_i), \quad i = 1, \dots, 4;$$

$$(4.26) \quad \widehat{DL}_{28+i}(\hat{\delta}) = \hat{v}(\hat{d}_i) - \hat{f}_1(\hat{d}_i), \quad i = 5, \dots, 8;$$

$$(4.27) \quad \left\{ \begin{array}{l} \widehat{DL}_{36+i}(\hat{\delta}) = -\frac{\partial \hat{v}}{\partial \hat{x}_1}(\hat{q}_i) + \frac{1}{|a_3 a_2|} < \frac{\partial F_K}{\partial \hat{x}_1}(\hat{q}_i), \quad t_{32} > \frac{d\hat{f}_2}{d\hat{x}_2}(\hat{q}_i) \\ + < \frac{\partial F_K}{\partial \hat{x}_1}(\hat{q}_i), \quad n_{32} > \hat{g}_2(\hat{q}_i), \quad i = 1, \dots, 5, \end{array} \right.$$

where for convenience \hat{q}_i means \hat{b}_1 for $i = 1$ and \hat{d}_{i-1} for $i = 2, \dots, 5$;

$$(4.28) \quad \left\{ \begin{array}{l} \widehat{DL}_{41+i}(\hat{\delta}) = -\frac{\partial \hat{v}}{\partial \hat{x}_2}(\hat{q}_i) + \frac{1}{|a_3 a_1|} < \frac{\partial F_K}{\partial \hat{x}_2}(\hat{q}_i), \quad t_{31} > \frac{d\hat{f}_1}{d\hat{x}_2}(\hat{q}_i) \\ + < \frac{\partial F_K}{\partial \hat{x}_2}(\hat{q}_i), \quad n_{31} > \hat{g}_1(\hat{q}_i), \quad i = 1, \dots, 5, \end{array} \right.$$

where for convenience \hat{q}_i means \hat{b}_2 for $i = 1$ and \hat{d}_{i+3} for $i = 2, \dots, 5$;

(resp. BELL : replace \hat{g}_1 and \hat{g}_2 by \hat{h}_1 and \hat{h}_2 , respectively).

Relations (3.16) to (3.18) show that \hat{f}_1 is the P_5 -Hermite polynomial which interpolates \hat{v} over the side $\hat{a}_3\hat{a}_1$, from the given degrees of freedom

$$(4.29) \quad \left\{ \begin{array}{l} \{\hat{v}(\hat{a}_1), \hat{v}(\hat{a}_3), D\hat{v}(\hat{a}_1)(\hat{a}_3 - \hat{a}_1), D\hat{v}(\hat{a}_3)(\hat{a}_1 - \hat{a}_3), \\ D^2\hat{v}(\hat{a}_1)(\hat{a}_3 - \hat{a}_1)^2, D^2\hat{v}(\hat{a}_3)(\hat{a}_1 - \hat{a}_3)^2\}. \end{array} \right.$$

In the sequel, we denote by the same letter the functions defined on a triangle K and their traces along the boundary ∂K . From the definition of function \hat{f}_1 , we get for any $\hat{v} \in W^{4,\infty}(\hat{a}_3\hat{a}_1)$

$$|\hat{v}(\hat{d}_i) - \hat{f}_1(\hat{d}_i)| \leq |\hat{v} - \hat{f}_1|_{0,\infty,\hat{a}_3\hat{a}_1} \leq c|\hat{v}|_{4,\infty,\hat{a}_3\hat{a}_1}, \quad i = 5, \dots, 8$$

and with Sobolev's theorem (see [29])

$$|\hat{v}(\hat{d}_i) - \hat{f}_1(\hat{d}_i)| \leq c\|\hat{v}\|_{4,\infty,\hat{K}} \leq c\|\hat{v}\|_{6,\hat{K}}, \quad \forall \hat{v} \in H^6(\hat{K}), \quad i = 5, \dots, 8.$$

Thus, the linear form

$$\mathcal{F} : \forall \hat{v} \in H^6(\hat{K}) \longrightarrow \mathcal{F}(\hat{v}) = \max_{i=5, \dots, 8} |\hat{v}(\hat{d}_i) - \hat{f}_1(\hat{d}_i)|$$

is continuous. Since it vanishes for any $\hat{v} \in P_5$, the lemma of Bramble-Hilbert [38] implies the existence of a constant c such that

$$|\hat{v}(\hat{d}_i) - \hat{f}_1(\hat{d}_i)| \leq c|\hat{v}|_{6, \hat{K}}, \quad \forall \hat{v} \in H^6(\hat{K}), \quad i = 5, \dots, 8,$$

or, with relation (4.16)

$$(4.30) \quad |\widehat{DL}_{28+i}(\hat{\delta})| = |\hat{v}(\hat{d}_i) - \hat{f}_1(\hat{d}_i)| \leq ch_K^5 \|v\|_{6, K}, \quad \forall \hat{v} \in H^6(\hat{K}), \quad i = 5, \dots, 8.$$

Similarly, we can prove that

$$(4.31) \quad \left\{ \begin{array}{l} \text{(resp. BELL :} \\ |\widehat{DL}_{28+i}(\hat{\delta})| = |\hat{v}(\hat{d}_i) - \hat{f}_1(\hat{d}_i)| \leq ch_K^4 \|v\|_{5, K}, \quad \forall \hat{v} \in H^5(\hat{K}), \quad i = 5, \dots, 8, \end{array} \right.$$

$$(4.32) \quad |\widehat{DL}_{28+i}(\hat{\delta})| = |\hat{v}(\hat{d}_i) - \hat{f}_2(\hat{d}_i)| \leq ch_K^5 \|v\|_{6, K}, \quad \forall \hat{v} \in H^6(\hat{K}), \quad i = 1, \dots, 4,$$

$$(4.33) \quad \left\{ \begin{array}{l} \text{(resp. BELL :} \\ |\widehat{DL}_{28+i}(\hat{\delta})| = |\hat{v}(\hat{d}_i) - \hat{f}_2(\hat{d}_i)| \leq ch_K^4 \|v\|_{5, K}, \quad \forall \hat{v} \in H^5(\hat{K}), \quad i = 1, \dots, 4. \end{array} \right.$$

Now, let us examine the case of degrees of freedom (4.28). The correspondences $\hat{v} = v \circ F_K$ and $q_i = F_K(\hat{q}_i)$ imply

$$\left\{ \begin{array}{l} \frac{\partial \hat{v}}{\partial \hat{x}_2}(\hat{q}_i) = D\hat{v}(\hat{q}_i)\vec{e}_2 = Dv(q_i)DF_K(\hat{q}_i)\vec{e}_2 = Dv(q_i)\frac{\partial F_K}{\partial \hat{x}_2}(\hat{q}_i) \\ = Dv(q_i) \left[\left\langle \frac{\partial F_K}{\partial \hat{x}_2}(\hat{q}_i), t_{31} \right\rangle t_{31} + \left\langle \frac{\partial F_K}{\partial \hat{x}_2}(\hat{q}_i), n_{31} \right\rangle n_{31} \right], \\ = \frac{1}{|a_3 a_1|} \left\langle \frac{\partial F_K}{\partial \hat{x}_2}(\hat{q}_i), t_{31} \right\rangle \frac{\partial \hat{v}}{\partial \hat{x}_1}(\hat{q}_i) + \left\langle \frac{\partial F_K}{\partial \hat{x}_2}(\hat{q}_i), n_{31} \right\rangle Dv(q_i)n_{31}, \end{array} \right.$$

and hence,

$$(4.34) \quad \left\{ \begin{array}{l} \widehat{DL}_{41+i}(\hat{\delta}) = \frac{1}{|a_3 a_1|} \left\langle \frac{\partial F_K}{\partial \hat{x}_2}(\hat{q}_i), t_{31} \right\rangle \left[\frac{d\hat{f}_1}{d\hat{x}_1}(\hat{q}_i) - \frac{\partial \hat{v}}{\partial \hat{x}_1}(\hat{q}_i) \right] \\ + \left\langle \frac{\partial F_K}{\partial \hat{x}_2}(\hat{q}_i), n_{31} \right\rangle (\hat{g}_1(\hat{q}_i) - Dv(q_i)n_{31}), \quad i = 1, \dots, 5. \end{array} \right.$$

A proof similar to that of estimate (4.32) shows that

$$(4.35) \quad \left| \frac{1}{|a_3 a_1|} < \frac{\partial F_K}{\partial \hat{x}_2}(\hat{q}_i), t_{31} > \left[\frac{d\hat{f}_1}{d\hat{x}_1}(\hat{q}_i) - \frac{\partial \hat{v}}{\partial \hat{x}_1}(\hat{q}_i) \right] \right| \leq ch_K^5 \|v\|_{6,K}, \quad \forall \hat{v} \in H^6(\hat{K}).$$

Now, let us consider the last term of relation (4.34). Firstly according to the proof of Theorem 3.2, we have since \hat{q}_1 means \hat{b}_2

$$(4.36) \quad \hat{g}_1(\hat{q}_1) - Dv(q_1)n_{31} = \hat{g}_1(\hat{b}_2) - Dv(b_2)n_{31} = 0.$$

The next term is $\hat{g}_1(\hat{q}_2) - Dv(q_2)n_{31} = \hat{g}_1(\hat{d}_5) - Dv(d_5)n_{31}$ where, according to (3.17) (3.19) and (3.21), the function $g_1 = \hat{g}_1 \circ F_K^{-1}$ is the one-dimensional P_4 -interpolate of the function $z(\cdot) = Dv(\cdot)n_{31}$ along the side $a_3 a_1$ by using the set of degrees of freedom :

$$\{z(a_i), Dz(a_i)t_{31}, i = 1, 3 ; z(b_2)\}.$$

Then the interpolation properties and Sobolev's theorem imply :

$$\begin{cases} |z(d_5) - g_1(d_5)| \leq |z - g_1|_{0,\infty,a_3 a_1} = |\hat{z} - \hat{g}_1|_{0,\infty,\hat{a}_3 \hat{a}_1} \leq c|\hat{z}|_{3,\infty,\hat{a}_3 \hat{a}_1} \\ \leq c\|\hat{z}\|_{3,\infty,\hat{K}} \leq c\|\hat{z}\|_{5,\hat{K}}, \quad \forall \hat{z} = z \circ F_K \in H^5(\hat{K}). \end{cases}$$

Thus, the linear form

$$\mathcal{G} : \forall \hat{z} \in H^5(K) \longmapsto |\hat{z}(\hat{d}_5) - \hat{g}_1(\hat{d}_5)| \text{ is continuous.}$$

Since it vanishes for any $\hat{z} \in P_4$, the Bramble-Hilbert lemma implies the existence of a constant c such that

$$(4.37) \quad |\hat{z}(\hat{d}_5) - \hat{g}_1(\hat{d}_5)| = |z(d_5) - g_1(d_5)| \leq c|\hat{z}|_{5,\hat{K}}, \quad \forall \hat{z} \in H^5(\hat{K}).$$

By analogy with (4.15) and (4.16) we obtain

$$(4.38) \quad |\hat{z}|_{5,\hat{K}} \leq Ch_K^4 \|z\|_{5,K}, \quad \forall \hat{z} \in H^5(\hat{K})$$

and since $z(\cdot) = Dv(\cdot)n_{31}$, we get

$$(4.39) \quad \|z\|_{5,K} \leq C\|v\|_{6,K},$$

so that relations (4.37) to (4.39) imply

$$(4.40) \quad |\hat{g}_1(\hat{d}_5) - Dv(d_5)n_{31}| \leq Ch_K^4 \|v\|_{6,K}, \quad \forall v \in H^6(K),$$

and three other relations at points \hat{d}_6 , \hat{d}_7 and \hat{d}_8 . Then, relations (2.17) (4.34) (4.35) (4.36) and (4.40) imply

$$(4.41) \quad |\widehat{DL}_{41+i}(\hat{\delta})| \leq Ch_K^5 \|v\|_{6,K}, \quad \forall v \in H^6(K), \quad i = 1, \dots, 5.$$

By the same way, we could get :

$$(4.42) \quad (\text{resp. BELL : } |\widehat{DL}_{41+i}(\hat{\delta})| \leq ch_K^4 \|v\|_{5,K}, \quad \forall v \in H^5(K), \quad i = 1, \dots, 5 ;$$

$$(4.43) \quad |\widehat{DL}_{36+i}(\hat{\delta})| \leq ch_K^5 \|v\|_{6,K}, \quad \forall v \in H^6(K), \quad i = 1, \dots, 5 ;$$

$$(4.44) \quad (\text{resp. BELL : } |\widehat{DL}_{36+i}(\hat{\delta})| \leq ch_K^4 \|v\|_{5,K}, \quad \forall v \in H^5(K), \quad i = 1, \dots, 5.$$

Step 3 : Estimate of $\widehat{DL}_i(\hat{\delta}), i = 47, \dots, 55$

From definitions (3.13) and (3.34) we obtain

$$(4.45) \quad \widehat{DL}_{46+i}(\hat{\delta}) = \hat{v}(\hat{d}_{8+i}) - \hat{f}_3(\hat{d}_{8+i}), \quad i = 1, \dots, 4 ;$$

$$(4.46) \quad \begin{cases} \widehat{DL}_{51}(\hat{\delta}) = \frac{\sqrt{2}}{2} \left[\frac{\partial \hat{v}}{\partial \hat{x}_1} + \frac{\partial \hat{v}}{\partial \hat{x}_2} \right] (\hat{b}_3) + \sqrt{2} \hat{g}_3(\hat{b}_3), \\ \widehat{DL}_{51+i}(\hat{\delta}) = \frac{\sqrt{2}}{2} \left[\frac{\partial \hat{v}}{\partial \hat{x}_1} + \frac{\partial \hat{v}}{\partial \hat{x}_2} \right] (\hat{d}_{8+i}) + \sqrt{2} \hat{g}_3(\hat{d}_{8+i}), \quad i = 1, \dots, 4. \end{cases}$$

By using the definition (3.31) of (\hat{f}_3) , a proof similar to that of (4.30) shows that

$$(4.47) \quad |\widehat{DL}_{46+i}(\hat{\delta})| = |\hat{v}(\hat{d}_{8+i}) - \hat{f}_3(\hat{d}_{8+i})| \leq ch_K^5 \|v\|_{6,K}, \quad \forall \hat{v} \in H^6(\hat{K}), \quad i = 1, \dots, 4,$$

$$(4.48) \quad \begin{cases} (\text{resp. BELL :} \\ |\widehat{DL}_{46+i}(\hat{\delta})| = |\hat{v}(\hat{d}_{8+i}) - \hat{f}_3(\hat{d}_{8+i})| \leq ch_K^4 \|v\|_{5,K}, \quad \forall \hat{v} \in H^5(\hat{K}), \quad i = 1, \dots, 4). \end{cases}$$

Now, consider the first term (4.46), i.e., $\widehat{DL}_{51}(\hat{\delta}) = -\sqrt{2}[\hat{g}_3(\hat{b}_3) - D\hat{v}(\hat{b}_3)(\hat{a}_3 - \hat{b}_3)]$. With definition (3.32) of \hat{g}_3 , Sobolev's theorem and Bramble-Hilbert lemma, we get

$$\begin{cases} |\widehat{DL}_{51}(\hat{\delta})| \leq \sqrt{2} |\hat{g}_3(\cdot) - D\hat{v}(\cdot)(\hat{a}_3 - \hat{b}_3)|_{0,\infty,\hat{a}_1\hat{a}_2} \\ \leq c |D\hat{v}(\cdot)(\hat{a}_3 - \hat{b}_3)|_{3,\infty,\hat{a}_1\hat{a}_2} \\ \leq c |\hat{v}|_{4,\infty,\hat{a}_1\hat{a}_2} \leq c \|\hat{v}\|_{6,\hat{K}} \leq ch_K^5 \|v\|_{6,K}, \quad \forall \hat{v} \in H^6(\hat{K}). \end{cases}$$

And, by similarity, we finally obtain

$$(4.49) \quad |\widehat{DL}_{50+i}(\hat{\delta})| \leq ch_K^5 \|v\|_{6,K}, \quad \forall \hat{v} \in H^6(\hat{K}), \quad i = 1, \dots, 5 ;$$

$$(4.50) \quad (\text{resp. BELL : } |\widehat{DL}_{50+i}(\hat{\delta})| \leq ch_K^4 \|v\|_{5,K}, \quad \forall \hat{v} \in H^5(\hat{K}), \quad i = 1, \dots, 5).$$

Note that for a curved finite element \mathcal{C}^1 -compatible with an ARGYRIS triangle we have in fact $\widehat{DL}_{51}(\hat{\delta}) = 0$. This property is not generally true for a curved finite element \mathcal{C}^1 -compatible with a BELL triangle.

Finally, relations (4.22) (4.23) imply

$$|\hat{\delta}|_{j,\hat{K}} \leq |\hat{\pi}v - \widehat{\pi}_K v|_{j,\hat{K}} \leq c \sum_{i=1}^{55} |\widehat{DL}_i(\hat{\delta})|.$$

Then, estimates (4.24) (4.30) (4.32) (4.41) (4.43) (4.47) and (4.49) [resp. BELL : (4.24) (4.31) (4.33) (4.42) (4.44) (4.48) and (4.50)] give estimate (4.20) [resp. BELL : (4.21)]. We obtain estimates (4.7) [resp. BELL : (4.8)] by combining estimates (4.18) and (4.20) [resp. BELL : (4.19) and (4.21)]. ■

4.2 Estimates of the interpolation error for curved finite elements \mathcal{C}^1 -compatible with ARGYRIS or BELL triangles in case of $F_K \in (P_3)^2$

In this case, we can prove similar results than in Theorem 4.1 :

Theorem 4.2 : There exists a constant c , independent of h_K , such that for all curved finite elements \mathcal{C}^1 -compatible with ARGYRIS (resp. BELL) triangles, we have :

$$(4.51) \quad \begin{cases} |v - \pi_K v|_{m,K} \leq ch_K^{k+1-m} \|v\|_{k+1,K}, \quad \forall v \in H^{k+1}(K), \\ k = 3, \dots, 5, \quad 0 \leq m \leq k+1, \end{cases}$$

$$(4.52) \quad \begin{cases} (\text{resp. BELL : } |v - \pi_K v|_{m,K} \leq ch_K^{k+1-m} \|v\|_{k+1,K}, \quad \forall v \in H^{k+1}(K), \\ k = 3, 4, \quad 0 \leq m \leq k+1), \end{cases}$$

where $\pi_K v$ is the interpolating function of v defined in section 3.3. ■

4.3 Estimates of the interpolation error for curved finite elements \mathcal{C}^0 -compatible with Hermite triangle of type (3) ($F_K \in (P_n)^2$, $n = 3$ or 5)

For the curved finite elements introduced in section 3.4, it is easy to prove the theorem :

Theorem 4.3 : There exists a constant c , independent of h_K , such that for all curved finite elements \mathcal{C}^0 -compatible with Hermite triangle of type (3), we have

$$|v - \pi_K v|_{m,K} \leq ch_K^{4-m} \|v\|_{4,K}, \quad 0 \leq m \leq 4, \quad \forall v \in H^4(K),$$

where $\pi_K v$ is the interpolating function of v defined in section 3.4. ■

5 CONCLUDING REMARKS

To conclude it is worth to emphasize some very interesting properties of these interpolation methods on curved triangles :

i) they allow to construct C^1 -interpolating functions from just a set of degrees of freedom which is completely compatible with those of the associate straight finite elements. In this way they used a mapping $F_K : \hat{K} \rightarrow K$ which is entirely symmetric and polynomial ;

ii) corresponding functions on the reference triangle are of polynomial type so that the study of the effect of numerical integration is straightforward (see Part 2, i.e. [4]) ;

iii) for a given degree of regularity, the asymptotic interpolation error estimates have the same order for the straight and for the associate curved finite elements ;

iv) thus, these curved finite elements constitute a really powerful tool to approximate fourth order problems set on plane curved boundary domains.

Acknowledgements

The author is indebted to P.G. CIARLET for suggesting this study and for his constant interest during the investigation. Also we take this opportunity to thank I. BABUSKA who has incited us to implement such curved C^1 -finite elements and J.M. BOISSERIE with whom we have had a fruitful collaboration to realize the implementation and some numerical experiments (see [5]).

References

- [1] I. BABUSKA, J. PITKARANTA, The plate paradox for hard and soft simple support, SIAM J. Math. Anal., Vol. 21, n° 3, pp. 551-576, May 1990.
- [2] M. BERNADOU, J.M. BOISSERIE, The Finite Element Method in Thin Shell Theory : Application to Arch Dam Simulations, Birkhäuser, Boston, 1982.
- [3] M. BERNADOU, On the approximation of junctions between plates or between shells, Computational Mechanics, Cheung, Lee and Leung (Eds.), pp. 35-45, Balkema, Rotterdam, 1991.
- [4] M. BERNADOU, C^1 -curved finite elements with numerical integration for thin plate and thin shell problems. Part 2 : Approximation of thin plate and thin shell problems (to appear).
- [5] M. BERNADOU, J.M. BOISSERIE, Curved finite elements of class C^1 : implementation and numerical experiments (to appear).

- [6] J.H. ARGYRIS, I. FRIED, D.W. SCHARPF, The TUBA family of plate elements for the matrix displacement method, *The Aeronautical Journal of the Royal Aeronautical Society*, 72 (1968), pp.701-709.
- [7] K. BELL, A refined triangular plate bending element, *Internat. J. Num. Meth. Eng.*, 1 (1969), pp. 101-122.
- [8] M. BERNADOU, Convergence of conforming finite element methods for general shell problems, *Internat. J. Engng. Sci.*, 18 (1980), pp. 249-276.
- [9] M. ZLÁMAL, The finite element method in domains with curved boundaries, *Internat. J. Numer. Methods Engng.*, 5 (1973), pp. 367-373.
- [10] M. ZLÁMAL, Curved elements in the finite element method, Part I, *SIAM J. Numer. Anal.*, 10 (1973), pp. 229-240.
- [11] M. ZLÁMAL, Curved elements in the finite element method, Part II, *SIAM J. Numer. Anal.*, 11 (1974), pp. 347-362.
- [12] R. SCOTT, Interpolated boundary conditions in the finite element method, *SIAM J. Numer. Anal.* 12, n° 3 (1975), pp. 404-427.
- [13] R. SCOTT, A survey of displacement methods for the plate bending problem (to appear in *Proceedings of the U.S.-Germany Symposium on Formulations and Computational Algorithms in Finite Element Analysis*, Massachusetts Institute of Technology, Cambridge, August 9-13, 1976).
- [14] L.E. MANSFIELD, Approximation of the boundary in the finite element solution of fourth order problems, *SIAM J. Numer. Anal.*, 15 (1978), pp. 568-579.
- [15] L.E. MANSFIELD, A Clough-Tocher type element useful for fourth order problems over non-polygonal domains, *Math. Comp.* 32 (1978), pp. 135-142.
- [16] A. ŽENIŠEK, A general theorem on triangular finite $C^{(m)}$ -elements, *Rev. Française Automat. Informat. Recherche Opérationnelle*, R-2 (1974), pp. 119-127.
- [17] A. ŽENIŠEK, Curved triangles finite $C^{(m)}$ -elements, *Apl. Mat.*, 23 (1978), pp. 346-377.
- [18] M. LENOIR, Optimal isoparametric finite elements and error estimates for domains involving curved boundaries. *SIAM J. Numer. Anal.*, Vol. 23, n° 3, pp. 562-580, June 1986.
- [19] C. BERNARDI, Optimal finite-element interpolation on curved domains, *SIAM J. Numer. Anal.* 26, n° 5 (1989), pp. 1212-1240.
- [20] J. KRIZKOVA, Special exact curved finite elements, *Applications of Mathematics*, 36, n° 2 (1991), pp. 81-95.

- [21] V.G. KORNEEV, S.E. PONOMAREV, Use of curvilinear finite elements in schemes for solving linear elliptic equations of order $2n$, Part I ; Chislennyye metody mekhaniki sploshnoi sredy Novosibirsk, 5, n° 5 (1974), pp. 78-97 (english translation : UCRL-trans. 10852).
- [22] V.G. KORNEEV, S.E. PONOMAREV, Use of curvilinear finite elements in schemes for solving linear elliptic equations of order $2n$, Part II ; Chislennyye metody mekhaniki sploshnoi sredy Novosibirsk, 6, n° 1 (1975), pp. 47-64 (in Russian).
- [23] J.H. ARGYRIS, N. LOCHNER, On the application of the SHEBA shell element. Comput. Meth. Appl. Mech. Engng. 1 (1972), pp. 317-347.
- [24] M.W. CHERNUKA, G.R. COWPER, G.M. LINDBERG, M.D. OLSON, Finite element analysis of plates with curved edges, Internat. J. Numer. Methods Engng., 4 (1972), pp. 49-65.
- [25] J. DOUGLAS, T. DUPONT, P. PERCELL, R. SCOTT, A family of C^1 -finite elements with optimal approximation properties for various Galerkin methods for 2nd and 4th order problems, Rev. Française Automat. Informat. Recherche Opérationnelle, Analyse Numérique, 13 (1979), pp. 227-256.
- [26] T.M. HRUDEY, A finite element procedure for plates with curved boundaries, Comput. Struct., 7 (1977), pp. 553-564.
- [27] P.G. CIARLET, The Finite Element Method for Elliptic Problems, North-Holland, Amsterdam, 1978.
- [28] P.G. CIARLET, Basic error estimates for elliptic problems. In : Handbook of Numerical Analysis (P.G. Ciarlet and J.L. Lions Editors). Vol. II : Finite Element Methods, Part 1, pp. 17-351, North-Holland, Amsterdam, 1991.
- [29] R.A. ADAMS, Sobolev Spaces, Academic Press, New York, 1975.
- [30] J.L. LIONS, E. MAGENES, Problèmes aux Limites non Homogènes et Applications, Vol. 1, Dunod, Paris, 1968.
- [31] J. NEČAS, Les Méthodes Directes en Théorie des Equations Elliptiques, Masson, Paris, 1967.
- [32] M. BERNADOU, Sur l'Analyse Numérique du Modèle Linéaire de Coques Minces de W.T. KOITER. Thèse d'Etat, Université Pierre et Marie Curie, 1978.
- [33] G.H. MEISTERS, C. OLECH, Locally one-to-one mappings and a classical theorem on Schlicht functions, Duke Math. J., 30 (1963), pp. 63-80.
- [34] C. CARATHEODORY, Theory of Functions of a Complex Variable, Volume 1, Chelsea Publishing Company, New-York, 1964.

- [35] H. CARTAN, Calcul Différentiel, Hermann, Paris, 1967.
- [36] J.M. BOISSERIE, A new C^1 finite element : full heptic, Internat. J. Num. Meth. Eng., 28 (1989), pp. 667-677.
- [37] P.G. CIARLET, P.A. RAVIART, Interpolation theory over curved elements, with applications to finite element methods, Comput. Methods Appl. Mech. Engng., 1 (1972), pp. 217-249.
- [38] J.H. BRAMBLE, S.R. HILBERT, Estimation of linear functionals on Sobolev spaces with application to Fourier transforms and spline interpolation, SIAM J. Numer. Anal. 7, n° 1 (1970), pp. 112-124.

ISSN 0249 - 6399