



Pancyclic arcs and connectivity in tournaments

Frédéric Havet

► **To cite this version:**

Frédéric Havet. Pancyclic arcs and connectivity in tournaments. RR-4378, INRIA. 2002. inria-00072210

HAL Id: inria-00072210

<https://hal.inria.fr/inria-00072210>

Submitted on 23 May 2006

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Pancyclic arcs and connectivity in tournaments

Frédéric Havet

N° 4378

Février 2002

THÈME 1

 ***rapport
de recherche***

Pancyclic arcs and connectivity in tournaments

Frédéric Havet

Thème 1 — Réseaux et systèmes
Projet Mascotte

Rapport de recherche n° 4378 — Février 2002 — 19 pages

Abstract: A *tournament* is an orientation of the edges of a complete graph. An arc is *pancyclic* in a digraph D if it is contained in a cycle of length l , for every $3 \leq l \leq |D|$. In [4], Moon showed that every strong tournament contains at least three pancyclic arcs and characterized the tournaments with exactly three pancyclic arcs. All these tournaments are not 2-strong. In this paper, we are interested in the minimum number $p_k(n)$ of pancyclic arcs in a k -strong tournament of order n . We conjecture that (for $k \geq 2$) there exists a constant $\alpha_k > 0$ such that $p_k(n) \geq \alpha_k n$. After proving that every 2-strong tournament has a hamiltonian cycle containing at least five pancyclic arcs, we deduce that for $k \geq 2$, $p_k(n) \geq 2k + 3$. We then characterize the tournaments having exactly four pancyclic arcs and those having exactly five pancyclic arcs.

Key-words: pancyclic, tournament, strong connectivity, strongness

Arcs pancycliques et connexité dans les tournois

Résumé : Un *tournoi* est une orientation des arêtes du graphe complet. Un arc est *pancyclique* dans un graphe orienté D s'il est contenu dans un cycle de longueur l , pour toute longueur $3 \leq l \leq |D|$. Dans [4], Moon a montré que tout tournoi fortement connexe contenait au moins trois arcs pancycliques et il a caractérisé les tournois ayant exactement trois arcs pancycliques. Tous ces tournois ne sont pas 2-fortement connexes. Dans ce rapport, nous étudions le nombre minimum $p_k(n)$ d'arcs pancycliques dans un tournoi k -fortement connexe. Nous conjecturons que (pour $k \geq 2$), il existe une constante α_k telle que $f_k(n) \geq \alpha_k n$. Nous prouvons d'abord que tout tournoi 2-fortement connexe possède un circuit hamiltonien contenant au moins cinq arcs pancycliques. Nous en déduisons ensuite que pour $k \geq 2$, $p_k(n) \geq 2k + 3$. Enfin, nous caractérisons les tournois ayant exactement quatre arcs pancycliques et ceux ayant exactement cinq arcs pancycliques.

Mots-clés : pancyclique, tournoi, forte connexité

1 Introduction

A *tournament* is an orientation of the arcs of a complete graph. Paths and cycles are always directed. A l -cycle is a cycle of length l .

An arc or a vertex is *pancyclic* in a digraph D if, for every $3 \leq l \leq |D|$, it is contained in an l -cycle.

A tournament is *strong* (or *strongly connected*) if for any two vertices x and y there exists a path beginning in x and terminating at y . A nonstrong tournament is said to be *reducible*. A tournament is k -strong, if $T - Y$ is strong for any set Y of $k - 1$ vertices. A tournament is $(= k)$ -strong or *exactly k -strong*, if it is k -strong and not $(k + 1)$ -strong.

To contain a pancyclic arc or vertex, a tournament must contain a hamiltonian cycle. Therefore, it must be strong according to the well known theorem of Camion [2]: *A tournament has a hamiltonian cycle if and only if it is strong*. Moon [3] gave an alternative proof of Camion's theorem by proving that every vertex of a strong tournament is pancyclic.

Analogously, one may wonder whether there are pancyclic arcs in tournament and how many. Moon [4] showed that every strong tournament contains at least three pancyclic arcs. Actually, he proved a somewhat stronger result : indeed, instead of considering the number $p(T)$ of pancyclic arcs in the tournament T , he proved that $h(T)$ the maximum number of pancyclic arcs contained in some hamiltonian cycle of T is at least 3.

Theorem 1 (Moon, [4]) *Let T be a strong tournament with $n \geq 3$ vertices.*

$$h(T) \geq 3$$

with equality holding only if $T \in \mathcal{P}_3$.

A tournament is in \mathcal{P}_3 if there is a vertex v such that $T - v$ is the transitive tournament $TT[t_1, t_2, \dots, t_m]$ ((t_i, t_j) is an arc if and only if $i < j$), and an integers $1 < i_1 \leq m$ such that $v \rightarrow t_j$ if and only if $1 \leq j < i_1$.

Let $p_k(n)$ be minimum number $p_k(n)$ of pancyclic arcs in a k -strong tournament of order n and let $h_k(n) := \min\{h(T); T \text{ } k\text{-strong of order } n\}$.

Because the tournaments of \mathcal{P}_3 are $(= 1)$ -strong, we have $p_1(n) = h_1(n) = 3$ and if $k \geq 2$, $p_k(n) \geq h_k(n) \geq 4$. However we consider that this lower bound 4 is not tight.

In this paper, we show Section 3 sufficient conditions for an arc to be pancyclic in a tournament. Using these conditions, we give an easy alternative proof of Theorem 1. Moreover our method allow us to go further. In Section 4, we prove that for $k \geq 2$, $h_k(n) \geq 5$ and then deduce that $p_k(n) \geq 2k + 3$. Finally, we characterize the tournaments with exactly four pancyclic arcs and those with exactly five pancyclic arcs.

However, our lower bound for h_k and p_k seems to be still far from the exact value. We conjecture that for $k \geq 2$, $p_k(n)$ tends linearly to infinity :

Conjecture 1 For $k \geq 2$, there exists a constant $\alpha_k > 0$ such that $p_k(n) \geq \alpha_k n$.

We cannot expect to have more pancyclic arcs since there are k -strong tournaments having less than $2kn$ pancyclic arcs.

Proposition 1 $p_k(n) \leq 2kn - 2k^2 - k$

Proof. Let T_n be the k -strong tournament obtained from the rotative tournament on $2k + 1$ vertices by blowing up a vertex with a transitive tournament TT of order $n - 2k$. Every arc in TT is not pancyclic in T since it is contained in no 3-cycle. Thus $(n - 2k)(n - 2k - 1)/2$ arcs are not pancyclic. ■

Proposition 2

$$h_k(n) \leq 3k$$

Proof. If $n \leq 3k$, we have trivially the answer. Suppose now that $n > 3k$. Consider the k -strong tournament T_n obtained from two TT_k , A and B , and one TT_{n-2k} C such that $A \rightarrow B \rightarrow C \rightarrow A$. It is easy to see that every arc contained in one of the three subtournaments A , B and C is not pancyclic because it is contained in no 3-cycle. It follows that $h(T_n) \leq 3k$. ■

The bound $3k$ is not tight because of the 2-strong tournaments of order $2k + 1 \leq n < 3k$. However, we think that if n is large enough, the above example are extremal.

Conjecture 2 For n sufficiently large, $h_k(n) = 3k$.

Alsopach [1] showed that every arc of a regular tournament is pancyclic. This implies that $h_k(2k + 1) = 2k + 1$. In order to avoid the exception of small order in Conjecture 2, it would make sense to first try to prove that $h_k(n) \geq 2k + 1$.

2 Definition and preliminaries

Let T be a tournament. Let x and y be two vertices of T . We write $x \rightarrow y$ if (x, y) is an arc of T . Likewise, let X and Y be two subdigraphs of T . We write $X \rightarrow Y$ if $x \rightarrow y$ for all pairs $(x, y) \in V(X) \times V(Y)$.

Let A_1, A_2, \dots, A_k be a family of subdigraphs of T . We denote by $T[A_1, A_2, \dots, A_k]$ the subtournament induced by T on the set of vertices $\bigcup_{1 \leq i \leq k} V(A_i)$ and by $T - [A_1, A_2, \dots, A_k]$

the subtournament induced by T on the set of vertices $V(T) \setminus \bigcup_{1 \leq i \leq k} V(A_i)$.

$A(X, Y)$ denotes the set of arc (x, y) with $x \in X$ and $y \in Y$. $A^+(X)$ is the set of arcs outgoing from X , that is $A(X, T - X)$ and $A^-(X)$ is the set of arcs ingoing into X , that is $A(T - X, X)$.

The *dual* of a digraph D is the digraph $-D$ on the same set of vertices such that $x \rightarrow y$ is an arc of $-D$ if and only if $y \rightarrow x$ is an arc of D .

An (x, y) -path is a path that begins in x and terminates at y .

One can easily show the following result (whose proof is left to the reader) that allows to extend an (x, y) path to a longer one under condition.

Proposition 3 *Let $P = (v_1, v_2, \dots, v_m)$ be a path in a tournament T and x a vertex of $T - P$.*

If there exist $1 \leq i < j \leq m$ such that $v_i \rightarrow x \rightarrow v_j$, then in T there is a path with $m + 1$ vertices starting in v_1 and ending in v_m .

A non-strong tournament T is said to be *reducible*. It admits a *reduction* into two subtournaments T_1 and T_2 such that $V(T_1) \cup V(T_2) = V(T)$ and $T_1 \rightarrow T_2$; in this case, we write $T = T_1 \rightarrow T_2$.

A (strong) *component* of T is a strong subtournament of T which is maximal by inclusion. Let T_1, T_2, \dots, T_m be the components of T . Then $(V(T_1), V(T_2), \dots, V(T_m))$ is a partition of $V(T)$ and without loss of generality, we may suppose that $T_i \rightarrow T_j$ whenever $i < j$. In this case we say that $T_1 \rightarrow T_2 \rightarrow \dots \rightarrow T_m$ is the *decomposition* of T . The component T_1 (resp. T_m) is called the *outsection* (resp. *insection*) of T , denoted by $out(T)$ (resp. $In(T)$) and its vertices are called the *outgenerators* of T (resp. *ingenerators* of T).

Proposition 4 *In a tournament, a vertex is the beginning of a hamiltonian path if and only if it is an outgenerator.*

Proof. Let T be a tournament. Then $T = Out(T) \rightarrow T - Out(T)$ (with $T - Out(T)$ empty when T is strong). $T[Out(T)]$ is strong, and thus by Camion's theorem admits a hamiltonian cycle C . Then every outgenerator v is the origin of a hamiltonian path Q of $T[Out(T)]$. And $T - Out(T)$ has a hamiltonian path P . So (P, Q) is hamiltonian path of T with origin v . ■

Proposition 5 *Let T be a reducible tournament of order n . Let u be an ingenerator and t an outgenerator of T . For any $1 \leq l \leq n - 1$, there is a (t, u) -path of length l .*

Proof. Let $T_1 \rightarrow T_2$ be a reduction of T and for $i = 1, 2$, let n_i be the order of T_i . Clearly, t is an ingenerator of T_2 and u an outgenerator of T_1 . Therefore, by Proposition 4, t is the end of a hamiltonian path $(t_{n_2-1}, t_{n_2-2}, \dots, t_1, t)$ of T_2 and u is the origin of a hamiltonian path $(u, u_1, \dots, u_{n_1-2}, u_{n_1-1})$ of T_1 . Now, for any $1 \leq l \leq n - 1$, pick $0 \leq l_1 \leq n_1 - 1$ and $0 \leq l_2 \leq n_2 - 1$ such that $l_1 + l_2 = l - 1$. Then $(u, u_1, \dots, u_{l_1}, t_{l_2}, t_{l_2-1}, \dots, t)$ is a path of length l . ■

A *reductor* of a tournament is a smallest subtournament X such that $T - X$ is reducible. If T is $(= k)$ -strong then a reductor has k vertices.

Proposition 6 *Let X be a reductor of a tournament T . Every element of X dominates an outgenerator of $T - X$ and is dominated by an ingenerator of $T - X$.*

Proof. Let x be an element of X . Let Y be the set of vertices that are not outgenerator of $T - X$. If $Out(T - X) \rightarrow x$, then $T - [X \setminus x]$ is reducible with reduction $Out(T - X) \rightarrow T[Y, x]$. This contradicts that X is a reductor.

Analogously, we prove that x is dominated by an ingenerator of $T - X$. ■

Proposition 7 *Let x and y be two vertices of a reductor X of a 2-tournament T . If $in(T - X) \geq 3$, there are two distinct vertices z_x and z_y of $In(T - X)$ such that $z_x \rightarrow x$ and $z_y \rightarrow y$.*

Proof. Suppose that two such vertices do not exist, then by Proposition 6, there is a vertex $u \in In(T - X)$ such that $u \rightarrow \{x, y\}$ and $In(T - X) \setminus u \leftarrow \{x, y\}$. Then $X \setminus \{x, y\} \cup \{u\}$ is a reductor of T , which is a contradiction. ■

3 Lower bounds for $p(T)$ and $h(T)$

3.1 Sufficient conditions for an arc to be pancyclic

Lemma 1 *Let X be a subtournament such that $T - X$ is reducible and every vertex of X dominates an outgenerator of $T - X$. Let v be an outgenerator of X and u an ingenerator of $T - X$. If $u \rightarrow v$, then the arc (u, v) is pancyclic.*

Proof. By Proposition 4, $v = v_0$ is the origin of a hamiltonian path $(v_0, v_1, \dots, v_{k-2}, v_{k-1})$ of $T[X]$. For $0 \leq i \leq k - 1$, let t_i be an outgenerator of $T - X$ dominated by v_i . For $3 \leq l \leq n$, take $0 \leq k' \leq k - 1$ and $1 \leq l \leq n - k - 1$ such that $k' + l + 2 = l$. Then by Proposition 5, in $T - X$, there is a $(t_{k'}, u)$ path Q of length l . Thus $(v_0, v_1, \dots, v_{k'}, Q, v_0)$ is a cycle of length l going through $(u, v) = (u, v_0)$. ■

Corollary 1 *Let X be the reductor of a tournament T . Let v be an outgenerator of X and u an ingenerator of $T - X$. If $u \rightarrow v$, then the arc (u, v) is pancyclic.*

Corollary 2 *Let T be a strong tournament.*

$$p(T) \geq h(T) \geq 2$$

Proof. Let X be a reductor of T . Let P be a hamiltonian path of X with beginning v and end w . Clearly, v is an outgenerator of X and w is an ingenerator of X . Then by Corollary 1 (and its dual), (u, v) and (w, t) are pancyclic.

And by Proposition 5, there is a (t, u) -path Q that is hamiltonian in $T - X$, thus (P, Q, v) is a hamiltonian cycle containing (u, v) and (w, t) . ■

Let T be a strong tournament T with reductor X . Let $X_1 \rightarrow X_2 \rightarrow \dots \rightarrow X_l$ be a decomposition of X and $T_1 \rightarrow T_2 \rightarrow \dots \rightarrow T_m$ be a decomposition of $T - X$.

In the remaining of this section, we examine the number of pancyclic arcs in the different parts of T .

Lemma 2 For $1 \leq i \leq l$, if an arc is pancyclic in X_i , then it is also pancyclic in T .

Proof. Let e be a pancyclic arc in one of the X_i . Then e is contained in a 3-cycle. It is also contained in a hamiltonian cycle and then a hamiltonian path of X_i . This hamiltonian path may be extended to a hamiltonian path $(v_0, v_1, \dots, v_{k-1})$ of X using hamiltonian paths of the $X_{i'}$, for $i' \neq i$. Let j be the index such that $e = (v_j, v_{j+1})$. Let $4 \leq l \leq n$. Choose $0 \leq l_1 \leq j$, $0 \leq l_2 \leq k - j - 2$ and $1 \leq l_3 \leq n - k - 1$ such that $l_1 + l_2 + l_3 + 3 = l$. By Proposition 6, there is an ingenerator u of $T - X$ dominating v_{j-l_1} and an outgenerator t of $T - X$ dominated by v_{j+1+l_2} . Then by Proposition 5, in $T - X$ there is a (t, u) -path of length l_3 . Hence $(v_{j-l_1}, v_{j-l_1+1}, \dots, v_{j+1+l_2}, P, v_{j-l_1})$ is an l -cycle containing e . ■

Lemma 3 Suppose that $T - X$ is the transitive tournament $TT(t_1, t_2, \dots, t_m)$.

For every $1 \leq i \leq m - 1$, the arc (t_i, t_{i+1}) is pancyclic if and only if it is contained in a 3-cycle.

Proof. By Proposition 6, $X \rightarrow t_1$ and $t_m \rightarrow X$. Let $(v_0, v_1, \dots, v_{k-2}, v_{k-1})$ be a hamiltonian path of X .

Suppose first that $i = 1$. Let $4 \leq l \leq n$. Take $0 \leq k' < k$, and $1 \leq l' < m - 2$ such that $k' + l' + 3 = l$. By Proposition 5, there is a (t_2, t_m) -path P of length l' in $T[t_2, \dots, t_m]$. Then $(t_1, P, v_0, v_1, \dots, v_{k'}, t_1)$ is a cycle of length l . Thus if (t_1, t_2) is contained in a 3-cycle then it is pancyclic.

By duality, we have the result if $i = m - 1$.

Suppose now that $1 < i < m - 1$. Let v be the vertex such that (t_i, t_{i+1}, v, t_i) is a 3-cycle. Necessarily, v is in X . Then $(t_i, t_{i+1}, v, t_1, t_i)$ is a 4-cycle. Let $5 \leq l \leq n$. The arc (t_i, t_{i+1}) is contained in a cycle of length l . Indeed, take $0 \leq k' < k$, $0 < l_1 < i$ and $0 \leq l_2 < m - i$ such that $k' + l_1 + l_2 + 3 = l$. By Proposition 5, there is a (t_1, t_i) -path P_1 of length l_1 in $T[t_1, \dots, t_i]$, and a (t_{i+1}, t_m) -path P_2 of length l_2 in $T[t_{i+1}, \dots, t_m]$. Then $(t_i, P_2, v_0, v_1, \dots, v_{k'}, P_1)$ is an l -cycle containing (t_i, t_{i+1}) . Thus (t_i, t_{i+1}) is pancyclic. ■

Lemma 4 For $1 < i < m$, if $|T_i| \geq 4$, then every T_i -pancyclic arc is pancyclic in T .

Proof. Let $(v_0, v_1, \dots, v_{k-2}, v_{k-1})$ be a hamiltonian path of X .

Let (a_1, a_2) be a pancyclic arc in T_i . It is contained in a hamiltonian cycle of T_i , $C_i = (a_1, a_2, \dots, a_{n_i}, a_1)$.

Let us prove that (a_1, a_2) is contained in a cycle of length l for all $5 \leq l \leq n$. Let $S_1 = T[T_1, T_2, \dots, T_{i-1}, a_1]$ and $S_2 = T - [X, S_1]$. Clearly, a_1 is an ingenerator of S_1 and a_2 an outgenerator of S_2 . There exists three integers $1 \leq l_1 \leq |S_1| - 1$, $1 \leq l_2 \leq |S_2| - 1$ and $0 \leq k' \leq k - 1$ such that $l_1 + l_2 + k' = l - 3$. By Proposition 6, there is an ingenerator u_0 of $T - X$ and then also of S_2 which dominates v_0 and an outgenerator $t_{k'}$ of S_1 which is dominated by $v_{k'}$. By Proposition 5, there is a $(t_{k'}, a_1)$ -path P_1 of length l_1 in S_1 and a (a_2, u_0) -path P_2 of length l_2 in S_2 . Then $(P_1, P_2, v_0, v_1, \dots, v_{k'}, t_{k'})$ is the desired l -cycle.

And because it is pancyclic in T_i , (a_1, a_2) is contained in an l -cycle, for $3 \leq l \leq n_i$. ■

Lemma 5 For any $1 < i < m$, if T_i is a 3-cycle, then two arcs of T_i are pancyclic in T .

Proof. Let $(v_0, v_1, \dots, v_{k-2}, v_{k-1})$ be a hamiltonian path of X . The component T_i is the 3-cycle (a_1, a_2, a_3, a_1) .

In the same way as in the proof of Lemma 4 the arc (a_1, a_2) , is contained in cycle of length l for all $5 \leq l \leq n$. Analogously, (a_2, a_3) and (a_3, a_1) are contained in cycle of length l for all $5 \leq l \leq n$.

Hence, it suffice to prove that two arcs of T_i is contained in a 4-cycle. Without loss of generality, we may assume that $\{a_1, a_2\} \leftarrow v_0$ or $\{a_1, a_2\} \rightarrow v_0$. If $\{a_1, a_2\} \leftarrow v_0$, then $(a_1, a_2, u_0, v_0, a_1)$ and $(a_2, a_3, u_0, v_0, a_2)$ are 4-cycles. If $\{a_1, a_2\} \rightarrow v_0$, then $(a_3, a_1, v_0, t_0, a_3)$ and $(a_1, a_2, v_0, t_0, a_1)$ are 4-cycles. ■

Lemma 6 *Let $C = (a_1, a_2, \dots, a_{n_1}, a_1)$ be a hamiltonian cycle of T_1 such that a_1 is dominated by an ingenerator of X .*

If (a_j, a_{j+1}) is pancyclic in T_1 and $1 \leq j \leq n_1 - 2$ then (a_j, a_{j+1}) is pancyclic in T .

Proof. By T_1 -pancyclicity, (a_j, a_{j+1}) is contained in a cycle of length l for $3 \leq l \leq n_1$.

Let now l be an integer of $[n_1 + 1, n]$.

Let v_{k-1} be an ingenerator of X dominating a_1 . By Proposition 4, there exists a hamiltonian path $(v_0, v_1, \dots, v_{k-2}, v_{k-1})$ of $T[X]$. By Proposition 6, for every $0 \leq i \leq k-1$, there is an ingenerator u_i of $T - X$ dominating v_i .

Let $1 \leq l_1 \leq n - k - j - 1$ and $0 \leq k' \leq k - 1$ such that $k' + l_1 = l - j - 2$. Obviously, a_{j+1} is an outgenerator of $T' = T - [X, a_1, a_2, \dots, a_j]$. And $u_{k-1-k'}$ is an ingenerator of T' . Thus, by Proposition 5 there is an $(a_{j+1}, u_{k-1-k'})$ path P of length l_1 . Hence $(P, v_{k-1-k'}, v_{k-1-k'}, \dots, v_{k-1}, a_1, a_2, \dots, a_{j+1})$ is a cycle of length l . ■

Lemma 7 *There are at least $h(T_1) - 1$ pancyclic arcs in T_1 .*

Proof. Let $C = (a_1, a_2, \dots, a_{n_1}, a_1)$ be a hamiltonian cycle of T_1 containing $h(T_1)$ T_1 -pancyclic arcs. Without loss of generality, we may suppose that a_1 is dominated by an ingenerator v of X .

Since $j_1 \leq n_1 - 2$, by Lemma 6, every pancyclic arc in $P = (a_1, a_2, \dots, a_{n_1-1})$ is also pancyclic in T . If P contains $h(T_1) - 1$ T_1 -pancyclic arcs, we have the result.

Hence we may assume that (a_{n_1-1}, a_{n_1}) and (a_{n_1}, a_1) are pancyclic in T_1 .

If $v \rightarrow a_{n_1}$, again by Lemma 6, the arc (a_{n_1}, a_1) is pancyclic in T .

If $v \leftarrow a_{n_1}$ then let us prove that $e_2 = (a_{n_1-1}, a_{n_1})$ is pancyclic in T . By pancyclicity in T_1 , e_2 is contained in a cycle of length l for $3 \leq l \leq n_1$. And $(a_1, a_2, \dots, a_{n_1}, v, a_1)$ is an $(n_1 + 1)$ -cycle containing e_2 .

Let now l be an integer of $[n_1 + 2, n]$.

By Proposition 4, there is a hamiltonian path such that $(v_0, v_1, \dots, v_{k-2}, v_{k-1})$ be a hamiltonian path of X . By Proposition 6, for every $0 \leq i \leq k-1$, there is an ingenerator u_i of $T - X$ dominating v_i .

Let $1 \leq l_1 \leq n - k - n_1 - 1$ and $0 \leq k' \leq k - 1$ such that $k' + l_1 = l - n_1 - 1$. Obviously, a_{n_1} is an outgenerator of $T' = T - [X, Out(X) \setminus a_{n_1}]$. And $u_{k-1-k'}$ is an ingenerator of T' . Thus, by Proposition 5 there is a path P of length l_1 with start a_{n_1} and end $u_{k-1-k'}$. Hence $(P, v_{k-1-k'}, v_{k-1-k'}, \dots, v_{k-1}, a_1, a_2, \dots, a_{n_1})$ is a cycle of length l . ■

3.2 The lower bounds

Using the above lemmas, we derive lower bounds for $p(T)$ and $h(T)$.

Definition 1 Let k be an integer and T a tournament. $\epsilon(k, T) = 1$ if $|T| \geq k$ and 0 otherwise.

Theorem 2 If $T - X$ is transitive then

$$p(T) \geq |A(\text{In}(T - X); \text{Out}(X))| + |A(\text{In}(X); \text{Out}(T - X))| + \sum_{j=1}^l p(X_j) + 1 \quad (1)$$

Otherwise

$$\begin{aligned} p(T) \geq & |A(\text{In}(T - X); \text{Out}(X))| + |A(\text{In}(X); \text{Out}(T - X))| + \sum_{j=1}^l p(X_j) \\ & + \sum_{i=2}^{m-1} \{2\epsilon(3, T_i) + (p(T_i) - 2)\epsilon(4, T_i)\} \\ & + \epsilon(3, T_1)(h(T_1) - 1) + \epsilon(3, T_m)(h(T_m) - 1) \end{aligned} \quad (2)$$

Proof. Since every outgenerator of $T - X$ is dominated by an ingenerator of $T - X$, according to Corollary 1, every arc of $A(\text{In}(T - X); \text{Out}(X))$ is pancyclic. By duality, every arc of $A(\text{In}(X); \text{Out}(T - X))$ is pancyclic. According to Lemma 2, there are at least $\sum_{j=1}^l p(X_j)$ T -pancyclic arcs in X .

Suppose now that $T - X$ is a transitive tournament. Then since $v_0 \rightarrow t_1$ and $v_0 \rightarrow t_m$ then there is an index i such that (t_i, t_{i+1}, v_0, t_i) is a 3-cycle. So by Lemma 3, this arc is pancyclic. So we obtain Equation 1.

To obtain Equation 2, let us now count the number of pancyclic arcs contained in each T_i such that $|T_i| \geq 3$. If $i = 1$ or $i = m$, by Lemma 7 (or its dual), $h(T_i) - 1$ arc are T -pancyclic.

If $1 < i < m$, then, if T_i is a 3-cycle then by Lemma 5, 2 arcs of T_i are pancyclic in T ; and if $|T_i| \geq 4$, by Lemma 4, each T_i -pancyclic arc is pancyclic in T . ■

Theorem 3 If $T - X$ is transitive then

$$h(T) \geq 3 + \sum_{j=1}^l \epsilon(3, X_j) \cdot \min\{h(X_j); |X_j| - 1\} \quad (3)$$

otherwise

$$\begin{aligned} h(T) \geq & 2 + \sum_{j=1}^l \epsilon(3, X_j) \cdot \min\{h(X_j); |X_j| - 1\} + \sum_{i=2}^{m-1} \epsilon(3, T_i) \cdot \min\{h(T_i); |T_i| - 1\} \\ & + \epsilon(3, T_1)(h(T_1) - 1) + \epsilon(3, T_m)(h(T_m) - 1) \end{aligned} \quad (4)$$

Proof. For $1 \leq j \leq l$, let P_j be a hamiltonian path of X_j defined as follows :

- If X_j is reduced to a single vertex x_j then $P_j = (x_j)$.
- if $|X_j| \leq 3$, let P_j is obtained from a hamiltonian cycle of X_j containing $h(X_j)$ X_j -pancyclic arcs by removing a non X_j -pancyclic arc if $h(X_j) < |X_j|$ or any arc if $h(X_j) = |X_j|$.

By Lemma 2, each P_j contains $\epsilon(3, X_j) \cdot \min\{h(X_j); |X_j| - 1\}$ T -pancyclic arcs. Let v be the beginning of P_1 and w the end of P_l .

Suppose first that $T - X$ is the transitive tournament $TT(t_1, t_2, \dots, t_m)$. By Lemma 3, there exists i such that (t_i, t_{i+1}) is pancyclic and by Corollary 1, (w, t_1) and (t_m, v) are pancyclic. Thus, the hamiltonian cycle $((t_1, t_2, \dots, t_m, P_1, P_2, \dots, P_l, t_1))$ contains $3 + \sum_{j=1}^l \epsilon(3, X_j) \cdot \min\{h(X_j); |X_j| - 1\}$.

Suppose now that $T - X$ is not transitive.

For $1 < i < m$, let Q_i be a hamiltonian path of T_i defined as follows :

- If T_i is reduced to a single vertex t_i then $Q_i = (t_i)$.
- If $|T_i| = 3$, then Q_i is a path formed by two arcs of T_i that are pancyclic in T . (Such arcs exists according to Lemma 5.)
- If $|T_i| \leq 4$, let Q_i is obtained from a hamiltonian cycle of T_i containing $h(T_i)$ T_i -pancyclic arcs by removing a non T_i -pancyclic arc if $h(T_i) < |T_i|$ or any arc if $h(T_i) = |T_i|$.

By Lemma 4, each Q_i contains $\epsilon(3, T_i) \cdot \min\{h(T_i); |T_i| - 1\}$ T -pancyclic arcs.

Let $C_1 = (a_1, a_2, \dots, a_{n_1}, a_1)$ be a hamiltonian path of T_1 containing $h(T_1)$ T_1 -pancyclic arcs. Then set $Q_1 := ((a_1, a_2, \dots, a_{n_1}))$. Analogously define Q_m . By (the proof of) Lemma 6 Q_1 (resp. Q_m) contains at least $h(T_1) - 1$ (resp. $h(T_m) - 1$) pancyclic arcs in T .

Then the hamiltonian cycle $(w, Q_1, Q_2, \dots, t_m, P_1, P_2, \dots, P_l)$ gives Equation 4. ■

3.3 Tournaments with $h(T) = 3$

From the lower bounds, one can easily derive Theorem 1 :

Proof of Theorem 1 Let us prove the result by induction on the order n of T . If $n = 3$, it is obviously true.

Suppose now that it is true for strong tournaments of order less than n . Let X be a reductor of T and $T_1 \rightarrow T_2 \rightarrow \dots \rightarrow T_m$ be a decomposition of $T - X$.

If $T - X$ is transitive, Equation 3 yields the result and if it is not then Equation 4 gives $h(T) \geq 4$.

Suppose now T is a tournament such that $h(T) = 3$. Let X be a reductor of T . By Equation 4 then X and $T - X$ are a transitive tournaments. Set $X = TT[v_0, v_1, \dots, v_{k-1}]$ and $T - X = TT[t_1, t_2, \dots, t_{n-k}]$. By Corollary 1, (t_{n-k}, v_0) and (v_{k-1}, t_1) are pancyclic. Therefore there is at most one pancyclic arc on the hamiltonian path of $T - X$. Thus by

Lemma 3, there is an index i such that $X \rightarrow t_j$ if and only if $j \leq i$. And (t_i, t_{i+1}) is pancyclic.

Suppose that T is 2-strong. Then $T - t_1$ and $T - t_{n-k}$ are strong, so $2 \leq i \leq n - k - 2$. By Lemma 1, (t_{n-k-1}, v_0) is pancyclic in $T - t_{n-k}$ and (t_{n-k}, v_1) is pancyclic in $T - v_0$. Thus two arcs are T -pancyclic because they are contained in the n -cycle $C_3 = (t_{n-k-1}, v_0, v_{k-1}, t_2, t_3, \dots, t_{n-k-2}, t_{n-k}, v_1, v_2, \dots, v_{k-2}, t_1, t_{n-k-1})$ if $k \geq 3$ or $C_2 = (t_{n-3}, v_0, t_1, t_{n-2}, v_1, t_2, t_3, \dots, t_{n-3})$ if $k = 2$. Analogously, (v_{k-1}, t_2) and (v_{k-2}, t_1) are also T -pancyclic. Hence the cycle C_2 or C_3 contains four pancyclic arcs. This is a contradiction.

Thus T is $(= 1)$ -strong. Then by Lemma 3 it is in \mathcal{P}_3 . ■

Note that in the proof of Theorem 1, we also show the following :

Proposition 8 *Let X be a reductor of a strong tournament. There is at least one T -pancyclic arc in $T - X$.*

4 Number of pancyclic arcs in k -strong tournaments

Theorem 4 (Yao, Guo and Zhang, [5]) *Every tournament contains a vertex x such that every outgoing arc is pancyclic.*

From this result, we derive lower bounds on the number of pancyclic arcs in a strong tournament :

Lemma 8

$$p(T) \geq h(T) + \delta^+ - 1 \tag{5}$$

$$p(T) \geq h(T) + \delta^+ + \delta^- - 3 \tag{6}$$

Proof. By Theorem 4 and its dual, there is a vertex x such that every outgoing arc is pancyclic and a vertex y such that every ingoing arc is pancyclic. Obviously, $|A^+(x) \cap A^-(y)| \leq 1$. There are $h(T)$ pancyclic arcs on a hamiltonian cycle. At most one of them is in $A^+(X)$ and at most two of them are in $A^+(x) \cup A^-(y)$. Hence, $p(T) \geq h(T) + d^+(x) - 1 \geq h(T) + \delta^+ - 1$, and

$$p(T) \geq d^+(x) + d^-(y) - |A^+(x) \cap A^-(y)| + h(T) - 2 \tag{7}$$

$$\geq d^+(x) + d^-(y) + h(T) - 3 \tag{8}$$

■

Lemma 9 *Let X be the reductor of a 2-strong tournament. Suppose that X is the transitive tournament $TT(v_0, v_1, \dots, v_{k-1})$.*

For every $0 \leq j \leq k - 2$, the arc (v_j, v_{j+1}) is pancyclic if and only if it is contained in a 3-cycle.

Proof. Let $0 \leq j \leq k-2$ and $4 \leq l \leq n$. Let us prove that (v_j, v_{j+1}) is contained in an l -cycle. There exist $0 \leq l_1 \leq j$, $0 \leq l_2 \leq k-j-2$ and $1 \leq l_3 \leq n-k$ such that $l_1 + l_2 + l_3 + 3 = l$. By Proposition 6, there is an ingenerator u of $T-X$ dominating v_{j-l_1} and an outgenerator t of $T-X$ dominated by v_{j+1+l_2} . And by Proposition 5, in $T-X$, there is an (t, u) -path P of length l_3 . Therefore, $(P, v_{j-l_1}, v_{j-l_1+1}, \dots, v_{j+1+l_2}, t)$ is the desired l -cycle. \blacksquare

Theorem 5 For every a 2-strong tournament T , $h(T) \geq 5$.

Proof. Let T be a 2-strong tournament such that $h(T) = 4$. Let X be a reductor of T and $T_1 \rightarrow T_2 \rightarrow \dots \rightarrow T_m$ be a decomposition of $T-X$. By Equations 3 and 4, we may assume that X is a transitive tournament, say $TT(v_0, v_1, \dots, v_{k-1})$, and that at most one of the T_i is not reduced to a single vertex t_i .

I) Suppose that for some $2 \leq i \leq m-1$, T_i is not reduced to a single vertex. Then by Equation 4, we may assume that T_i is a 3-cycle (a, b, c, a) . By Lemma 10, for $1 \leq j \leq i-1$, $X \rightarrow t_j$ and for $i+1 \leq j \leq m$, $X \leftarrow t_j$. Without loss of generality, we may assume that both a and b dominate a vertex in X . Then by Lemma 10, (t_{i-1}, a) and (t_{i-1}, b) are pancyclic and by (the proof of) Lemma 5, the arcs (a, b) and (c, a) are pancyclic. If c dominates a vertex in X then (b, c) is pancyclic, and if c is dominated a vertex in X then (c, t_{i+1}) is pancyclic. In any case, the hamiltonian cycle $(t_1, t_2, \dots, t_{i-1}, a, b, c, t_{i+1}, \dots, t_m, v_0, v_1, \dots, v_{k-1}, t_1)$ contains five pancyclic arcs.

II) Suppose that $|T_1| \geq 3$.

By Equation 4, we may assume that $h(T_1) = 3$. So $T_1 \in \mathcal{P}_3$ according to Theorem 1. Let w be the reductor of T_1 such that $T_1 - w$ is the transitive tournament.

If for some $2 \leq j < m$, t_j is dominated by a vertex of X , then by Lemma 10 (dual), there is a pancyclic arc in the path $(t_j, t_{j+1}, \dots, t_m)$. Let P be a hamiltonian path of T_1 beginning at an outneighbour of V_{k-1} . The hamiltonian cycle $C = (v_{k-1}, P, t_2, \dots, t_{i-1}, t_i, t_{i+1}, \dots, t_m, v_0, v_1, \dots, v_{k-1})$ contains five pancyclic arcs. Hence we may assume that $X \leftarrow t_j$, for $2 \leq j \leq m$.

If there is a pancyclic arc in the path $(v_0, v_1, \dots, v_{k-1})$, the cycle C contains five pancyclic arcs. So, by Lemma 9, we may assume that $N_{T_1}^+(v_{k-1}) \subseteq N_{T_1}^+(v_{k-2}) \subseteq \dots \subseteq N_{T_1}^+(v_0)$. There are two distinct vertices r_1 and r_2 of T_1 that are dominated by v_{k-1} . Let C_1 be a hamiltonian cycle of T_1 containing three T_1 -pancyclic arcs. Let r_1^- (resp. r_2^+) be the vertex dominating r_1 (resp. r_2) in C . Let P_1 (resp. P_2) be the subpaths of C_1 with beginning r_1 (resp. r_2) and end r_2^- (resp. r_1^-).

1) Suppose first that $m \geq 3$.

Let C' be the cycle defined as follows:

- If $k \geq 3$, then $C' := (v_0, v_{k-1}, P_1, t_m, v_1, v_2, \dots, v_{k-2}, P_2, t_2, t_3, \dots, t_{m-1}, v_0)$;
- if $k = 2$, then $C' := (v_0, P_2, t_m, v_1, P_1, t_2, t_2, t_3, \dots, t_{m-1}, v_0)$.

Since C_1 contains at least three T_1 -pancyclic arcs, one of them, say e is contained in P_1 or P_2 . So, by Lemma 6, e is pancyclic in T . Moreover C' contains the four arcs (t_m, v_1) , (t_{m-1}, v_0) , (v_{k-2}, r_2) and (v_{k-1}, r_1) . By Lemma 1, these arcs are pancyclic in respectively in $T - [v_0]$, $T - [t_m]$, $T - [v_{k-1}]$ and T . Hence because they are contained in C' , these four arcs are pancyclic. Thus C' contains five pancyclic arcs.

- 2) Suppose now that $m = 2$. Then v_0 (and then every v_j , $0 \leq j \leq k-1$) has an inneighbour in T_1 which dominates X . Without loss of generality we may suppose that this inneighbour s_2^- is between r_1 and r_2 in C_1 , that s_2 the successor of s_2^- along C_1 dominates v_0 . Let Q_1 (resp. Q_2) be the path beginning in r_1 (resp. s_2) and ending in s_2^- (resp. r_1^-).

Set $C_0 := (v_0, Q_2, t_2, v_1, v_2, \dots, v_{k-1}, Q_1, v_0)$. It is easy to check that (v_0, s_2) and (t_2, v_1) and (v_{k-1}, r_1) are pancyclic in T . If $P_1 \cup P_2$ contains two T_1 -pancyclic arcs then these two are also pancyclic in T by Lemma 6. So C_0 contains five pancyclic arcs.

Hence we may assume that both (r_1^-, r_1) and (s_2^-, s_2) are pancyclic. Let e be the third T_1 -pancyclic arc. Let x be a vertex of Q_1 such that $v_0 \rightarrow x$ and its successor x^+ in Q_1 dominates v_0 . Such a vertex exists since $v_0 \rightarrow r_1$ and $s_2^- \rightarrow v_0$. The arc (x, x^+) of Q_1 is pancyclic. Indeed ...

Thus if $e = (x, x^+)$, so $e \in Q_1$.

Three cases may arise:

- a) Suppose that $w = s_2 = r_1^-$. Then $T_1 - [s_2^-, s_2] \rightarrow s_2^-$, thus (s_2^-, v_0, r_1, s_2^-) is a 3-cycle. For $4 \leq l \leq n_1 + 1$, (s_2^-, v_0) is contained in the l -cycle obtained by replacing the arc (s_2^-, s_2) by (s_2^-, v_0, s_2) in the $l-1$ -cycle. Now since $v_0 \rightarrow X - v_0$ and $X \rightarrow r_1$, by Proposition 3 (applied for every vertex of $X - v_0$ one after another), (s_2^-, v_0) is contained in an l -cycle for $n_1 + 2 \leq l \leq n - 1$. And it is contained in C_0 . Hence, (s_2^-, v_0) is pancyclic and C_0 contains five pancyclic arcs.
- b) Suppose that $w = r_1$. Then $r_1 \rightarrow T_1 - Q_2$ and $Q_2 \rightarrow r_1$. Then (s_2^-, v_0, r_1, s_2^-) is a 3-cycle. For $4 \leq l \leq n_1 + 1$, pick Q'_1 a subpath of Q_1 ending in s_2^- of length $l_1 < |Q_1| - 2$ and Q'_2 a subpath of Q_2 beginning in s_2 of length $l_2 < |Q_2| - 2$ such that $l_1 + l_2 + 4 = l$, then $(s_2^-, v_0, Q'_2, r_1, Q'_1)$ is an l -cycle. Analogously to the end of Case a), we obtain that (s_2^-, v_0) is pancyclic and C_0 contains five pancyclic arcs.
- Suppose that $w = s_2^- = x^+$. Then $r_1 \rightarrow w$, so (s_2^-, v_0, r_1, s_2^-) is a 3-cycle. And $T_1 - s_2^-$ is a transitive tournament with outgenerator s_2 and ingenerator x . Thus for $4 \leq l \leq n_1 + 1$, in $T_1 - s_2^-$, there is an (s_2, x) -path P of length $l - 2$. Hence, (Q, s_2^-, v_0, s_2) is an l -cycle. Again, analogously to the end of Case a), we obtain that (s_2^-, v_0) is pancyclic and C_0 contains five pancyclic arcs.

- III) Suppose now that $T - X$ is a transitive tournament.

If $v_0 \rightarrow t_{n-k-1}$, then $X' = T[t_{n-k}, v_1, v_2, \dots, v_{k-1}]$ is a reductor of T and $T - X'$ is not a transitive. Indeed v_0 is dominated by some vertex t_i and then (t_i, v_0, t_1, t_i) is a 3-cycle. So we have the result by one of the previous case.

Analogously, we obtain the result if $v_{k-2} \leftarrow t_2$. Thus we may assume that $v_0 \leftarrow t_{n-k-1}$ and $v_{k-2} \rightarrow t_2$.

By Lemma 1, (t_{n-k-1}, v_0) is pancyclic in $T - t_{n-k}$ and (t_{n-k}, v_1) is pancyclic in $T - v_0$. Thus two arcs are T -pancyclic because they are contained in the n -cycle $C_3 = (t_{n-k-1}, v_0, v_{k-1}, t_2, t_3, \dots, t_{n-k-2}, t_{n-k}, v_1, t_{n-k-1})$ if $k \geq 3$ or $C_2 = (t_{n-3}, v_0, t_1, t_{n-2}, v_1, t_2, t_3, \dots, t_{n-3})$ if $k = 2$. Analogously, (v_{k-1}, t_2) and (v_{k-2}, t_1) are also T -pancyclic.

Suppose that $k \geq 3$, then the 4 pancyclic arcs (t_{n-k-1}, v_0) , (t_{n-k}, v_1) , (v_{k-1}, t_2) and (v_{k-2}, t_1) are contained in the two n -cycles C_3 and $C'_3 = (t_{n-k-1}, v_0, v_{k-1}, t_2, t_{n-k}, v_1, v_2, \dots, v_{k-2}, t_1, t_3, t_4, \dots, t_{n-k-1})$. Therefore no arcs in $\{(t_i, t_{i+1}), 2 \leq i \leq n-k-2\}$ is T pancyclic. By Lemma 3, $t_i \rightarrow v_0$ for $2 \leq i \leq n-k$ and (t_1, t_2) is pancyclic, and $v_{k-1} \rightarrow t_i$ for $1 \leq i \leq n-k-1$ and (t_{n-k-1}, t_{n-k}) is pancyclic. Thus at least one of the arcs (v_i, v_{i+1}) is in a 3-cycle is pancyclic by Lemma 9. Hence the cycle $C = (v_0, v_1, \dots, v_{k-1}, t_1, t_2, \dots, t_{n-k}, v_0)$ contains five pancyclic arcs.

Thus we may assume that $k = 2$. We may assume that there is no pancyclic arcs in $\{(t_i, t_{i+1}), 2 \leq i \leq n-4\}$, otherwise C_2 contains five pancyclic arcs. Then by Lemma 3, $t_i \rightarrow v_0$ for $2 \leq i \leq n-2$ and (t_1, t_2) is pancyclic, and $v_1 \rightarrow t_i$ for $1 \leq i \leq n-3$ and (t_{n-3}, t_2) is pancyclic. Hence (v_0, v_1) is in the 3-cycle (v_0, v_1, t_2, v_0) , so by Lemma 9, it is pancyclic. Hence the cycle $C = (v_0, v_1, t_1, t_2, \dots, t_{n-2}, v_0)$ contains five pancyclic arcs. ■

This result is best possible since the regular tournament R_5 on five vertices is 2-strong and obviously satisfies $h(R_5) = 5$.

It follows directly from Theorems 5 and Equation 6, that for $k \geq 2$, every k -strong tournament has at least $2k + 2$ pancyclic arcs.

Corollary 3 *Every k -strong tournament has at least $2k + 2$ pancyclic arcs.*

We now prove a slightly better result.

Theorem 6 *Every k -strong ($k \geq 2$) tournament has at least $2k + 3$ pancyclic arcs.*

Proof. Let x and y be vertices such that the arcs of $A^+(x) \cup A^-(y)$ are pancyclic. By Equation 7, we have the result, if $d^+(x) \geq k + 1$ or $d^-(y) \geq k + 1$ or $A^+(x) \cap A^-(y) = \emptyset$. Thus we may assume that $x \rightarrow y$, and $d^+(x) = d^-(y) = k$.

Then $X = N^+(x)$ is a reductor of T containing y . We have $2k - 1$ pancyclic arcs in $A^+(x) \cup A^-(y)$ and by Corollary 1, there is at least one pancyclic arc e_1 in $A(X, Out(T - X))$.

If X is not transitive, then according to Lemma 2, there are at least three pancyclic arcs in X , with at most one of them in $A^-(y)$. And by Proposition 8, there is at most one pancyclic arc in $T - X$. It follows that $p(t) \geq 2k + 3$. Hence we may assume that X is the transitive tournament $TT(v_0, v_1, \dots, v_{k-1})$.

Let $T_1 \rightarrow T_2 \rightarrow \dots \rightarrow T_m$ be a decomposition of $T - X$. By Lemmas 4, 5 and 6, at most one of the T_i is not reduced to the vertex t_i .

Suppose that for some $2 \leq i \leq m-1$, T_i is not reduced to a single vertex. If $|T_i| \geq 4$ then by Lemma 4, there are at least three pancyclic arcs in T_i . Thus $p(T) \geq 2k+3$. Assume now that T_i is a 3-cycle. Two arcs of T_i are pancyclic by Lemma 5. Let t_i be a vertex of T_i . By Lemma 10, if $t_i \rightarrow v_0$, then (t_{i-1}, t_i) is pancyclic and if $t_i \leftarrow v_0$, then (t_i, t_{i+1}) is pancyclic. Hence, $p(T) \geq 2k+3$.

Suppose now that T_1 is not reduced to a single vertex. By Lemma 7, we have the result if $h(T_1) \geq 4$. So we may assume that $T_1 \in \mathcal{P}_3$. Also by Lemma 7, there are two pancyclic arcs in T_1 . According to Proposition 7 (dual), there are two distinct vertices w_1 and w_2 of T_1 such that $v_{k-1} \rightarrow w_1$ and $v_{k-2} \rightarrow w_2$. Let C_1 be a hamiltonian cycle of T_1 .

- Suppose first that $m \geq 3$. If a vertex t_i with $2 \leq i \leq m-1$ dominates an element of X then by Lemma 10 one of the arcs (t_j, t_{j+1}) with $i \leq j \leq m-1$ is pancyclic. Hence $p(T) \geq 2k+3$. So we may assume that $T - [T_1, X] \rightarrow X$. Let us prove that (v_{k-2}, w_2) is pancyclic : it is contained in the hamiltonian cycle $(P_2, t_m, v_{k-1}, P_1, t_2, t_3, \dots, t_{m-1}, v_0, v_1, \dots, v_{k-2}, w_2)$, where P_1 (resp. P_2) is the subpath along C_1 of T_1 starting in w_1 (resp. w_2) and ending in the predecessor of w_2 (resp. w_1). For $3 \leq l \leq n-1$, let $0 \leq k' \leq k-2$, $0 \leq m' \leq m-2$ and $0 \leq l_1 \leq n_1-1$ such that $l_1+k'+m'+3 = l$. There is a path Q_1 of length l_1 starting in w_2 . Thus (v_{k-2}, w_2) is contained in the l -cycle $(v_{k-2}, Q_2, t_{m-m'}, t_{m-m'+1}, \dots, t_m, v_{k-2-k'}, v_{k-1-k'}, \dots, v_{k-2})$. Hence T has at least $2k+3$ pancyclic arcs.

- Suppose now that $m = 2$. Since $d_X^-(v_0) \leq k-2$, v_0 is dominated by at least one vertex in T_1 . Hence there exists a vertex w_0 of T_1 dominated by v_0 such that its predecessor w_0^- along C_1 dominates v_0 .

Because $d_X^+(v_{k-1}) \leq k-2$, v_{k-1} dominates at least two vertices of T_1 , so at least one, say w_1 , distinct of w_0 .

Let us now prove that (v_0, w_0) is pancyclic. For $3 \leq l \leq n_1+1$, let Q_0 be a path in T_1 starting at w_0 of length $l-3$. Then (v_0, Q_0, t_2, v_0) is an l -path. Let Q_1 (resp. Q_2) be the subpath of C_1 starting at w_1 (resp. w_0) and ending in w_0^- (resp. the predecessor of w_1) along C_1 . Then for $n_1+2 \leq l \leq n$, $(Q_1, v_0, Q_2, t_2, v_{n+1-l}, v_{n-l}, \dots, v_{k-1}, w_1)$ is an l -path containing (v_0, w_0) .

Hence T has at least $2k+3$ pancyclic arcs.

Suppose now that $T - X$ is the transitive tournament $TT(t_1, t_2, \dots, t_{n-k})$.

Let i_0 be the smallest integer $i > 1$ such that $v_{k-1} \rightarrow t_i$.

- Suppose that $i_0 = 2$. Since T is 2-strong, v_0 is dominated by vertex t_{i_1} distinct from t_{n-k} . By Lemma 1, (v_{k-1}, t_2) and (v_{k-2}, t_1) are respectively $(T - t_1)$ - and $(T - v_{k-1})$ -pancyclic, thus they are contained in l -cycle for any $3 \leq l \leq n-1$. And they both are in the following hamiltonian cycle : $(v_{k-1}, t_2, t_3, \dots, t_{i_1}, v_0, v_1, \dots, v_{k-2}, t_1, t_{i_1+1}, t_{i_1+2}, \dots, t_{n-k}, v_{k-1})$. Thus they both are pancyclic in T . Moreover, by Proposition 8, there is a T -pancyclic arc in $T - X$. Hence, $p(T) \geq 2k+3$.

- Suppose now that $i_0 > 2$. Then $X' = T[X - v_{-1}, t_1]$ is a reductor of X . And by Lemma 3, (t_1, t_2) is T -pancyclic.

The subtournament $T' = T[t_{i_0}, t_{i_0+1}, \dots, t_{n-k}, v_{k-1}]$ is a strong component of $T - X'$. By Lemma 7, there are $h(T') - 1$ T' -pancyclic arcs in T' . And at most one of them is in $A^+(x)$. Thus, if $h(T') \geq 4$, we obtain $p(T) \geq 2k + 3$. So we may assume that $h(T') = 3$, so $T' \in \mathcal{P}_3$. Since $T' - v_{k-1}$ is transitive, (t_{n-k}, v_{k-1}) and (v_{k-1}, t_{i_0}) are T' -pancyclic. Let e_2 be the third T' -pancyclic arc. By Lemma 6, e_2 is also T -pancyclic because $t_{n-k} = x$ dominates v_0 .

If v_0 is dominated by a vertex of $T' - [x, v_{k-1}]$ then by Lemma 6, (v_{k-1}, t_{i_0}) is pancyclic in T and so $p(T) \geq 2k + 3$.

If v_0 dominates $T' - [x, v_{k-1}]$, it must have an in-neighbour t_{i_2} with $2 \leq i_2 \leq i_0 - 1$. Hence, by Proposition 10, there is a pancyclic arc (t_i, t_{i+1}) with $1 \leq i \leq i_0 - 2$. So, $p(T) \geq 2k + 3$. ■

5 Tournaments with exactly four pancyclic arcs

We now prove a generalization of Lemma 3.

Lemma 10 *Let T be a strong tournament, X a reductor of T and $T_1 \rightarrow T_2 \rightarrow \dots \rightarrow T_m$ be a decomposition of $T - X$.*

- i) For $1 < i < m - 1$, if there exist $t_i \in T_i$ and $t_{i+1} \in T_{i+1}$ such that (t_i, t_{i+1}) is in a 3-cycle then (t_i, t_{i+1}) is pancyclic.*
- ii) If $m \geq 3$ and $T_1 = \{t_1\}$ and there is a vertex $t_2 \in T_2$ such that (t_1, t_2) is in a 3-cycle then (t_1, t_2) is pancyclic.*

Proof. Let $(v_0, v_1, \dots, v_{k-1})$ be a hamiltonian path of X and for $1 \leq i \leq m$, set $n_i = |T_i|$.

i) Let v be the vertex of X such that (t_i, t_{i+1}, v, t_i) is a 3-cycle. Then v belongs to X and then by Proposition 6, dominates a vertex $t \in T_1$. Then $(t_i, t_{i+1}, v, t, t_i)$ is a 4-cycle. Let t_1 be an element of T_1 that is dominated by v_{k-1} and t_m be an element of T_m dominating v_0 . Let $5 \leq l \leq n$. The arc (t_i, t_{i+1}) is contained in a cycle of length l . Indeed, take $0 \leq k' < k$, $0 < l_1 < \sum_{j=1}^i n_j$ and $0 \leq l_2 < \sum_{j=i+1}^m n_j$ such that $k' + l_1 + l_2 + 3 = l$. By Proposition 5, there is a (t_1, t_i) -path P_1 of length l_1 in $T[T_1, \dots, T_i]$, and a (t_{i+1}, t_m) -path P_2 of length l_2 in $T[T_{i+1}, \dots, T_m]$. Then $(t_i, P_2, v_0, v_1, \dots, v_{k'}, P_1)$ is an l -cycle containing (t_i, t_{i+1}) . Hence (t_i, t_{i+1}) is pancyclic.

ii) Let $4 \leq l \leq n$. Let us prove that (t_1, t_2) is contained in an l -cycle. By Proposition 6, $v_{k-1} \rightarrow t_1$. Let t_m be an element of T_m dominating v_0 . Since $m \geq 3$, $T - [X, t_1]$ is reducible and t_2 is one of its outgenerators and t_m one of its ingenerators. Therefore, by Proposition 5, there is a (t_1, t_m) -path P of length $l - 3$ in $T - [X, t_1]$. Then $(P, v_0, v_1, \dots, v_{k-1}, t_1, t_2)$ is the desired l -cycle. ■

Definition 2 A tournament is in \mathcal{P}_4 if there is a vertex v such that $T - v$ is the transitive tournament $TT[t_1, t_2, \dots, t_m]$ and three integers $1 < i_1 < i_2 < i_3 \leq m$ such that $v \rightarrow t_j$ if and only if $1 \leq j < i_1$ or $i_2 \leq j < i_3$.

Theorem 7 A tournament has exactly 4 pancyclic arcs if and only if it is in \mathcal{P}_4 .

Proof. It is easy to check that every tournament of \mathcal{P}_4 has exactly four pancyclic arcs.

Let T be a tournament with exactly 4 pancyclic arcs. By Corollary 3, T is (= 1)-strong. Let $\{v\}$ be a reductor of X and $T_1 \rightarrow T_2 \rightarrow \dots \rightarrow T_m$ be a decomposition of $T - v$. By Equation 2, at most one of the $T - i$, $1 \leq i \leq m$ is not reduced to a single vertex.

Let us prove that there exists a decomposition such that each T_i is reduced to a single vertex.

1. If $T - v$ is a transitive tournament, then by Lemma 3, T is in \mathcal{P}_4 .
2. Suppose that there exist $1 < i < m$, such that T_m is not reduced to a vertex. By Corollary 1, (t_m, v) and (v, t_1) are pancyclic. By Equation 2, then $|T_i| = 3$. If $T_i \rightarrow v$ or $v \rightarrow T_i$ then the three arcs of T_i are pancyclic, then T has 5 pancyclic arcs. This is a contradiction. Then there is a vertex $t_i \in T_i$ such that $t_i \rightarrow v$. Thus there is an arc (t_j, t_{j+1}) with $1 \leq j < i$ such that (v, t_j, t_{j+1}, v) is a 3-cycle. So by Lemma 10, (t_j, t_{j+1}) is pancyclic. Again T has 5 pancyclic arcs which is a contradiction.
3. Suppose that T_m is not reduced to a vertex. By Equation 2, v has a unique inneighbour u in T_m . Set $T_i = \{t_i\}$ for $1 \leq i \leq m-1$. By Corollary 1, (u, v) and (v, t_1) are pancyclic and by Lemma 6 dual, there are two pancyclic arcs in T_m . Therefore, there is no other pancyclic arcs in T . Thus according to Lemma 10, $v \rightarrow \{t_i, 1 \leq i \leq m-1\}$. Then u is a reductor. And there are 3 pancyclic arcs which are not in $T_m - u$. Then by Lemmas 4, 5 and 7, $T - u$ is a transitive tournament. So we have the result by Case 1.

■

6 Tournaments with exactly five pancyclic arcs

Definition 3 A tournament is in \mathcal{P}_5 if there is a vertex v such that $T - v$ is the transitive tournament $TT[t_1, t_2, \dots, t_m]$ and three integers $1 < i_1 < i_2 < i_3 < i_4 < i_5 \leq m$ such that $v \rightarrow t_j$ if and only if $1 \leq j < i_1$ or $i_2 \leq j < i_3$ or $i_4 \leq j < i_5$.

The tournament $Q(n)$ ($n \geq 5$) is the tournament constructed from the transitive tournament $TT_{n-4} = TT(t_1, t_2, \dots, t_{n-4})$ and four vertices a, b, c and v such that (a, b, c, a) is a 3-cycle, $v \rightarrow TT_{n-4}$, $TT_{n-4} \rightarrow \{a, b, c\}$, $\{a, b\} \rightarrow v$ and $c \leftarrow v$.

Proposition 9

$$p(Q(5)) = 5$$

For $n \geq 6$, $p(Q(n)) = 6$.

Proof. v is a reductor of T . By Corollary 1, (a, v) , (b, v) and (v, t_1) are pancyclic. And By Lemma 6, (a, b) and (c, a) are pancyclic. Thus $p(Q(n)) \geq 5$.

Now an arc in TT_{n-4} is not pancyclic in T because it is contained in no 3-cycle ; (b, c) is not pancyclic because it is contained in no 4-cycle ; and (v, c) is not pancyclic because it is contained in no l -cycle for $l \geq 5$. For $1 \leq i < n - 4$ the arc (t_i, x) with $x \in \{a, b, c\}$ is not pancyclic because it is contained in no hamiltonian cycle. The arc (t_{n-4}, a) is not pancyclic since it is contained in no hamiltonian cycle and (t_{n-4}, c) is not pancyclic since it is contained in no 3-cycle

For $3 \leq l \leq n - 2$, the arc (t_{n-4}, b) is contained in the l -cycle $(t_{n-4}, b, v, t_{n-1-l}, \dots, t_{n-4})$. For $5 \leq l \leq n$, the arc (t_{n-4}, b) is contained in the l -cycle $(t_{n-4}, b, c, a, v, t_{n+1-l}, \dots, t_{n-4})$. Thus, if $n \geq 6$, $p(Q(n)) = 6$. It is easy to check that if $n = 5$, the arc (t_1, b) is contained in no 4-cycle. Thus $p(Q(5)) = 5$. \blacksquare

Theorem 8 *A tournament has exactly 5 pancyclic arcs if and only if it is in \mathcal{P}_5 or is $Q(5)$.*

Proof. It is easy to check that every tournament of $\mathcal{P}_5 \cup \{Q(5)\}$ has exactly five pancyclic arcs.

Let T be a tournament with exactly 5 pancyclic arcs. By Corollary 3, T is $(= 1)$ -strong. By Equation 2, at most one of the T_i , $1 \leq i \leq m$ is not reduced to a single vertex.

- If $T - v$ is a transitive tournament. Then by Lemma 3, T is in \mathcal{P}_5 .
- Suppose that there exist $1 < i < m$, such that T_m is not reduced to a vertex. By Corollary 1, (t_m, v) and (v, t_1) are pancyclic. By directionnal duality, we may suppose that there is a vertex $t_i \in T_i$ such that $t_i \rightarrow v$. Then by Lemma 10, there is a pancyclic arc in $T[t_1, t_2, \dots, t_i]$. Thus, there are at most two pancyclic arc in T_i . Then by Lemma 4 and the proof of Lemma 5, T_i is a 3-cycle containing 2 pancyclic arcs and there is a vertex $s_i \in T_i$ dominated by v . Then by Lemma 10, there is a pancyclic arc in $T[s_i, t_{i+1}, \dots, t_m]$. So $p(T) \geq 6$ which is a contradiction.
- Suppose that T_m is not reduced to a vertex.

Then $\delta^+ \geq 2$. So by Equation 5, $h(T) \leq 4$ thus by Theorem 1, $T_m \in \mathcal{P}_3$. Let w be a reductor of T_m such that $T_m - w$ is the transitive tournament $TT[a_1, \dots, a_p]$. Let i be the index such that $w \rightarrow a_i$ and $w \leftarrow a_{i+1}$.

$v \rightarrow T_m \setminus \{a_i, w, a_p\}$, otherwise by Lemma 6, $h(T) \geq 5$ which is a contradiction. Also $v \rightarrow T - [v, T_m]$ otherwise by Lemma 10, there is a pancyclic arc in $\{(t_j, t_{j+1}), 1 \leq j \leq m - 2\}$.

By Equation 2, $h_{T_m}^-(v) \leq 2$.

If $h_{T_m}^-(v) = 1$, then let u be the inneighbour of v in T_m . The vertex u is a reductor. And $out(T - u) < out(T - v)$. Iterating the process, we find a reductor v' such that $T - v'$ is transitive or $h_{T_m}^-(v') = 2$. So we may assume that $h_{T_m}^-(v) = 2$.

- Suppose that $N_{T_m}^-(v) = \{a_i, w\}$. If $i \neq 1$, then by Lemma 6 dual, the three arcs (w, a_1) , (a_p, w) and (a_i, a_{i+1}) are pancyclic. So $p(T) \geq 6$ which is a contradiction. Thus $i = 1$. It is easy to see that (a_1, a_2) is contained in every cycle of length l , for $5 \leq l \leq n$. It follows that $|T_m| = 3$. Then $T = Q(n)$ and by Proposition 9, $T = Q(5)$.
- If $N_{T_m}^-(v) = \{w, a_p\}$, then by Lemma 6 dual, (a_p, w) and (a_i, a_{i+1}) are pancyclic. It is easy to see that (w, a_1) is contained in every cycle of length l , for $5 \leq l \leq n$. And because (w, a_1) is pancyclic in T_m , it follows that $|T_m| = 3$. Thus $T = Q(n)$ and by Proposition 9, $T = Q(5)$.
- Suppose that $N_{T_m}^-(v) = \{a_i, a_p\}$. If $i \neq p - 1$, then by Lemma 6 dual, the three arcs (w, a_1) , (a_p, w) and (a_i, a_{i+1}) are pancyclic. So $p(T) \geq 6$ which is a contradiction. Thus $i = p - 1$. It is easy to see that (a_p, w) is contained in every cycle of length l , for $5 \leq l \leq n$. And because (a_p, w) is pancyclic in T_m , it follows that $|T_m| = 3$. So $T = Q(n)$ and by Proposition 9, $T = Q(5)$.

■

References

- [1] B. Alspach, Cycles of each length in regular tournaments, *Canad. Math. Bull.* **10** (1967), 283–286.
- [2] P. Camion, Chemins et circuits hamiltoniens des graphes complets, *C.R. Acad. Sci. Paris* **249** (1959), 2151–2152. , *J. Combin. Inform. System Sci.*, **19** (3-4)(1994), 207–214.
- [3] J. W. Moon, *Topics on Tournaments*, (Holt, Rinehart and Winston, New York) (1968).
- [4] J. W. Moon, On k -cyclic and pancyclic arcs in strong tournaments, *J. Combin. Inform. System Sci.*, **19** (3-4)(1994), 207–214.
- [5] T. Yao, Y. Guo and K. Zhang, Pancyclic out-arcs of a vertex in tournaments, *Discrete Appl. Math.*, **99** (1-3)(2000), 245–249.



Unité de recherche INRIA Sophia Antipolis

2004, route des Lucioles - BP 93 - 06902 Sophia Antipolis Cedex (France)

Unité de recherche INRIA Lorraine : LORIA, Technopôle de Nancy-Brabois - Campus scientifique
615, rue du Jardin Botanique - BP 101 - 54602 Villers-lès-Nancy Cedex (France)

Unité de recherche INRIA Rennes : IRISA, Campus universitaire de Beaulieu - 35042 Rennes Cedex (France)

Unité de recherche INRIA Rhône-Alpes : 655, avenue de l'Europe - 38330 Montbonnot-St-Martin (France)

Unité de recherche INRIA Rocquencourt : Domaine de Voluceau - Rocquencourt - BP 105 - 78153 Le Chesnay Cedex (France)

Éditeur

INRIA - Domaine de Voluceau - Rocquencourt, BP 105 - 78153 Le Chesnay Cedex (France)

<http://www.inria.fr>

ISSN 0249-6399