



# Improving the Robustness of Difference of Convex Algorithm in the Research of a Global Optimum of a Nonconvex Differentiable Function Defined on a Bounded Closed Interval

Leslous Fadila, Philippe Marthon, Ouanes Mohand

## ► To cite this version:

Leslous Fadila, Philippe Marthon, Ouanes Mohand. Improving the Robustness of Difference of Convex Algorithm in the Research of a Global Optimum of a Nonconvex Differentiable Function Defined on a Bounded Closed Interval. Applied Mathematical Sciences, 2014, vol. 8 (n 1), pp. 1-12. <10.12988/ams.2014.35258>. <hal-01137376v2>

**HAL Id: hal-01137376**

**<https://hal.archives-ouvertes.fr/hal-01137376v2>**

Submitted on 31 Mar 2015

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.





## Open Archive TOULOUSE Archive Ouverte (OATAO)

OATAO is an open access repository that collects the work of Toulouse researchers and makes it freely available over the web where possible.

This is an author-deposited version published in : <http://oatao.univ-toulouse.fr/>  
Eprints ID : 12886

**To link to this article** : DOI :10.12988/ams.2014.35258  
URL : <http://dx.doi.org/10.12988/ams.2014.35258>

**To cite this version** : Fadila, Leslous and Marthon, Philippe and Mohand, Ouanes *[Improving the Robustness of Difference of Convex Algorithm in the Research of a Global Optimum of a Nonconvex Differentiable Function Defined on a Bounded Closed Interval.](#)* (2014) Applied Mathematical Sciences, vol. 8 (n° 1). pp. 1-12. ISSN 1314-7552

Any correspondance concerning this service should be sent to the repository administrator: [staff-oatao@listes-diff.inp-toulouse.fr](mailto:staff-oatao@listes-diff.inp-toulouse.fr)

# Improving the Robustness of Difference of Convex Algorithm in the Research of a Global Optimum of a Non convex Differentiable Function Defined on a Bounded Closed Interval

**Leslous Fadila**

Laboratory LAROMAD  
Faculty of Science, University Mouloud Mammeri Tizi-Ouzou, Algeria  
famafarach@yahoo.fr

**Philippe Marthon**

ENSEEIH-IRIT, University of Toulouse, France  
Philippe.Marthon@enseeiht.fr

**Ouanes Mohand**

Laboratory LAROMAD  
Faculty of Science, University Mouloud Mammeri Tizi-Ouzou, Algeria  
ouanes-mohand@yahoo.fr

## **Abstract**

In this paper we present an algorithm for solving a DC problem non convex on an interval  $[a, b]$  of  $\mathbb{R}$ . We use the DCA (Difference of Convex Algorithm) and the minimum of the average of two approximations of the function from  $a$  and  $b$ .

This strategy has the advantage of giving in general a minimum to be situated in the attraction zone of the global minimum searched. After applying the DCA from this minimum we certainly arrive at the global minimum searched.

**Keywords:** Optimization DC and DCA, global optimization, non convex optimization

## 1 Introduction

The `fminbnd` function from MATLAB is a standard method for resolution of a real function minimization defined on a bounded closed interval  $[a, b] \subset R$ . It realizes a golden section search and parabolic interpolation. It provides us with only a local minimum, not necessarily global if the function is not unimodal [2], [3], [7].

In this paper, we propose an alternate method based on the decomposition of the function in a difference of convex functions (DC) and the application DCA algorithm [6], [8].

The DCA also generally provides a local minimum not necessarily global (or even a critical point) [9], [10], [11].

As the DCA is very sensitive to the choice of the initial point [5], [12], [1], [4], we propose not to choose a point of departure. Instead, we minimize the average of two approximations of the function from  $a$  and  $b$ .

This strategy has the advantage of giving generally a minimum to be located in the attraction zone of the global minimum searched.

We apply the DCA from the minimum found [12], we arrive certainly to the global minimum searched.

## 2 Problem Formulation

Let us consider the optimization DC problem:

$$(P_{dc}) \iff \min\{f(x) = g(x) - h(x), x \in [a, b]\}$$

$f : R^n \longrightarrow R$  nonconvex

$g : R^n \longrightarrow R$  convex

$h : R^n \longrightarrow R$  convex

We want to solve the problem  $P_{dc}$  by applying the DCA to the minimum of the average of the two approximations of  $f$  from  $a$  and  $b$  (MDC).

### 2.1 Principle of the (MDC) methode

The DCA is very sensitive to the choice of the starting point. [5], [6]

In the case of a minimization a real function defined on  $[a, b]$ , the minimum found when starting from  $a$  will generally be different from that found when




Figure 1: Principle of MDC

starting from b.

We propose not to choose a starting point. Instead we want to minimize the average of two approximations to  $f$  from  $a$  and  $b$  (MDC) let:

$$\min \frac{1}{2}(f_k(x, a) + f_k(x, b))$$

with:

$$f_k(x, a) = g(x) - h'(a)(x - a) - h(a)$$

$$f_k(x, b) = g(x) - h'(b)(x - b) - h(b)$$

This strategy has the advantage of providing in general a minimum to be located in the attraction zone of the minimum global searched as illustrated by the following example:

## 2.2 The principle of DCA

Note that DCA works only with DC components  $g$  and  $h$  [8] , [12].

At the  $k$ -th iteration of DCA,  $h$  is replaced by its affine minorant  $h_k(x) = h(x^k) + \langle x - x^k, y^k \rangle$  in the neighborhood of  $x^k$ .

Knowing that  $h$  is a convex function, we have therefore  $h(x) \geq h_k(x), \forall x \in X$ .

As a result,  $g(x) - [h(x^k) + \langle x - x^k, y^k \rangle] \geq g(x) - h(x), \forall x \in X$ .

That is to say,  $g(x) - [h(x^k) + \langle x - x^k, y^k \rangle]$  is a majorant function of function  $f(x)$ .

Indeed, the surface of  $f^k$  can be imagined as a bowl being placed directly above the surface of  $f$ ; Moreover, the two surfaces touch at point  $(x^k, f(x^k))$




Figure 2: Principle of DCA

### 2.3 Proposition 1:

Let  $g, h$  two convex functions, differentiable on  $X$ . Putting  $f^k(x) = g(x) - [h(x^k) + \langle x - x^k, y^k \rangle]$ . we have [10] :

- $f^k(x) \geq f(x), \forall x \in X$ .
- $f^k(x^k) = f(x^k)$ .
- $\nabla f^k(x^k) = \nabla f(x^k)$ .

### 2.4 Proof:

Knowing that  $h$  is convex,  $h(x) \geq h_k(x), \forall x \in X$ .

Consequently,  $f^k(x) = g(x) - [h(x^k) + \langle x - x^k, y^k \rangle] \geq g(x) - h(x) = f(x), \forall x \in X$ , that is to say:  $f^k(x) = g(x) - [h(x^k) + \langle x - x^k, y^k \rangle] \geq g(x) - h(x) = f(x), \forall x \in X$ , that is to say  $f^k(x) \geq f(x), \forall x \in X$ .

$$f^k(x) = g(x^k) - [h(x^k) + \langle x^k - x^k, y^k \rangle] = g(x^k) - h(x^k) + 0 = f(x^k).$$

$$\nabla f^k(x) = \nabla g(x) - y^k.$$

Since  $g$  is differentiable, we have:  $y^k = \nabla h(x^k)$ . This is why  $\nabla f^k(x) = \nabla g(x) - \nabla h(x^k)$ .

Finally, we obtain  $\nabla f^k(x^k) = \nabla g(x^k) - \nabla h(x^k) = \nabla f(x^k)$ .

## 2.5 Proposition 2:

1. Sequences  $\{g(x^k) - h(x^k)\}$  and  $\{h^*(y^k) - g^*(y^k)\}$  decrease and tend to the same limit  $\beta$  which is higher than or equal to the global optimal value  $\alpha$ .
2. If  $(g - h)(x^{k+1}) = (g - h)(x^k)$  the algorithm stops at iteration  $k+1$ , and the point  $x^k$  (respectively  $y^k$ ) is a critical point of  $g-h$  (resp.  $h^* - g^*$ ).
3. If the optimal value of Problem (P) is finite and if Sequences  $\{x^k\}$  and  $\{y^k\}$  are bounded, then any value of adherence  $x^*$  of Sequence  $\{x^k\}$  (respectively  $y^*$  of Sequence  $\{y^k\}$ ) is a critical point of  $g-h$  (resp.  $h^* - g^*$ ). [10]

### 2.5.1 Remark:

Thanks to Proposition 2, DCA stops if at least one of Sequences  $\{(g - h)(x^k)\}$ ,  $\{(h^* - g^*)(y^k)\}$ ,  $\{x^k\}$ ,  $\{y^k\}$  converges. In practice, we often use the following stop conditions:

- $|(g - h)(x^{k+1}) - (g - h)(x^k)| \leq \epsilon$ .
- $\|x^{k+1} - x^k\| \leq \epsilon$ .

## 2.6 Properties of DCA

**1-** The DCA constructs a sequence  $\{x^k\}$  such that Sequence  $\{f(x^k)\}$  is decreasing. This can be easily verified on the figure 2, because  $x^{k+1}$  is a minimum of  $f^k$  ( therefore  $f^k(x^k) \geq f^k(x^{k+1})$  and  $f(x^{k+1}) \leq f^k(x^{k+1})$ , as  $f^k(x^k) = f(x^k)$ . Finally, we have  $f(x^k) \geq f^k(x^{k+1}) \geq f(x^{k+1})$ . This shows that Sequence  $\{f(x^{k+1})\}$  is decreasing.

**2-** For the boundedness and convergence of DCA, knowing that  $f: R^n \rightarrow R \cup \{+\infty\}$  is bounded from below in  $R^n$ , Sequence  $\{f(x^k)\}$  is also bounded from below. We know that a sequence  $\{f(x^k)\}$  that is decreasing and bounded from below is convergent.

**3-** When DCA converges to a point  $x^*$ , this point must be a generalized KKT point. this can be easily understood with the help for Figure.2. If we start DCA from a generalized KKT point of  $f$  ( $x^*$  the figure), then DCA stops immediately at this point because it is also a minimum of  $f^*$  ( the convex majorant function defined at point  $x^*$  by  $f(x^*) = g(x) - [h(x^*) + \langle x - x^*, y^* \rangle]$ ,  $y^* \in \partial h(x^*)$ ).

**4-** It can be seen, thanks to the figure that DCA has the option to skip certain neighborhoods of local minima. For example, DCA jumps a local minimum between  $x^k$  and  $x^{k+1}$  then arrives at a neighborhood of the global solution. Though we can not always ensure that this phenomenon accure, we can understand that the performance of DCA is likely to depend on the DC decomposition DC and on the position of the initial point.



### 3 DCA (DC Algorithm)

Initial Step:  $x^0$  given,  $k=0$ .

Step 1: We search  $y^k \in \partial h(x^k)$ .

Step 2: We determine  $x^{k+1} \in \partial g^*(y^k)$ .

Step 3: If the stop conditions are satisfied then DCA is terminated; Otherwise  $k=k+1$  and we repeat the Step 1.

### 4 Application of DCA:

#### 4.1 Example 1:

$$(P) \iff \begin{cases} f(x) = (x-1)(x-2)(x+1)(x+2) \longrightarrow \text{Min} \\ x \in [-3, +3] \end{cases}$$

DCA is applied from Point 3 and Point -3, Next we look for the minimum of the two minima in order to find the global minimum.

$$g(x) = x^4 + 4, h(x) = 5x^2, h'(x) = 10x, y^k = \nabla h(x^k)$$

$$\text{If } x^0 = 3, k = 0, y^0 = 30$$

Step 1

$$x^1 = 1,96, y^1 = 19,6$$

Step 2

$$x^2 = 1,7, y^2 = 17$$

Step 3

$$x^3 = 1,58, y^3 = 15,8$$

$$\text{such as } |x^3 - x^2| = |1,58 - 1,7| < \epsilon$$

Then  $x=1,58$  is the optimal solution of the problem (global minimum) with:

$$f(1,58) = -2,25$$

$$\text{If } x^0 = -3, k = 0, y^0 = -30$$

Step 1

$$x^1 = -1,96, y^1 = -19,6$$

Step 2

$$x^2 = -1,7, y^2 = -17$$

Step 3

$$x^3 = -1,58, y^3 = -15,8$$

$$\text{As we have } |x^3 - x^2| = |-1,58 + 1,7| < \epsilon$$

Then  $x=-1,58$  is the optimal solution of the problem (global minimum) with:

$$f(-1,58) = -2,25.$$




Figure 3: Graphical Solution of Example 1 (DCA)

## 4.2 Remark:

Since  $f(-1,58)=f(1,58)$ , applying the DCA from -3 or from 3, we arrive at the global minimum because function  $f$  is symmetrical.

The standard function `fminbnd` of MATLAB gives the same solution  $x=1,58$  (global minimum)

## 4.3 Exmpl 2:

$$(P) \iff \begin{cases} f(x) = (x - 1, 2)(x - 1, 8)(x + 1)(x + 2) \longrightarrow Min \\ x \in [-3, +3] \end{cases}$$

We apply the DCA from Point 3 and Point -3. Next, we look for the minimum of the two minima in order to find the global minimum.

$$g(x) = x^4 + 0, 48x + 4, 32, h(x) = 4, 84x^2, h'(x) = 9, 68x, y^k = \nabla h(x^k)$$

If  $x^0 = 3, k = 0$

Step 3 of DCA gives:

$$x^3 = +1, 58(\text{nonglobal, localminimum})$$

If  $x^0 = -3, k = 0$

Step 3 of DCA gives:

$$x^3 = -1, 58(\text{global minimum}).$$

`Fminbnd` function of MATLAB gives the solution  $x=1.58$  (non global, local minimum)




Figure 4: Graphical Solution of Example 2 (DCA)

#### 4.4 remark:

In this example we have two minima: One is global and the other is local. So we are Compiled to take the minimum of the two, which is a global minimum.

The solution of example 2 is:

$x = -1,58$  (minimum global)

The solution of Example 2 with the standard function `fminbnd` of MATLAB is:

$x = +1,58$  (local minimum).

#### 4.5 Example 3:

$$(P) \iff \begin{cases} f(x) = (x - 1)(x - 2)(x + 1.8)(x + 1.2) \longrightarrow Min \\ x \in [-3, +3] \end{cases}$$

$$g(x) = x^4 - 0,48x + 4,32, \quad h(x) = 4,84x^2, \quad h'(x) = 9,68x$$

$$y^k = \nabla h(x^k)$$

$$\text{If } x^0 = +3$$

Step 3 of DCA gives the solution  $x = +1,58$  (global minimum).

$$\text{If } x^0 = -3$$

Step 3 of DCA gives the solution  $x = -1,58$  (local minimum).




Figure 5: Graphical Solution of Example 3 with DCA

The standard function `fminbnd` of MATLAB gives the solution  $x=-1,58$  (local minimum).

#### 4.6 Remark:

DCA is very sensitive to the starting point. Then we propose in Example 4 to solve the problem of Example 3 with the proposed method, and to obtain the global solution with a single iteration of DCA.

### 5 Algorithm of the proposed Method (MDCA):

Step 0:  $x^0 = \min_{\frac{1}{2}}(f_k(x, a) + f_k(x, b)), k = 0$ .

Step 1: Application of DCA from  $x^0$ .

### 6 Application of the proposed algorithm (MDCA)

#### 6.1 Example 4:

$$(P) \iff \begin{cases} f(x) = (x - 1)(x - 2)(x + 1.8)(x + 1.2) \longrightarrow Min \\ x \in [-3, +3] \end{cases}$$

$$g(x) = x^4 - 0,48x + 4,32, h(x) = 4,84x^2, h'(x) = 9,68x$$

$$y^k = \nabla h(x^k)$$

We aim to minimize the average of the two approximations of  $f$  from  $-3$  and from  $+3$ ,  $(f_k(x, -3), f_k(x, +3))$  with:

$$f_k(x, -3) = g(x) - h'(-3)(x + 3) - h(-3)$$

$$f_k(x, +3) = g(x) - h'(3)(x - 3) - h(3)$$

Therefore:

$$f_k(x, -3) = x^4 + 28,56x + 47,88$$

$$f_k(x, +3) = x^4 - 29,52x + 47,88$$

Step:0

Solve the problem convex  $(P')$  following:

$$(P') \iff \min \frac{1}{2} [f_k(x, -3) + f_k(x, +3)]$$

$x=0,48$  is the solution of Problem  $(P')$

Step:1

Application of DCA from  $x=0,48$

$$x^0 = 0,48, k = 0, y^0 = 4,64$$

$x^1$  is the solution of the convex problem:

$$\min \{x^4 - 16,11x + 5,44\}$$

$x=1,58$  is the optimal solution (global minimum) of Problem  $(P)$

While the standard function `fminbnd` of MATLAB gives a local minimum ( $x=-1,58$ ).

## 6.2 Remark:

Using the proposed method we did not chose a starting point, instead we want to minimize the average of two approximations of  $f$  from  $3$  and from  $-3$  which will provide a minimum located in the attraction zone of the global minimum. DCA is applied from this minimum, which gives the global minimum searched.

## 7 Conclusion

We dealt in our work with a particular class of optimization problems, namely: non convex problems (DC).

The strategy of minimizing the average followed by the standard application of DCA has led to the production of the global minimum of function  $f$ , while the standard function `fminbnd` of MATLAB found a non global, local minimum.

It now remains to test other examples to better evaluate the pertinence of this strategy, reinforcing the importance of DCA in solving non convex problems.




Figure 6: Solution of Example 4 (DCA)




Figure 7: Solution of Example 4 (MDCA)

## References

- [1] H.A. Le Thi, N.V. Vinh, et T. Phan Din: A Combined DCA and New Cutting Plane Techniques for Globally Solving Mixed Linear Programming, SIAM Conference on Optimization, Stockholm (2005).
- [2] Hartman P.: On functions representable as a difference of convex functions. Pacific J. Maths. 9, 707-713, (1959).
- [3] Horst R., Hoang T.: Global Optimization : Deterministic Approaches, 2nd revised edition. Springer, Berlin,(1993).
- [4] Le The H.A: Solving large scale molecular distance geometry problems by a smoothing technique via the gaussian transform and d.c. programming, Journal of Global Optimization, Vol. 27(4), 375-397, (2003)
- [5] Nguyen Canh Nam, Le Thi Hoai An, Phan Din Tao : A Branch and Bound Algorithm Based on DC Programming and DCA for Strategic Capacity Planning in supply Chain Design for a New Market Opportunity. Operations rechearch proceedings (2006).
- [6] Niu Y.S., Pham Din T.: A DC Programming Approach for Mixed-Integer Linear Programs. Computation and Optimization in Information Systems and Management Sciences, Book Series: Communications in Computer and Information Science, Springer Berlin Heidelberg, 244-253, (2008).
- [7] Parpas P., Rustem B.: Global optimization of the scenario generation and portfolio selection problems, Computational Science and Its Applications,

Lecture folio selection problems, Computational Science and Its Applications, Lecture Notes in Computer Science, Vol. 3982, 908-917, (2006).

- [8] Pham Dinh T.: Algorithms for solving a class of non convex optimization problems. Methods of subgradients. Fermat days 85. Mathematics for optimization, J.B. Hiriart Urruty (ed.), Elsevier Science Publishers, B.V. North-Holland, (1986).
- [9] Pham Din T.: Duality in d.c (difference of convex functions) optimization. Subgradient methods. Trends in Mathematical Optimization, K.H. Hoffmann et al. (ed.), International Series of Numer Math.84, Birkhauser, (1988)
- [10] Pham Dinh Tao and Le Thi Hoai An, Stabilité de la dualité lagrangienne en optimisation d.c. (différence de deux fonctions convexes). C.R.Acad. Paris, t.318,SerieI, (1994).
- [11] Pham Dinh T., Le Thi H.A., Convex analysis approach to D.C programming: Theory, Algorithms and Application. Acta Mathematica Vietnamica, Vol.22 NO .1, 289-355, (1997).
- [12] Pham Dinh T., Le Thi H.A.: DC Programming. Theory, Algorithms, Applications: The state of the Art. First International Whorkshop on global Constrained Optimization and Constraint Satisfaction, Nice, October 2-4, (2002)