



# Uniqueness in a class of Hamilton-Jacobi equations with constraints

Sepideh Mirrahimi, Jean-Michel Roquejoffre

## ► To cite this version:

Sepideh Mirrahimi, Jean-Michel Roquejoffre. Uniqueness in a class of Hamilton-Jacobi equations with constraints. 2015. <hal-01116483>

**HAL Id: hal-01116483**

**<https://hal.archives-ouvertes.fr/hal-01116483>**

Submitted on 13 Feb 2015

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Uniqueness in a class of Hamilton-Jacobi equations with constraints

Sepideh Mirrahimi <sup>a</sup>, Jean-Michel Roquejoffre <sup>a</sup>

<sup>a</sup>*Institut de Mathématiques (UMR CNRS 5219), Université Paul Sabatier, 118 Route de Narbonne, 31062 Toulouse Cedex, France*

Received \*\*\*\*\*; accepted after revision +++++

Presented by

---

## Abstract

In this note, we discuss a class of time-dependent Hamilton-Jacobi equations depending on a function of time, this function being chosen in order to keep the maximum of the solution to the constant value 0. The main result of the note is that the full problem has a unique classical solution. The motivation is a selection-mutation model which, in the limit of small diffusion, exhibits concentration on the zero level set of the solution of the Hamilton-Jacobi equation. The uniqueness result that we prove implies strong convergence and error estimates for the selection-mutation model. *To cite this article: S. Mirrahimi, J.-M. Roquejoffre, C. R. Acad. Sci. Paris, Ser. I 340 (2015).*

## Résumé

**Unicité pour une classe d'équations de Hamilton-Jacobi avec contraintes.** Dans cette note, on discute une classe d'équations de Hamilton-Jacobi dépendant du temps, et d'une fonction inconnue du temps choisie pour que le maximum de la solution de l'équation de Hamilton-Jacobi prenne tout le temps la valeur 0. Le résultat principal de cette note est que le problème complet admet une unique solution classique. La motivation est un modèle de sélection-mutation qui, dans la limite d'une diffusivité nulle, présente une concentration sur la ligne de niveau 0 de la solution de l'équation de Hamilton-Jacobi. Le résultat d'unicité que nous démontrons implique une convergence forte avec estimations d'erreur pour le modèle de sélection-mutation. *Pour citer cet article : S. Mirrahimi, J.-M. Roquejoffre, C. R. Acad. Sci. Paris, Ser. I 340 (2015).*

---

*Email addresses:* [sepideh.mirrahimi@math.univ-toulouse.fr](mailto:sepideh.mirrahimi@math.univ-toulouse.fr) (Sepideh Mirrahimi),  
[jean-michel.roquejoffre@mth.univ-toulouse.fr](mailto:jean-michel.roquejoffre@mth.univ-toulouse.fr) (Jean-Michel Roquejoffre).

## Version française abrégée

On présente dans cette note un résultat d'unicité pour le problème de Hamilton-Jacobi suivant, d'inconnues  $(u(t, x), I(t))$  :

$$\begin{cases} \partial_t u = |\nabla u|^2 + R(x, I), & (t > 0, x \in \mathbb{R}^d), \quad \max_x u(t, x) = 0, \\ I(0) = I_0 > 0, \quad u(0, x) = u_0(x). \end{cases} \quad (1)$$

La donnée  $R(x, I)$  vérifie des hypothèses de stricte concavité par rapport à  $x$  et de monotonie par rapport à  $I$  explicites plus bas. La donnée initiale  $u_0$  vérifie aussi des hypothèses spéciales de concavité. Ainsi, la fonction  $I(t)$  doit être choisie pour que la solution  $u(t, x)$  de l'équation de Hamilton-Jacobi ait, à chaque instant, un maximum égal à 0.

Nous avons alors le

**Theorem 0.1** *On choisit  $R \in C^2$ , et on suppose l'existence de  $I_M > 0$  tel que  $\max_{x \in \mathbb{R}^d} R(x, I_M) = 0 = R(0, I_M)$ . De plus,  $R$  est supposée strictement concave et, pour  $|x|$  grand, comprise entre deux paraboles. La donnée initiale  $u_0$  est également à décroissance quadratique et strictement concave.*

*Le problème (1) a une unique solution  $(u, I)$ , où  $u$  est une solution classique de l'équation de Hamilton-Jacobi. De plus  $u \in L_{loc}^\infty(\mathbb{R}^+; W_{loc}^{3,\infty}(\mathbb{R}^d)) \cap W_{loc}^{1,\infty}(\mathbb{R}^+; L_{loc}^\infty(\mathbb{R}^d)) \times W^{1,\infty}(\mathbb{R})$ .*

L'existence pour (1) a été démontrée en plusieurs endroits, voir par exemple [8] ou [1]. L'unicité est donc notre résultat principal, c'était un problème ouvert. L'unicité était en effet connue seulement pour un cas très particulier (voir [8]).

Le modèle (1) intervient dans la limite  $\varepsilon \rightarrow 0$  des solutions de

$$\partial_t n_\varepsilon - \varepsilon \Delta n_\varepsilon = \frac{n_\varepsilon}{\varepsilon} R(x, I_\varepsilon(t)) \quad (t > 0, x \in \mathbb{R}^d), \quad I_\varepsilon(t) = \int_{\mathbb{R}^d} \psi(x) n_\varepsilon(t, x) dx, \quad (2)$$

où  $n_\varepsilon(t, x)$  est la densité d'une population caractérisée par un trait biologique  $x$   $d$ -dimensionnel. La compétition pour une ressource unique est représentée par  $I_\varepsilon(t)$ ,  $\psi > 0$  régulière donnée. Le terme  $R(x, I)$  est le taux de reproduction. Les hypothèses de concavité sont des hypothèses techniques, mais pertinentes au plan biologique. Par une transformation de Hopf-Cole  $n_\varepsilon = \exp(u_\varepsilon/\varepsilon)$  on se ramène à l'équation sur  $u_\varepsilon$  suivante :

$$\partial_t u_\varepsilon = \varepsilon \Delta u_\varepsilon + |\nabla u_\varepsilon|^2 + R(x, I_\varepsilon) \quad (3)$$

qui, dans la limite  $\varepsilon \rightarrow 0$ , donne l'équation sur  $u$ . On s'attend alors à ce que  $n_\varepsilon$  se concentre aux points où  $u$  est proche de 0. Et, dans cette limite,  $I_\varepsilon$  apparaît comme une sorte de multiplicateur de Lagrange.

La convergence de (3) vers (1) à une sous-suite près est connue depuis [8]. Le Théorème 0.1 donne la convergence de toute la famille, ainsi que des estimations d'erreur. Soit  $x_\varepsilon(t)$  le point où  $u_\varepsilon(t, \cdot)$  atteint son maximum. On suppose l'existence de  $I^0$  tel que  $0 < I^0 \leq I_\varepsilon(0) := \int_{\mathbb{R}^d} \psi(x) n_\varepsilon^0(x) dx < I_M$ , et on suppose

$$n_\varepsilon^0 = e^{u_\varepsilon^0/\varepsilon} = \frac{r}{\varepsilon^{d/2}} e^{u_0/\varepsilon}, \quad \text{with } u_0 \in C^2(\mathbb{R}^d) \quad \text{and} \quad \max_{x \in \mathbb{R}^d} u_0(x) = 0.$$

Le résultat est alors le

**Theorem 0.2** *Soit  $n_\varepsilon$  la solution de (2) et  $u_\varepsilon$  définie par (3). Nous avons les développements asymptotiques suivants*

$$I_\varepsilon = I + \varepsilon I_1 + o(1), \quad x_\varepsilon = \bar{x} + \varepsilon \bar{x}_1 + o(1), \quad u_\varepsilon = u + \varepsilon \log\left(\frac{r}{\varepsilon^{d/2}}\right) + \varepsilon u_1 + o(1).$$

Les termes  $I_1$ ,  $\bar{x}_1$  et  $u_1$  vont être présentés dans [7]. Ce résultat implique le corollaire

**Corollary 0.3** *Nous avons l'approximation suivante pour  $n_\varepsilon$  :*

$$n_\varepsilon(t, x) = \frac{r}{\varepsilon^{\frac{d}{2}}} \left( \exp(u_1 + \frac{u}{\varepsilon}) + o(1) \right).$$

*En particulier, lorsque  $\varepsilon \rightarrow 0$ , toute la suite  $(n_\varepsilon)_\varepsilon$  converge :*

$$n_\varepsilon(t, x) \longrightarrow \bar{\rho}(t) \delta(x - \bar{x}(t)) \quad \text{au sens des mesures,}$$

$$\text{avec } \rho(t) = \frac{I(t)}{\psi(\bar{x}(t))}.$$

En d'autres termes, la population se concentre sur un trait dominant qui évolue avec le temps. On note que la convergence de  $n_\varepsilon$  à une sous-suite prés était déjà établie dans [5].

Tous ces résultats seront détaillés dans [6] et [7].

## 1. Introduction

The purpose of this note is to discuss uniqueness in the following problem, with unknowns  $(I(t), u(t, x))$ :

$$\begin{cases} \partial_t u = |\nabla u|^2 + R(x, I) & (t > 0, x \in \mathbb{R}^d), \quad \max_x u(t, x) = 0, \\ I(0) = I_0 > 0, \quad u(0, x) = u_0(x), \end{cases} \quad (4)$$

where  $I_0 > 0$  and  $u_0$  is a concave, quadratic function:

$$-\underline{L}_0 - \underline{L}_1|x|^2 \leq u_0(x) \leq \bar{L}_0 - \bar{L}_1|x|^2, \quad -2\underline{L}_1 \leq D^2 u_0 \leq -2\bar{L}_1, \quad D^3 u_0 \in L^\infty(\mathbb{R}^d).$$

The constraint on the maximum of  $u(t, \cdot)$  makes the problem nonstandard. Our main result is

**Theorem 1.1** *Choose  $R \in C^2$ , and suppose that there is  $I_M > 0$  such that  $\max_{x \in \mathbb{R}^d} R(x, I_M) = 0 = R(0, I_M)$ .*

*Also assume the following concavity and regularity properties for  $R$ :*

$$\begin{aligned} -\underline{K}_1|x|^2 \leq R(x, I) \leq \bar{K}_0 - \bar{K}_1|x|^2, & \quad \text{for } 0 \leq I \leq I_M, \\ -2\underline{K}_1 \leq D^2 R(x, I) \leq -2\bar{K}_1 < 0 \text{ and } D^3 R(\cdot, I) \in L^\infty(\mathbb{R}^d) & \quad \text{for } 0 \leq I \leq I_M, \\ -\underline{K}_2 \leq \partial_I R \leq -\bar{K}_2, \end{aligned}$$

$$|\partial_{I x_i}^2 R(x, I)| + |\partial^3 R_{I x_i x_j}(x, I)| \leq K_3, \quad \text{for } 0 \leq I \leq I_M, \text{ and } i, j = 1, 2, \dots, d.$$

*Problem (4) has a unique solution  $(u, I)$ , where  $u$  solves the Hamilton-Jacobi equation in the classical sense. Moreover  $u \in L_{loc}^\infty(R^+; W_{loc}^{3, \infty}(\mathbb{R}^d)) \cap W_{loc}^{1, \infty}(R^+; L_{loc}^\infty(\mathbb{R}^d)) \times W^{1, \infty}(\mathbb{R})$ .*

Existence to (4) has been proved in various contexts (see [8, 1, 5]). Thus, our contribution is uniqueness, which has up to now been an open problem. The uniqueness has indeed been known only for a very particular case (see [8]).

The rest of the note is organized as follows. In Section 2, we explain the motivation and, in particular, the meaning of the various assumptions. In Section 3, we revisit existence for (4), which will entail an unconventional ODE formulation for uniqueness. Section 4, which is the main part of the note, provides a fairly complete sketch of the uniqueness proof. In Section 5, we give an application.

## 2. Background and motivation

Model (4) arises in the limit  $\varepsilon \rightarrow 0$  of the solutions to the problem

$$\partial_t n_\varepsilon - \varepsilon \Delta n_\varepsilon = \frac{n_\varepsilon}{\varepsilon} R(x, I_\varepsilon(t)) \quad (t > 0, x \in \mathbb{R}^d), \quad I_\varepsilon(t) = \int_{\mathbb{R}^d} \psi(x) n_\varepsilon(t, x) dx, \quad (5)$$

where  $n_\varepsilon(t, x)$  is the density of a population characterized by a  $d$ -dimensional biological trait  $x$ . The population competes for a single resource, this is represented by  $I_\varepsilon(t)$ , where  $\psi$  is a given positive smooth function. The term  $R(x, I)$  is the reproduction rate; it is, as can be expected, very negative for large  $x$  and decreases as the competition increases. Such models can be derived from individual based stochastic processes in the limit of large populations (see [2]). The concavity assumption on  $R$  is a technical one, although biologically relevant. The Hopf-Cole transformation  $n_\varepsilon = \exp(u_\varepsilon/\varepsilon)$  yields the equation

$$\partial_t u_\varepsilon = \varepsilon \Delta u_\varepsilon + |\nabla u_\varepsilon|^2 + R(x, I_\varepsilon) \quad (6)$$

which, in the limit  $\varepsilon \rightarrow 0$ , yields the equation for  $u$ . Now,  $I_\varepsilon$  being uniformly positive and bounded in  $\varepsilon$ , the Hopf-Cole transformation leads to the constraint on  $u$ . Moreover, one expects that  $n_\varepsilon$  concentrates at the points where  $u$  is close to 0 and the function  $I_\varepsilon$  appears, in the limit, as a sort of Lagrange multiplier.

This approach, based on the Hopf-Cole transformation, to study (5) has been introduced in [4] and then developed in different contexts (see for instance [8,1,3,5]). Long time asymptotics of such models have also been studied in [9] and the references therein.

### 3. Existence

Existence to a solution to (4) is obtained by letting  $\varepsilon \rightarrow 0$  in (6). The main step is the

**Theorem 3.1** (uniform estimates for  $u_\varepsilon$ , [5]) *There exists  $I_m > 0$  such that  $0 < I_m \leq I_\varepsilon(t) \leq I_M + C\varepsilon^2$ . Moreover we have the following estimates on  $u_\varepsilon$*

$$\begin{cases} -\underline{L}_0 - \underline{L}_1 |x|^2 - \varepsilon 2d \underline{L}_1 t \leq u_\varepsilon(t, x) \leq \bar{L}_0 - \bar{L}_1 |x|^2 + (\bar{K}_0 + 2d\varepsilon \bar{L}_1) t, \\ \underline{L}_1 - 2t \underline{K}_1 \leq D^2 u_\varepsilon(t, x) \leq -2\bar{L}_1, \quad \|D^3 u_\varepsilon(t, \cdot)\|_{L^\infty} \leq C(T), \quad \text{for } t \in [0, T]. \end{cases} \quad (7)$$

The bounds for  $u_\varepsilon$  can be obtained for any uniformly bounded function  $I_\varepsilon$ , not only for that of (5). This remark will be an important ingredient of the uniqueness proof.

### 4. Uniqueness

For a given continuous function  $I(t)$  such that  $0 < I(t) < I_M$ , one may construct a solution of  $\partial_t u = |\nabla u|^2 + R(x, I)$  with initial datum  $u_0$ . Just as in Theorem 3.1, this solution satisfies estimates (7). And so,  $u(t, \cdot)$  being strictly concave and quadratically decreasing, there exists a unique function  $\bar{x}(t)$  such that  $u(t, \bar{x}(t)) = \max_{x \in \mathbb{R}^d} u(t, x)$ . Assume that  $I(t)$  is chosen such that  $u(t, \bar{x}(t)) = 0$ . Then, from the equation on  $u$  we deduce that  $R(\bar{x}(t), I(t)) = 0$ . Notice also that, because  $\partial_I R < 0$ , we have  $R(\bar{x}(t), 0) > 0$ . Finally, differentiating  $\nabla u(t, \bar{x}(t)) = 0$  and plugging in the equation for  $u$  we obtain an ODE for  $\bar{x}$ :  $\dot{\bar{x}}(t) = (-D^2 u(t, \bar{x}(t)))^{-1} \nabla_x R(\bar{x}(t), \bar{I}(t))$ .

The idea is thus to change the constrained problem (4) by the following slightly nonstandard differential system:

$$\begin{cases} R(\bar{x}(t), I(t)) = 0, & \text{for } t \in [0, T], \\ \dot{\bar{x}}(t) = (-D^2 u(t, \bar{x}(t)))^{-1} \nabla_x R(\bar{x}(t), \bar{I}(t)), & \text{for } t \in [0, T], \\ \partial_t u = |\nabla u|^2 + R(x, I), & \text{in } [0, T] \times \mathbb{R}^d, \end{cases} \quad (8)$$

with initial conditions

$$I(0) = I_0, \quad u(0, \cdot) = u_0(\cdot), \quad \bar{x}(0) = \bar{x}_0, \quad \text{such that } R(x_0, I_0) = 0. \quad (9)$$

Note that (8) is really a differential system because the assumptions on  $R$  imply that  $I(t)$  can implicitly be expressed in terms of  $\bar{x}(t)$ . And it is slightly nonstandard because  $\bar{x}$  solves an ODE whose nonlinearity depends on  $u$ . Finally, note that, as soon as  $u$  satisfies the concavity and regularity estimates (7), system (8) is equivalent to the constrained problem (4).

This suggests to use a simple fixed point argument to prove uniqueness to (8) (and so, to (4)). Which in turn suggests to set up the following scheme: starting from  $x(t) \in C([0, T], \mathbb{R}^d)$ , such that  $x(0) = x_0$ , where  $R(x_0, 0) > 0$ . Let  $I(t)$  solve  $R(x(t), I(t)) = 0$  on  $[0, T]$  with  $R(x_0, I_0) = 0$ . Let  $v(t, \cdot)$  be the unique solution to

$$\partial_t v = |\nabla v|^2 + R(x, I), \quad v(0, x) = u_0(x). \quad (10)$$

Let  $y(t)$  solve  $\dot{y}(t) = (-D^2 v(t, x(t)))^{-1} \nabla_x R(x(t), I(t))$  on  $[0, T]$  with initial datum  $y(0) = x_0$ . Setting  $y := \Phi(x)$ , we notice that uniqueness is proved as soon as we have proved that  $\Phi$  has a unique fixed point. One additional feature about a solution  $(\bar{I}, u, \bar{x})$  of (8):

**Lemma 4.1** *The function  $\bar{I}(t)$  is increasing.*

We claim that our problem reduces to proving the

**Theorem 4.2** *There exists  $C > 0$  universal and  $\delta > 0$ , which is small as  $R(x_0, 0)$  tends to 0, such that  $\Phi$  is a contraction from  $C([0, \delta], \overline{B}_C(x_0))$  to itself; here  $B_r(a)$  denotes the ball of centre  $a \in \mathbb{R}^d$  and radius  $r > 0$ .*

Note indeed that, by Lemma 4.1, we have, because  $\partial_I R < 0$ :

$$R(\bar{x}(\delta), 0) = R(\bar{x}(\delta), 0) - R(\bar{x}(\delta), \bar{I}(\delta)) \geq c\bar{I}(\delta) \geq cI_0,$$

for some universal  $c > 0$ . Hence Theorem 4.2 can be iterated to yield global existence and uniqueness.

Let us give an overview of the proof of Theorem 4.2. For  $I \in C([0, \delta]; [0, I_M])$ , let  $V(I)$  be the (unique) solution of (10). The main step is the following

**Lemma 4.3** *Let  $I_1, I_2 \in C([0, \delta]; [0, I_M])$ . Then*

$$\|V(I_1) - V(I_2)\|_{W^{2,\infty}([0,\delta] \times \mathbb{R}^d)} \leq C\|I_1 - I_2\|_{L^\infty([0,\delta])}\delta.$$

This lemma, once proved, opens the way to Theorem 4.2. Indeed the equation  $R(x, I) = 0$  yields a smooth mapping  $x \mapsto I$ , and  $I \mapsto V$  is a Lipschitz mapping thanks to Lemma 4.3. Moreover, the equation  $\dot{y}(t) = (-D^2 v(t, x(t)))^{-1} \nabla_x R(x(t), I(t))$  yields a Lipschitz mapping  $v \mapsto y$  by the estimates for  $v$  given by Theorem 3.1.

Lemma 4.3 is more involved. If  $I_1$  and  $I_2$  are as in the assumptions of the lemma, the function  $r = V(I_1) - V(I_2)$  solves

$$\begin{cases} \partial_t r = (\nabla v_1 + \nabla v_2) \cdot \nabla r + R(x, I_2) - R(x, I_1), & \text{in } [0, \delta] \times \mathbb{R}^d \\ r(0, x) = 0, & \text{for all } x \in \mathbb{R}^d. \end{cases} \quad (11)$$

with  $v_i = V(I_i)$ . Note that the above equation has a unique classical solution which can be computed by the method of characteristics. The characteristics solve

$$\dot{\gamma}(t) = -\nabla v_1(t, \gamma) - \nabla v_2(t, \gamma), \quad (12)$$

and, due to the estimates of Theorem 3.1, they exist globally. So, one may successively express  $r$  given by integration along characteristics, and estimate its derivatives recursively.

## 5. Application

Convergence of (6) to (4) had already been proved in [8,1], along subsequences. The uniqueness part of Theorem 1.1 yields the convergence of the full family of solutions  $u_\varepsilon$  of (6), instead of convergence along a subsequence. Moreover it allows an expansion of  $I_\varepsilon$ ,  $u_\varepsilon$  and  $x_\varepsilon$  (the maximum point of  $u_\varepsilon$  at each time) in terms of  $\varepsilon$ . Here are the results.

Assume that there is  $I^0$  such that  $0 < I^0 \leq I_\varepsilon(0) := \int_{\mathbb{R}^d} \psi(x)n_\varepsilon^0(x)dx < I_M$ , and that

$$n_\varepsilon^0 = e^{u_\varepsilon^0/\varepsilon} = \frac{r}{\varepsilon^{d/2}} e^{u^0/\varepsilon}, \quad \text{with } u^0 \in C^2(\mathbb{R}^d) \quad \text{and} \quad \max_{x \in \mathbb{R}^d} u^0(x) = 0.$$

The result is the

**Theorem 5.1** *Let  $n_\varepsilon$  be the solution of (5) and  $u_\varepsilon$  be defined by (6). We have the following asymptotic expansions*

$$I_\varepsilon = I + \varepsilon I_1 + o(1), \quad x_\varepsilon = \bar{x} + \varepsilon \bar{x}_1 + o(1), \quad u_\varepsilon = u + \varepsilon \log\left(\frac{r}{\varepsilon^{d/2}}\right) + \varepsilon u_1 + o(1).$$

The terms  $I_1$ ,  $\bar{x}_1$  and  $u_1$  will be provided in [7]. This yields the corollary

**Corollary 5.2** *We have the following approximation for  $n_\varepsilon$ :*

$$n_\varepsilon(t, x) = \frac{r}{\varepsilon^{d/2}} \left( \exp\left(u_1(t, x) + \frac{u(t, x)}{\varepsilon}\right) + o(1) \right).$$

In particular, as  $\varepsilon \rightarrow 0$ , the whole sequence  $(n_\varepsilon)_\varepsilon$  converges:

$$n_\varepsilon(t, x) \longrightarrow \bar{\rho}(t) \delta(x - \bar{x}(t)), \quad \text{weakly in the sense of measures,}$$

with  $\rho(t) = \frac{I(t)}{\psi(\bar{x}(t))}$ .

In other words, the population density concentrates on a dominant trait which evolves in time. We note that the convergence of  $n_\varepsilon$  along subsequences was already established in [5].

The above results will be detailed in [6] and [7].

## Acknowledgements

S. Mirrahimi was partially funded by the french ANR projects KIBORD ANR-13-BS01-0004 and MOD-EVOL ANR-13-JS01-0009. J.-M. Roquejoffre has received funding from the European Research Council under the European Union's Seventh Framework Programme (FP/2007-2013) / ERC Grant Agreement n. 321186 - ReaDi - "Reaction-Diffusion Equations, Propagation and Modelling" held by Henri Berestycki. He was also partially supported by the French National Research Agency (ANR), within the project NONLOCAL ANR-14-CE25-0013. Both authors are grateful to the Labex CIMI for organizing a PDE-probability quarter in Toulouse, which provided quite a stimulating environment to this collaboration.

## References

- [1] G. BARLES, S. MIRRAHIMI, AND B. PERTHAME, *Concentration in Lotka-Volterra parabolic or integral equations: a general convergence result*. Methods Appl. Anal., **16(3)** (2009) pp.321–340.
- [2] N. CHAMPAGNAT, R. FERRIÈRE, AND S. MÉLÉARD, *Individual-based probabilistic models of adaptive evolution and various scaling approximations*, **59** (2008) Progress in Probability. Birkhäuser.

- [3] N. CHAMPAGNAT AND P.-E. JABIN, *The evolutionary limit for models of populations interacting competitively via several resources*, Journal of Differential Equations, **261** (2011), pp.179–195.
- [4] O. DIEKMANN, P.-E. JABIN, S. MISCHLER, AND B. PERTHAME, *The dynamics of adaptation: an illuminating example and a Hamilton-Jacobi approach*, Th. Pop. Biol., **67(4)** (2005) pp.257–271.
- [5] A. LORZ, S. MIRRAHIMI, B PERTHAME, *Dirac mass dynamics in a multidimensional nonlocal parabolic equation*. Communications in Partial Differential Equations, **36** (2011) pp.1071–1098.
- [6] S. MIRRAHIMI, J.-M. ROQUEJOFFRE, *Uniqueness in a Hamilton-Jacobi equation with constraints*, in preparation.
- [7] S. MIRRAHIMI, J.-M. ROQUEJOFFRE, *Approximation of solutions of selection-mutation models and error estimates*, in preparation.
- [8] B. PERTHAME, G. BARLES, *Dirac concentrations in Lotka-Volterra parabolic PDEs*, Indiana Univ. Math. J. **57** (2008), pp. 3275–3301.
- [9] G. RAOUL, *Etude qualitative et numérique d'équations aux dérivées partielles issues des sciences de la nature*, PhD thesis, ENS Cachan, 2009.