



Means in complete manifolds: uniqueness and approximation

Marc Arnaudon, Laurent Miclo

► To cite this version:

Marc Arnaudon, Laurent Miclo. Means in complete manifolds: uniqueness and approximation. ESAIM: Probability and Statistics, EDP Sciences, 2014, 18, pp.185-206. <hal-00717677>

HAL Id: hal-00717677

<https://hal.archives-ouvertes.fr/hal-00717677>

Submitted on 13 Jul 2012

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

MEANS IN COMPLETE MANIFOLDS: UNIQUENESS AND APPROXIMATION

MARC ARNAUDON AND LAURENT MICLO

ABSTRACT. Let M be a complete Riemannian manifold, $N \in \mathbb{N}$ and $p \geq 1$. We prove that almost everywhere on $x = (x_1, \dots, x_N) \in M^N$ for Lebesgue measure in M^N , the measure $\mu(x) = \frac{1}{N} \sum_{k=1}^N \delta_{x_k}$ has a unique p -mean $e_p(x)$.

As a consequence, if $X = (X_1, \dots, X_N)$ is a M^N -valued random variable with absolutely continuous law, then almost surely $\mu(X(\omega))$ has a unique p -mean. In particular if $(X_n)_{n \geq 1}$ is an independent sample of an absolutely continuous law in M , then the process $e_{p,n}(\omega) = e_p(X_1(\omega), \dots, X_n(\omega))$ is well-defined.

Assume M is compact and consider a probability measure ν in M . Using partial simulated annealing, we define a continuous semimartingale which converges to the set of minimizers of the integral of distance at power p with respect to ν . When the set is a singleton, it converges to the p -mean.

1. INTRODUCTION

Finding the mean of the median or more generally the p -mean e_p of a probability measure in a manifold (the point which minimizes integral with respect to this measure of distance at power p) has numerous applications. There is not much to say for the mean in \mathbb{R}^d , almost the only case where there is a closed formula, and the most important case as the most useful estimator in statistics when the measure is uniform law on a sample. For medians in \mathbb{R}^d the situation is more complicated. Uniqueness holds as soon as the support of the probability measure is not carried by a line. The first algorithm for computing e_1 is due to Weisfeld in [24]. As for the computation of e_∞ (the center of the smallest ball containing the support of the measure), Badöiu and Clarkson gave a fast and simple algorithm in [6]. For many applications in biology, signal processing, information geometry, extension to other spaces is necessary. The median in Hilbert space is computed in [9]. In nonlinear spaces with convexity assumptions, uniqueness has been established in [18] for the mean, [1] for the p -mean. Many algorithms of computation now exist. As far as deterministic algorithms are concerned, one can cite [19], [12], [13], [2] for the mean in Riemannian manifolds, [3] for the mean in Finsler manifolds, [11] and more generally [25] for the median, [5] for e_∞ . Stochastic algorithms avoid to compute the gradient of the functional to minimize. They can be found in [23], [4]. For other functionals to minimize, see [8].

In this paper we investigate the case of non necessarily convex, complete Riemannian manifolds. Our first result (Theorem 2.1) concerns uniqueness of the p -mean of the uniform measure on a finite set $\{x_1, \dots, x_n\}$ of points, almost everywhere on $x = (x_1, \dots, x_n)$ for the Lebesgue measure. This generalizes Bhattacharya and Patangreanu result on the circle ([7], case $p = 2$). See also [10] for more general uniqueness criterions on the circle.

For computation of the p -mean, usual deterministic algorithms are not possible any more, due to the fact that the functional to minimize may have many local minima. So restricting to symmetric spaces we use a simulated annealing method with a continuous stochastic process, together with an estimation of the gradient to minimize via a drift moving faster and faster. With this method we are able to define a process which converges in distribution to the p -mean for $p \in [1, \infty)$ (Theorem 4.3, and Theorem 3.2 for more general but smooth functionals).

The main applications are in signal processing with polarimetric signal, but also for the group of rotations of \mathbb{R}^n , so as to determine averages on rotations. Also this solves many problems of optimization which may arise in economy, decision support, operation research. Notice that on the circle, fast computation of the mean has been performed in [17]. In fact this is a case where a closed formula can be found. For general case the situation is much more complicated and the convergence of our processes is slower and weaker. Jump processes and algorithms related to the continuous processes presented here will be investigated in a forthcoming paper.

2. UNIQUENESS OF p -MEANS FOR UNIFORM MEASURES WITH FINITE SUPPORT

Let M be a d -dimensional complete Riemannian manifold with Riemannian distance denoted by ρ . For ν a probability measure on M and $p \geq 1$, we define

$$(2.1) \quad \begin{aligned} H_{p,\nu} : M &\rightarrow \mathbb{R}_+ \cup \{+\infty\}, \\ y &\mapsto \int_M \rho^p(y, z) \nu(dz). \end{aligned}$$

Either $H_{p,\nu} \equiv \infty$ or for all $y \in M$, $H_{p,\nu}(y) < \infty$. In the latter case we denote by $Q_{p,\nu}$ the set of minimizers of $H_{p,\nu}$. When $Q_{p,\nu}$ has only one element we denote it by $e_{p,\nu}$ and call it the p -mean of ν . When there is no possible confusion we let $e_p = e_{p,\nu}$. For $x = (x_1, \dots, x_N) \in M^N$, we let

$$(2.2) \quad \mu(x) = \frac{1}{N} \sum_{k=1}^N \delta_{x_k}.$$

Clearly $H_{p,\mu(x)}$ is finite.

Theorem 2.1. *Assume $p > 1$ or $\{d > 1 \text{ and } N > 2\}$. For almost all $x \in M^N$, $Q_{p,\mu(x)}$ has a unique element $e_{p,\mu(x)}$*

Remark 2.2. This theorem extends Theorem 4.15 in [26] where the same result has been established for $p = 1$ and M compact.

Proof. We begin with the case $p > 1$.

Since $\mu(x)$ has a finite support, we can assume that M is a compact Riemannian manifold. For this a smooth modification outside a large ball is sufficient. For instance we can choose a radius so that the boundary is smooth, double the ball and finally smoothen the metric locally around the place where the pasting has been performed.

So in the sequel we will assume that M is compact, with diameter L . For $y \in M$ we denote by $S_y M \subset T_y M$ the set of unit tangent vectors above y . Let

$$(2.3) \quad \tilde{V} = \{(y, n), y \in M, n = (n_1, \dots, n_N), n_j \in S_y M, j = 1, \dots, N\} \times [0, 2L]^N.$$

Note \tilde{V} is a compact smooth $(N+1)d$ -dimensional manifold with boundary.

Define

$$(2.4) \quad \begin{aligned} \tilde{\phi} : \tilde{V} &\rightarrow M^N \\ (y, n, r) &\mapsto (\exp_y(n_1 r_1), \dots, \exp_y(n_N r_N)). \end{aligned}$$

The map $\tilde{\phi}$ is onto. If $x = (x_1, \dots, x_N) \in M^N$, consider $y \in M$ minimizing $H_{p,\mu(x)}$. Then among all (n, r) such that

$$(2.5) \quad \tilde{\phi}(y, n, r) = x$$

we can choose one so that for all $k = 1, \dots, N$ the map $s \mapsto \exp_y(s n_k)$ is a minimal geodesic for $s \in [0, r_k]$. For this choice we have

$$(2.6) \quad H_{p,\mu(x)}(y) = \frac{1}{N} \sum_{k=1}^N r_k^p.$$

Now since y minimizes $H_{p,\mu(x)}$, from equation (2.6) at y and variation of arc length formula, we have for all $u \in T_y M$

$$(2.7) \quad \left\langle \sum_{k=1}^N r_k^{p-1} n_k, u \right\rangle \leq 0$$

and this implies

$$(2.8) \quad \sum_{k=1}^N r_k^{p-1} n_k = 0.$$

So letting

$$(2.9) \quad \tilde{W}_p = \left\{ (y, n, r) \in \tilde{V}, \sum_{k=1}^N r_k^{p-1} n_k = 0 \right\}$$

and $\tilde{\phi}_p = \phi|_{\tilde{W}_p}$ the restriction of $\tilde{\phi}$ to \tilde{W}_p , $\tilde{\phi}_p$ is onto, on M^N by (2.5) and (2.8).

By Sard's theorem, the set $C_1 \subset M^N$ of singular values of $\tilde{\phi}_p$ has measure 0. It is closed since \tilde{W}_p is compact.

Let us prove that the set

$$(2.10) \quad C_2 := \{(x_1, \dots, x_N) \in M^N, \{x_1, \dots, x_N\} \cap Q_{p,\mu(x_1, \dots, x_N)} \neq \emptyset\}$$

has Lebesgue-measure 0: we can assume that for $i \neq j$, $x_i \neq x_j$ since we exclude 0-measure sets. So the elements we consider are images by $\tilde{\phi}_p$ of

$$(2.11) \quad \hat{W}_p = \left\{ (y, n, r) \in \tilde{W}_p, r_1 = 0, \forall k \geq 2 \ r_k > 0 \right\}.$$

The set \hat{W}_p is a submanifold of codimension 1 of \tilde{W}_p . Now $\dim \tilde{W}_p = Nd = \dim M^N$ so $\dim \hat{W}_p = \dim M^N - 1$ and its image by $\tilde{\phi}_p$ is of measure 0 in M^N . As a conclusion, C_2 has measure 0.

Define

$$(2.12) \quad C_3 := \{(x_1, \dots, x_N) \in M^N, \exists i \neq j \text{ s.t. } x_i = x_j\}$$

and $C = C_1 \cup C_2 \cup C_3$. The set C is closed in M^N and has measure 0. Letting

$$(2.13) \quad W_p = \left\{ (y, n, r) \in \tilde{W}_p, \forall k = 1, \dots, N, r_k \in (0, 2L) \right\},$$

we proved that $\tilde{\phi}_p|_{W_p}$ is onto on $M^N \setminus C$. Denote $\phi_p = \tilde{\phi}_p|_{W_p}$. Since W_p has same dimension as M^N and \tilde{W}_p is compact, every point x of $M^N \setminus C$ has a neighbourhood V_x such that $\phi_p^{-1}(V_x) = U_{1,x} \cup \dots \cup U_{m_x,x}$ where the $U_{j,x}$ are disjoint open subsets of W_p and

$$(2.14) \quad \phi_p|_{U_{j,x}} : U_{j,x} \rightarrow \phi_p(U_{j,x})$$

is a diffeomorphism. Now since $M^N \setminus C$ is second countable we can cover it by a countable number of such sets V_x . So to prove that the p -mean is almost everywhere unique it is sufficient to prove it on V_x .

For $x' \in V_x$ denote $x' = (x'_1, \dots, x'_N)$, and for $i \in \{1 \dots m_x\}$, write

$$(\phi|_{U_{i,x}})^{-1}(x') = (y_i(x'), n_1^i(x'), \dots, n_d^i(x'), r_1^i(x'), \dots, r_d^i(x')).$$

Let $i, j \in \{1 \dots m_x\}$ satisfy $i \neq j$. If $y_i(x'), y_j(x') \in Q_{p,\mu(x')}$ then we have

$$(2.15) \quad H_{p,\mu(x')} \circ y_i(x') = H_{p,\mu(x')} \circ y_j(x').$$

We can assume with the same argument as for (2.5) and (2.6) that the maps

$$(2.16) \quad \gamma_{i,k,x'} : s \mapsto \exp_{y_i(x')}(sn_k^i(x')) \quad \text{and} \quad \gamma_{j,k,x'} : s \mapsto \exp_{y_j(x')}(sn_k^j(x'))$$

are minimal geodesics respectively on $[0, r_k^i(x')]$ and $[0, r_k^j(x')]$. So letting $h_p : W_p \rightarrow \mathbb{R}$, $(y, n, r) \mapsto \sum_{k=1}^N r_k^p$, we have

$$\frac{1}{N} h_p \circ (\phi_p|_{U_{i,x}})^{-1}(x') = H_{p,\mu(x')} \circ y_i(x'), \quad \frac{1}{N} h_p \circ (\phi_p|_{U_{j,x}})^{-1}(x') = H_{p,\mu(x')} \circ y_j(x').$$

It is sufficient to prove that for all $x' \in V_x$,

$$(2.17) \quad h_p \circ (\phi_p|_{U_{i,x}})^{-1}(x') = h_p \circ (\phi_p|_{U_{j,x}})^{-1}(x')$$

implies

$$(2.18) \quad \text{grad}_{x'} (h_p \circ (\phi_p|_{U_{i,x}})^{-1}) \neq \text{grad}_{x'} (h_p \circ (\phi_p|_{U_{j,x}})^{-1}).$$

Indeed with (2.18) we will be able to deduce that the set

$$(2.19) \quad \{x' \in V_x, h_p \circ (\phi_p|_{U_{i,x}})^{-1} = h_p \circ (\phi_p|_{U_{j,x}})^{-1}\}$$

has codimension ≥ 1 in V_x and this will imply that

$$(2.20) \quad \{x' \in V_x, H_{p,\mu(x')} \circ y_i(x') = H_{p,\mu(x')} \circ y_j(x')\}$$

has codimension ≥ 1 in V_x .

Let us prove (2.18). For $k = 1, \dots, N$ let

$$m_k^i(x') = -\dot{\gamma}_{i,k,x'}(r_k^i(x')) \quad \text{and} \quad m_k^j(x') = -\dot{\gamma}_{j,k,x'}(r_k^j(x')).$$

These unit vectors satisfy

$$\exp_{x'_k}(r_k^i(x')m_k^i(x')) = y_i(x') \quad \text{and} \quad \exp_{x'_k}(r_k^j(x')m_k^j(x')) = y_j(x').$$

Then noting that $(h_p \circ (\phi_p|_{U_{i,x}})^{-1})(x') = \sum_{k=1}^N (r_k^i)^p(x')$ we get

$$\begin{aligned} & d_{x'} (h_p \circ (\phi_p|_{U_{i,x}})^{-1})(\cdot) \\ &= \left\langle -p \sum_{k=1}^N (r_k^i)^{p-1}(x') n_k^i(x'), T_{x'} y_i(\cdot) \right\rangle_{T_{y_i(x')} M} \\ & - p \langle ((r_1^i(x'))^{p-1} m_1^i(x'), \dots, (r_N^i(x'))^{p-1} m_N^i(x')), \cdot \rangle_{T_{x'} M^N}. \end{aligned}$$

Due to the fact that $(y_i(x'), n^i(x'), r^i(x')) \in W_p$, the first term in the right vanishes. So

$$(2.21) \quad \text{grad}_{x'} (h_p \circ (\phi_p|_{U_{i,x}})^{-1}) = -p \left((r_1^i(x'))^{p-1} m_1^i(x'), \dots, (r_N^i(x'))^{p-1} m_N^i(x') \right)$$

and similarly

$$(2.22) \quad \text{grad}_{x'} (h_p \circ (\phi_p|_{U_{j,x}})^{-1}) = -p \left((r_1^j(x'))^{p-1} m_1^j(x'), \dots, (r_N^j(x'))^{p-1} m_N^j(x') \right).$$

Since $y_i(x') \neq y_j(x')$ we have $(r_1^i(x'), m_1^i(x')) \neq (r_1^j(x'), m_1^j(x'))$, so $(r_1^i(x'))^{p-1} m_1^i(x') \neq (r_1^j(x'))^{p-1} m_1^j(x')$, from which we conclude that

$$\text{grad}_{x'} (h_p \circ (\phi_p|_{U_{i,x}})^{-1}) \neq \text{grad}_{x'} (h_p \circ (\phi_p|_{U_{j,x}})^{-1}).$$

This achieves the proof for the case $p > 1$.

Let us now consider the case $p = 1$. The result is due to Yang in [26], we give the proof here for completeness.

The main difference is that the subset of M^N of points $x = (x_1, \dots, x_N)$ so that $x_i \in Q_{1,\mu(x)}$ for some i has positive measure.

First consider the open subset U of M^N of points x such that for all $i = 1, \dots, N$, $x_i \notin Q_{1,\mu(x)}$.

Consider the closed subset C_0 of M^N of points $(x_1, \dots, x_N) = \tilde{\phi}(y, n, r)$, with $(y, n, r) \in \tilde{V}$ such that for all $j, k = 1, \dots, N$, $n_j = \pm n_k$. Since $d > 1$ and $N > 2$ this subset has Lebesgue measure 0.

Replacing M^N by U and C by $C_0 \cup C$, the argument is similar until (2.17). But now we will be able to prove that (2.17) implies (2.18) only in some neighbourhoods $V_{x,x'}$ to be precised later, of $x' \in V_x$ such that the geodesics

$$s \mapsto \exp_{y_i(x')}(sn_k^i(x')) \quad \text{and} \quad s \mapsto \exp_{y_j(x')}(sn_k^j(x'))$$

are minimal respectively on $[0, r_k^i(x')]$ and $[0, r_k^j(x')]$. But this will be sufficient since every compact subset of V_x can be covered by a finite number of these neighbourhoods $V_{x,x'}$.

Making the above assumption on x' , the proof is similar until (2.21) and (2.22). Then we have

$$(2.23) \quad \text{grad}_{x'} (h_1 \circ (\phi_1|_{U_{i,x}})^{-1}) = - (m_1^i(x'), \dots, m_N^i(x'))$$

and

$$(2.24) \quad \text{grad}_{x'} (h_1 \circ (\phi_1|_{U_{j,x}})^{-1}) = - (m_1^j(x'), \dots, m_N^j(x')).$$

Assume

$$\text{grad}_{x'} (h_1 \circ (\phi_1|_{U_{i,x}})^{-1}) = \text{grad}_{x'} (h_1 \circ (\phi_1|_{U_{j,x}})^{-1}).$$

Then for all $k = 1, \dots, N$, $m_k^i(x') = m_k^j(x')$. In particular for $k = 1$ this implies (possibly by exchanging i and j) that $y_i(x')$ lies in the minimizing geodesic from x'_1 to $y_j(x')$. Now since $x' \notin C_0$ there exists $k \in \{1, \dots, N\}$ such that $x'_k \notin \{\exp_{y_i(x')}(sn_1^i(x')), s \in [-2L, 2L]\}$. On the other hand since $m_k^i(x') = m_k^j(x')$, $y_j(x')$ (or $y_i(x')$) lies on the minimizing geodesic from x'_k to $y_i(x')$ (or $y_j(x')$). As a consequence there are two minimizing geodesics from $y_i(x')$ to $y_j(x')$. But this is impossible since the geodesic from x'_1 to $y_j(x')$ is minimizing, contains

$y_i(x')$ and $x'_1 \neq y_i(x')$ by the fact that we have supposed that $x'_1 \notin Q_{1,\mu(x')}$ and $y_i(x') \in Q_{1,\mu(x')}$. So

$$\text{grad}_{x'}(h_1 \circ (\phi_1|_{U_{i,x}})^{-1}) \neq \text{grad}_{x'}(h_1 \circ (\phi_1|_{U_{j,x}})^{-1}),$$

and by continuity this is true in a neighbourhood $V_{x,x'}$ of x' .

Now we consider the case where $x'_1 \in Q_{1,\mu(x')}$ and $x'_2 \notin Q_{1,\mu(x')}$. We follow the same lines as in the previous part with the difference that now $y_i(x') = x'_1$ and for the definition of $U_{i,x}$ W_1 is replaced by

$$W_1^i = \{(y, n, r) \in V, r_1 = 0\}.$$

The definition of $U_{j,x}$ remains unchanged. By [25] Theorem 1

$$\left\| \frac{1}{N} \sum_{k=2}^N n_k^i(x') \right\| \leq \mu_N(x')(\{x'_1\})$$

which gives

$$(2.25) \quad \left\| \sum_{k=2}^N n_k^i(x') \right\| \leq 1.$$

Since $d > 1$ and $N > 2$, the submanifolds of V_x images of

$$\left\{ (y, n, r) \in U_{i,x}, \left\| \sum_{k=2}^N n_k \right\| = 1 \right\}$$

and

$$\left\{ (y, n, r) \in U_{i,x}, \sum_{k=2}^N n_k = 0 \right\}$$

by ϕ_1 have measure 0, so we can exclude them. On the subset

$$\left\{ (y, n, r) \in U_{i,x}, 0 < \left\| \sum_{k=2}^N n_k \right\| < 1 \right\},$$

the function h_1 is smooth and on its image by ϕ_1 ,

$$(2.26) \quad \text{grad}_{x'}(h_1 \circ (\phi_1|_{U_{i,x}})^{-1}) = - (0, m_2^i(x'), \dots, m_N^i(x')).$$

Again

$$(2.27) \quad \text{grad}_{x'}(h_1 \circ (\phi_1|_{U_{j,x}})^{-1}) = - (m_1^j(x'), \dots, m_N^j(x')).$$

They are not equal, and this achieves the proof for this case by the same argument as before.

Finally we consider the case where $x'_1, x'_2 \in Q_{1,\mu(x')}$ with $x'_1 = y_i(x')$ and $x'_2 = y_j(x')$. We follow the same line as in the previous case, but now for the definition of $U_{j,x}$, W_1 is replaced by

$$W_1^j = \{(y, n, r) \in V, r_2 = 0\}.$$

Again we can exclude the submanifolds of V_x images of

$$\left\{ (y, n, r) \in U_{j,x}, \left\| \sum_{k \in \{1, \dots, N\}, k \neq 2} n_k \right\| = 1 \right\}$$

and

$$\left\{ (y, n, r) \in U_{j,x}, \sum_{k \in \{1, \dots, N\}, k \neq 2} n_k = 0 \right\}$$

by ϕ_1 and work on

$$\begin{aligned} & \phi_1 \left(\left\{ (y, n, r) \in U_{j,x}, 0 < \left\| \sum_{k \in \{1, \dots, N\}, k \neq 2} n_k \right\| < 1 \right\} \right) \\ & \cap \phi_1 \left(\left\{ (y, n, r) \in U_{i,x}, 0 < \left\| \sum_{k=2}^N n_k \right\| < 1 \right\} \right). \end{aligned}$$

On this set $h_1 \circ (\phi_1|_{U_{i,x}})^{-1}$ and $h_1 \circ (\phi_1|_{U_{j,x}})^{-1}$ are smooth and

$$(2.28) \quad \text{grad}_{x'} (h_1 \circ (\phi_1|_{U_{i,x}})^{-1}) = - (0, m_2^i(x'), \dots, m_N^i(x')).$$

$$(2.29) \quad \text{grad}_{x'} (h_1 \circ (\phi_1|_{U_{j,x}})^{-1}) = - (m_1^j(x'), 0, m_3^j(x'), \dots, m_N^j(x')).$$

They are not equal, and this achieves the proof. \square

Corollary 2.3. *Let $p \in [1, \infty)$ and $X = (X_1, \dots, X_N)$ a random variable with values in M^N , which has an absolutely continuous law. Then almost-surely $\mu(X(\omega))$ has a unique p -mean $e_p(X(\omega))$.*

Corollary 2.4. *Let $p \in [1, \infty)$ and $(X_n)_{n \geq 1}$ a sequence of i.i.d. M -valued random variables with absolutely continuous laws. Then the process of empirical p -means*

$$\left(e_{p,n}(\omega) := e_p(X_1(\omega), \dots, X_n(\omega)) \right)_{n \geq 1}$$

is well-defined.

Remark 2.5. For $p = 2$ and M a circle, it has been proved in [7] that the assumption can be weakened: the same result holds if the law has no atom.

We believe that it would be interesting to study the behaviour of the process $(e_{p,n})_{n \geq 1}$ in many situations. For instance when the law of X_1 is uniform on a compact symmetric space (even the case of the circle is highly non trivial) one would observe a recurrent but irregular and slower and slower process. Again on a compact symmetric space, when the law ν of X_1 has a finite number of p -means due to a finite group of symmetries, one would observe an almost stationary behaviour, and at increasingly spaced times jumps between smaller and smaller neighbourhoods of the p -means of ν .

3. FINDING THE MINIMIZERS OF SOME INTEGRATED FUNCTIONALS WITH SIMULATED ANNEALING

Let M be a compact Riemannian manifold. For simplicity and without loss of generality we assume that M has Lebesgue volume 1. On M consider a probability law ν with a density with respect to Lebesgue measure, also denoted by ν . Assume we are given a continuous function $\kappa : M \times M \rightarrow \mathbb{R}_+$, where $\kappa(\theta, y)$ is interpreted as some kind of cost for going from θ to y . Assume furthermore that for all $y \in M$

the function $\theta \mapsto \kappa(\theta, y)$ is smooth and that its first and second derivative in θ are uniformly bounded in (θ, y) . Consider on M the functional

$$(3.1) \quad \begin{aligned} U : M &\rightarrow \mathbb{R}_+ \\ \theta &\mapsto \int_M \kappa(\theta, y) \nu(dy) \end{aligned}$$

Denote by \mathcal{M} the set of minimizers of U . The aim of this section is to find a continuous semimartingale which converges in law to \mathcal{M} . Also we try to avoid using the gradient of U , which in many cases is difficult or impossible to compute.

For this we will use a sequence $(P_k)_{k \geq 0}$ of independent random variables with law ν , a Poisson process N_t on \mathbb{N} with intensity γ_t^{-1} where

$$(3.2) \quad \gamma_t = (1 + t)^{-1}.$$

Define

$$(3.3) \quad c(U) = 2 \sup_{\theta, y \in M} \left(\inf_{\phi \in \mathcal{C}_{\theta, y}} e(\phi) \right),$$

$\mathcal{C}_{\theta, y}$ denoting the set of continuous paths $[0, 1] \rightarrow M$ and for $\phi \in \mathcal{C}_{\theta, y}$, the elevation $e(\phi)$ being defined as

$$(3.4) \quad e(\phi) = \sup_{0 \leq t \leq 1} U(\phi(t)) - U(\theta) - U(y) + \inf_{z \in M} U(z).$$

Let

$$(3.5) \quad \beta_t = \frac{1}{k} \ln(1 + t),$$

the constant k satisfying $k > c(U)$.

We assume that $(N_t)_{t \geq 0}$ is independent of the sequence $(P_k)_{k \geq 0}$. We let $(B_t)_{t \geq 0}$ be a Brownian motion with values in \mathbb{R}^r for some $r \in \mathbb{N}$, independent of $(N_t)_{t \geq 0}$ and $(P_k)_{k \geq 0}$, and σ a smooth section of $TM \otimes (\mathbb{R}^r)^*$: for all $\theta \in M$, $\sigma(\theta)$ is a linear map $\mathbb{R}^r \rightarrow T_\theta M$. We assume that for all $\theta \in M$, we have $\sigma(\theta)\sigma(\theta)^* = \text{id}_{T_\theta M}$. We fix $\theta_0 \in M$ and let Θ_t be the solution started at θ_0 of the Itô equation

$$(3.6) \quad d\Theta_t = \sigma(\Theta_t) dB_t - \beta_t \text{grad}_{\Theta_t} \kappa(\cdot, Y_t) dt \quad \text{with } Y_t = P_{N_t}.$$

Recall that if $P(\Theta_t) : T_{\theta_0} M \rightarrow T_{\Theta_t} M$ is the parallel transport map along (Θ_t) , then

$$(3.7) \quad d\Theta_t = P(\Theta_t) d \left(\int_0^\cdot P(\Theta_s)^{-1} \circ d\Theta_s \right)_t.$$

Also define Θ_t^0 the solution started at θ_0 of the Itô equation

$$(3.8) \quad d\Theta_t^0 = \sigma(\Theta_t^0) dB_t - \beta_t \left(\int_M \text{grad}_{\Theta_t^0} \kappa(\cdot, y) \nu(y) dy \right) dt.$$

Note (3.8) rewrites as

$$(3.9) \quad d\Theta_t^0 = \sigma(\Theta_t^0) dB_t - \beta_t \text{grad}_{\Theta_t^0} U dt,$$

so that the same equation with fixed β instead of β_t has an invariant law with density

$$(3.10) \quad \mu_\beta(\theta) = \frac{1}{Z_\beta} e^{-2\beta U(\theta)}, \quad \text{with } Z_\beta = \int_M e^{-2\beta U(\theta')} d\theta'.$$

The process Θ_t^0 is an inhomogeneous diffusion with generator

$$(3.11) \quad L_t^0(\theta) = \frac{1}{2}\Delta(\theta) - \beta_t \operatorname{grad}_\theta U.$$

Denote by $m_t(\theta)$ the density of Θ_t .

The process (Θ_t, Y_t) is Markovian with generator L_t given by

$$(3.12) \quad \begin{aligned} L_t f(\theta, y) &= \left(\frac{1}{2}\Delta(\theta) - \beta_t \operatorname{grad}_\theta \kappa(\cdot, y) \right) f(\cdot, y) + \gamma_t^{-1} \int_M (f(\theta, z) - f(\theta, y)) \nu(dz) \\ &= L_{1,t} f(\cdot, y)(\theta) + L_{2,t} f(\theta, \cdot)(y). \end{aligned}$$

We know that for all neighbourhood \mathcal{N} of \mathcal{M} , $\int_{\mathcal{N}} \mu_\beta(\theta) d\theta$ converges to 1 as $\beta \rightarrow \infty$. So to prove that $\int_{\mathcal{N}} m_t(\theta) d\theta$ converges to 1 it is sufficient to prove the following proposition:

Proposition 3.1. *The entropy*

$$(3.13) \quad J_t := \int_M \ln \left(\frac{m_t(\theta)}{\mu_{\beta_t}(\theta)} \right) m_t(\theta) d\theta$$

converges to 0 as $t \rightarrow \infty$.

Proof. Let us compute

$$(3.14) \quad \frac{dJ_t}{dt} = \int_M \frac{dm_t(\theta)}{dt} d\theta - \int_M \frac{d \ln \mu_{\beta_t}(\theta)}{dt} m_t(\theta) d\theta + \int_M \ln \left(\frac{m_t(\theta)}{\mu_{\beta_t}(\theta)} \right) \frac{dm_t(\theta)}{dt} d\theta.$$

Since for all t $m_t(\theta)$ is a probability density, the first term in the right vanishes. So we get

$$(3.15) \quad \frac{dJ_t}{dt} = 2\beta'_t \int_M U(\theta)(m_t(\theta) - \mu_{\beta_t}(\theta)) d\theta + \int_M L_t \left[\ln \left(\frac{m_t(\theta)}{\mu_{\beta_t}(\theta)} \right) \right] m_t(\theta) d\theta$$

where the last term comes from Dynkin formula. For the first term in the right we have using (3.5)

$$(3.16) \quad 2\beta'_t \int_M U(\theta)(m_t(\theta) - \mu_{\beta_t}(\theta)) d\theta \leq 4\|U\|_\infty |\beta'_t| \leq \frac{4\|\kappa\|_\infty}{k(1+t)}.$$

Now we split the second term in the right of (3.15) into

$$(3.17) \quad \begin{aligned} & \int_M L_t \left[\ln \left(\frac{m_t(\theta)}{\mu_{\beta_t}(\theta)} \right) \right] m_t(\theta) d\theta \\ &= \int_M L_t^0 \left[\ln \left(\frac{m_t(\theta)}{\mu_{\beta_t}(\theta)} \right) \right] m_t(\theta) d\theta + \int_M R_t(\theta, y) \left[\ln \left(\frac{m_t(\theta)}{\mu_{\beta_t}(\theta)} \right) \right] m_t(\theta) d\theta. \end{aligned}$$

We have

$$(3.18) \quad \begin{aligned} & \int_M L_t^0 \left[\ln \left(\frac{m_t(\theta)}{\mu_{\beta_t}(\theta)} \right) \right] m_t(\theta) d\theta \\ &= \int_M L_t^0 \left[\left(\frac{m_t(\theta)}{\mu_{\beta_t}(\theta)} \right) \right] \mu_{\beta_t}(\theta) d\theta - \frac{1}{2} \int_M \left\| \nabla \ln \left(\frac{m_t(\theta)}{\mu_{\beta_t}(\theta)} \right) \right\|^2 \mu_{\beta_t}(\theta) d\theta \\ &= -2 \int_M \left\| \nabla \sqrt{\frac{m_t(\theta)}{\mu_{\beta_t}(\theta)}} \right\|^2 \mu_{\beta_t}(\theta) d\theta \\ &\leq -2c_2(\beta_t \vee 1)^{-p} \exp(-c(U)\beta_t) J_t \end{aligned}$$

for some $c_2 > 0$ and integer $p > 0$ by logarithmic Sobolev inequality ([15] and [16], for more details see [20]). Note we used again Dynkin formula to prove the vanishing of the first term in the right of the second line.

As for the second term we have

$$\begin{aligned}
& \int_M R_t(\theta, y) \left[\ln \left(\frac{m_t(\theta)}{\mu_{\beta_t}(\theta)} \right) \right] m_t(\theta) d\theta \\
&= \int_M -\beta_t \left\langle d \ln \left(\frac{m_t(\theta)}{\mu_{\beta_t}(\theta)} \right), \text{grad}_\theta \kappa(\cdot, y) - \int_M \text{grad}_\theta \kappa(\cdot, z) \nu(dz) \right\rangle m_t(\theta, y) d\theta dy \\
&= -\beta_t \int_M \left\langle d \ln \left(\frac{m_t(\theta)}{\mu_{\beta_t}(\theta)} \right), \int_M \text{grad}_\theta \kappa(\cdot, y) (m_t(y|\theta) - \nu(y)) \right\rangle m_t(\theta) d\theta dy \\
&= 2\beta_t \int_M \sqrt{\frac{\mu_{\beta_t}(\theta)}{m_t(\theta)}} \left\langle d \sqrt{\frac{m_t(\theta)}{\mu_{\beta_t}(\theta)}}, R_t(\theta) \right\rangle m_t(\theta) d\theta
\end{aligned}$$

with

$$R_t(\theta) = - \int_M \text{grad}_\theta \kappa(\cdot, y) (m_t(y|\theta) - \nu(y)) dy.$$

So by Cauchy-Schwartz inequality

$$\begin{aligned}
& \int_M R_t(\theta, y) \left[\ln \left(\frac{m_t(\theta)}{\mu_{\beta_t}(\theta)} \right) \right] m_t(\theta) d\theta \\
&\leq 2\beta_t \left(\int_M \left\| \nabla \sqrt{\frac{m_t(\theta)}{\mu_{\beta_t}(\theta)}} \right\|^2 \mu_{\beta_t}(\theta) d\theta \right)^{1/2} \left(\int_M \|R_t(\theta)\|^2 m_t(\theta) d\theta \right)^{1/2} \\
&\leq \beta_t^2 \int_M \|R_t(\theta)\|^2 m_t(\theta) d\theta + \int_M \left\| \nabla \sqrt{\frac{m_t(\theta)}{\mu_{\beta_t}(\theta)}} \right\|^2 \mu_{\beta_t}(\theta) d\theta.
\end{aligned}$$

Replacing $2c_2$ by c_2 in (3.18) we can after summing get rid of the second term in the right. Defining

$$(3.19) \quad K = \sup_{\theta, y \in M} \|\text{grad}_\theta \kappa(\cdot, y)\|,$$

let us now bound

$$\begin{aligned}
\int_M \|R_t(\theta)\|^2 m_t(\theta) d\theta &= \int_M \left\| \int_M \text{grad}_\theta \kappa(\cdot, y) (m_t(y|\theta) - \nu(y)) dy \right\|^2 m_t(\theta) d\theta \\
&\leq \int_M \left\| K \int_M |m_t(y|\theta) - \nu(y)| dy \right\|^2 m_t(\theta) d\theta \\
&\leq 32K^2 \int_M \left(\int_M \ln \left(\frac{m_t(y|\theta)}{\nu(y)} \right) m_t(y|\theta) dy \right) m_t(\theta) d\theta \\
&= 32K^2 I_t
\end{aligned}$$

where we have defined

$$(3.20) \quad I_t = \int_{M \times M} \ln \left(\frac{m_t(y|\theta)}{\nu(y)} \right) m_t(y, \theta) dy.$$

We also used classical bound of total variation by entropy ([16]):

$$\int_M |m_t(y|\theta) - \nu(y)| dy \leq 4\sqrt{2} \left(\int_M \ln \left(\frac{m_t(y|\theta)}{\nu(y)} \right) m_t(y|\theta) dy \right)^{1/2}$$

At this stage we proved that

$$(3.21) \quad \frac{dJ_t}{dt} \leq \frac{4\|\kappa\|_\infty}{k(1+t)} - c_2(\beta_t \vee 1)^{-p} \exp(-c(U)\beta_t) J_t + \beta_t^2 32K^2 I_t.$$

The next step is to find a suitable bound for $\frac{dI_t}{dt}$. As before

$$(3.22) \quad \begin{aligned} \frac{dI_t}{dt} &= \int_{M \times M} L_t \left[\ln \left(\frac{m_t(y|\theta)}{\nu(y)} \right) \right] m_t(y, \theta) d\theta dy \\ &= \int_{M \times M} (L_{2,t} + L_{1,t}) \left[\ln \left(\frac{m_t(y|\theta)}{\nu(y)} \right) \right] m_t(y, \theta) d\theta dy. \end{aligned}$$

We begin with the first term:

$$\begin{aligned} &\int_{M \times M} L_{2,t} \left[\ln \left(\frac{m_t(y|\theta)}{\nu(y)} \right) \right] m_t(y, \theta) d\theta dy \\ &= \gamma_t^{-1} \int_{M \times M} \int_M \left[\ln \left(\frac{m_t(z|\theta)}{\nu(z)} \right) - \ln \left(\frac{m_t(y|\theta)}{\nu(y)} \right) \right] \nu(dz) m_t(y, \theta) d\theta dy \\ &= \gamma_t^{-1} \int_{M \times M} \ln \left(\frac{m_t(y|\theta)}{\nu(y)} \right) (\nu(y) - m_t(y|\theta)) m_t(\theta) d\theta. \end{aligned}$$

By Jensen inequality we have

$$\begin{aligned} &\int_{M \times M} \ln \left(\frac{m_t(y|\theta)}{\nu(y)} \right) \nu(y) m_t(\theta) dy d\theta \\ &= \int_M \left(\int_M \ln \left(\frac{m_t(y|\theta)}{\nu(y)} \right) \nu(y) dy \right) m_t(\theta) d\theta \\ &\leq \int_M \ln \left(\int_M \frac{m_t(y|\theta)}{\nu(y)} \nu(y) dy \right) m_t(\theta) d\theta \\ &= \int_M \ln(1) m_t(\theta) d\theta = 0 \end{aligned}$$

Consequently

$$\begin{aligned} &\int_{M \times M} L_{2,t} \left[\ln \left(\frac{m_t(y|\theta)}{\nu(y)} \right) \right] m_t(y, \theta) d\theta dy \\ &\leq -\gamma_t^{-1} \int_{M \times M} \ln \left(\frac{m_t(y|\theta)}{\nu(y)} \right) m_t(y|\theta) m_t(\theta) d\theta \end{aligned}$$

which rewrites as

$$(3.23) \quad \int_{M \times M} L_{2,t} \left[\ln \left(\frac{m_t(y|\theta)}{\nu(y)} \right) \right] m_t(y, \theta) d\theta dy \leq -\gamma_t^{-1} I_t.$$

Let us now consider the second term in the right of (3.22). Since

$$\ln \left(\frac{m_t(y|\theta)}{\nu(y)} \right) = \ln \left(\frac{m_t(\theta|y)}{m_t(\theta)} \right)$$

it rewrites as

$$(3.24) \quad \begin{aligned} &\int_{M \times M} L_{1,t} \left[\ln \left(\frac{m_t(\theta|y)}{m_t(\theta)} \right) \right] m_t(y, \theta) d\theta dy \\ &= \int_{M \times M} L_{1,t} [\ln(m_t(\theta|y)) - \ln(m_t(\theta))] m_t(y, \theta) d\theta dy. \end{aligned}$$

But

$$\begin{aligned}
(3.25) \quad & \int_{M \times M} L_{1,t} \ln(m_t(\theta|y)) m_t(y, \theta) d\theta dy \\
&= \frac{1}{2} \int_{M \times M} \Delta \ln(m_t(\theta|y)) - \beta_t \int_M \langle d_\theta \ln m_t(\cdot|y), \text{grad}_\theta \kappa(\cdot, y) \rangle m_t(y, \theta) d\theta dy \\
&= -2 \int_{M \times M} \left\| \nabla \sqrt{m_t(\theta|y)} \right\|^2 d\theta \nu(dy) \\
&\quad - \beta_t \int_{M \times M} \langle d_\theta m_t(\cdot|y), \text{grad}_\theta \kappa(\cdot, y) \rangle \nu(y) d\theta dy.
\end{aligned}$$

Let us bound the absolute value of the last term:

$$\begin{aligned}
(3.26) \quad & \left| -\beta_t \int_{M \times M} \langle d_\theta m_t(\cdot|y), \text{grad}_\theta \kappa(\cdot, y) \rangle \nu(y) d\theta dy \right| \\
&= \left| 2\beta_t \int_{M \times M} \langle d_\theta \sqrt{m_t(\cdot|y)}, \text{grad}_\theta \kappa(\cdot, y) \rangle \sqrt{m_t(\theta|y)} \nu(dy) d\theta dy \right| \\
&\leq 2\beta_t K \int_{M \times M} \left\| \nabla \sqrt{m_t(\theta|y)} \right\| \sqrt{m_t(\theta|y)} \nu(y) d\theta dy \\
&\leq \int_{M \times M} \left(\frac{1}{2} \beta_t^2 K^2 m_t(\theta|y) + 2 \left\| \nabla \sqrt{m_t(\theta|y)} \right\|^2 \right) \nu(y) d\theta dy \\
&= \frac{1}{2} \beta_t^2 K^2 + 2 \int_{M \times M} \left\| \nabla \sqrt{m_t(\theta|y)} \right\|^2 \nu(y) d\theta dy.
\end{aligned}$$

This yields

$$(3.27) \quad \int_{M \times M} L_{1,t} \ln(m_t(\theta|y)) m_t(y, \theta) d\theta dy \leq \frac{1}{2} \beta_t^2 K^2$$

We also have to bound the last term in (3.24):

$$(3.28) \quad -L_{1,t} \ln(m_t(\theta)) = -L_{1,t} \ln \left(\frac{m_t(\theta)}{\mu_{\beta_t}(\theta)} \right) - L_{1,t} \ln(\mu_{\beta_t}(\theta)).$$

We already know that

$$(3.29) \quad - \int_M L_{1,t} \ln \left(\frac{m_t(\theta)}{\mu_{\beta_t}(\theta)} \right) m_t(\theta) d\theta \leq -\frac{dJ_t}{dt} + 4\|\kappa\|_\infty \beta_t'.$$

For the second term we have

$$\begin{aligned}
(3.30) \quad & -L_{1,t} \ln(\mu_{\beta_t}(\theta)) = 2L_{1,t} U(\theta) \\
&= \beta_t \Delta U(\theta) + \beta_t^2 \langle dU, \text{grad}_\theta \kappa(\cdot, y) \rangle \\
&\leq K'(\beta_t \vee 1) \beta_t
\end{aligned}$$

with

$$(3.31) \quad K' = \sup_{\theta, y \in M} |\Delta_\theta \kappa(\cdot, y)| + K^2.$$

Finally we obtain

$$(3.32) \quad \frac{dI_t}{dt} \leq 4\|\kappa\|_\infty \beta_t' + K'(\beta_t \vee 1) \beta_t - \frac{dJ_t}{dt} - \gamma_t^{-1} I_t$$

together with (3.21):

$$(3.33) \quad \frac{dJ_t}{dt} \leq \frac{4\|\kappa\|_\infty}{k(1+t)} - c_2(\beta_t \vee 1)^{-p} \exp(-c(U)\beta_t) J_t + 2\beta_t^2 32K^2 I_t.$$

At this stage we can use the end of the proof of theorem 1 in [21] to obtain that under assumptions (3.5) and (3.3) then

$$(3.34) \quad \lim_{t \rightarrow \infty} J_t = 0$$

(notice that in Section 4 we will prove this in a more general context). \square

Theorem 3.2. *Assume*

$$(3.35) \quad \beta_t = \frac{1}{k} \ln(1+t), \quad \text{and} \quad \gamma_t = (1+t)^{-1},$$

where $k > c(U)$, ($c(U)$ defined in (3.3)). Then for any neighbourhood \mathcal{N} of \mathcal{M} ,

$$(3.36) \quad \lim_{t \rightarrow \infty} \mathbb{P}[\Theta_t \in \mathcal{N}] = 1.$$

Proof. We use Proposition 3.1 together with the fact that

$$\|m_t - \mu_{\beta_t}\| \leq 4\sqrt{2J_t}$$

and

$$\lim_{t \rightarrow \infty} \mu_{\beta_t}(\mathcal{N}) = 1.$$

\square

4. APPLICATION TO LOCATION OF p -MEANS IN SYMMETRIC SPACES

In this section we assume that M is a compact symmetric space endowed with the canonical Riemannian metric of volume 1. Denote by ρ the Riemannian distance in M , D its diameter. We fix $p \geq 1$ and consider a probability measure ν on M . We aim to find at least one element of $Q_{p,\nu}$ by using the result of the previous section. In particular if ν has a unique p -mean e_p , then we will be able to construct a process which converges in probability to e_p as $t \rightarrow \infty$.

Denote by $p(s, x, y)$ the heat kernel on M , and for $s > 0$ let ν_s be the probability measure with density

$$(4.1) \quad \nu_s(y) = \int_M p(s, y, z) \nu(dz),$$

and let

$$(4.2) \quad \begin{aligned} \kappa_s &: M \times M \rightarrow \mathbb{R}_+ \\ (\theta, y) &\mapsto \int_M p(s, \theta, z) \rho^p(z, y) dz, \end{aligned}$$

and

$$(4.3) \quad \begin{aligned} U_{s_1, s_2} &: M \rightarrow \mathbb{R}_+ \\ \theta &\mapsto \int_M \kappa_{s_1}(\theta, y) \nu_{s_2}(y) dy. \end{aligned}$$

Also let $U = H_{p,\nu}$. Clearly ν_{s_1} and κ_{s_2} satisfy the assumption of the previous section. Moreover, denoting by \mathcal{M}_{s_1, s_2} the set of minimizers of U_{s_1, s_2} then as $s_1, s_2 \rightarrow 0$ we have $\mathcal{M}_{s_1, s_2} \rightarrow Q_{p,\nu}$ in the sense that for any neighbourhood \mathcal{N} of $Q_{p,\nu}$, we have $\mathcal{M}_{s_1, s_2} \subset \mathcal{N}$ for all s_1, s_2 sufficiently small. This is due to the fact that as $s_1, s_2 \rightarrow 0$, $U_{s_1, s_2}(\theta) \rightarrow U(\theta)$ uniformly in θ .

Lemma 4.1. *For all $s_1, s_2 > 0$ we have*

$$(4.4) \quad U_{s_1, s_2}(\theta) = U_{0, s_1 + s_2}(\theta) = \int_M \rho(\theta, y) \nu_{s_1 + s_2}(y) dy.$$

Proof. Fix $\theta, y \in M$, let m be the middle point of a minimal geodesic from θ to y and i_m the symmetry centered at m . We have

$$\begin{aligned} \int_M p(s_1, \theta, z) \rho^p(z, y) dz &= \int_M p(s_1, i_m(\theta), i_m(z)) \rho^p(i_m(z), i_m(y)) dz \\ &= \int_M p(s_1, i_m(\theta), z') \rho^p(z', i_m(y)) dz' \\ &= \int_M p(s_1, y, z') \rho^p(z', \theta) dz' \\ &= \int_M \rho^p(\theta, z') p(s_1, z', y) dz' \end{aligned}$$

where we first used the invariance by isometry of the heat kernel and then did the change of variable $z' = i_m(z)$ in the integral and finally used the symmetry of the heat kernel. To finish the proof we are left to use the convolution property of the heat semigroup. \square

Corollary 4.2. *Defining*

$$(4.5) \quad K'' = \sup_{\theta, y \in M} \|\text{grad}_\theta \rho(\cdot, y)\|,$$

we have for all $s_1, s_2 > 0, \theta, y \in M$,

$$(4.6) \quad \|\text{grad}_\theta \kappa_{s_1}(\cdot, y)\| \leq p D^{p-1} K'' =: K \quad \text{and} \quad \|\text{grad}_\theta U_{s_1, s_2}\| \leq K.$$

With all these properties we would like to find $s_1(t) \searrow 0$ and $s_2(t) \searrow 0$ such that the process Θ_t started at θ_0 and solution to

$$(4.7) \quad d\Theta_t = \sigma(\Theta_t) dB_t - \beta_t \text{grad}_{\Theta_t} \kappa_{s_1(t)}(\cdot, Y_t^{s_2(t)}) dt$$

converges in law to e_p , $(N_t, Y_t^{s_2(t)})$ being a Poisson point process in $[0, \infty) \times M$ with intensity $\gamma(t)^{-1} \nu_{s_2(t)}(y) dt dy$, independent of (B_t) . This is the object of the next theorem in which we will take

$$s_1(t) = s_2(t) = s_t = (\ln(1+t))^{-1}.$$

So define Θ_t^0 the solution started at θ_0 of the Itô equation

$$(4.8) \quad d\Theta_t^0 = \sigma(\Theta_t^0) dB_t - \beta_t \left(\int_M \text{grad}_{\Theta_t^0} \kappa_{s_t}(\cdot, y) \nu_{s_t}(y) dy \right) dt.$$

Notice that using Lemma 4.1, (4.8) rewrites as

$$(4.9) \quad d\Theta_t^0 = \sigma(\Theta_t^0) dB_t - \beta_t \text{grad}_{\Theta_t^0} U_{2s_t} dt,$$

where $U_{2s_t} := U_{0, 2s_t}$, so that the same equation with fixed (β, s) instead of (β_t, s_t) has an invariant law with density

$$(4.10) \quad \mu_{\beta, s}(\theta) = \frac{1}{Z_{\beta, s}} e^{-2\beta U_{2s}(\theta)}, \quad \text{with} \quad Z_{\beta, s} = \int_M e^{-2\beta U_{2s}(\theta')} d\theta'.$$

The process Θ_t^0 is an inhomogeneous diffusion with generator

$$(4.11) \quad L_t^0(\theta) = \frac{1}{2} \Delta(\theta) - \beta_t \text{grad}_\theta U_{2s_t}.$$

Denote by $m_t(\theta)$ the density of Θ_t .

Let $Y_t := Y_t^{s_t}$. The process (Θ_t, Y_t) is Markovian with generator L_t given by

$$(4.12) \quad \begin{aligned} L_t f(\theta, y) &= \left(\frac{1}{2} \Delta(\theta) - \beta_t \operatorname{grad}_\theta \kappa_{s_t}(\cdot, y) \right) f(\cdot, y) + \gamma_t^{-1} \int_M (f(\theta, z) - f(\theta, y)) \nu_{s_t}(dz) \\ &= L_{1,t} f(\cdot, y)(\theta) + L_{2,t} f(\theta, \cdot)(y). \end{aligned}$$

We know that for all neighbourhood \mathcal{N} of $Q_{p,\nu}$, $\int_{\mathcal{N}} \mu_{\beta,s}(\theta) d\theta$ converges to 1 as $\beta \rightarrow \infty$, uniformly in s sufficiently small. Again define

$$(4.13) \quad J_t := \int_M \ln \left(\frac{m_t(\theta)}{\mu_{\beta_t, s_t}(\theta)} \right) m_t(\theta) d\theta.$$

Theorem 4.3. *Assume*

$$(4.14) \quad \beta_t = \frac{1}{k} \ln(1+t), \quad \gamma_t = (1+t)^{-1}, \quad s_1(t) = s_2(t) = s(t) = (\ln(1+t))^{-1}.$$

where $k > c(U)$, ($c(U)$ defined in (3.3)). Then for any neighbourhood \mathcal{N} of $Q_{p,\nu}$, the process Θ_t defined in equation (4.7) satisfies

$$(4.15) \quad \lim_{t \rightarrow \infty} \mathbb{P}[\Theta_t \in \mathcal{N}] = 1.$$

Proof. We use Proposition 4.4 below together with the fact that

$$\|m_t - \mu_{\beta_t, s_t}\| \leq 4\sqrt{2J_t}$$

and

$$\lim_{t \rightarrow \infty} \mu_{\beta_t, s_t}(\mathcal{N}) = 1.$$

□

Proposition 4.4. *The entropy*

$$(4.16) \quad J_t = \int_M \ln \left(\frac{m_t(\theta)}{\mu_{\beta_t, s_t}(\theta)} \right) m_t(\theta) d\theta$$

converges to 0 as $t \rightarrow \infty$.

Proof. Let us compute as before

$$(4.17) \quad \frac{dJ_t}{dt} = - \int_M \partial_t \ln(\mu_{\beta_t, s_t}(\theta)) m_t(\theta) d\theta + \int_M L_t \left[\ln \left(\frac{m_t(\theta)}{\mu_{\beta_t, s_t}(\theta)} \right) \right] m_t(\theta) d\theta.$$

For the first term in the right we have using (4.10)

$$(4.18) \quad \begin{aligned} &\partial_t \ln(\mu_{\beta_t, s_t}(\theta)) \\ &= -2\beta'_t U_{2s_t} - 2\beta_t \int_{M \times M} 2s'_t \partial_s \ln p(2s_t, \theta, z) p(2s_t, \theta, z) \rho^p(z, y) \nu(dy) dz \\ &\quad + 2\beta'_t \int_M U_{2s_t}(\theta') \mu_{\beta_t, s_t}(\theta') d\theta' \\ &\quad + 2\beta_t \int_M \left(\int_{M \times M} 2s'_t \partial_s \ln p(2s_t, \theta', z) p(2s_t, \theta', z) \rho^p(z, y) dz \nu(dy) \right) \mu_{\beta_t, s_t}(\theta') d\theta'. \end{aligned}$$

It is known that there exists $C_0 > 0$ such that $\forall s \in (0, 1]$

$$(4.19) \quad |\partial_s \ln p(s, \theta, z)| \leq \frac{C_0}{s^2},$$

see e.g. [14] and [22] where bounds of the type $|\nabla_\theta \ln p(s, \theta, z)| \leq \frac{C_1}{s}$ and $|\nabla_\theta^2 \ln p(s, \theta, z)| \leq \frac{C_2}{s^2}$ are given. Here we use

$$|\partial_s \ln p(s, \theta, z)| = \frac{1}{2} \left| \frac{\Delta_\theta p(s, \theta, z)}{p(s, \theta, z)} \right| \leq \frac{\dim M}{2} (|\nabla_\theta^2 \ln p(s, \theta, z)| + |\nabla_\theta \ln p(s, \theta, z)|^2).$$

So (4.18) and (4.19) yield

$$(4.20) \quad |\partial_t \ln(\mu_{\beta_t, s_t}(\theta))| \leq D^p \left(4\beta'_t + \frac{C_0 \beta_t |s'_t|}{s_t^2} \right).$$

which implies

$$(4.21) \quad |\partial_t \ln(\mu_{\beta_t, s_t}(\theta))| \leq C \left(\beta'_t + \frac{\beta_t |s'_t|}{s_t^2} \right).$$

with

$$(4.22) \quad C = D^p(4 + C_0).$$

Evaluating with (4.14) and integrating on M we get

$$(4.23) \quad \left| - \int_M \partial_t \ln(\mu_{\beta_t, s_t}(\theta)) m_t(\theta) d\theta \right| \leq \frac{C}{(1+t)k} (1 + \ln(1+t)).$$

Now we split the second term in the right of (4.17) into

$$(4.24) \quad \begin{aligned} & \int_M L_t \left[\ln \left(\frac{m_t(\theta)}{\mu_{\beta_t, s_t}(\theta)} \right) \right] m_t(\theta, y) d\theta dy \\ &= \int_M L_t^0 \left[\ln \left(\frac{m_t(\theta)}{\mu_{\beta_t, s_t}(\theta)} \right) \right] m_t(\theta) d\theta + \int_M R_t(\theta, y) \left[\ln \left(\frac{m_t(\theta)}{\mu_{\beta_t, s_t}(\theta)} \right) \right] m_t(\theta, y) d\theta dy. \end{aligned}$$

We have as for (3.18)

$$(4.25) \quad \begin{aligned} \int_M L_t^0 \left[\ln \left(\frac{m_t(\theta)}{\mu_{\beta_t, s_t}(\theta)} \right) \right] m_t(\theta) d\theta &= -2 \int_M \left\| \nabla \sqrt{\frac{m_t(\theta)}{\mu_{\beta_t, s_t}(\theta)}} \right\|^2 \mu_{\beta_t, s_t}(\theta) d\theta \\ &\leq -2c_2(\beta_t \vee 1)^{-p} \exp(-c(U_{2s_t})\beta_t) J_t \end{aligned}$$

for some $c_2 > 0$ and integer $p > 0$ by logarithmic Sobolev inequality ([20]).

The computation for the second term is similar to the one after (3.18) and we get

$$\begin{aligned} & \int_M R_t(\theta, y) \left[\ln \left(\frac{m_t(\theta)}{\mu_{\beta_t, s_t}(\theta)} \right) \right] m_t(\theta, y) d\theta dy \\ &= 2\beta_t \int_M \sqrt{\frac{\mu_{\beta_t, s_t}(\theta)}{m_t(\theta)}} \left\langle d \sqrt{\frac{m_t(\theta)}{\mu_{\beta_t, s_t}(\theta)}}, R_t(\theta) \right\rangle m_t(\theta) d\theta \end{aligned}$$

with

$$R_t(\theta) = - \int_M \text{grad}_\theta \kappa_{s_t}(\cdot, y) (m_t(y|\theta) - \nu_{s_t}(y)) dy,$$

and again

$$\begin{aligned} & \int_M R_t(\theta, y) \left[\ln \left(\frac{m_t(\theta)}{\mu_{\beta_t, s_t}(\theta)} \right) \right] m_t(\theta) d\theta \\ & \leq \beta_t^2 \int_M \|R_t(\theta)\|^2 m_t(\theta) d\theta + \int_M \left\| \nabla \sqrt{\frac{m_t(\theta)}{\mu_{\beta_t, s_t}(\theta)}} \right\|^2 \mu_{\beta_t, s_t}(\theta) d\theta. \end{aligned}$$

Replacing $2c_2$ by c_2 in (4.25) we can after summing get rid of the second term in the right.

Here again

$$\int_M \|R_t(\theta)\|^2 m_t(\theta) d\theta \leq 32K^2 I_t$$

where we have defined

$$(4.26) \quad I_t = \int_{M \times M} \ln \left(\frac{m_t(y|\theta)}{\nu_{s_t}(y)} \right) m_t(y, \theta) dy.$$

At this stage we proved that

$$(4.27) \quad \begin{aligned} \frac{dJ_t}{dt} & \leq \frac{C}{(1+t)k} (1 + \ln(1+t)) \\ & - c_2(\beta_t \vee 1)^{-p} \exp(-c(U_{2s_t})\beta_t) J_t + \beta_t^2 32K^2 I_t. \end{aligned}$$

The next step is to find a suitable bound for $\frac{dI_t}{dt}$. As before

$$(4.28) \quad \begin{aligned} \frac{dI_t}{dt} & = - \int_M \partial_t \ln(\nu_{s_t}(y)) m_t(\theta, y) d\theta dy \\ & + \int_{M \times M} L_t \left[\ln \left(\frac{m_t(y|\theta)}{\nu_{s_t}(y)} \right) \right] m_t(y, \theta) d\theta dy \end{aligned}$$

and

$$(4.29) \quad \begin{aligned} & \left| \int_M \partial_t \ln(\nu_{s_t}(y)) m_t(\theta, y) d\theta dy \right| \\ & = \left| s'_t \int_M \partial_s \ln p(s_t, y, z) p(s_t, y, z) \nu(dz) \right| \\ & \leq \frac{|s'_t| C_0}{s_t^2} = \frac{C_0}{1+t}. \end{aligned}$$

Now

$$(4.30) \quad \begin{aligned} & \int_{M \times M} L_t \left[\ln \left(\frac{m_t(y|\theta)}{\nu_{s_t}(y)} \right) \right] m_t(y, \theta) d\theta dy \\ & = \int_{M \times M} (L_{2,t} + L_{1,t}) \left[\ln \left(\frac{m_t(y|\theta)}{\nu_{s_t}(y)} \right) \right] m_t(y, \theta) d\theta dy. \end{aligned}$$

We begin with the first term:

$$\begin{aligned} & \int_{M \times M} L_{2,t} \left[\ln \left(\frac{m_t(y|\theta)}{\nu_{s_t}(y)} \right) \right] m_t(y, \theta) d\theta dy \\ & = \gamma_t^{-1} \int_{M \times M} \ln \left(\frac{m_t(y|\theta)}{\nu_{s_t}(y)} \right) (\nu_{s_t}(y) - m_t(y|\theta)) m_t(\theta) d\theta \end{aligned}$$

and estimate it as for (3.23):

$$(4.31) \quad \int_{M \times M} L_{2,t} \left[\ln \left(\frac{m_t(y|\theta)}{\nu_{s_t}(y)} \right) \right] m_t(y, \theta) d\theta dy \leq -\gamma_t^{-1} I_t.$$

For the second term in the right of (4.28) we need to introduce the density $f_t(y)$ of $Y_t^{s_t}$. Since

$$\ln \left(\frac{m_t(y|\theta)}{\nu_{s_t}(y)} \right) = \ln \left(\frac{m_t(\theta|y)}{m_t(\theta)} \right) + \ln \left(\frac{f_t(y)}{\nu_{s_t}(y)} \right)$$

it rewrites as

$$(4.32) \quad \begin{aligned} & \int_{M \times M} L_{1,t} \left[\ln \left(\frac{m_t(\theta|y)}{m_t(\theta)} \right) \right] m_t(y, \theta) d\theta dy \\ &= \int_{M \times M} L_{1,t} [\ln(m_t(\theta|y)) - \ln(m_t(\theta))] m_t(y, \theta) d\theta dy. \end{aligned}$$

Similarly to (3.25)

$$(4.33) \quad \begin{aligned} & \int_{M \times M} L_{1,t} \ln(m_t(\theta|y)) m_t(y, \theta) d\theta dy \\ &= \frac{1}{2} \int_{M \times M} \Delta \ln(m_t(\theta|y)) - \beta_t \int_M \langle d_\theta \ln m_t(\cdot|y), \text{grad}_\theta \kappa_{s_t}(\cdot, y) \rangle m_t(y, \theta) d\theta dy \\ &= -2 \int_{M \times M} \left\| \nabla \sqrt{m_t(\theta|y)} \right\|^2 d\theta f_t(y) dy \\ &\quad - \beta_t \int_{M \times M} \langle d_\theta m_t(\cdot|y), \text{grad}_\theta \kappa_{s_t}(\cdot, y) \rangle f_t(y) d\theta dy. \end{aligned}$$

For the absolute value of the last term:

$$(4.34) \quad \begin{aligned} & \left| -\beta_t \int_{M \times M} \langle d_\theta m_t(\cdot|y), \text{grad}_\theta \kappa(\cdot, y) \rangle \nu(y) f_t(y) d\theta dy \right| \\ & \leq \frac{1}{2} \beta_t^2 K^2 + 2 \int_{M \times M} \left\| \nabla \sqrt{m_t(\theta|y)} \right\|^2 f_t(y) d\theta dy. \end{aligned}$$

We get as in (3.27)

$$(4.35) \quad \int_{M \times M} L_{1,t} \ln(m_t(\theta|y)) m_t(y, \theta) d\theta dy \leq \frac{1}{2} \beta_t^2 K^2$$

Then we bound the last term in (4.32):

$$(4.36) \quad -L_{1,t} \ln(m_t(\theta)) = -L_{1,t} \ln \left(\frac{m_t(\theta)}{\mu_{\beta_t, s_t}(\theta)} \right) - L_{1,t} \ln(\mu_{\beta_t, s_t}(\theta)).$$

We already know by (4.17) and (4.23) that

$$(4.37) \quad - \int_M L_{1,t} \ln \left(\frac{m_t(\theta)}{\mu_{\beta_t, s_t}(\theta)} \right) m_t(\theta) d\theta \leq -\frac{dJ_t}{dt} + \frac{C}{(1+t)k} (1 + \ln(1+t)).$$

For the second term we have

$$(4.38) \quad \begin{aligned} L_{1,t} \ln(\mu_{\beta_t, s_t}(\theta)) &= -2\beta_t L_{1,t} U_{2s_t}(\theta) \\ &= -\beta_t \Delta U_{2s_t}(\theta) + 2\beta_t^2 \langle dU_{2s_t}, \text{grad}_\theta \kappa_{s_t}(\cdot, y) \rangle \\ &\leq K'(\beta_t \vee 1) \beta_t s_t^{-2} \end{aligned}$$

for some $K' > 0$, where we used

$$\Delta U_{2s} = \int_M (\Delta_\theta \ln p(2s, \theta, y) + \|\nabla_\theta \ln p(2s, \theta, y)\|^2) p(2s, \theta, y) \rho^p(y, z) \nu(dz)$$

and standard bound for the first and second derivatives of the heat kernel ([14] and [22]).

Finally we obtain

$$(4.39) \quad \frac{dI_t}{dt} \leq \frac{C_0}{(1+t)} + \frac{K'}{k^2} (\ln(1+t) \vee k) (\ln(1+t))^3 - \frac{dJ_t}{dt} - (1+t)I_t$$

together with (4.27):

$$(4.40) \quad \frac{dJ_t}{dt} \leq \frac{C}{(1+t)k} (1 + \ln(1+t)) - c_2(\beta_t \vee 1)^{-p} \exp(-c(U_{2s_t})\beta_t) J_t + 2\beta_t^2 32K^2 I_t$$

which rewrites as

$$(4.41) \quad \frac{dI_t}{dt} \leq k_1 (\ln(1+t))^4 - \frac{dJ_t}{dt} - (1+t)I_t$$

and

$$(4.42) \quad \frac{dJ_t}{dt} \leq c_1 \left(\frac{\ln(1+t)}{1+t} + (\ln(1+t))^2 I_t \right) - c_2 (\ln(1+t))^{-p} (1+t)^{-\frac{c(U_{2s_t})}{k}} J_t$$

for some constants $c_1, k_1 > 0$, as soon as $t \geq 2$. At this stage we can use a similar computation to the end of the proof of theorem 1 in [21] to obtain that under assumptions (4.14) and (3.3) then

$$(4.43) \quad \lim_{t \rightarrow \infty} J_t = 0.$$

However we will do the calculation for completeness, and because there are some small differences. Recall $U_s \rightarrow U$ uniformly as $s \rightarrow 0$. Moreover $2s_t \rightarrow 0$ as $t \rightarrow \infty$, so we get

$$\limsup_{t \rightarrow \infty} c(U_{2s_t}) \leq c(U).$$

As a consequence, for t sufficiently large we have

$$(4.44) \quad \frac{c(U_{2s_t})}{k} \leq 1 - \varepsilon$$

for some $\varepsilon > 0$. Let

$$(4.45) \quad \ell_t = \frac{c_1 (\ln(1+t))^2}{1+t + c_1 (\ln(1+t))^2 - c_2 (\ln(1+t))^{-p} (1+t)^{-(1-\varepsilon)}}$$

where $\varepsilon > 0$ is defined in (4.44). It is easily checked that for t sufficiently large ℓ_t is positive and decreasing, and that it converges to 0 as $t \rightarrow \infty$. Define

$$(4.46) \quad K_t = J_t + \ell_t I_t.$$

We will prove that $K_t \rightarrow 0$ as $t \rightarrow \infty$ and from this we will get (4.43).

for t sufficiently large,

$$(4.47) \quad \frac{dK_t}{dt} \leq \frac{dJ_t}{dt} + \ell_t \frac{dI_t}{dt}$$

and this yields with (4.41) and (4.42)

$$\begin{aligned} \frac{dK_t}{dt} &\leq (1 - \ell_t)c_1 \frac{\ln(1+t)}{1+t} + c_1(\ln(1+t))^2 I_t \\ &\quad - \ell_t c_1 (\ln(1+t))^2 I_t - (1 - \ell_t)c_2 (\ln(1+t))^{-p} (1+t)^{-\frac{c(U_{2s_t})}{k}} J_t \\ &\quad + \ell_t k_1 (\ln(1+t))^4 - (1+t)\ell_t I_t. \end{aligned}$$

Replacing $c_1(\ln(1+t))^2$ at the end of the first line by

$$\ell_t \left(1+t + c_1(\ln(1+t))^2 - c_2(\ln(1+t))^{-p} (1+t)^{-(1-\varepsilon)} \right)$$

by the help of (4.45) we obtain

$$\begin{aligned} \frac{dK_t}{dt} &\leq c_1 \frac{\ln(1+t)}{1+t} - c_2 \ell_t (\ln(1+t))^{-p} (1+t)^{-\frac{c(U_{2s_t})}{k}} I_t \\ &\quad - (1 - \ell_t)c_2 (\ln(1+t))^{-p} (1+t)^{-\frac{c(U_{2s_t})}{k}} J_t + \ell_t k_1 (\ln(1+t))^4 \end{aligned}$$

and this yields using $-(1+t)^{-\frac{c(U_{2s_t})}{k}} \leq -(1+t)^{-(1-\varepsilon)}$:

$$(4.48) \quad \frac{dK_t}{dt} \leq A_t - B_t K_t$$

with

$$(4.49) \quad A_t = c_1 \frac{\ln(1+t)}{1+t} + \ell_t k_1 \ln(1+t)^4$$

and

$$(4.50) \quad B_t = (1 - \ell_t)c_2 (\ln(1+t))^{-p} (1+t)^{-(1-\varepsilon)}.$$

A sufficient condition for K_t to converge to 0 as $t \rightarrow \infty$ is

$$(4.51) \quad \int_0^\infty B_t dt = +\infty$$

and

$$(4.52) \quad \lim_{t \rightarrow \infty} \frac{A_t}{B_t} = 0.$$

Condition (4.51) clearly is realized. As for condition (4.52) we easily see that

$$\frac{c_1 \frac{\ln(1+t)}{1+t}}{(1 - \ell_t)c_2 (\ln(1+t))^{-p} (1+t)^{-(1-\varepsilon)}} \rightarrow 0$$

and also

$$\frac{\ell_t k_1 (\ln(1+t))^4}{(1 - \ell_t)c_2 (\ln(1+t))^{-p} (1+t)^{-(1-\varepsilon)}} \rightarrow 0$$

from the fact that

$$\ell_t \leq \frac{c(\ln(1+t))^2}{1+t}$$

for some $c > 0$. □

REFERENCES

- [1] B. Afsari, *Riemannian L^p center of mass : existence, uniqueness, and convexity*, Proceedings of the American Mathematical Society, S 0002-9939(2010)10541-5, Article electronically published on August 27, 2010.
- [2] B. Afsari, R. Tron and R. Vidal, *On the convergence of gradient descent for finding the Riemannian center of mass* arXiv:1201.0925
- [3] M. Arnaudon, F. Nielsen, *Medians and means in Finsler geometry*, LMS Journal of Computation and Mathematics volume 15, pp. 23-37 (2012)
- [4] Arnaudon, M., Dombry, C., Phan, A., Yang, L., *Stochastic algorithms for computing means of probability measures* Stoch. Proc. Appl. 122 (2012), pp. 1437-1455.
- [5] M. Arnaudon, F. Nielsen, *On computing the Riemannian 1-Center*, Computational Geometry: Theory and Applications, to appear
- [6] Bădoiu, M., Clarkson, K. L., 2003, *Smaller core-sets for balls*, Proceedings of the fourteenth annual ACM-SIAM symposium on Discrete algorithms. Society for Industrial and Applied Mathematics, Philadelphia, PA, USA, pp. 801–802.
- [7] R. Bhattacharya and V. Patangreanu, *Large sample theory of intrinsic and extrinsic sample means on manifolds (i)*, Annals of Statistics 31 (1), pp. 1–29 (2003)
- [8] S. Bonnabel, *Convergence des méthodes de gradient stochastique sur les variétés riemanniennes* In GRETSI. Bordeaux. 2011
- [9] H. Cardot, P. Cnac, P.-A. Zitt, *Efficient and fast estimation of the geometric median in Hilbert spaces with an averaged stochastic gradient algorithm*, Bernoulli.
- [10] B. Charlier, *Necessary and sufficient condition for the existence of a Fréchet mean on the circle* arXiv:1109.1986.
- [11] P.T. Fletcher, S. Venkatasubramanian, S. Joshi, *The geometric median on Riemannian manifolds with application to robust atlas estimation*, NeuroImage, 45 (2009), pp. S143–S152
- [12] D. Groisser, *Newton’s method, zeroes of vector fields, and the Riemannian center of mass* Adv. in Appl. Math. 33 (2004), no. 1, 95135.
- [13] D. Groisser, *On the convergence of some Procrustean averaging algorithms* Stochastics 77 (2005), no. 1, 3160.
- [14] E.P. Hsu, *Estimates of derivatives of the heat kernel on a compact Riemannian manifold*, Proceedings of the American Mathematical Society, Vol. 127 (1999), n. 12, pp 3739–3744.
- [15] R. Holley, S. Kusuoka and D. Stroock, *Asymptotics of the spectral gap with applications to the theory of simulated annealing*, J. Funct. Anal. 83 (1989) 333–347.
- [16] R. Holley and D. Stroock, *Annealing via Sobolev inequalities*, Comm. Math. Phys. 115 (1988) 553–569.
- [17] T. Hotz, S. Huckemann, *Intrinsic mean on the circle: Uniqueness, Locus and Asymptotics*, arXiv.org, 1108:2141
- [18] W.S. Kendall, *Probability, convexity and harmonic maps with small image I: uniqueness and fine existence*, Proc. London Math. Soc. (3) 61 no. 2 (1990) pp. 371–406
- [19] H. Le, *Estimation of Riemannian barycentres*, LMS J. Comput. Math. 7 (2004), pp. 193–200
- [20] L. Miclo, *Recuit simulé sans potentiel sur une variété compacte*, Stochastics and Stochastic Reports 41 (1992) 23–56.
- [21] L. Miclo, *Recuit simulé partiel*, Stochastic Processes and their Applications 65 (1996) 281–298
- [22] S.J. Sheu, *Some estimates of the transition density function of a nondegenerate diffusion Markov process*, The Annals of Probability 19, n. 2 (1991), 538–561.
- [23] K.T. Sturm, *Probability measures on metric spaces of nonpositive curvature*, Heat kernels and analysis on manifolds, graphs, and metric spaces (Paris, 2002), 357390, Contemp. Math., 338, Amer. Math. Soc., Providence, RI, 2003.
- [24] E. Weiszfeld, *Sur le point pour lequel la somme des distances de n points donnés est minimum*, Tohoku Math. J. 43 (1937), pp. 355–386
- [25] L.Yang, *Riemannian median and its estimation*, LMS Journal of Computation and Mathematics, Vol 13 (2010), pp 461–479.
- [26] L.Yang, *Some properties of Frechet medians in Riemannian manifolds*, preprint

LABORATOIRE DE MATHÉMATIQUES ET APPLICATIONS
CNRS: UMR 7348
UNIVERSITÉ DE POITIERS, TÉLÉPORT 2 - BP 30179
F-86962 FUTUROSCOPE CHASSENEUIL CEDEX, FRANCE
E-mail address: `marc.arnaudon@math.univ-poitiers.fr`

INSTITUT DE MATHÉMATIQUE DE TOULOUSE
CNRS: UMR 5219
118, ROUTE DE NARBONNE
F-31062 TOULOUSE CEDEX 9, FRANCE
E-mail address: `laurent.miclo@math.univ-toulouse.fr`