



A gradient flow approach to a thin film approximation of the Muskat problem

Philippe Laurencot, Bogdan-Vasile Matioc

► To cite this version:

Philippe Laurencot, Bogdan-Vasile Matioc. A gradient flow approach to a thin film approximation of the Muskat problem. *Calculus of Variations and Partial Differential Equations*, Springer Verlag, 2013, 47, pp.319-341. <10.1007/s00526-012-0520-5>. <hal-00636647>

HAL Id: hal-00636647

<https://hal.archives-ouvertes.fr/hal-00636647>

Submitted on 28 Oct 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A GRADIENT FLOW APPROACH TO A THIN FILM APPROXIMATION OF THE MUSKAT PROBLEM

PHILIPPE LAURENÇOT AND BOGDAN-VASILE MATIOC

ABSTRACT. A fully coupled system of two second-order parabolic degenerate equations arising as a thin film approximation to the Muskat problem is interpreted as a gradient flow for the 2-Wasserstein distance in the space of probability measures with finite second moment. A variational scheme is then set up and is the starting point of the construction of weak solutions. The availability of two Liapunov functionals turns out to be a central tool to obtain the needed regularity to identify the Euler-Lagrange equation in the variational scheme.

1. INTRODUCTION

The Muskat model is a free boundary problem describing the motion of two immiscible fluids with different densities and viscosities in a porous medium (such as intrusion of water into oil). Assuming that the thickness of the two fluid layers is small, a thin film approximation to the Muskat problem has been recently derived in [10] for the space and time evolution of the thickness $f = f(t, x) \geq 0$ and $g = g(t, x) \geq 0$ of the two fluids ($f + g$ being then the total height of the layer) and reads

$$\begin{cases} \partial_t f &= (1 + R)\partial_x (f\partial_x f) + R\partial_x (f\partial_x g), \\ \partial_t g &= R_\mu\partial_x (g\partial_x f) + R_\mu\partial_x (g\partial_x g), \end{cases} \quad (t, x) \in (0, \infty) \times \mathbb{R}, \quad (1.1a)$$

supplemented with the initial conditions

$$f(0) = f_0, \quad g(0) = g_0, \quad x \in \mathbb{R}. \quad (1.1b)$$

Here, R and R_μ are two positive real numbers depending on the densities and the viscosities of the fluids. Since f and g may vanish, (1.1a) is a strongly coupled degenerate parabolic system with a full diffusion matrix due to the terms $\partial_x(f\partial_x g)$ and $\partial_x(g\partial_x f)$. There is however an underlying structure which results in the availability of an energy functional

$$\mathcal{E}(f, g) := \frac{1}{2} \int_{\mathbb{R}} [f^2 + R(f + g)^2] dx, \quad (1.2)$$

which decreases along the flow. More precisely, a formal computation reveals that

$$\frac{d}{dt} \mathcal{E}(f, g) = - \int_{\mathbb{R}} \left[f \left((1 + R)\partial_x f + R\partial_x g \right)^2 + R R_\mu g \left(\partial_x f + \partial_x g \right)^2 \right] dx. \quad (1.3)$$

A similar property is actually valid when (1.1a) is set on a bounded interval $(0, L)$ with homogeneous Neumann boundary conditions: in that setting, the stationary solutions are constants and the principle

Date: October 28, 2011.

2010 Mathematics Subject Classification. 35K65, 35K40, 47J30, 35Q35.

Key words and phrases. thin film, degenerate parabolic system, gradient flow, Wasserstein distance.

Partially supported by the french-german PROCOPE program 20190SE.

of linearized stability is used in [10] to construct global classical solutions which stay in a small neighbourhood of positive constant stationary states. Local existence and uniqueness of classical solutions (with positive components) are also established in [10] by using the general theory for nonlinear parabolic systems developed in [2]. Weak solutions have been subsequently constructed in [9] by a compactness method: the first step is to study a regularized system in which the cross-diffusion terms are “weakened” and to show that it has global strong solutions, the proof combining the theory from [2] for the local well-posedness and suitable estimates for the global existence. Some of these estimates turn out to be independent of the regularisation parameter and provide sufficient information to pass to the limit as the regularisation parameter goes to zero and obtain a weak solution to (1.1a) in a second step. A key argument in the analysis of [9] was to notice that there is another Liapunov functional for (1.1a) given by

$$\mathcal{H}(f, g) := \int_{\mathbb{R}} \left[f \ln f + \frac{R}{R_\mu} g \ln g \right] dx, \quad (1.4)$$

which evolves along the flow as follows:

$$\frac{d}{dt} \mathcal{H}(f, g) = - \int_{\mathbb{R}} [|\partial_x f|^2 + R |\partial_x f + \partial_x g|^2] dx.$$

The basic idea behind the above computation is to notice that an alternative formulation of (1.1a) is

$$\begin{cases} \partial_t f &= \partial_x [f \partial_x ((1+R)f + Rg)], \\ \partial_t g &= R_\mu \partial_x [g \partial_x (f+g)], \end{cases} \quad (t, x) \in (0, \infty) \times \mathbb{R},$$

so that it is rather natural to multiply the f -equation by $\ln f$ and the g -equation by $\ln g$ and find nice cancellations after integrating by parts. In this note, we go one step further and observe that a concise formulation of (1.1a) is actually

$$\begin{cases} \partial_t f &= \partial_x \left[f \partial_x \left(\frac{\delta \mathcal{E}}{\delta f}(f, g) \right) \right], \\ \frac{R}{R_\mu} \partial_t g &= \partial_x \left[g \partial_x \left(\frac{\delta \mathcal{E}}{\delta g}(f, g) \right) \right], \end{cases} \quad (t, x) \in (0, \infty) \times \mathbb{R}, \quad (1.5)$$

which is strongly reminiscent of the interpretation of second-order parabolic equations as gradient flows with respect to the 2-Wasserstein distance, see [3, Chapter 11] and [18, Chapter 8]. Indeed, since the pioneering works [12] on the linear Fokker-Planck equation and [15, 16] on the porous medium equation, several equations have been interpreted as gradient flows with respect to some Wasserstein metrics, including doubly degenerate parabolic equations [1], a model for type-II semiconductors [4], the Smoluchowski-Poisson equation [5], some kinetic equations [6, 8], and some fourth-order degenerate parabolic equations [14], to give a few examples, see also [3] for a general approach. As far as we know, the system (1.5) seems to be the first example of a system of parabolic partial differential equations which can be interpreted as a gradient flow for Wasserstein metrics. Let us however mention that the parabolic-parabolic Keller-Segel system arising in the modeling of chemotaxis has a mixed Wasserstein- L^2 gradient flow structure [7].

The purpose of this note is then to show that the heuristic argument outlined previously can be made rigorous and to construct weak solutions to (1.1) by this approach. More precisely, let \mathcal{K} be the convex

subset of the Banach space $L^1(\mathbb{R}, (1+x^2)dx) \cap L^2(\mathbb{R})$ defined by

$$\mathcal{K} := \left\{ h \in L^1(\mathbb{R}, (1+x^2)dx) \cap L^2(\mathbb{R}) : h \geq 0 \text{ a.e. and } \int_{\mathbb{R}} h(x) dx = 1 \right\}, \quad (1.6)$$

and consider initial data $(f_0, g_0) \in \mathcal{K}_2 := \mathcal{K} \times \mathcal{K}$. We next denote the set of Borel probability measures on \mathbb{R} with finite second moment by $\mathcal{P}_2(\mathbb{R})$ and the 2–Wasserstein distance on $\mathcal{P}_2(\mathbb{R})$ by W_2 . Recall that, given two Borel probability measures μ and ν in $\mathcal{P}_2(\mathbb{R})$,

$$W_2^2(\mu, \nu) := \inf_{\pi \in \Pi(\mu, \nu)} \int_{\mathbb{R}^2} |x - y|^2 d\pi(x, y),$$

where $\Pi(\mu, \nu)$ is the set of all probability measures $\pi \in \mathcal{P}(\mathbb{R}^2)$ which have marginals μ and ν , that is $\pi[A \times \mathbb{R}] = \mu[A]$ and $\pi[\mathbb{R} \times B] = \nu[B]$ for all measurable subsets A and B of \mathbb{R} . Alternatively, $\pi \in \Pi(\mu, \nu)$ is equivalent to

$$\int_{\mathbb{R}^2} (\phi(x) + \psi(y)) d\pi(x, y) = \int_{\mathbb{R}} \phi(x) d\mu(x) + \int_{\mathbb{R}} \psi(y) d\nu(y) \quad \text{for all } (\phi, \psi) \in L^1(\mathbb{R}; \mathbb{R}^2).$$

With these notation, our result reads:

Theorem 1.1. *Assume that $R > 0$, $R_\mu > 0$. Given $\tau > 0$ and $(f_0, g_0) \in \mathcal{K}_2$, the sequence $(f_\tau^n, g_\tau^n)_{n \geq 0}$ obtained recursively by setting*

$$(f_\tau^0, g_\tau^0) := (f_0, g_0), \quad (1.7)$$

$$\mathcal{F}_\tau^n (f_\tau^{n+1}, g_\tau^{n+1}) := \inf_{(u, v) \in \mathcal{K}_2} \mathcal{F}_\tau^n(u, v), \quad (1.8)$$

with

$$\mathcal{F}_\tau^n(u, v) := \frac{1}{2\tau} \left(W_2^2(u, f_\tau^n) + \frac{R}{R_\mu} W_2^2(v, g_\tau^n) \right) + \mathcal{E}(u, v), \quad (u, v) \in \mathcal{K}_2, \quad (1.9)$$

is well-defined. Introducing the interpolation (f_τ, g_τ) defined by

$$f_\tau(t) := f_\tau^n \text{ and } g_\tau(t) := g_\tau^n \text{ for } t \in [n\tau, (n+1)\tau) \text{ and } n \geq 0, \quad (1.10)$$

there exist a sequence $(\tau_k)_{k \geq 1}$ of positive real numbers, $\tau_k \searrow 0$, and functions $(f, g) : [0, \infty) \rightarrow \mathcal{K}_2$ such that

$$(f_{\tau_k}, g_{\tau_k}) \longrightarrow (f, g) \text{ in } L^2((0, T) \times \mathbb{R}; \mathbb{R}^2) \text{ for all } T > 0. \quad (1.11)$$

Moreover,

- (i) $(f, g) \in L^\infty(0, \infty; L^2(\mathbb{R}; \mathbb{R}^2))$, $(\partial_x f, \partial_x g) \in L^2(0, t; H^1(\mathbb{R}; \mathbb{R}^2))$,
- (ii) $(f, g) \in C([0, \infty); H^{-3}(\mathbb{R}; \mathbb{R}^2))$ with $(f, g)(0) = (f_0, g_0)$,

and the pair (f, g) is a weak solution of (1.1) in the sense that

$$\begin{cases} \int_{\mathbb{R}} f(t) \xi dx - \int_{\mathbb{R}} f_0 \xi dx + \int_0^t \int_{\mathbb{R}} f(\sigma) [(1+R)\partial_x f + R\partial_x g](\sigma) \partial_x \xi dx d\sigma = 0, \\ \int_{\mathbb{R}} g(t) \xi dx - \int_{\mathbb{R}} g_0 \xi dx + R_\mu \int_0^t \int_{\mathbb{R}} g(\sigma) (\partial_x f + \partial_x g)(\sigma) \partial_x \xi dx d\sigma = 0, \end{cases} \quad (1.12)$$

for all $\xi \in C_0^\infty(\mathbb{R})$ and $t \geq 0$. In addition, (f, g) satisfy the following estimates

$$(a) \quad \mathcal{H}(f(T), g(T)) + \int_0^T \int_{\mathbb{R}} [|\partial_x f|^2 + R|\partial_x(f+g)|^2] dx dt \leq \mathcal{H}(f_0, g_0),$$

$$(b) \quad \mathcal{E}(f(T), g(T)) + \frac{1}{2} \int_0^T \int_{\mathbb{R}} [f((1+R)\partial_x f + R\partial_x g)^2 + RR_\mu g(\partial_x f + \partial_x g)^2] dx dt \leq \mathcal{E}(f_0, g_0),$$

for a.e. $T \in (0, \infty)$, \mathcal{E} and \mathcal{H} being the functionals defined by (1.2) and (1.4), respectively.

Let us briefly outline the proof of Theorem 1.1: in the next section, we study the variational problem (1.8) and the properties of its minimizers. A key argument here is to note that the availability of the Liapunov functional (1.4) allows us to apply an argument from [14] which guarantees that the minimizers are not only in $L^2(\mathbb{R}; \mathbb{R}^2)$ but also in $H^1(\mathbb{R}; \mathbb{R}^2)$. This property is crucial in order to derive the Euler-Lagrange equation in Section 2.2. The latter is then used to obtain additional regularity on the minimizers, adapting an argument from [15]. Convergence of the variational approximation is established in Section 3. Finally, three technical results are collected in the Appendix.

As a final comment, let us point out that we have assumed for simplicity that the initial data f_0 and g_0 are probability measures but that the case of initial data having different masses may be handled in the same way after a suitable rescaling: more precisely, let $(f_0, g_0) \in L^2(\mathbb{R}) \cap L^1(\mathbb{R}, (1+x^2)dx)$ and denote a solution to (1.1) by (f, g) . Setting $F := f/\|f_0\|_1$ and $G := g/\|g_0\|_1$ and recalling that $\|f(t)\|_1 = \|f_0\|_1$ and $\|g(t)\|_1 = \|g_0\|_1$ for all $t \geq 0$, we realize that (F, G) solves

$$\begin{cases} \frac{1}{\|g_0\|_1} \partial_t F &= \partial_x [F \partial_x ((1+R)\eta^2 F + RG)], \\ \frac{R}{R_\mu \|f_0\|_1} \partial_t G &= \partial_x [G \partial_x (RF + R\eta^{-2}G)], \end{cases} \quad (t, x) \in (0, \infty) \times \mathbb{R},$$

with $\eta^2 := \|f_0\|_1/\|g_0\|_1$ and initial data $(F_0, G_0) := (f_0/\|f_0\|_1, g_0/\|g_0\|_1) \in \mathcal{K}_2$. The corresponding variational scheme then involves the functional

$$\frac{1}{2\tau} \left(\frac{1}{\|g_0\|_1} W_2^2(u, F_0) + \frac{R}{R_\mu \|f_0\|_1} W_2^2(v, G_0) \right) + \frac{\eta^2}{2} \|u\|_2^2 + \frac{R}{2} \|\eta u + \eta^{-1} v\|_2^2, \quad (u, v) \in \mathcal{K}_2,$$

to which the analysis performed below (with $\eta = 1$) also applies.

2. A VARIATIONAL SCHEME

Given $\tau > 0$ and $(f_0, g_0) \in \mathcal{K}_2$, we introduce the functional

$$\mathcal{F}_\tau(u, v) := \frac{1}{2\tau} \left(W_2^2(u, f_0) + \frac{R}{R_\mu} W_2^2(v, g_0) \right) + \mathcal{E}(u, v), \quad (u, v) \in \mathcal{K}_2, \quad (2.1)$$

and consider the minimization problem

$$\inf_{(u, v) \in \mathcal{K}_2} \mathcal{F}_\tau(u, v). \quad (2.2)$$

2.1. Existence and properties of minimizers. Let us start by proving that, for each $(f_0, g_0) \in \mathcal{K}_2$, the minimization problem (2.2) has a unique solution in \mathcal{K}_2 .

Lemma 2.1. *Given $(f_0, g_0) \in \mathcal{K}_2$ and $\tau > 0$, there exists a unique minimizer $(f, g) \in \mathcal{K}_2$ of (2.2). Additionally, $(f, g) \in H^1(\mathbb{R}; \mathbb{R}^2)$ with*

$$\|\partial_x f\|_2^2 + R\|\partial_x(f+g)\|_2^2 \leq \frac{1}{\tau} \left[H(f_0) - H(f) + \frac{R}{R_\mu} (H(g_0) - H(g)) \right], \quad (2.3)$$

where

$$H(h) := \int_{\mathbb{R}} h \ln(h) dx \quad \text{for } h \in L^1(\mathbb{R}) \text{ such that } h \geq 0 \text{ a.e. and } h \ln(h) \in L^1(\mathbb{R}). \quad (2.4)$$

Recall that, if $h \in \mathcal{K}$, then $h \ln h \in L^1(\mathbb{R})$ (see Lemma A.1 below) so that the right-hand side of (2.3) is well-defined.

Proof. The uniqueness of the minimizer follows from the convexity of \mathcal{K}_2 and W_2^2 and the strict convexity of the energy functional \mathcal{E} .

We next prove the existence of a minimizer. To this end, pick a minimizing sequence $(u_k, v_k)_{k \geq 1} \in \mathcal{K}_2$. There exists a constant $C > 0$ such that

$$\|u_k\|_2 + \|v_k\|_2 \leq C, \quad k \geq 1, \quad (2.5)$$

$$W_2(u_k, f_0) + W_2(v_k, g_0) \leq C, \quad k \geq 1. \quad (2.6)$$

From (2.5) we obtain at once that there exist $(f, g) \in L^2(\mathbb{R}; \mathbb{R}^2)$ and a subsequence of $(u_k, v_k)_{k \geq 1}$ (denoted again by $(u_k, v_k)_{k \geq 1}$) such that

$$u_k \rightharpoonup f \quad \text{and} \quad v_k \rightharpoonup g \quad \text{in } L^2(\mathbb{R}). \quad (2.7)$$

Let us first check that $(f, g) \in \mathcal{K}_2$. Indeed, the nonnegativity of f and g readily follows from that of u_k and v_k by (2.7) while integrating the inequality $x^2 \leq 2y^2 + 2|x-y|^2$ with respect to an arbitrary $\pi \in \Pi(u_k, f_0)$ yields

$$\begin{aligned} \int_{\mathbb{R}} u_k(x) x^2 dx &= \int_{\mathbb{R}^2} x^2 d\pi(x, y) \leq 2 \int_{\mathbb{R}^2} y^2 d\pi(x, y) + 2 \int_{\mathbb{R}^2} |x-y|^2 d\pi(x, y) \\ &\leq 2 \int_{\mathbb{R}^2} f_0(y) y^2 dy + 2 \int_{\mathbb{R}^2} |x-y|^2 d\pi(x, y), \end{aligned}$$

which implies by virtue of (2.6) that

$$\int_{\mathbb{R}} u_k(x) x^2 dx \leq 2 \int_{\mathbb{R}} f_0(x) x^2 dx + 2W_2^2(u_k, f_0) \leq C, \quad k \geq 1. \quad (2.8)$$

Similarly,

$$\int_{\mathbb{R}} v_k(x) x^2 dx \leq C, \quad k \geq 1. \quad (2.9)$$

Owing to (2.5), (2.8), and (2.9), we deduce from the Dunford-Pettis theorem that $(u_k)_{k \geq 1}$ and $(v_k)_{k \geq 1}$ are weakly sequentially compact in $L^1(\mathbb{R})$. We may thus assume (after possibly extracting a further subsequence) that $u_k \rightharpoonup f$ and $v_k \rightharpoonup g$ in $L^1(\mathbb{R})$, whence

$$\int_{\mathbb{R}} f(x) dx = \lim_{k \rightarrow \infty} \int_{\mathbb{R}} u_k(x) dx = 1 \quad \text{and} \quad \int_{\mathbb{R}} g(x) dx = \lim_{k \rightarrow \infty} \int_{\mathbb{R}} v_k(x) dx = 1.$$

Finally, combining (2.5), (2.8), and (2.9) with a truncation argument ensure that f and g both belong to $L^1(\mathbb{R}, (1+x^2)dx)$. Summarising, we have shown that $(f, g) \in \mathcal{K}_2$.

The next step is to prove that

$$\mathcal{F}_\tau(f, g) = \inf_{(u, v) \in \mathcal{K}_2} \mathcal{F}_\tau(u, v).$$

Indeed, on the one hand, the weak convergence (2.7) implies that

$$\mathcal{E}(f, g) \leq \liminf_{k \rightarrow \infty} \mathcal{E}(u_k, v_k).$$

On the other hand, we recall that the 2-Wasserstein metric W_2 is lower semicontinuous with respect to the narrow convergence of probability measures in each of its arguments, see [3, Proposition 7.1.3], and the weak convergence of $(u_k, v_k)_{k \geq 1}$ in $L^1(\mathbb{R}; \mathbb{R}^2)$ ensures that

$$W_2^2(f, f_0) \leq \liminf_{k \rightarrow \infty} W_2^2(u_k, f_0) \quad \text{and} \quad W_2^2(g, g_0) \leq \liminf_{k \rightarrow \infty} W_2^2(v_k, g_0).$$

Consequently,

$$\mathcal{F}_\tau(f, g) \leq \liminf_{k \rightarrow \infty} \mathcal{F}_\tau(u_k, v_k) \quad \text{with} \quad (f, g) \in \mathcal{K}_2,$$

so that (f, g) is a minimizer of \mathcal{F}_τ in \mathcal{K}_2 .

As a final step, we show that f and g belong to $H^1(\mathbb{R})$. To this end, we follow the approach developed in [14] and take advantage of the availability of another Liapunov function as already discussed in the Introduction. More precisely, denote the heat semigroup by $(G_t)_{t \geq 0}$, that is,

$$(G_t h)(x) := \frac{1}{\sqrt{4\pi t}} \int_{\mathbb{R}} \exp\left(-\frac{|x-y|^2}{4t}\right) h(y) dy, \quad (t, x) \in [0, \infty) \times \mathbb{R},$$

for $h \in L^1(\mathbb{R})$. Since $(f, g) \in \mathcal{K}_2$, classical properties of the heat semigroup ensure that $(G_t f, G_t g) \in \mathcal{K}_2$ for all $t \geq 0$. Consequently, $\mathcal{F}_\tau(f, g) \leq \mathcal{F}_\tau(G_t f, G_t g)$ and we deduce that

$$\mathcal{E}(f, g) - \mathcal{E}(G_t f, G_t g) \leq \frac{1}{2\tau} \left[(W_2^2(G_t f, f_0) - W_2^2(f, f_0)) + \frac{R}{R_\mu} (W_2^2(G_t g, g_0) - W_2^2(g, g_0)) \right] \quad (2.10)$$

for all $t \geq 0$. Moreover, for all $t > 0$, we have

$$\frac{d}{dt} \mathcal{E}(G_t f, G_t g) = \int_{\mathbb{R}} [G_t f \partial_t G_t f + R (G_t f + G_t g) \partial_t (G_t f + G_t g)] dx = -\|\partial_x G_t f\|_2^2 - R \|\partial_x G_t (f + g)\|_2^2,$$

and by integration with respect to time we find that

$$\frac{1}{t} \int_0^t [\|\partial_x G_s f\|_2^2 + R \|\partial_x G_s (f + g)\|_2^2] ds \leq \frac{\mathcal{E}(f, g) - \mathcal{E}(G_t f, G_t g)}{t} \quad \text{for all } t > 0.$$

Since $s \mapsto \|\partial_x G_s h\|_2$ is non-increasing for $h \in L^1(\mathbb{R})$ we end up with

$$\|\partial_x G_t f\|_2^2 + R \|\partial_x G_t (f + g)\|_2^2 \leq \frac{\mathcal{E}(f, g) - \mathcal{E}(G_t f, G_t g)}{t} \quad \text{for all } t > 0. \quad (2.11)$$

We recall now some properties of the heat flow in connection with the 2-Wasserstein distance W_2 , see [3, 8, 16], these properties being actually collected in [14, Theorem 2.4]. The heat flow is the gradient flow of the entropy functional H given by (2.4) for W_2 and, for all $(h, \tilde{h}) \in \mathcal{K}_2$, we have [3, Theorem 11.1.4]

$$\frac{1}{2} \frac{d}{dt} W_2^2(G_t h, \tilde{h}) + H(G_t h) \leq H(\tilde{h}) \quad \text{for a.e. } t \geq 0. \quad (2.12)$$

Choosing $(h, \tilde{h}) = (f, f_0)$ and $(h, \tilde{h}) = (g, g_0)$ in (2.12), we obtain

$$\frac{1}{2} \frac{d}{dt} \left[W_2^2(G_t f, f_0) + \frac{R}{R_\mu} W_2^2(G_t g, g_0) \right] \leq H(f_0) - H(G_t f) + \frac{R}{R_\mu} (H(g_0) - H(G_t g))$$

for a.e. $t \geq 0$. Integrating the above inequality with respect to time and using the time monotonicity of $s \mapsto H(G_s f)$ and $s \mapsto H(G_s g)$ give

$$\begin{aligned} & \frac{1}{2} \left[W_2^2(G_t f, f_0) - W_2^2(f, f_0) + \frac{R}{R_\mu} (W_2^2(G_t g, g_0) - W_2^2(g, g_0)) \right] \\ & \leq \int_0^t \left[H(f_0) - H(G_s f) + \frac{R}{R_\mu} (H(g_0) - H(G_s g)) \right] ds \\ & \leq t \left[H(f_0) - H(G_t f) + \frac{R}{R_\mu} (H(g_0) - H(G_t g)) \right]. \end{aligned} \quad (2.13)$$

Gathering (2.10), (2.11), and (2.13), we find

$$\|\partial_x G_t f\|_2^2 + R \|\partial_x G_t(f + g)\|_2^2 \leq \frac{1}{\tau} \left[H(f_0) - H(G_t f) + \frac{R}{R_\mu} (H(g_0) - H(G_t g)) \right] \quad (2.14)$$

for $t > 0$. As a direct consequence of (2.14) and the boundedness from below (A.2) of H in \mathcal{K} , $(\partial_x G_t f)_{t>0}$ and $(\partial_x G_t g)_{t>0}$ are bounded in $L^2(\mathbb{R})$ and converge to $\partial_x f$ and $\partial_x g$, respectively, in the sense of distributions as $t \rightarrow 0$. This implies that both f and g belongs to $H^1(\mathbb{R})$ and we can pass to the limit as $t \rightarrow 0$ in (2.14) to obtain the desired estimate (2.3) and finish the proof. \square

2.2. The Euler-Lagrange equation. We now identify the Euler-Lagrange equation corresponding to the minimization problem (2.2).

Lemma 2.2. *Given $(f_0, g_0) \in \mathcal{K}_2$ and $\tau > 0$, the minimizer (f, g) of \mathcal{F}_τ in \mathcal{K}_2 satisfies*

$$\left| \frac{1}{\tau} \int_{\mathbb{R}} \xi (f - f_0) dx + \int_{\mathbb{R}} [(1 + R) f \partial_x f + R f \partial_x g] \partial_x \xi dx \right| \leq \frac{\|\partial_x^2 \xi\|_\infty}{2\tau} W_2^2(f, f_0), \quad (2.15)$$

$$\left| \frac{1}{\tau} \int_{\mathbb{R}} \xi (g - g_0) dx + R_\mu \int_{\mathbb{R}} [g \partial_x f + g \partial_x g] \partial_x \xi dx \right| \leq \frac{\|\partial_x^2 \xi\|_\infty}{2\tau} W_2^2(g, g_0), \quad (2.16)$$

for $\xi \in C_0^\infty(\mathbb{R})$.

Proof. To derive (2.15)-(2.16) we follow the general strategy outlined in [18, Chapter 8]. According to Brenier's theorem [18, Theorem 2.12], there are two convex functions $\varphi : \mathbb{R} \rightarrow \mathbb{R}$ and $\psi : \mathbb{R} \rightarrow \mathbb{R}$ which are uniquely determined up to an additive constant such that

$$W_2^2(f, f_0) = \int_{\mathbb{R}} |x - \partial_x \varphi(x)|^2 f_0(x) dx = \inf_{T \# f_0 = f} \int_{\mathbb{R}} |x - T(x)|^2 f_0(x) dx, \quad (2.17a)$$

where the infimum is taken over all measurable functions $T : \mathbb{R} \rightarrow \mathbb{R}$ pushing f_0 forward to f ($f = T \# f_0$), i.e. satisfying

$$\int_B f(x) dx = \int_{T^{-1}(B)} f_0(x) dx \quad \text{for all Borel sets } B \text{ of } \mathbb{R},$$

and

$$W_2^2(g, g_0) = \int_{\mathbb{R}} |x - \partial_x \psi(x)|^2 g_0(x) dx = \inf_{S \# g_0 = g} \int_{\mathbb{R}} |x - S(x)|^2 g_0(x) dx. \quad (2.17b)$$

We pick now two test functions η and ξ in $C_0^\infty(\mathbb{R})$ and define

$$f_\varepsilon := ((\text{id} + \varepsilon\xi) \circ \partial_x \varphi) \# f_0 = (\text{id} + \varepsilon\xi) \# f \quad \text{and} \quad g_\varepsilon := ((\text{id} + \varepsilon\eta) \circ \partial_x \psi) \# g_0 = (\text{id} + \varepsilon\eta) \# g \quad (2.18)$$

for each $\varepsilon \in [0, 1]$, where id is the identity function on \mathbb{R} . To ease notation we set

$$T_\varepsilon := \text{id} + \varepsilon\xi \quad \text{and} \quad S_\varepsilon := \text{id} + \varepsilon\eta, \quad (2.19)$$

and observe that there is ε_0 small enough (depending on both ξ and η) such that, for $\varepsilon \in [0, \varepsilon_0]$, T_ε and S_ε are C^∞ -diffeomorphisms in \mathbb{R} . Then, by (2.18), we find the identities

$$f_\varepsilon = \frac{f \circ T_\varepsilon^{-1}}{\partial_x T_\varepsilon \circ T_\varepsilon^{-1}} \quad \text{and} \quad g_\varepsilon = \frac{g \circ S_\varepsilon^{-1}}{\partial_x S_\varepsilon \circ S_\varepsilon^{-1}}, \quad \varepsilon \in (0, \varepsilon_0]. \quad (2.20)$$

Observing that $\|f_\varepsilon\|_1 = \|f\|_1 = \|g_\varepsilon\|_1 = \|g\|_1 = 1$ and

$$\|f_\varepsilon\|_2^2 = \int_{\mathbb{R}} \frac{|f(x)|^2}{\partial_x T_\varepsilon(x)} dx \quad \text{and} \quad \|g_\varepsilon\|_2^2 = \int_{\mathbb{R}} \frac{|g(x)|^2}{\partial_x S_\varepsilon(x)} dx, \quad (2.21)$$

we clearly have $(f_\varepsilon, g_\varepsilon) \in \mathcal{K}_2$ for all $\varepsilon \in (0, \varepsilon_0]$ and thus $\mathcal{F}_\tau(f, g) \leq \mathcal{F}_\tau(f_\varepsilon, g_\varepsilon)$. Consequently,

$$0 \leq \frac{1}{2\tau} \left[W_2^2(f_\varepsilon, f_0) - W_2^2(f, f_0) + \frac{R}{R_\mu} (W_2^2(g_\varepsilon, g_0) - W_2^2(g, g_0)) \right] + \mathcal{E}(f_\varepsilon, g_\varepsilon) - \mathcal{E}(f, g). \quad (2.22)$$

Concerning the energy \mathcal{E} , it follows from (2.21) that

$$2(\mathcal{E}(f_\varepsilon, g_\varepsilon) - \mathcal{E}(f, g)) = (1 + R) I_1^\varepsilon + R I_2^\varepsilon + 2R I_3^\varepsilon, \quad (2.23)$$

with

$$I_1^\varepsilon := \int_{\mathbb{R}} \left(\frac{1}{\partial_x T_\varepsilon(x)} - 1 \right) |f(x)|^2 dx, \quad I_2^\varepsilon := \int_{\mathbb{R}} \left(\frac{1}{\partial_x S_\varepsilon(x)} - 1 \right) |g(x)|^2 dx, \\ I_3^\varepsilon := \int_{\mathbb{R}} (f_\varepsilon g_\varepsilon - f g)(x) dx.$$

We now consider the three integrals in the right-hand side of the relation (2.23) separately: since

$$I_1^\varepsilon = -\varepsilon \int_{\mathbb{R}} \frac{\partial_x \xi}{1 + \varepsilon \partial_x \xi} f^2 dx \quad \text{and} \quad I_2^\varepsilon = -\varepsilon \int_{\mathbb{R}} \frac{\partial_x \eta}{1 + \varepsilon \partial_x \eta} g^2 dx,$$

it readily follows from Lebesgue's dominated convergence theorem that

$$\lim_{\varepsilon \rightarrow 0} \frac{I_1^\varepsilon}{\varepsilon} = - \int_{\mathbb{R}} \partial_x \xi f^2 dx \quad \text{and} \quad \lim_{\varepsilon \rightarrow 0} \frac{I_2^\varepsilon}{\varepsilon} = - \int_{\mathbb{R}} \partial_x \eta g^2 dx. \quad (2.24)$$

We next turn to the term I_3^ε involving both f and g and split it in two terms $2I_3^\varepsilon = I_{31}^\varepsilon + I_{32}^\varepsilon$ with

$$I_{31}^\varepsilon := \int_{\mathbb{R}} (f_\varepsilon + f)(g_\varepsilon - g) dx \quad \text{and} \quad I_{32}^\varepsilon := \int_{\mathbb{R}} (g_\varepsilon + g)(f_\varepsilon - f) dx.$$

By (2.20),

$$I_{31}^\varepsilon = \int_{\mathbb{R}} \left(\frac{f \circ T_\varepsilon^{-1}}{\partial_x T_\varepsilon \circ T_\varepsilon^{-1}} + f \right) \left(\frac{g \circ S_\varepsilon^{-1}}{\partial_x S_\varepsilon \circ S_\varepsilon^{-1}} - g \right) dx \\ = \int_{\mathbb{R}} \left(\frac{f \circ T_\varepsilon^{-1} \circ S_\varepsilon}{\partial_x T_\varepsilon \circ T_\varepsilon^{-1} \circ S_\varepsilon} + f \circ S_\varepsilon \right) (g - (g \circ S_\varepsilon) \partial_x S_\varepsilon) dx \\ = \int_{\mathbb{R}} \left(\frac{f \circ T_\varepsilon^{-1} \circ S_\varepsilon}{\partial_x T_\varepsilon \circ T_\varepsilon^{-1} \circ S_\varepsilon} + f \circ S_\varepsilon \right) (g - g \circ S_\varepsilon) dx - \varepsilon \int_{\mathbb{R}} \partial_x \eta (g \circ S_\varepsilon) \left(\frac{f \circ T_\varepsilon^{-1} \circ S_\varepsilon}{\partial_x T_\varepsilon \circ T_\varepsilon^{-1} \circ S_\varepsilon} + f \circ S_\varepsilon \right) dx.$$

On the one hand, invoking Lemma A.2 (with $(h, \zeta) = (g, \eta)$), we know that $(g - g \circ S_\varepsilon)/\varepsilon \rightharpoonup -\eta \partial_x g$ in $L^2(\mathbb{R})$ as $\varepsilon \rightarrow 0$. On the other hand, using again Lemma A.2 as well as Lemma A.3, we have that $f \circ S_\varepsilon \rightarrow f$ and $f \circ T_\varepsilon^{-1} \circ S_\varepsilon \rightarrow f$ in $L^2(\mathbb{R})$ as $\varepsilon \rightarrow 0$, and so does $f \circ T_\varepsilon^{-1} \circ S_\varepsilon / (\partial_x T_\varepsilon \circ T_\varepsilon^{-1} \circ S_\varepsilon)$ owing to the uniform convergence of $(\partial_x T_\varepsilon)_\varepsilon$ to 1. Consequently,

$$\lim_{\varepsilon \rightarrow 0} \frac{I_{31}^\varepsilon}{\varepsilon} = -2 \int_{\mathbb{R}} f \partial_x(\eta g) dx, \quad (2.25)$$

and similarly

$$\lim_{\varepsilon \rightarrow 0} \frac{I_{32}^\varepsilon}{\varepsilon} = -2 \int_{\mathbb{R}} g \partial_x(\xi f) dx. \quad (2.26)$$

Gathering (2.24)-(2.26), it follows from (2.23) that

$$\lim_{\varepsilon \rightarrow 0} \frac{\mathcal{E}(f_\varepsilon, g_\varepsilon) - \mathcal{E}(f, g)}{\varepsilon} = -(1+R) \int_{\mathbb{R}} \frac{f^2}{2} \partial_x \xi dx - R \int_{\mathbb{R}} \frac{g^2}{2} \partial_x \eta dx - R \int_{\mathbb{R}} [f \partial_x(\eta g) + g \partial_x(\xi f)] dx. \quad (2.27)$$

To handle the terms of (2.22) involving the Wasserstein distance, we argue as in [18, Section 8.4] and write

$$\begin{aligned} W_2^2(f_\varepsilon, f_0) &\leq \int_{\mathbb{R}} |\text{id} - T_\varepsilon \circ \partial_x \varphi|^2 f_0 dx = \int_{\mathbb{R}} |\text{id} - \partial_x \varphi - \varepsilon \xi \circ \partial_x \varphi|^2 f_0 dx \\ &= \int_{\mathbb{R}} |\text{id} - \partial_x \varphi|^2 f_0 dx - 2\varepsilon \int_{\mathbb{R}} (\text{id} - \partial_x \varphi) (\xi \circ \partial_x \varphi) f_0 dx + \varepsilon^2 \int_{\mathbb{R}} |\xi \circ \partial_x \varphi|^2 f_0 dx, \end{aligned}$$

from which we deduce, according to the definition of $\partial_x \varphi$,

$$W_2^2(f_\varepsilon, f_0) \leq W_2^2(f, f_0) - 2\varepsilon \int_{\mathbb{R}} (\text{id} - \partial_x \varphi) (\xi \circ \partial_x \varphi) f_0 dx + \varepsilon^2 \int_{\mathbb{R}} |\xi \circ \partial_x \varphi|^2 f_0 dx, \quad (2.28)$$

and similarly

$$W_2^2(g_\varepsilon, g_0) \leq W_2^2(g, g_0) - 2\varepsilon \int_{\mathbb{R}} (\text{id} - \partial_x \psi) (\eta \circ \partial_x \psi) g_0 dx + \varepsilon^2 \int_{\mathbb{R}} |\eta \circ \partial_x \psi|^2 g_0 dx. \quad (2.29)$$

Summing (2.27), (2.28), and (2.29), we obtain by dividing (2.22) by ε and letting $\varepsilon \rightarrow 0$ that

$$\begin{aligned} &\frac{1}{\tau} \left[\int_{\mathbb{R}} (\text{id} - \partial_x \varphi) (\xi \circ \partial_x \varphi) f_0 dx + \frac{R}{R_\mu} \int_{\mathbb{R}} (\text{id} - \partial_x \psi) (\eta \circ \partial_x \psi) g_0 dx \right] \\ &+ (1+R) \int_{\mathbb{R}} \partial_x \xi \frac{f^2}{2} dx + R \int_{\mathbb{R}} \partial_x \eta \frac{g^2}{2} dx + R \int_{\mathbb{R}} [f \partial_x(\eta g) + g \partial_x(\xi f)] dx \leq 0. \end{aligned}$$

Since the relation is valid for (ξ, η) as well as for $(-\xi, -\eta)$, we end up with

$$\begin{aligned} &\frac{1}{\tau} \left[\int_{\mathbb{R}} (\text{id} - \partial_x \varphi) (\xi \circ \partial_x \varphi) f_0 dx + \frac{R}{R_\mu} \int_{\mathbb{R}} (\text{id} - \partial_x \psi) (\eta \circ \partial_x \psi) g_0 dx \right] \\ &+ (1+R) \int_{\mathbb{R}} \partial_x \xi \frac{f^2}{2} dx + R \int_{\mathbb{R}} \partial_x \eta \frac{g^2}{2} dx + R \int_{\mathbb{R}} [f \partial_x(\eta g) + g \partial_x(\xi f)] dx = 0 \end{aligned} \quad (2.30)$$

for all $(\xi, \eta) \in C_0^\infty(\mathbb{R}; \mathbb{R}^2)$.

Consider now $\Xi \in C_0^\infty(\mathbb{R})$. For $x \in \mathbb{R}$, we have

$$\begin{aligned} |\Xi(x) - \Xi(\partial_x \varphi(x)) - \partial_x \Xi(\partial_x \varphi(x)) (x - \partial_x \varphi(x))| &= \left| \int_{\partial_x \varphi(x)}^x (x-y) \partial_x^2 \Xi(y) dy \right| \\ &\leq \|\partial_x^2 \Xi\|_\infty \frac{(x - \partial_x \varphi(x))^2}{2}. \end{aligned}$$

Multiplying the above inequality by $f_0(x)$, integrating over \mathbb{R} , and using the definition of $\partial_x \varphi$ yield

$$\left| \int_{\mathbb{R}} [\Xi(x) - \Xi(\partial_x \varphi(x)) - \partial_x \Xi(\partial_x \varphi(x)) (x - \partial_x \varphi(x))] f_0(x) dx \right| \leq \|\partial_x^2 \Xi\|_\infty \frac{W_2^2(f, f_0)}{2}. \quad (2.31)$$

Owing to (2.30) with $(\xi, \eta) = (\partial_x \Xi, 0)$ and the property $f = \partial_x \varphi \# f_0$, we deduce that

$$\left| \frac{1}{\tau} \int_{\mathbb{R}} (f - f_0) \Xi dx - (1+R) \int_{\mathbb{R}} \frac{f^2}{2} \partial_x^2 \Xi dx - R \int_{\mathbb{R}} g \partial_x (f \partial_x \Xi) dx \right| \leq \frac{1}{2} \|\partial_x^2 \Xi\|_\infty \frac{W_2^2(f, f_0)}{\tau}.$$

Taking into account that $(f, g) \in H^1(\mathbb{R}; \mathbb{R}^2)$ by Lemma 2.1, we arrive, after integrating by parts once, to (2.15). A similar argument leads to (2.16). \square

We next develop further an argument from the proof of [15, Proposition 2] which allows us to gain regularity on f and g by using the Euler-Lagrange equation.

Corollary 2.3. *The functions $\sqrt{f} \partial_x((1+R)f + Rg)$ and $\sqrt{g} \partial_x(f+g)$ both belong to $L^2(\mathbb{R})$ and*

$$\tau \left\| \sqrt{f} \partial_x[(1+R)f + Rg] \right\|_2 \leq W_2(f, f_0), \quad (2.32a)$$

$$\tau R_\mu \left\| \sqrt{g} \partial_x(f+g) \right\|_2 \leq W_2(g, g_0). \quad (2.32b)$$

It is worth mentioning here that the estimates (2.32) match exactly the regularity of (f, g) given by the dissipation in the energy inequality (1.3).

Proof. Consider $\xi \in C_0^\infty(\mathbb{R})$. We infer from (2.30) with $\eta = 0$ that, after integrating by parts,

$$\int_{\mathbb{R}} [(1+R) f \partial_x f + R f \partial_x g] \xi dx = \frac{1}{\tau} \int_{\mathbb{R}} (x - \partial_x \varphi(x)) (\xi \circ \partial_x \varphi)(x) f_0(x) dx.$$

Since $f = \partial_x \varphi \# f_0$, it follows from the Cauchy-Schwarz inequality and (2.17a) that

$$\begin{aligned} &\left| \int_{\mathbb{R}} (x - \partial_x \varphi(x)) (\xi \circ \partial_x \varphi)(x) f_0(x) dx \right| \\ &\leq \left(\int_{\mathbb{R}} (x - \partial_x \varphi(x))^2 f_0(x) dx \right)^{1/2} \left(\int_{\mathbb{R}} (\xi \circ \partial_x \varphi)^2(x) f_0(x) dx \right)^{1/2} \\ &\leq W_2(f, f_0) \left(\int_{\mathbb{R}} \xi^2(x) f(x) dx \right)^{1/2}. \end{aligned}$$

Therefore,

$$\left| \int_{\mathbb{R}} [(1+R) f \partial_x f + R f \partial_x g] \xi dx \right| \leq \frac{W_2(f, f_0)}{\tau} \left(\int_{\mathbb{R}} \xi^2(x) f(x) dx \right)^{1/2}. \quad (2.33)$$

Consider next a nonnegative function $\chi \in C_0^\infty(\mathbb{R})$ such that $\|\chi\|_1 = 1$ and define $\chi_m(x) := m\chi(mx)$ for $m \geq 1$ and $x \in \mathbb{R}$. Then, $(\chi_m)_{m \geq 1}$ is a sequence of mollifiers in \mathbb{R} and, given $\vartheta \in C_0^\infty(\mathbb{R})$ and $m \geq 1$,

the function $\vartheta/(m^{-1/4} + \chi_m * f)^{1/2}$ belongs to $C_0^\infty(\mathbb{R})$. Taking $\xi = \vartheta/(m^{-1/4} + \chi_m * f)^{1/2}$ in (2.33), we obtain

$$\left| \int_{\mathbb{R}} \frac{f \partial_x[(1+R)f + Rg]}{\sqrt{m^{-1/4} + \chi_m * f}} \vartheta dx \right| \leq \frac{W_2(f, f_0)}{\tau} \left\| \frac{f}{m^{-1/4} + \chi_m * f} \right\|_\infty^{1/2} \|\vartheta\|_2.$$

The previous inequality being valid for all $\vartheta \in C_0^\infty(\mathbb{R})$, a duality argument yields

$$\left\| \frac{f \partial_x[(1+R)f + Rg]}{\sqrt{m^{-1/4} + \chi_m * f}} \right\|_2 \leq \frac{W_2(f, f_0)}{\tau} \left\| \frac{f}{m^{-1/4} + \chi_m * f} \right\|_\infty^{1/2}. \quad (2.34)$$

Now, since $f \in H^1(\mathbb{R})$ by Lemma 2.1, we have $\|\chi_m * f - f\|_\infty \leq C_\chi \|\partial_x f\|_2 m^{-1/2}$ for some constant $C_\chi > 0$ depending only on χ from which we deduce that

$$\left\| \frac{f}{m^{-1/4} + \chi_m * f} \right\|_\infty \leq \left\| \frac{f - \chi_m * f}{m^{-1/4} + \chi_m * f} \right\|_\infty + \left\| \frac{\chi_m * f}{m^{-1/4} + \chi_m * f} \right\|_\infty \leq 1 + \frac{C_\chi \|\partial_x f\|_2}{m^{1/4}}. \quad (2.35)$$

In particular, for $x \in \mathbb{R}$,

$$\left| \frac{f(x)}{\sqrt{m^{-1/4} + \chi_m * f(x)}} \right| \leq \left(1 + \sqrt{C_\chi \|\partial_x f\|_2} \right) \sqrt{f(x)} \in L^2(\mathbb{R})$$

and

$$\lim_{m \rightarrow \infty} \frac{f(x)}{\sqrt{m^{-1/4} + \chi_m * f(x)}} = \begin{cases} 0 = \sqrt{f(x)} & \text{if } f(x) = 0, \\ \sqrt{f(x)} & \text{if } f(x) > 0, \end{cases}$$

so that

$$\frac{f}{\sqrt{m^{-1/4} + \chi_m * f}} \longrightarrow \sqrt{f} \quad \text{in } L^2(\mathbb{R})$$

by the Lebesgue dominated convergence theorem. Since $(1+R)f + Rg$ belongs to $H^1(\mathbb{R})$ by Lemma 2.1, we conclude that

$$\frac{f}{\sqrt{m^{-1/4} + \chi_m * f}} \partial_x[(1+R)f + Rg] \longrightarrow \sqrt{f} \partial_x[(1+R)f + Rg] \quad \text{in } L^1(\mathbb{R}). \quad (2.36)$$

Owing to (2.35) and (2.36), we may let $m \rightarrow \infty$ in (2.34) and deduce that $\sqrt{f} \partial_x[(1+R)f + Rg] \in L^2(\mathbb{R})$ and satisfies (2.32a). The proof of (2.32b) is similar. \square

2.3. Interpolation. Thanks to the results established in the previous sections, we are now in a position to set up a variational scheme to approximate the solution to (1.1). More precisely, given $(f_0, g_0) \in \mathcal{K}_2$ and $\tau \in (0, 1)$, we define inductively a sequence $(f_\tau^n, g_\tau^n)_{n \geq 0}$ as follows:

$$(f_\tau^0, g_\tau^0) := (f_0, g_0), \quad (2.37)$$

$$\mathcal{F}_\tau^n(f_\tau^{n+1}, g_\tau^{n+1}) := \inf_{(u,v) \in \mathcal{K}_2} \mathcal{F}_\tau^n(u, v), \quad (2.38)$$

with

$$\mathcal{F}_\tau^n(u, v) := \frac{1}{2\tau} \left(W_2^2(u, f_\tau^n) + \frac{R}{R_\mu} W_2^2(v, g_\tau^n) \right) + \mathcal{E}(u, v), \quad (u, v) \in \mathcal{K}_2,$$

the existence and uniqueness of $(f_\tau^{n+1}, g_\tau^{n+1})$ being guaranteed by Lemma 2.1 for each $n \geq 0$. We next define two interpolation functions f_τ and g_τ by (1.10), i.e. $f_\tau(t) := f_\tau^n$ and $g_\tau(t) := g_\tau^n$ for $t \in [n\tau, (n+1)\tau)$ and $n \geq 0$. By Lemma 2.2, we have

$$\begin{cases} \left| \int_{\mathbb{R}} (f_\tau^n - f_\tau^{n-1}) \xi dx + \tau \int_{\mathbb{R}} f_\tau^n \partial_x ((1+R) f_\tau^n + R g_\tau^n) \partial_x \xi dx \right| \leq \frac{\|\partial_x^2 \xi\|_\infty}{2} W_2^2(f_\tau^n, f_\tau^{n-1}), \\ \left| \int_{\mathbb{R}} (g_\tau^n - g_\tau^{n-1}) \xi dx + \tau R_\mu \int_{\mathbb{R}} g_\tau^n \partial_x (g_\tau^n + g_\tau^n) \partial_x \xi dx \right| \leq \frac{\|\partial_x^2 \xi\|_\infty}{2} W_2^2(g_\tau^n, g_\tau^{n-1}), \end{cases} \quad (2.39)$$

for all $n \geq 1$ and $\xi \in C_0^\infty(\mathbb{R})$. Given $T > 0$ arbitrary, we set $N := [T/\tau]$. Summing both equations of (2.39) from $n = 1$ to $n = N$, we find

$$\left| \int_{\mathbb{R}} (f_\tau(T) - f_0) \xi dx + \int_\tau^{(N+1)\tau} \int_{\mathbb{R}} f_\tau \partial_x ((1+R) f_\tau + R g_\tau) \partial_x \xi dx dt \right| \leq \frac{\|\partial_x^2 \xi\|_\infty}{2} \sum_{n=1}^N W_2^2(f_\tau^n, f_\tau^{n-1}), \quad (2.40)$$

$$\left| \int_{\mathbb{R}} (g_\tau(T) - g_0) \xi dx + R_\mu \int_\tau^{(N+1)\tau} \int_{\mathbb{R}} g_\tau \partial_x (f_\tau + g_\tau) \partial_x \xi dx dt \right| \leq \frac{\|\partial_x^2 \xi\|_\infty}{2} \sum_{n=1}^N W_2^2(g_\tau^n, g_\tau^{n-1}). \quad (2.41)$$

3. CONVERGENCE

We gather in the next lemma various properties of the interpolations (f_τ, g_τ) defined in Section 2.3 which are consequences of Lemma 2.1 and Corollary 2.3.

Lemma 3.1. *There exists a positive constant C_1 depending only on $R, R_\mu, f_0,$ and g_0 such that, for all $t \geq 0$ and $\tau \in (0, 1)$, we have*

$$(i) \quad \int_{\mathbb{R}} f_\tau(t) dx = \int_{\mathbb{R}} g_\tau(t) dx = 1, \quad (3.1)$$

$$(ii) \quad \sum_{n=1}^{\infty} [W_2^2(f_\tau^n, f_\tau^{n-1}) + W_2^2(g_\tau^n, g_\tau^{n-1})] \leq C_1 \tau, \quad (3.2)$$

$$(iii) \quad \mathcal{E}(f_\tau(t), g_\tau(t)) \leq \mathcal{E}(f_\tau(s), g_\tau(s)), \quad s \in [0, t], \quad (3.3)$$

$$(iv) \quad \int_{\mathbb{R}} (f_\tau + g_\tau)(t, x) x^2 dx \leq C_1 (1+t), \quad (3.4)$$

$$(v) \quad \int_\tau^t [\|\partial_x f_\tau(s)\|_2^2 + \|\partial_x g_\tau(s)\|_2^2] ds \leq C_1 (1+t), \quad (3.5)$$

$$(vi) \quad \int_\tau^\infty \int_{\mathbb{R}} f_\tau |\partial_x [(1+R) f_\tau + R g_\tau]|^2 dx ds \leq C_1, \quad (3.6)$$

$$(vii) \quad \int_\tau^\infty \int_{\mathbb{R}} g_\tau |\partial_x (f_\tau + g_\tau)|^2 dx ds \leq C_1. \quad (3.7)$$

Proof. The property (3.1) readily follows from the fact that $(f_\tau^n, g_\tau^n) \in \mathcal{K}_2$ for all $n \geq 0$ and $\tau > 0$. Next, for $\tau > 0$ and $n \geq 1$, the minimizing property of (f_τ^n, g_τ^n) ensures that

$$\mathcal{E}(f_\tau^n, g_\tau^n) + \frac{1}{2\tau} \left[W_2^2(f_\tau^n, f_\tau^{n-1}) + \frac{R}{R_\mu} W_2^2(g_\tau^n, g_\tau^{n-1}) \right] \leq \mathcal{E}(f_\tau^{n-1}, g_\tau^{n-1}). \quad (3.8)$$

Given $t \in (0, \infty)$ and $s \in [0, t]$, we set $N := [t/\tau]$, $\nu := [s/\tau]$, and sum (3.8) from $n = \nu + 1$ up to $n = N$ to obtain, since $(f_\tau, g_\tau)(t) = (f_\tau^N, g_\tau^N)$ and $(f_\tau, g_\tau)(s) = (f_\tau^\nu, g_\tau^\nu)$,

$$\mathcal{E}(f_\tau(t), g_\tau(t)) + \frac{1}{2\tau} \sum_{n=\nu+1}^N \left[W_2^2(f_\tau^n, f_\tau^{n-1}) + \frac{R}{R_\mu} W_2^2(g_\tau^n, g_\tau^{n-1}) \right] \leq \mathcal{E}(f_\tau(s), g_\tau(s)). \quad (3.9)$$

The monotonicity property (3.3) is a straightforward consequence of (3.9) while the nonnegativity of \mathcal{E} and (3.9) with $s = \nu = 0$ give

$$\sum_{n=1}^N \left[W_2^2(f_\tau^n, f_\tau^{n-1}) + \frac{R}{R_\mu} W_2^2(g_\tau^n, g_\tau^{n-1}) \right] \leq 2\mathcal{E}(f_0, g_0) \tau.$$

Since the right-hand side of the above inequality does not depend on N , we obtain (3.2). In order to prove (3.4), we combine (2.8) and (3.2) and obtain for $t \geq 0$ with $N := [t/\tau]$

$$\begin{aligned} \int_{\mathbb{R}} f_\tau(t, x) x^2 dx &= \int_{\mathbb{R}} f_\tau^N(x) x^2 dx \leq 2 \int_{\mathbb{R}} f_0(x) x^2 dx + 2W_2^2(f_\tau^N, f_0) \\ &\leq 2 \int_{\mathbb{R}} f_0(x) x^2 dx + 2N \sum_{n=1}^N W_2^2(f_\tau^n, f_\tau^{n-1}) \\ &\leq 2 \int_{\mathbb{R}} f_0(x) x^2 dx + 4N\tau \mathcal{E}(f_0, g_0) \leq C(1+t). \end{aligned}$$

We next infer from (2.3) that, for $n \geq 1$,

$$\tau \left(\|\partial_x f_\tau^n\|_2^2 + R \|\partial_x(f_\tau^n + g_\tau^n)\|_2^2 \right) \leq H(f_\tau^{n-1}) - H(f_\tau^n) + \frac{R}{R_\mu} \left(H(g_\tau^{n-1}) - H(g_\tau^n) \right).$$

Let $N \geq 1$. Summation from $n = 1$ to N yields

$$\int_{\tau}^{(N+1)\tau} \left(\|\partial_x f_\tau(s)\|_2^2 + R \|\partial_x(f_\tau + g_\tau)(s)\|_2^2 \right) ds \leq H(f_0) - H(f_\tau(N\tau)) + \frac{R}{R_\mu} \left(H(g_0) - H(g_\tau(N\tau)) \right). \quad (3.10)$$

It now follows from Lemma A.1, (3.1), (3.4), and (3.10) that

$$\begin{aligned} &\int_{\tau}^{(N+1)\tau} \left(\|\partial_x f_\tau(s)\|_2^2 + R \|\partial_x(f_\tau + g_\tau)(s)\|_2^2 \right) ds \\ &\leq H(f_0) + \frac{R}{R_\mu} H(g_0) + \frac{(R + R_\mu)C_\ell}{R_\mu} + \int_{\mathbb{R}} (1+x^2) \left(f_\tau(N\tau) + \frac{R}{R_\mu} g_\tau(N\tau) \right) \leq C(1+N\tau), \end{aligned}$$

which entails the validity of (3.5) for $t \in [N\tau, (N+1)\tau)$.

We finish the proof by showing (3.6) and (3.7). By Corollary 2.3, we have for $n \geq 1$

$$\tau^2 \left\| \sqrt{f_\tau^n} \partial_x [(1+R) f_\tau^n + R g_\tau^n] \right\|_2^2 \leq W_2^2(f_\tau^n, f_\tau^{n-1}).$$

Summing over $n \geq 1$ and using (3.2) give

$$\sum_{n=1}^{\infty} \tau^2 \left\| \sqrt{f_\tau^n} \partial_x [(1+R) f_\tau^n + R g_\tau^n] \right\|_2^2 \leq \sum_{n=1}^{\infty} W_2^2(f_\tau^n, f_\tau^{n-1}) \leq C_1\tau,$$

whence (3.6). The proof of (3.7) also relies on Corollary 2.3 and is similar. \square

3.1. Compactness. We now turn to the compactness properties of $(f_\tau)_{\tau>0}$ and $(g_\tau)_{\tau>0}$ and point out that the nonlinearity of (1.1a) requires strong compactness. We first observe that the compactness with respect to the space variable x is granted by (3.5) thanks to the following lemma.

Lemma 3.2. *The spaces $H^1(\mathbb{R}) \cap L^1(\mathbb{R}, (1+x^2) dx)$ and $L^2(\mathbb{R}) \cap L^1(\mathbb{R}, (1+x^2) dx)$ are compactly embedded in $L^2(\mathbb{R})$ and $H^{-3}(\mathbb{R})$, respectively.*

Proof. Let us first consider a bounded sequence $(h_i)_{i \geq 1}$ in $H^1(\mathbb{R}) \cap L^1(\mathbb{R}, (1+x^2) dx)$. On the one hand, since $H^1(\mathbb{R})$ is continuously embedded in $L^\infty(\mathbb{R})$ and $C^{1/2}(\mathbb{R})$, the Arzelà-Ascoli theorem implies that there are $h \in H^1(\mathbb{R})$ and a subsequence of $(h_i)_{i \geq 1}$ (not relabeled), such that $(h_i)_{i \geq 1}$ converges to h in $C([-R, R])$ for all $R > 0$. On the other hand, using once more the embedding of $H^1(\mathbb{R})$ in $L^\infty(\mathbb{R})$, we have for $R > 1$

$$\begin{aligned} \int_{\mathbb{R}} |h_i(x) - h(x)|^2 dx &\leq \int_{\{|x| \leq R\}} |h_i(x) - h(x)|^2 dx + \int_{\{|x| > R\}} |h_i(x) - h(x)|^2 dx \\ &\leq 2R \|h_i - h\|_{C([-R, R])}^2 + \frac{1}{R^2} \|h_i - h\|_\infty \int_{\mathbb{R}} x^2 |h_i(x) - h(x)| dx \\ &\leq 2R \|h_i - h\|_{C([-R, R])}^2 + \frac{2}{R^2} \sup_{i \geq 1} \left\{ \|h_i\|_\infty \int_{\mathbb{R}} x^2 |h_i(x)| dx \right\}. \end{aligned}$$

Letting first $i \rightarrow \infty$ and then $R \rightarrow \infty$ shows that $(h_i)_{i \geq 1}$ converges to h in $L^2(\mathbb{R})$.

Next, let $(h_i)_{i \geq 1}$ be a bounded sequence in $L^2(\mathbb{R}) \cap L^1(\mathbb{R}, (1+x^2) dx)$ and denote the Fourier transform of h_i by $\mathcal{F}h_i$ for $i \geq 1$. A straightforward consequence of the bounds for $(h_i)_{i \geq 1}$ is that $(\mathcal{F}h_i)_{i \geq 1}$ is bounded in $L^2(\mathbb{R}) \cap W^{2, \infty}(\mathbb{R})$. Arguing as above, this implies that $(\mathcal{F}h_i)_{i \geq 1}$ is relatively compact in $L^2(\mathbb{R}, (1+x^2)^{-3} dx)$. Coming back to the original variable, $(h_i)_{i \geq 1}$ is relatively compact in $H^{-3}(\mathbb{R})$ as claimed. \square

We next turn to the compactness in time and prove the following result:

Lemma 3.3. *There is a positive constant C_2 depending only on R , R_μ , f_0 , and g_0 such that, for $\tau \in (0, 1)$ and $(t, s) \in [0, \infty) \times [0, \infty)$,*

$$\|f_\tau(t) - f_\tau(s)\|_{H^{-3}} + \|g_\tau(t) - g_\tau(s)\|_{H^{-3}} \leq C_2 \sqrt{|t-s| + \tau}. \quad (3.11)$$

Proof. Consider $t \in (0, \infty)$, $s \in [0, t]$, and define the integers $N := [t/\tau]$ and $\nu := [s/\tau]$. Either $N = \nu$ and $f_\tau(t) - f_\tau(s) = 0$ satisfies (3.11) or $N \geq \nu + 1$ and it follows from (2.39) that, for $n \in \{\nu + 1, \dots, N\}$ and $\xi \in C_0^\infty(\mathbb{R})$,

$$\begin{aligned} \left| \int_{\mathbb{R}} (f_\tau^n - f_\tau^{n-1}) \xi dx \right| &\leq \int_{n\tau}^{(n+1)\tau} \int_{\mathbb{R}} f_\tau(s) |\partial_x [(1+R) f_\tau + R g_\tau](s)| |\partial_x \xi| dx ds \\ &\quad + \frac{\|\partial_x^2 \xi\|_\infty}{2} W_2^2(f_\tau^n, f_\tau^{n-1}) \end{aligned}$$

Summing the above inequality from $n = \nu + 1$ to $n = N$ and using (3.1), (3.3), (3.6), and the Cauchy-Schwartz inequality, we are led to

$$\begin{aligned}
\left| \int_{\mathbb{R}} (f_{\tau}(t) - f_{\tau}(s)) \xi \, dx \right| &= \left| \int_{\mathbb{R}} (f_{\tau}^N - f_{\tau}^{\nu}) \xi \, dx \right| \leq \sum_{n=\nu+1}^N \left| \int_{\mathbb{R}} (f_{\tau}^n - f_{\tau}^{n-1}) \xi \, dx \right| \\
&\leq \int_{(\nu+1)\tau}^{(N+1)\tau} \int_{\mathbb{R}} f_{\tau}(s) |\partial_x [(1+R) f_{\tau} + R g_{\tau}](s)| |\partial_x \xi| \, dx \, ds \\
&\quad + \frac{\|\partial_x^2 \xi\|_{\infty}}{2} \sum_{n=\nu+1}^N W_2^2(f_{\tau}^n, f_{\tau}^{n-1}) \\
&\leq \|\partial_x \xi\|_{\infty} \int_{(\nu+1)\tau}^{(N+1)\tau} \|f_{\tau}(s)\|_1^{1/2} \left\| \sqrt{f_{\tau}} \partial_x [(1+R) f_{\tau} + R g_{\tau}](s) \right\|_2 \, ds \\
&\quad + C_1 \tau \|\partial_x^2 \xi\|_{\infty} \\
&\leq C \|\xi\|_{W^{2,\infty}} \left(\sqrt{(N-\nu)\tau} + \tau \right) \\
&\leq C \|\xi\|_{W^{2,\infty}} \left(\sqrt{t-s} + \tau + \tau \right).
\end{aligned}$$

Since $H^3(\mathbb{R})$ is continuously embedded in $W^{2,\infty}(\mathbb{R})$, the claimed estimate for $f_{\tau}(t) - f_{\tau}(s)$ follows by a density argument. A similar computation relying on (2.39), (3.1), (3.3), and (3.7) gives the same estimate for $g_{\tau}(t) - g_{\tau}(s)$ and completes the proof of Lemma 3.3. \square

We are now in a position to establish the strong compactness of $(f_{\tau}, g_{\tau})_{\tau>0}$ in $L^2((0, T) \times \mathbb{R})$ for all $T > 0$ as announced in (1.11).

Lemma 3.4. *There are a sequence $(\tau_k)_{k \geq 1}$, $\tau_k \rightarrow 0$, and functions f and g in $C([0, \infty); H^{-3}(\mathbb{R}))$ such that, for all $t \geq 0$,*

$$(f_{\tau_k}(t), g_{\tau_k}(t)) \longrightarrow (f(t), g(t)) \quad \text{in } H^{-3}(\mathbb{R}; \mathbb{R}^2), \quad (3.12)$$

$$(f_{\tau_k}, g_{\tau_k}) \longrightarrow (f, g) \quad \text{in } L^2((0, t) \times \mathbb{R}; \mathbb{R}^2), \quad (3.13)$$

$$(f_{\tau_k}, g_{\tau_k}) \longrightarrow (f, g) \quad \text{a.e. in } (0, \infty) \times \mathbb{R}. \quad (3.14)$$

Proof. The proof relies on [3, Proposition 3.3.1] and [17, Lemma 9]. Indeed, it first follows from (3.1), (3.3), (3.4), and Lemma 3.2 that $(f_{\tau}(t))_{\tau \in (0,1)}$ lies in a compact subset of $H^{-3}(\mathbb{R})$. This fact, together with Lemma 3.3 and a refined version of the Arzelà-Ascoli theorem [3, Proposition 3.3.1] ensures that there are a sequence $(\tau_k)_{k \geq 1}$, $\tau_k \rightarrow 0$, and a function $f \in C([0, \infty); H^{-3}(\mathbb{R}))$ such that $(f_{\tau_k}(t))$ converges towards $f(t)$ in $H^{-3}(\mathbb{R}; \mathbb{R}^2)$ for each $t \geq 0$. Since the same argument applies for $(g_{\tau})_{\tau \in (0,1)}$, we have established (3.12). We then infer from (3.3), the embedding of $L^2(\mathbb{R})$ in $H^{-3}(\mathbb{R})$, the convergence (3.12), and the Lebesgue dominated convergence theorem that

$$(f_{\tau_k}, g_{\tau_k}) \longrightarrow (f, g) \quad \text{in } L^2(0, T; H^{-3}(\mathbb{R}; \mathbb{R}^2)) \quad \text{for all } T > 0. \quad (3.15)$$

Now, given $\delta \in (0, 1)$ and $T > 1$, the estimates (3.1), (3.3) (with $s = 0$), (3.4), and (3.5) in Lemma 3.1 ensure that

$$(f_{\tau_k}, g_{\tau_k})_{k \geq 1} \quad \text{is bounded in } L^2(\delta, T; H^1(\mathbb{R}) \cap L^1(\mathbb{R}, (1+x^2) \, dx)). \quad (3.16)$$

Since $H^1(\mathbb{R}) \cap L^1(\mathbb{R}, (1+x^2) \, dx)$ is compactly embedded in $L^2(\mathbb{R})$ by Lemma 3.2 and $L^2(\mathbb{R})$ is continuously embedded in $H^{-3}(\mathbb{R})$, we are in a position to apply [17, Lemma 9] and deduce from (3.15) and (3.16) that

$(f_{\tau_k}, g_{\tau_k})_{k \geq 1}$ converges towards (f, g) in $L^2((\delta, T) \times \mathbb{R}; \mathbb{R}^2)$. Owing to (3.3), this convergence may actually be improved to (3.13). The a.e. convergence (3.14) then follows from (3.13) after possibly extracting a further subsequence. \square

Finally, (3.5) implies that, after possibly extracting a further subsequence, we may assume that

$$(\partial_x f_{\tau_k}, \partial_x g_{\tau_k}) \rightharpoonup (\partial_x f, \partial_x g) \quad \text{in} \quad L^2((\delta, T) \times \mathbb{R}) \quad \text{for all} \quad 0 < \delta < T. \quad (3.17)$$

Now, combining (3.6), (3.7), (3.13), and (3.17), we obtain

$$\begin{cases} \sqrt{f_{\tau_k}} \partial_x [(1+R) f_{\tau_k} + R g_{\tau_k}] & \rightharpoonup \sqrt{f} \partial_x [(1+R) f + R g] \\ \sqrt{g_{\tau_k}} \partial_x (f_{\tau_k} + g_{\tau_k}) & \rightharpoonup \sqrt{g} \partial_x (f + g) \end{cases} \quad \text{in} \quad L^2((\delta, T) \times \mathbb{R}) \quad (3.18)$$

for $0 < \delta < T$, while (3.13) and (3.17) imply that, for $0 < \delta < T$,

$$\begin{cases} f_{\tau_k} \partial_x [(1+R) f_{\tau_k} + R g_{\tau_k}] & \rightharpoonup f \partial_x [(1+R) f + R g] \\ g_{\tau_k} \partial_x (f_{\tau_k} + g_{\tau_k}) & \rightharpoonup g \partial_x (f + g) \end{cases} \quad \text{in} \quad L^1((\delta, T) \times \mathbb{R}). \quad (3.19)$$

3.2. Passing to the limit. Combining the convergence (3.12) with the estimates (3.1), (3.3) (with $s = 0$) and (3.4) in Lemma 3.1 ensures that $(f(t), g(t)) \in \mathcal{K}_2$ for all $t \geq 0$. Moreover, gathering (3.3), (3.5), (3.13), and (3.17), we conclude that (f, g) satisfies the integrability properties (i) of Theorem 1.1. In addition, it follows from (3.12) and Lemma 3.3 that

$$\|f(t) - f(s)\|_{H^{-3}} + \|g(t) - g(s)\|_{H^{-3}} \leq C_2 \sqrt{|t - s|}, \quad (t, s) \in [0, \infty) \times [0, \infty), \quad (3.20)$$

which proves assertion (ii) of Theorem 1.1.

In order to establish the estimate (b) of Theorem 1.1, we pick $T > 0$ and set $N_k := [T/\tau_k]$ for all integers $k \geq 1$. Then, we infer from Corollary 2.3 and (3.9) (with $s = 0$) that for all $k \geq 1$ we have

$$\begin{aligned} & \frac{1}{2} \int_{\tau_k}^T \left\{ \left\| \sqrt{f_{\tau_k}(\sigma)} \partial_x [(1+R)f_{\tau_k} + Rg_{\tau_k}](\sigma) \right\|_2^2 + RR_\mu \left\| \sqrt{g_{\tau_k}(\sigma)} \partial_x [f_{\tau_k} + g_{\tau_k}](\sigma) \right\|_2^2 \right\} d\sigma \\ & \leq \sum_{n=1}^{N_k} \left[\frac{W_2^2(f_{\tau_k}^n, f_{\tau_k}^{n-1})}{2\tau_k} + \frac{R}{R_\mu} \frac{W_2^2(g_{\tau_k}^n, g_{\tau_k}^{n-1})}{2\tau_k} \right] \leq \mathcal{E}(f_0, g_0) - \mathcal{E}(f_{\tau_k}(T), g_{\tau_k}(T)). \end{aligned}$$

Letting $k \rightarrow \infty$, the convergences (3.13) and (3.18) lead us to

$$\begin{aligned} & \frac{1}{2} \int_\delta^T \left\{ \left\| \sqrt{f(\sigma)} \partial_x [(1+R)f + Rg](\sigma) \right\|_2^2 + RR_\mu \left\| \sqrt{g(\sigma)} \partial_x [f + g](\sigma) \right\|_2^2 \right\} d\sigma \\ & \leq \mathcal{E}(f_0, g_0) - \mathcal{E}(f(T), g(T)). \end{aligned}$$

for all $\delta \in (0, 1)$, whence the desired assertion (b) of Theorem 1.1 after letting $\delta \rightarrow 0$.

Now, we identify the equations solved by f and g . To this end, fix $\xi \in C_0^\infty(\mathbb{R})$, $t \in (0, \infty)$, $s \in (0, t)$ and set $N := [t/\tau]$ and $\nu := [s/\tau]$. We infer from (2.40), (3.1), (3.2), and (3.6) that

$$\begin{aligned}
& \left| \int_{\mathbb{R}} (f_\tau(t) - f_\tau(s)) \xi dx + \int_s^t \int_{\mathbb{R}} f_\tau(\sigma) \partial_x [(1+R) f_\tau + R g_\tau](\sigma) \partial_x \xi dx d\sigma \right| \\
& \leq \left| \int_{\mathbb{R}} (f_\tau(t) - f_0) \xi dx + \int_\tau^{(N+1)\tau} \int_{\mathbb{R}} f_\tau(\sigma) \partial_x [(1+R) f_\tau + R g_\tau](\sigma) \partial_x \xi dx d\sigma \right| \\
& \quad + \left| \int_{\mathbb{R}} (f_\tau(s) - f_0) \xi dx + \int_\tau^{(\nu+1)\tau} \int_{\mathbb{R}} f_\tau(\sigma) \partial_x [(1+R) f_\tau + R g_\tau](\sigma) \partial_x \xi dx d\sigma \right| \\
& \quad + \left| \int_t^{(N+1)\tau} \int_{\mathbb{R}} f_\tau(\sigma) \partial_x [(1+R) f_\tau + R g_\tau](\sigma) \partial_x \xi dx d\sigma \right| \\
& \quad + \left| \int_s^{(\nu+1)\tau} \int_{\mathbb{R}} f_\tau(\sigma) \partial_x [(1+R) f_\tau + R g_\tau](\sigma) \partial_x \xi dx d\sigma \right| \\
& \leq \|\partial_x^2 \xi\|_\infty \sum_{n=1}^N W_2^2(f_\tau^n, f_\tau^{n-1}) \\
& \quad + \|\partial_x \xi\|_\infty \int_s^{(\nu+1)\tau} \|f_\tau(\sigma)\|_1^{1/2} \left\| \sqrt{f_\tau} \partial_x [(1+R) f_\tau + R g_\tau](\sigma) \right\|_2 d\sigma \\
& \quad + \|\partial_x \xi\|_\infty \int_t^{(N+1)\tau} \|f_\tau(\sigma)\|_1^{1/2} \left\| \sqrt{f_\tau} \partial_x [(1+R) f_\tau + R g_\tau](\sigma) \right\|_2 d\sigma \\
& \leq C \|\xi\|_{W^{2,\infty}} (\tau + \sqrt{\tau}).
\end{aligned}$$

Taking $\tau = \tau_k$ in the above inequality and letting $k \rightarrow \infty$ with the help of (3.12) and (3.19), we end up with the first identity in (1.12)

$$\int_{\mathbb{R}} (f(t) - f(s)) \xi dx + \int_s^t \int_{\mathbb{R}} f(\sigma) \partial_x [(1+R) f + R g](\sigma) \partial_x \xi dx d\sigma = 0.$$

The proof of the second one being similar, it remains to check the property (a) stated in Theorem 1.1. To this end, we first claim that

$$(f_{\tau_k} \ln f_{\tau_k}, g_{\tau_k} \ln g_{\tau_k}) \longrightarrow (f \ln f, g \ln g) \quad \text{in } L^1((0, T) \times \mathbb{R}), \quad T > 0. \quad (3.21)$$

Indeed, by (3.13) and the continuity of $r \mapsto r \ln r$ in $[0, \infty)$, we have for $T > 0$

$$(f_{\tau_k} \ln f_{\tau_k}, g_{\tau_k} \ln g_{\tau_k}) \longrightarrow (f \ln f, g \ln g) \quad \text{a.e. in } (0, T) \times \mathbb{R}. \quad (3.22)$$

Moreover, it readily follows from (3.3) (with $s = 0$) that

$$(f_{\tau_k} \ln f_{\tau_k}, g_{\tau_k} \ln g_{\tau_k})_{k \geq 1} \quad \text{is uniformly integrable in } L^1((0, T) \times \mathbb{R}; \mathbb{R}^2), \quad (3.23)$$

while (3.3), (3.4), and the inequality $|r \ln r| \leq 2\sqrt{r} \max\{r, 1\}$, $r \geq 0$, guarantee that, for $R > 1$,

$$\begin{aligned}
\int_0^T \int_{\{|x| \geq R\}} |f_{\tau_k} \ln f_{\tau_k}| dx dt &\leq 2 \int_0^T \int_{\{|x| \geq R\}} \sqrt{f_{\tau_k}} \mathbf{1}_{[0,1]}(f_{\tau_k}) dx dt \\
&+ 2 \int_0^T \int_{\{|x| \geq R\}} f_{\tau_k}^{3/2} \mathbf{1}_{(1,R)}(f_{\tau_k}) dx dt \\
&+ 2 \int_0^T \int_{\{|x| \geq R\}} f_{\tau_k}^{3/2} \mathbf{1}_{[R,\infty)}(f_{\tau_k}) dx dt \\
&\leq 2 \left(\int_0^T \int_{\{|x| \geq R\}} x^2 f_{\tau_k} dx dt \right)^{1/2} \left(\int_0^T \int_{\{|x| \geq R\}} \frac{dx dt}{x^2} \right)^{1/2} \\
&+ 2\sqrt{R} \int_0^T \int_{\{|x| \geq R\}} f_{\tau_k} dx dt + \frac{2}{\sqrt{R}} \int_0^T \int_{\{|x| \geq R\}} f_{\tau_k}^2 dx dt \\
&\leq C \frac{1+T}{\sqrt{R}} + \frac{2}{R^{3/2}} \int_0^T \int_{\{|x| \geq R\}} x^2 f_{\tau_k} dx dt + C \frac{1+T}{\sqrt{R}} \\
&\leq C \frac{1+T}{\sqrt{R}}. \tag{3.24}
\end{aligned}$$

Due to (3.22)-(3.24), we are in a position to apply Vitali's convergence theorem (see, e.g., [11, Theorem 2.24] or [13, Théorème I.4.13]) and deduce the claim (3.21) for $(f_{\tau_k})_{k \geq 1}$, the proof for $(g_{\tau_k})_{k \geq 1}$ being identical. Consequently, after possibly extracting a subsequence, we have also

$$(H(f_{\tau_k}(t)), H(g_{\tau_k}(t))) \longrightarrow (H(f(t)), H(g(t))) \quad \text{a.e. in } (0, \infty), \tag{3.25}$$

the functional H being defined in (2.4). We next infer from (3.17) and the Fatou lemma that, for $t > 0$,

$$\begin{aligned}
\int_0^t (\|\partial_x f(s)\|_2^2 + R \|\partial_x(f+g)(s)\|_2^2) ds &= \lim_{\delta \rightarrow 0} \int_\delta^t (\|\partial_x f(s)\|_2^2 + R \|\partial_x(f+g)(s)\|_2^2) ds \\
&\leq \lim_{\delta \rightarrow 0} \liminf_{k \rightarrow \infty} \int_\delta^t (\|\partial_x f_{\tau_k}(s)\|_2^2 + R \|\partial_x(f_{\tau_k} + g_{\tau_k})(s)\|_2^2) ds. \tag{3.26}
\end{aligned}$$

Owing to (3.25) and (3.26), we may pass to the limit as $k \rightarrow \infty$ in (3.10) to obtain the assertion (a) of Theorem 1.1, which completes its proof.

ACKNOWLEDGEMENTS

PhL warmly thanks Adrien Blanchet and Giuseppe Savaré for helpful discussions on the optimal transport approach to partial differential equations. This work was initiated during a visit of BVM at the Institut de Mathématiques de Toulouse, Université Paul Sabatier - Toulouse 3. He is grateful for the hospitality.

APPENDIX A. SOME TECHNICAL RESULTS

We first collect some well-known properties of the functional H defined by (2.4).

Lemma A.1. *Let h be a nonnegative function in $L^1(\mathbb{R}, (1+x^2)dx) \cap L^2(\mathbb{R})$. Then $h \ln h \in L^1(\mathbb{R})$ and there is a positive constant C_ℓ such that*

$$\int_{\mathbb{R}} h(x) |\ln h(x)| dx \leq C_\ell + \int_{\mathbb{R}} h(x) (1+x^2) dx + \|h\|_2^2, \quad (\text{A.1})$$

$$H(h) \geq -C_\ell - \int_{\mathbb{R}} h(x) (1+x^2) dx. \quad (\text{A.2})$$

Proof. Introducing the function $\omega(x) := e^{-(1+x^2)}$, $x \in \mathbb{R}$, and using the monotonicity of $r \mapsto r |\ln r|$ in $[0, 1/e]$, we have

$$\begin{aligned} \int_{\mathbb{R}} h(x) |\ln h(x)| dx &= \int_{\{h(x) < \omega(x)\}} h(x) |\ln h(x)| dx + \int_{\{\omega(x) \leq h(x) \leq 1\}} h(x) |\ln h(x)| dx \\ &\quad + \int_{\{h(x) > 1\}} h(x) |\ln h(x)| dx \\ &\leq \int_{\{h(x) < \omega(x)\}} e^{-(1+|x|^2)} (1+x^2) dx + \int_{\{\omega(x) \leq h(x) \leq 1\}} h(x)(1+x^2) dx \\ &\quad + \int_{\{h(x) > 1\}} h(x)(h(x) - 1) dx \\ &\leq \int_{\mathbb{R}} e^{-(1+|x|^2)} (1+x^2) dx + \int_{\mathbb{R}} h(x)(1+x^2) dx + \|h\|_2^2, \end{aligned}$$

whence (A.1). Similarly,

$$\begin{aligned} H(h) &\geq \int_{\{h(x) < \omega(x)\}} h(x) \ln h(x) dx + \int_{\{\omega(x) \leq h(x) \leq 1\}} h(x) \ln h(x) dx \\ &\geq - \int_{\{h(x) < \omega(x)\}} e^{-(1+|x|^2)} (1+x^2) dx - \int_{\{\omega(x) \leq h(x) \leq 1\}} h(x)(1+x^2) dx, \end{aligned}$$

from which (A.2) readily follows. \square

The next results allowed us to identify the limit of some terms arising in the derivation of the Euler-Lagrange equation in Lemma 2.2.

Lemma A.2. *Consider $h \in H^1(\mathbb{R})$ and $\zeta \in C_0^\infty(\mathbb{R})$. Setting $\zeta_\varepsilon := \text{id} + \varepsilon \zeta$ for $\varepsilon > 0$, we have*

$$h \circ \zeta_\varepsilon \xrightarrow{\varepsilon \rightarrow 0} h \quad \text{in } L^2(\mathbb{R}) \quad \text{and} \quad \frac{h \circ \zeta_\varepsilon - h}{\varepsilon} \xrightarrow{\varepsilon \rightarrow 0} \zeta \partial_x h \quad \text{in } L^2(\mathbb{R}). \quad (\text{A.3})$$

Proof. Since ζ_ε is a C^∞ -diffeomorphism from \mathbb{R} onto \mathbb{R} for ε small enough, its inverse ζ_ε^{-1} is well-defined and satisfies

$$|x - \zeta_\varepsilon^{-1}(x)| \leq \varepsilon \|\zeta\|_\infty, \quad x \in \mathbb{R}. \quad (\text{A.4})$$

It follows from the Cauchy-Schwarz inequality, the Fubini theorem, and (A.4) that

$$\begin{aligned} \|h \circ \zeta_\varepsilon - h\|_2^2 &= \int_{\mathbb{R}} \left(\int_x^{\zeta_\varepsilon(x)} \partial_x h(y) dy \right)^2 dx \leq \int_{\mathbb{R}} |x - \zeta_\varepsilon(x)| \left| \int_x^{\zeta_\varepsilon(x)} |\partial_x h(y)|^2 dy \right| dx \\ &\leq \varepsilon \|\zeta\|_\infty \int_{\mathbb{R}} |\partial_x h(y)|^2 |y - \zeta_\varepsilon^{-1}(y)| dy \leq \varepsilon^2 \|\zeta\|_\infty^2 \|\partial_x h\|_2^2, \end{aligned}$$

which gives the first assertion in (A.3) and the boundedness of $((h \circ \zeta_\varepsilon - h)/\varepsilon)_\varepsilon$ in $L^2(\mathbb{R})$. Next, since $h \in H^1(\mathbb{R})$, almost every $x \in \mathbb{R}$ is a Lebesgue point for $\partial_x h$ and, for such points,

$$\frac{h(x + \varepsilon\zeta(x)) - h(x)}{\varepsilon} = \frac{1}{\varepsilon\zeta(x)} \int_x^{x+\varepsilon\zeta(x)} \partial_x h(y) dy \xrightarrow{\varepsilon \rightarrow 0} \partial_x h(x) \zeta(x).$$

Therefore, $((h \circ \zeta_\varepsilon - h)/\varepsilon)_\varepsilon$ converges a.e. to $\zeta \partial_x h$ as $\varepsilon \rightarrow 0$ and is bounded in $L^2(\mathbb{R})$, and the second assertion in (A.3) readily follows from these two facts. \square

The first assertion of Lemma A.2 is actually true in a more general setting:

Lemma A.3. *Consider $h \in H^1(\mathbb{R})$ and a sequence $(\zeta_\varepsilon)_{\varepsilon>0}$ of functions in $C_0^\infty(\mathbb{R})$ such that $\omega_\varepsilon := \|\zeta_\varepsilon - \text{id}\|_\infty \rightarrow 0$ as $\varepsilon \rightarrow 0$. Then*

$$h \circ \zeta_\varepsilon \xrightarrow{\varepsilon \rightarrow 0} h \quad \text{in } L^2(\mathbb{R}).$$

Proof. As in the proof of Lemma A.2, it follows from the Cauchy-Schwarz inequality and Fubini's theorem that

$$\begin{aligned} \|h \circ \zeta_\varepsilon - h\|_2^2 &\leq \int_{\mathbb{R}} |x - \zeta_\varepsilon(x)| \left| \int_x^{\zeta_\varepsilon(x)} |\partial_x h(y)|^2 dy \right| dx \leq \omega_\varepsilon \int_{\mathbb{R}} \int_{x-\omega_\varepsilon}^{x+\omega_\varepsilon} |\partial_x h(y)|^2 dy dx \\ &\leq 2 \omega_\varepsilon^2 \|\partial_x h\|_2^2, \end{aligned}$$

and the right-hand side of the above inequality converges to zero as $\varepsilon \rightarrow 0$. \square

REFERENCES

- [1] M. Agueh. Existence of solutions to degenerate parabolic equations via the Monge-Kantorovich theory. *Adv. Differential Equations*, 10(3):309–360, 2005.
- [2] H. Amann. Nonhomogeneous linear and quasilinear elliptic and parabolic boundary value problems. In *Function spaces, differential operators and nonlinear analysis (Friedrichroda, 1992)*, volume 133 of *Teubner-Texte Math.*, pages 9–126. Teubner, Stuttgart, 1993.
- [3] L. Ambrosio, N. Gigli, and G. Savaré. *Gradient flows in metric spaces and in the space of probability measures*. Lectures in Mathematics ETH Zürich. Birkhäuser Verlag, Basel, second edition, 2008.
- [4] L. Ambrosio and S. Serfaty. A gradient flow approach to an evolution problem arising in superconductivity. *Comm. Pure Appl. Math.*, 61(11):1495–1539, 2008.
- [5] A. Blanchet, V. Calvez, and J. A. Carrillo. Convergence of the mass-transport steepest descent scheme for the subcritical Patlak-Keller-Segel model. *SIAM J. Numer. Anal.*, 46(2):691–721, 2008.
- [6] E. A. Carlen and W. Gangbo. Solution of a model Boltzmann equation via steepest descent in the 2-Wasserstein metric. *Arch. Ration. Mech. Anal.*, 172(1):21–64, 2004.
- [7] J. A. Carrillo and S. Lisini. A Wasserstein- L^2 mixed gradient flow approach to the fully parabolic Keller-Segel model. In preparation.
- [8] J. A. Carrillo, R. J. McCann, and C. Villani. Contractions in the 2-Wasserstein length space and thermalization of granular media. *Arch. Ration. Mech. Anal.*, 179(2):217–263, 2006.
- [9] J. Escher, Ph. Laurençot, and B.-V. Matioc. Existence and stability of weak solutions for a degenerate parabolic system modelling two-phase flows in porous media. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 28(4):583–598, 2011.
- [10] J. Escher, A.-V. Matioc, and B.-V. Matioc. Modelling and analysis of the Muskat problem for thin fluid layers. *J. Math. Fluid Mech.*, to appear.
- [11] I. Fonseca and G. Leoni. *Modern methods in the calculus of variations: L^p spaces*. Springer Monographs in Mathematics. Springer, New York, 2007.
- [12] R. Jordan, D. Kinderlehrer, and F. Otto. The variational formulation of the Fokker-Planck equation. *SIAM J. Math. Anal.*, 29(1):1–17, 1998.
- [13] O. Kavian. *Introduction à la théorie des points critiques et applications aux problèmes elliptiques*, volume 13 of *Mathématiques & Applications (Berlin) [Mathematics & Applications]*. Springer-Verlag, Paris, 1993.

- [14] D. Matthes, R. J. McCann, and G. Savaré. A family of nonlinear fourth order equations of gradient flow type. *Comm. Partial Differential Equations*, 34(10-12):1352–1397, 2009.
- [15] F. Otto. Dynamics of labyrinthine pattern formation in magnetic fluids: a mean-field theory. *Arch. Rational Mech. Anal.*, 141(1):63–103, 1998.
- [16] F. Otto. The geometry of dissipative evolution equations: the porous medium equation. *Comm. Partial Differential Equations*, 26(1-2):101–174, 2001.
- [17] J. Simon. Compact sets in the space $L^p(0, T; B)$. *Ann. Mat. Pura Appl. (4)*, 146:65–96, 1987.
- [18] C. Villani. *Topics in optimal transportation*, volume 58 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2003.

INSTITUT DE MATHÉMATIQUES DE TOULOUSE, CNRS UMR 5219, UNIVERSITÉ DE TOULOUSE, F-31062 TOULOUSE
CEDEX 9, FRANCE

E-mail address: `laurenco@math.univ-toulouse.fr`

INSTITUT FÜR MATHEMATIK, UNIVERSITÄT WIEN, NORDBERGSTRASSE 15, 1090 WIEN, ÖSTERREICH

E-mail address: `bogdan-vasile.matioc@univie.ac.at`