



Smooth type II blow up solutions to the four dimensional energy critical wave equation

Matthieu Hillairet, Pierre Raphaël

► To cite this version:

Matthieu Hillairet, Pierre Raphaël. Smooth type II blow up solutions to the four dimensional energy critical wave equation. *Analysis & PDE*, Mathematical Sciences Publishers, 2012, 5 (4), pp.777-829. <hal-00524838>

HAL Id: hal-00524838

<https://hal.archives-ouvertes.fr/hal-00524838>

Submitted on 8 Oct 2010

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

SMOOTH TYPE II BLOW UP SOLUTIONS TO THE FOUR DIMENSIONAL ENERGY CRITICAL WAVE EQUATION

MATTHIEU HILLAIRET AND PIERRE RAPHAËL

ABSTRACT. We exhibit C^∞ type II blow up solutions to the focusing energy critical wave equation in dimension $N = 4$. These solutions admit near blow up time a decomposition

$$u(t, x) = \frac{1}{\lambda^{\frac{N-2}{2}}(t)} (Q + \varepsilon(t)) \left(\frac{x}{\lambda(t)} \right) \quad \text{with} \quad \|\varepsilon(t), \partial_t \varepsilon(t)\|_{\dot{H}^1 \times L^2} \ll 1$$

where Q is the extremizing profile of the Sobolev embedding $\dot{H}^1 \rightarrow L^{2^*}$, and a blow up speed

$$\lambda(t) = (T - t) e^{-\sqrt{|\log(T-t)|}(1+o(1))} \quad \text{as} \quad t \rightarrow T.$$

1. Introduction

1.1. Setting of the problem. We deal in this paper with the energy critical focusing wave equation

$$\begin{cases} \partial_{tt} u - \Delta u - f(u) = 0 & \text{with } f(t) = t^{\frac{N+2}{N-2}}, \\ (u, \partial_t u)|_{t=0} = (u_0, u_1), & (t, x) \in \mathbb{R} \times \mathbb{R}^N. \end{cases} \quad (1.1)$$

in dimension

$$N = 4.$$

This is a special case of the nonlinear wave equation

$$\partial_{tt} u - \Delta u - f(u) = 0 \quad (1.2)$$

which since the pioneering works by Jörrens [10] has attracted a considerable amount of works. For the energy critical nonlinearity $f(u) = \pm t^{\frac{N+2}{N-2}}$, the Cauchy problem is locally well posed in the energy space $\dot{H}^1 \times L^2$ and the solution propagates regularity, see for example Sogge [35] and references therein. Recall that in this case, (1.2) admits a conserved energy

$$E(u(t)) = E(u_0, u_1) = \frac{1}{2} \int (\partial_t u)^2 + \frac{1}{2} \int |\nabla u|^2 \mp \frac{N-2}{2N} \int u^{\frac{2N}{N-2}}$$

which is left invariant by the scaling symmetry of the flow:

$$u_\lambda(t, x) = \frac{1}{\lambda^{\frac{N-2}{2}}} u\left(\frac{t}{\lambda}, \frac{x}{\lambda^2}\right).$$

Global existence in the defocusing case was proved by Struwe [38] for radial data and Grillakis [9] for general data. For focusing nonlinearities, a sharp threshold criterion of global existence and scattering or finite time blow up is obtained by Kenig and Merle [14] based on the solitonic solution to (1.1):

$$Q(r) = \left(\frac{1}{1 + \frac{r^2}{N(N-2)}} \right)^{\frac{N-2}{2}} \quad (1.3)$$

which is the extremizing profile of the Sobolev embedding $\dot{H}^1 \rightarrow L^{2^*}$. Indeed, for initial data (u_0, u_1) such that $E(u_0, u_1) < E(Q, 0)$, those with $\|\nabla u_0\|_{L^2} < \|\nabla Q\|_{L^2}$ have global solutions and scatter,

while those with $\|\nabla u_0\|_{L^2} > \|\nabla Q\|_{L^2}$ lead to finite time blow up.

Note that like in the works by Levine [20], see also Strauss [37], and as is standard in a nonlinear dispersive setting, blow up is derived through obstructive convexity arguments, see also Karageorgis and Strauss [11] for refined statements near the soliton Q . However, this approach gives very little insight into the description of the blow up mechanism and the description of the flow even just near the ground state soliton Q is still only at its beginning.

1.2. On the energy critical wave map problem. There is an important litterature devoted to the construction of blow up solutions for nonlinear wave equations, see e.g. Alinhac [1], and Merle and Zaag [28], [29] for the study of the ODE type of blow up for subcritical nonlinearities. For energy critical problems like (1.1), recent important progress has been made through the study of the two dimensional energy critical corotational wave map to the 2-sphere:

$$\partial_{tt}u - \partial_{rr}u - \frac{\partial_r u}{r} - \frac{k^2 \sin 2u}{2r^2} = 0, \quad (1.4)$$

where $k \in \mathbb{N}^*$ is the homotopy number. The ground state is given there by

$$Q(r) = 2 \tan^{-1}(r^k).$$

After the pioneering works by Christodoulou, Tahvildar-Zadeh [4], Shatah and Tahvildar-Zadeh [36] and Struwe [39] and their detailed study of the concentration of energy scenario, the first explicit description of singularity formation for the $k = 1$ case is derived by Krieger, Schlag and Tataru [17] who construct finite energy finite time blow up solutions of the form

$$u(t, x) = (Q + \varepsilon)(t, \frac{x}{\lambda(t)}) \quad \text{with} \quad \|\varepsilon(t), \partial_t \varepsilon(t)\|_{\dot{H}^1 \times L^2} \ll 1 \quad (1.5)$$

with a blow up speed given by

$$\lambda(t) = (T - t)^\nu, \quad \forall \nu > \frac{3}{2},$$

see also [19]. The spectacular feature of this result is to exhibit arbitrarily slow blow up regimes further and further from self similarity which would correspond to the –forbidden, see [39]– self similar law

$$\lambda(t) \sim T - t. \quad (1.6)$$

Numerics suggest [3] that this blow up scenario is non generic and corresponds to finite codimensional manifolds. After the pioneering works [34] for large homotopy number $k \geq 4$, Raphaël and Rodnianski [31] give a complete description of a *stable* blow up dynamics which originates from smooth data and for all homotopy number $k \geq 1$. The blow up speed obeys in this regime a *universal law which depends in an essential way on the rate of convergence* of the ground state Q to its asymptotic value

$$\pi - Q \sim \frac{1}{r^k} \quad \text{as} \quad r \rightarrow \infty,$$

and indeed the stable blow up regime corresponds to a decomposition (1.5) with the blow up speed

$$\lambda(t) \sim \begin{cases} c_k \frac{T-t}{|\log(T-t)|^{\frac{1}{2k-2}}} & \text{for } k \geq 2, \\ (T-t)e^{-\sqrt{|\log(T-t)|}} & \text{for } k = 1. \end{cases} \quad (1.7)$$

Note that this work draws an important analogy with another critical problem, the L^2 critical nonlinear Schrödinger equation, where a similar universality of the stable singularity formation near the ground state is proved by Merle and Raphaël in the series of papers [23], [24], [30], [25], [26], [27].

1.3. Statement of the result. For the power nonlinearity energy critical problem (1.1), there has been recent progress towards the understanding of the flow near the solitary wave Q . In [15], Krieger and Schlag construct in dimension $N = 3$ a codimension one manifold of initial data near Q which yield global solutions asymptotically converging to the soliton manifold. The strategy developed by Krieger, Schlag, Tataru in [17] for the wave map problem has been adapted in [18] to show in dimension $N = 3$ the existence of finite energy finite time blow up solutions of the form

$$u(t, x) = \frac{1}{\lambda^{\frac{N-2}{2}}(t)}(Q + \varepsilon)(t, \frac{x}{\lambda(t)}) \quad \text{with} \quad \|\varepsilon(t), \partial_t \varepsilon(t)\|_{\dot{H}^1 \times L^2} \ll 1$$

and with a blow up speed given by

$$\lambda(t) = (T - t)^\nu, \quad \forall \nu > \frac{3}{2}. \quad (1.8)$$

The quantization of the energy at blow up for small type II blow up solutions in dimension $N \in \{3, 5\}$ is proved in [6], [7] in the radial and non radial cases. In particular, for radial data, if $T < +\infty$ and

$$\sup_{t \in [0, T]} [|\nabla u(t)|_{L^2}^2 + \partial_t u|_{L^2}^2] \leq |\nabla Q|_{L^2}^2 + \alpha^*, \quad \alpha^* \ll 1,$$

then there exists a dilation parameter $\lambda(t) \rightarrow 0$ as $t \rightarrow T$ and asymptotic profiles $(u^*, v^*) \in H^1 \times L^2$ such that

$$\left(u(t, x) - \frac{1}{\lambda^{\frac{N-2}{2}}(t)} Q\left(\frac{x}{\lambda(t)}\right), \partial_t u(t) \right) \rightarrow (u^*, v^*) \quad \text{in} \quad \dot{H}^1 \times L^2 \quad \text{as} \quad t \rightarrow T,$$

see [27] for related classification results for the L^2 critical (NLS).

These works however leave open the question of the existence of *smooth* type II blow up solutions. We claim that such smooth type II blow up solutions can be constructed in dimension $N = 4$ as the formal analogue of the singular dynamics exhibited by Raphaël and Rodnianski [31] for the wave map problem in the least homotopy number class $k = 1$. The following theorem is the main result of this paper:

Theorem 1.1 (Existence of smooth type II blow up solutions in dimension $N = 4$). *Let $N = 4$. Then for all $\alpha^* > 0$, there exist C^∞ initial data (u_0, u_1) with*

$$E(u_0, u_1) < E(Q, 0) + \alpha^*$$

such that the corresponding solution to the energy critical focusing wave equation (1.1) blows up in finite time $T = T(u_0, u_1) < +\infty$ in a type II regime according to the following dynamics: there exist $(u^, v^*) \in \dot{H}^1 \times L^2$ such that*

$$\left(u(t, x) - \frac{1}{\lambda^{\frac{N-2}{2}}(t)} Q\left(\frac{x}{\lambda(t)}\right), \partial_t u(t) \right) \rightarrow (u^*, v^*) \quad \text{in} \quad \dot{H}^1 \times L^2 \quad \text{as} \quad t \rightarrow T \quad (1.9)$$

with a blow up speed given by

$$\lambda(t) = (T - t)e^{-\sqrt{|\log(T-t)|}(1+o(1))} \quad \text{as} \quad t \rightarrow T. \quad (1.10)$$

Comments on the result

1. On the smoothness of the initial data: An important feature of Theorem 1.1 is to exhibit a new blow up speed which is valid for C^∞ solutions. Indeed, while the Krieger, Schlag, Tataru [18] approach provides a continuum of blow up speeds, the exact regularity of the obtained solutions is not known, which is an unpleasant consequence of their construction scheme. In fact, it is expected that C^∞ initial data should lead to *quantize* blow up rates hence breaking the continuum of blow up speeds (1.8), we refer to [2] for a related discussion in the context of the energy critical harmonic

heat flow. Hence we expect the blow up rate (1.10) to correspond to the minimal type II blow up speed of smooth solutions with small super critical energy. Such a general lower bound on blow up rate in the spirit of the one obtained by Merle and Raphael for the L^2 critical NLS [30], [26] is an open problem. The construction of excited blow up solutions with other speeds and C^∞ regularity also remains to be done. This problematic is related to the understanding of the structure of the flow near Q which is still at its beginning.

2. On the codimension one manifold: The proof of Theorem 1.1 involves a detailed description of the set of initial data leading to the type II blow up with speed (1.10). Indeed, given a small enough parameter $b_0 > 0$ and a suitable deformation Q_{b_0} of the soliton with

$$Q_{b_0} \rightarrow Q \text{ as } b_0 \rightarrow 0$$

in some strong sense, we show that for any smooth and radially symmetric excess of energy

$$\|\eta_0, \eta_1\|_{H^2 \times H^1} \lesssim \frac{b_0^2}{|\log(b_0)|},$$

we can find $d_+(b_0, \eta_0, \eta_1) \in \mathbb{R}$ such that the solution to (1.1) with initial data

$$u_0 = Q_{b_0} + \eta_0 + d_+ \psi, \quad u_1 = b_0 \left(\frac{N-2}{2} Q_{b_0} + y \cdot \nabla Q_{b_0} \right) + \eta_1$$

blows up in finite time in the regime described by Theorem 1.1. Here ψ is the bound state of the linearized operator close to Q and generates the unstable mode, we refer to Definition 3.4 and Proposition 3.5 for precise statements. Hence the set of blow up solutions we construct live on a codimension one manifold in the radial class in some weak sense. Following [15], [16], the proof that this set is indeed a codimension one manifold relies on proving some Lipschitz regularity of the map $(b_0, \eta_0, \eta_1) \rightarrow d_+(b_0, \eta_0, \eta_1)$, and in particular some local uniqueness to begin with. The analysis in [16] shows that this may be a delicate step in some cases. Our solution is constructed using a soft continuous topological argument of Brouwer type coupled with suitable monotonicity properties in the spirit of Cote, Martel and Merle [5], and in other related settings, see e.g. Martel [21], Raphaël and Szeftel [32], this strategy has proved to be quite powerful to eventually achieve strong uniqueness results. This interesting question in our setting will require additional efforts and needs to be addressed separately in details.

3. Extension to higher dimensions: We focus onto the case of dimension $N = 4$ for the sake of simplicity, and our main objective is to provide a robust framework to construct C^∞ type II blow up solutions. However, following the heuristic developed in [31], the blow up speed (1.10) corresponds to the $k = 1$ case in (1.7), and we similarly conjecture in dimension $N \geq 5$ the existence type II finite type blow up solutions close to Q with blow up speed

$$\lambda(t) \sim c_N \frac{T-t}{|\log(T-t)|^{\frac{1}{N-4}}}.$$

Note from (1.3) that the higher the dimension, the fastest the decay of the ground state Q , and this should avoid some difficulties which occur only in low dimension like in [31] for large homotopy number $k \geq 4$. We expect the strategy developed in this paper to carry over to the case $N = 5, 6$, but the extension to large dimension will be confronted in particular to the difficulty of the lack of smoothness of the nonlinearity. Let us also insist onto the fact that the case $N = 4$ is in many ways the more delicate one in terms of the strong coupling of the main part of the solution and the outgoing tail due to the slow decay of Q , which results in the somewhat pathological blow up speed (1.10). This comment becomes even more dramatic in dimension $N = 3$ where we expect

our analysis to be applicable to construct C^∞ type II blow up solutions, but this seems to require a slightly different approach.

1.4. Strategy of the proof. Let us briefly summarize the strategy of the proof of Theorem 1.1.

step 1 Approximate self similar solution.

Let D, Λ denote the differential operators (1.18). Exact self similar solutions to (1.1) of the form

$$u(t, x) = \frac{1}{\lambda^{\frac{N-2}{2}}(t)} Q_b \left(\frac{x}{\lambda(t)} \right) \quad \text{with } b = -\lambda_t$$

where Q_b satisfies the self similar equation

$$\Delta Q_b - b^2 D \Lambda Q_b + Q_b^3 = 0 \tag{1.11}$$

are known to develop a singularity on the light cone $y = \frac{T-t}{\lambda(t)} = \frac{1}{b}$ leading to an unbounded Dirichlet energy $\|\nabla Q_b\|_{L^2} = +\infty$, see Kavian, Weissler [12]. We therefore assume $0 < b \ll 1$ and consider a one term expansion approximation

$$Q_b = Q + b^2 T_1$$

which injected into (1.11) yields at the order b^2 :

$$H T_1 = -D \Lambda Q. \tag{1.12}$$

Here H is the linearized operator close to Q given by

$$H = -\Delta - \frac{N+2}{N-2} Q^{\frac{4}{N-2}}. \tag{1.13}$$

The spectral structure of H is well known in connection to the fact that Q is an extremizer of the Sobolev embedding $\dot{H}^1 \rightarrow L^{2^*}$, and in the radial sector, H admits one non positive eigenvalue with well localized eigenvector ψ :

$$H \psi = -\zeta \psi, \quad \zeta > 0, \tag{1.14}$$

and a *resonance* at the boundary of the continuum spectrum generated by the scaling invariance of (1.1):

$$H(\Lambda Q) = 0, \quad \Lambda Q(r) \sim \frac{C}{r^{N-2}} \quad \text{as } r \rightarrow +\infty. \tag{1.15}$$

In order to solve (1.12), we first remove the leading order growth in the exact solution $T_1 = \frac{1}{4}|y|^2 Q$ which is consequence of the flux computation:

$$(D \Lambda Q, \Lambda Q) = \frac{1}{2} \lim_{y \rightarrow +\infty} y^4 |\Lambda Q|^2 > 0 \tag{1.16}$$

due to the slow decay of Q in dimension $N = 4$ from (1.3). For this, we solve

$$H T_1 = -D \Lambda Q + c_b \Lambda Q \mathbf{1}_{y \leq \frac{1}{b}} \quad \text{with } c_b = \frac{(D \Lambda Q, \Lambda Q)}{\int_{y \leq \frac{1}{b}} |\Lambda Q|^2} \sim \frac{1}{2|\log b|} \quad \text{as } b \rightarrow 0.$$

The purpose of this construction is to yield after a suitable localization process an $o(b^2)$ approximate solution to the self similar equation (1.11) which dominant term near and past the light cone is still given by Q itself in the sense that:

$$b^2 |T_1| \ll Q \quad \text{for } y \geq \frac{1}{b}.$$

This identifies Q as the leading order radiation term¹.

step 2 Bootstrap estimates.

We now roughly consider initial data of the form

$$u_0 = Q_{b_0} + d_+ \psi + \eta_0, \quad u_1 = b_0 \Lambda Q_{b_0} + \eta_1, \quad \text{with } |d_+| + \|\eta_0, \eta_1\|_{H^2 \times H^1} \ll b_0^2, \quad (1.17)$$

and introduce a modulated decomposition of the flow

$$u(t, x) = \frac{1}{\lambda^{\frac{N-2}{2}}(t)} (Q_{b(t)} + \varepsilon) \left(t, \frac{x}{\lambda(t)} \right), \quad b(t) = -\lambda_t.$$

Here we face the major difference between the power nonlinearity wave equation (1.1) and the critical wave map problem (1.4) which is the presence of a negative eigenvalue in the first case (1.14) for the linearized operator H close to Q . This induces an instability in the modulation equations for b, λ which is absent in the wave map case, leading to stable blow up dynamics. However, we claim that the ODE type instability generated by (1.14) is the *only* instability mechanism.

The situation is conceptually similar to the one studied in [5] where multisolitary wave solutions are constructed in the supercritical regime despite the presence of exponentially growing modes for the linearized operator which are absent in the subcritical regime. We adapt a similar scheme of proof which does not rely on a fixed point argument to solve the problem from infinity in time², but by directly following the flow for *any* initial data of the form (1.17). This reduces the full problem to a one dimensional dynamical system for which a classical clever continuity argument yields the existence of $d_+(b_0, \eta_0, \eta_1)$ such that the unstable mode is extinct, see section 5.

The key is hence to control the flow under the a priori control of the unstable mode, and here we adapt the technology developed in [31] which relies on *monotonicity properties* of the linearized Hamiltonian at the H^2 level of regularity. However, the analysis in [31] heavily relies on the existence of a decomposition of the Hamiltonian

$$H = A^* A, \quad A = -\partial_y + V(y)$$

which is central in the proof of the main monotonicity property and is lost in our setting. This forces us to revisit the approach in several ways, and to rely in particular on fine algebraic properties of the flow³ near Q and coercitivity properties of suitable quadratic forms in the spirit of [22], [23], see Lemma 4.7, which remarkably turn out to be almost explicit thanks to the formula (1.3). We are eventually able to find $d_+(b_0, \eta_0, \eta_1)$ for which to leading order

$$b_s \sim -c_b b^2 \sim -\frac{b^2}{2|\log b|}, \quad b = -\lambda_t, \quad \frac{ds}{dt} = \frac{1}{\lambda}, \quad |d_+| + \|\partial_{yy} \varepsilon\|_{L^2} \ll b^2$$

which reintegration in time yields finite time blow up in the regime described by Theorem 1.1.

1.5. Notations. We define the differential operators:

$$\Lambda f = \frac{N-2}{2} f + y \cdot \nabla f \quad (\dot{H}^1 \text{ scaling}), \quad Df = \frac{N}{2} f + y \cdot \nabla f \quad (L^2 \text{ scaling}). \quad (1.18)$$

Denoting

$$(f, g) = \int f g = \int_0^{+\infty} f(r) g(r) r^{N-1} dr$$

¹see [31] for a further discussion on this issue and the role played by the non vanishing Pohozaev integration (1.16)

²after renormalization of the time

³see in particular (4.23), (4.38)

the $L^2(\mathbb{R}^N)$ radial inner product, we observe the integration by parts formula:

$$(Df, g) = -(f, Dg), \quad (\Lambda f, g) + (\Lambda g, f) = -2(f, g). \quad (1.19)$$

Given f and $\lambda > 0$, we shall denote:

$$f_\lambda(t, r) = \frac{1}{\lambda^{\frac{N-2}{2}}} f\left(t, \frac{r}{\lambda}\right),$$

and the space rescaled variable will always be denoted by

$$y = \frac{r}{\lambda}.$$

We let χ be a smooth positive radial cut off function $\chi(r) = 1$ for $r \leq 1$ and $\chi(r) = 0$ for $r \geq 2$. For a given parameter $B > 0$, we let

$$\chi_B(r) = \chi\left(\frac{r}{B}\right). \quad (1.20)$$

Given $b > 0$, we set

$$B_0 = \frac{2}{b}, \quad B_1 = \frac{|\log b|}{b}. \quad (1.21)$$

To clarify the exposition we use the notation $a \lesssim b$ when there exists a constant C with no relevant dependency on (a, b) such that $a \leq Cb$. In particular, we do not allow constants C to depend on the parameter M except in Appendix A.

Acknowledgments: The authors would like to thank Igor Rodnianski for stimulating discussions about this work. P.R is supported by the French ANR Jeune Chercheur SWAP.

2. Computation of the modified self-similar profile

This section is devoted to the construction of an approximate self-similar solution Q_b which describes the dominant part of the blow up profile inside the backward light cone from the singular point $(0, T)$ and displays a slow decay at infinity which is eventually responsible for the modifications to the blow up speed with respect to the self similar law. The key to this construction is the fact that the structure of the linearized operator H close to Q is completely explicit in the radial sector thanks to the explicit formulas at hand for the elements of the kernel.

We introduce the direction

$$\Phi = D\Lambda Q \quad (2.1)$$

which displays the cancellation

$$|\Phi(y)| \lesssim \frac{1}{1+y^4} \quad (2.2)$$

and the crucial nondegeneracy which follows from the Pohozaev integration by parts formula:

$$(\Phi, \Lambda Q) = \lim_{y \rightarrow +\infty} \left(\frac{1}{2} y^4 |\Lambda Q|^2 \right) = 32 > 0. \quad (2.3)$$

Proposition 2.1 (Approximate self-similar solution). *Let M denote a large enough constant. Then there exists $b^*(M) > 0$ small enough such that for all $0 < b < b^*(M)$, there exists a smooth radially symmetric profile T_1 satisfying the orthogonality condition*

$$(T_1, \chi_M \Phi) = 0 \quad (2.4)$$

such that

$$P_{B_1} = Q + \chi_{B_1} b^2 T_1 \quad (2.5)$$

is an approximate self similar solution in the following sense. Let

$$\Psi_{B_1} = -\Delta P_{B_1} + b^2 D\Lambda P_{B_1} - f(P_{B_1}), \quad (2.6)$$

then for all $k \geq 0, 0 \leq y \leq \frac{1}{b^2}$,

$$\left| \frac{d^k T_1}{dy^k}(y) \right| \lesssim \frac{1}{1+y^k} \left[\frac{1 + |\log(by)|}{|\log b|} \mathbf{1}_{2 \leq y \leq \frac{B_0}{2}} + \frac{1}{b^2 y^2 |\log b|} \mathbf{1}_{y \geq \frac{B_0}{2}} + \frac{\log(M) + |\log(1+y)|}{1+y^2} \right], \quad (2.7)$$

$$\left| \frac{d^k}{dy^k} \frac{\partial P_{B_1}}{\partial b} \right| \lesssim \frac{b \mathbf{1}_{y \leq 2B_1}}{1+y^k} \left[\frac{1 + |\log(by)|}{|\log b|} \mathbf{1}_{2 \leq y \leq \frac{B_0}{2}} + \frac{1}{b^2 y^2 |\log b|} \mathbf{1}_{y \geq \frac{B_0}{2}} + \frac{\log(M) + |\log(1+y)|}{1+y^2} \right], \quad (2.8)$$

and, for all $k \geq 0, y \geq 0$,

$$\begin{aligned} & \left| \frac{d^k}{dy^k} (\Psi_{B_1} - c_b b^2 \chi_{\frac{B_0}{4}} \Lambda Q) \right| \\ & \lesssim \frac{b^4}{1+y^k} \left[\frac{1 + |\log(by)|}{|\log b|} \mathbf{1}_{2 \leq y \leq \frac{B_0}{2}} + \frac{1}{b^2 y^2 |\log b|} \mathbf{1}_{2B_1 \geq y \geq \frac{B_0}{2}} + \frac{\log(M) + |\log(1+y)|}{1+y^2} \mathbf{1}_{y \leq 2B_1} \right] \\ & + \frac{b^2}{(1+y^{4+k})} \mathbf{1}_{y \geq B_1/2}, \end{aligned} \quad (2.9)$$

for some constant

$$c_b = \frac{1}{2|\log b|} \left(1 + O\left(\frac{1}{|\log b|} \right) \right). \quad (2.10)$$

Proof of Proposition 2.1

Step 1 Inversion of H .

The first green function of H is given from scaling invariance by

$$\Lambda Q(y) = \frac{N-2}{2 \left(1 + \frac{y^2}{N(N-2)} \right)^{\frac{N}{2}}} \left(1 - \frac{y^2}{N(N-2)} \right), \quad (2.11)$$

which admits the following asymptotics:

$$\forall k \geq 0, \quad \frac{d^k (\Lambda Q)}{dy^k}(y) = \begin{cases} O(1) & \text{as } y \rightarrow 0, \\ O(y^{-(N-2+k)}) & \text{as } y \rightarrow \infty \end{cases} \quad (2.12)$$

Let now

$$\Gamma(y) = -\Lambda Q(y) \int_1^y \frac{ds}{s^{N-1} (\Lambda Q)^2(s)},$$

be another (singular at the origin⁴) element of the kernel of H which can be found from the Wronskian relation:

$$\Gamma' \Lambda Q - \Gamma (\Lambda Q)' = \frac{-1}{y^{N-1}}.$$

From this we easily find the asymptotics of $\Gamma^{(k)}$ for any integer k :

$$\frac{d^k \Gamma}{dy^k}(y) = \begin{cases} O(y^{-(N-2+k)}) & \text{as } y \rightarrow 0 \\ O(y^{-k}) & \text{as } y \rightarrow \infty. \end{cases} \quad (2.13)$$

A smooth solution to $Hw = F$ is given by:

$$w(y) = \Gamma(y) \int_0^y F(s) \Lambda Q(s) s^{N-1} ds - \Lambda Q(y) \int_0^y F(s) \Gamma(s) s^{N-1} ds. \quad (2.14)$$

⁴Note that Γ must be smooth at $y = \sqrt{N(N-2)}$ where ΛQ vanishes from the radial ODE $H\Gamma = 0$

We now look for a solution to the self similar equation in the form $Q + b^2 T_1$. This yields:

$$\begin{aligned}\Psi_b &= -\Delta Q_b + b^2 D\Lambda Q_b - f(Q_b) \\ &= b^2 (HT_1 + D\Lambda Q) + b^4 D\Lambda T_1 - [f(Q + b^2 T_1) - f(Q) - b^2 f'(Q)T_1].\end{aligned}\quad (2.15)$$

Step 2 Computation of T_1 .

Thanks to the anomalous decay (2.2), we chose T_1 solution to

$$\begin{cases} HT_1 &= F = -D\Lambda Q + c_b \chi_{\frac{B_0}{4}} \Lambda Q, \\ (T_1, \chi_M \Phi) &= 0, \end{cases}\quad (2.16)$$

with c_b chosen such that:

$$(F, \Lambda Q) = 0 \quad (2.17)$$

i.e. from Pohozaev integration by parts formula, see (1.21) and (2.3) ,

$$\begin{aligned}c_b &= \frac{(D\Lambda Q, \Lambda Q)}{(\chi_{\frac{B_0}{4}} \Lambda Q, \Lambda Q)} = \frac{1 \lim_{y \rightarrow +\infty} y^4 |\Lambda Q(y)|^2}{2 \int \chi_{\frac{B_0}{4}} |\Lambda Q|^2} \\ &= \frac{1}{2|\log b|} \left(1 + O\left(\frac{1}{|\log b|}\right) \right) \quad \text{as } b \rightarrow 0.\end{aligned}$$

This yields (2.10). Following (2.14), we first consider

$$\tilde{T}_1(y) = \Gamma(y) \int_0^y F(s) \Lambda Q(s) s^3 ds - \Lambda Q(y) \int_0^y F(s) \Gamma(s) s^3 ds \quad (2.18)$$

The smoothness of \tilde{T}_1 at the origin follows from (2.18) together with elliptic regularity from (2.16).

We now examine the behavior of \tilde{T}_1 at large y .

We first observe that, from the orthogonality (2.17):

$$\tilde{T}_1(y) = - \left[\Gamma(y) \int_y^{+\infty} F(s) \Lambda Q(s) s^3 ds + \Lambda Q(y) \int_0^y F(s) \Gamma(s) s^3 ds \right]$$

Hence, from the degeneracy $|D\Lambda Q| = O(y^{-4})$, this yields that, for $\frac{B_0}{2} \leq y \leq \frac{1}{b^2}$:

$$\begin{aligned}|\tilde{T}_1(y)| &\lesssim \int_y^{+\infty} \frac{s^3}{(1+s^4)(1+s^2)} ds + \frac{1}{y^2} \left[\int_0^y \frac{1+s^3}{1+s^4} ds + |c_b| \int_0^{B_0} \frac{s^3}{1+s^2} ds \right] \\ &\lesssim \frac{|\log(1+y)|}{1+y^2} + \frac{1}{b^2 y^2 |\log b|}.\end{aligned}\quad (2.19)$$

similarly, for $1 \leq y \leq \frac{B_0}{2}$,

$$\begin{aligned}|\tilde{T}_1(y)| &= \left| \Gamma(y) \int_y^{+\infty} F(s) \Lambda Q(s) s^3 ds + \Lambda Q(y) \int_0^y F(s) \Gamma(s) s^3 ds \right| \\ &\lesssim \int_y^{+\infty} \frac{s^3}{(1+s^4)(1+s^2)} ds + |c_b| \int_y^{B_0} \frac{s^3}{(1+s^2)^2} ds \\ &\quad + \frac{1}{1+y^2} \left[\int_0^y \frac{s^3}{1+s^4} ds + |c_b| \int_0^y \frac{s^3}{1+s^2} ds \right] \\ &\lesssim \frac{1 + |\log(by)|}{|\log b|} + \frac{|\log(1+y)|}{1+y^2}.\end{aligned}\quad (2.20)$$

We now choose thanks to (2.3):

$$T_1(y) = \tilde{T}_1(y) - c\Lambda Q \quad \text{with} \quad c = \frac{(\tilde{T}_1, \chi_M \Phi)}{(\chi_M \Phi, \Lambda Q)}$$

so that the orthogonality condition (2.4) is fulfilled. We note that so that the bounds (2.19) and (2.20) ensure that c remains bounded by $\log(M)$ uniformly in M and b , provided b is chosen sufficiently small w.r.t. M .

This yields (2.7) for $k = 0$, the other cases follow similarly.

Step 4 Estimate on Ψ_{B_1} and $\partial_b \Psi_{B_1}$

We now cut off the slow decaying tail T_1 according to (2.5) and estimate the corresponding error to self similarity Ψ_{B_1} given by (2.6).

We compute:

$$\begin{aligned} \Psi_{B_1} &= b^2 \chi_{B_1} (HT_1 + D\Lambda Q) \\ &\quad + b^2 [-2\chi'_{B_1} T_1' - T_1 \Delta \chi_{B_1} + (1 - \chi_{B_1}) D\Lambda Q + b^2 D\Lambda(\chi_{B_1} T_1)] \\ &\quad - [f(Q + b^2 \chi_{B_1} T_1) - f(Q) - \chi_{B_1} f'(Q) T_1]. \end{aligned}$$

Outside the support of χ_{B_1} we have thus $\Psi_{B_1} = b^2 D\Lambda Q$. On the other hand, in dimension $N = 4$, we have the Taylor expansion :

$$f(Q + b^2 \chi_{B_1} T_1) - f(Q) - \chi_{B_1} f'(Q) T_1 = b^4 \chi_{B_1}^2 T_1^2(y) \int_0^1 (1 - \tau) (Q(y) + \tau b^2 \chi_{B_1} T_1(y)) d\tau.$$

We thus estimate from (2.7), (2.15), (2.16) and the degeneracy (2.2) for $y \leq 2B_1$:

$$\begin{aligned} \left| \Psi_{B_1} - b^2 c_b \chi_{\frac{B_0}{4}} \Lambda Q \right| &\lesssim b^2 \mathbf{1}_{y \geq B_1/2} \left(\frac{T_1'}{1+y} + \frac{T_1}{1+y^2} + \frac{1}{1+y^4} \right) \\ &\quad + b^4 |D\Lambda(\chi_{B_1} T_1)| + b^4 |T_1^2(y)| \int_0^1 (1 - \tau) |Q(y) + \tau b^2 T_1(y)| d\tau. \end{aligned}$$

(2.7) now yields (2.9) for $k = 0$. Further derivatives are estimated similarly thanks to the smoothness of the nonlinearity. We emphasize here that, given $B > 0$ large, we have $1/(1+y) \lesssim 1/B \lesssim 1/(1+y)$ on the support of χ'_B , so that differentiating χ_B acts as a multiplication by $1/(1+y)$. Furthermore, there holds $1/B_1 = o(b)$ so that we can always dominate $1/(1+y)$ by b on the support of χ'_{B_1} .

Finally, we compute $\partial_b P_{B_1}$ from (2.5).

To this end, we note that $\partial_b c_b = O(1/b |\log(b)|^2)$ when $b \rightarrow 0$ so that the source term for T_1 in (2.16) satisfies

$$\partial_b F = \left[O\left(\frac{1}{b|\log b|}\right) \chi_{B_0/4} + O\left(\frac{1}{b|\log b|}\right) \rho_{B_0/4} \right] \Lambda Q$$

where $\rho(z) = z\chi'(z) \in \mathcal{C}_c^\infty(0, \infty)$ and we keep the convention for function dilation. Hence, the same arguments as for T_1 enable to show that $\partial_b \tilde{T}_1$ and then $\partial_b T_1$ satisfy the estimates:

$$\left| \frac{d^k \partial_b T_1}{dy^k}(y) \right| \lesssim \frac{1}{b(1+y^k)} \left[\frac{1 + |\log(by)|}{|\log b|} \mathbf{1}_{2 \leq y \leq \frac{B_0}{2}} + \frac{1}{b^2 y^2 |\log b|} \mathbf{1}_{y \geq \frac{B_0}{2}} + \frac{1 + |\log(1+y)|}{1+y^2} \right]. \quad (2.21)$$

Finally, we compute from (2.5)

$$\partial_b P_{B_1} = 2b \chi_{B_1} T_1 + b^2 \partial_b \log(B_1) \rho_{B_1} T_1 + b^2 \chi_{B_1} \partial_b T_1. \quad (2.22)$$

This decomposition together with (2.7) and the previous computation yield (2.8).

This concludes the proof of Proposition 2.1.

3. Description of the trapped regime

We display in this section the regime which leads to the blow up dynamics described by Theorem 1.1.

3.1. Modulation of solutions to (1.1). Let us start with describing the set of solutions among which the finite time blow up scenario described by Theorem 1.1 is likely to arise. We recall from (1.14) that ψ denotes the bound state of H with eigenvalue $-\zeta < 0$. The following lemma is a standard consequence of the implicit function theorem and the smoothness of the flow, see Appendix A.

Lemma 3.1 (Modulation theory). *Let M be a sufficiently large constant to be chosen later and $0 < b_0 < b_0^*(M)$ small enough. Let (η_0, η_1, d_+) satisfying the smallness condition:*

$$|d_+| + \|\eta_0, \nabla \eta_0, \eta_1 + b_0(1 - \chi_{B_1(b_0)})\Lambda Q, \nabla \eta_1\|_{\dot{H}^1 \times \dot{H}^1 \times L^2 \times L^2} \lesssim \frac{b_0^2}{|\log b_0|}, \quad (3.1)$$

then, there exists a time T_0 such that the unique solution $u \in \mathcal{C}^2([0, T_0]; L^2(\mathbb{R}^N)) \cap \mathcal{C}([0, T_0]; H^2(\mathbb{R}^N))$ to (1.1) with initial data :

$$u_0 = P_{B_1(b_0)} + \eta_0 + d_+ \psi, \quad u_1 = b_0 \Lambda P_{B_1(b_0)} + \eta_1, \quad (3.2)$$

admits on $[0, T_0]$ a unique decomposition

$$u(t) = (P_{B_1(b(t))} + \varepsilon(t))_{\lambda(t)} \quad (3.3)$$

with

1. $\lambda \in \mathcal{C}^2([0, T_0], \mathbb{R}_+^*)$ with

$$\forall t \in [0, T_0], \quad (\varepsilon(t), \chi_M \Phi) = 0 \quad \text{and} \quad b(t) = -\lambda_t; \quad (3.4)$$

2. there holds the smallness:

$$\|\nabla \varepsilon(t)\|_{L^2} \lesssim b_0 |\log b_0| \quad |b(t) - b_0| + |\lambda(t) - 1| + \|\nabla^2 \varepsilon(t)\|_{L^2} \lesssim \frac{b_0^2}{|\log b_0|} \quad \forall t \in [0, T_0]. \quad (3.5)$$

Remark 3.2. Recall that the slow decay of Q and the choice of P_{B_1} induces an unbounded tail of P_{B_1} in the energy norm, and more specifically $\|\Lambda Q\|_{L^2} = +\infty$, hence the need for the compensation in the norm for the time derivative in (3.1).

3.2. Decomposition of the flow and modulation equations. Considering initial data satisfying the assumption of the above lemma, we now write the evolution equation induced by (1.1) in terms of the decomposition (3.3). Let

$$u(t, r) = \frac{1}{[\lambda(t)]^{\frac{N}{2}-1}} (P_{B_1(b(t))} + \varepsilon) \left(t, \frac{r}{\lambda(t)}\right) = (P_{B_1(b(t))})_{\lambda(t)} + w(t, r) \quad (3.6)$$

where $b = -\lambda_t$. Let us derive the equations for w and ε . Let

$$s(t) = \int_0^t \frac{d\tau}{\lambda(\tau)} \quad (3.7)$$

be the rescaled time. We shall make an intensive use of the following rescaling formulas:

$$u(t, r) = \frac{1}{\lambda^{N/2-1}} v(s, y), \quad y = \frac{r}{\lambda}, \quad \frac{ds}{dt} = \frac{1}{\lambda}, \quad (3.8)$$

$$\partial_t u = \frac{1}{\lambda} (\partial_s v + b \Lambda v)_\lambda, \quad (3.9)$$

$$\partial_{tt} u = \frac{1}{\lambda^2} [\partial_s^2 v + b(\partial_s v + 2\Lambda \partial_s v) + b^2 D \Lambda v + b_s \Lambda v]_\lambda. \quad (3.10)$$

In particular, we derive from (1.1) the equation for ε :

$$\begin{aligned} \partial_s^2 \varepsilon + H_{B_1} \varepsilon &= -\Psi_{B_1} - b_s \Lambda P_{B_1} - b(\partial_s P_{B_1} + 2\Lambda \partial_s P_{B_1}) - \partial_s^2 P_{B_1} \\ &\quad - b(\partial_s \varepsilon + 2\Lambda \partial_s \varepsilon) - b_s \Lambda \varepsilon + N(\varepsilon) \end{aligned} \quad (3.11)$$

where, implicitly, $B_1 = B_1(b(t))$ and H_{B_1} is the linear operator associated to the profile P_{B_1}

$$H_{B_1} \varepsilon = -\Delta \varepsilon + b^2 D \Lambda \varepsilon - f'(P_{B_1}) \varepsilon, \quad (3.12)$$

and the nonlinearity:

$$N(\varepsilon) = f(P_{B_1} + \varepsilon) - f(P_{B_1}) - f'(P_{B_1}) \varepsilon. \quad (3.13)$$

Alternatively, the equation for w takes the form:

$$\partial_t^2 w + \tilde{H}_{B_1} w = -[\partial_t^2 (P_{B_1})_\lambda - \Delta (P_{B_1})_\lambda - f((P_{B_1})_\lambda)] + N_\lambda(w)$$

with

$$\tilde{H}_{B_1} w = -\Delta w - f'((P_{B_1})_\lambda) w, \quad (3.14)$$

$$N_\lambda(w) = f((P_{B_1})_\lambda + w) - f((P_{B_1})_\lambda) - f'((P_{B_1})_\lambda) w. \quad (3.15)$$

We then expand using (3.9), (3.10):

$$\begin{aligned} \partial_t^2 (P_{B_1})_\lambda - \Delta (P_{B_1})_\lambda - f((P_{B_1})_\lambda) &= \frac{1}{\lambda^2} [\partial_{ss} P_{B_1} + b(\partial_s P_{B_1} + 2\Lambda \partial_s P_{B_1}) + b_s \Lambda P_{B_1} + \Psi_B]_\lambda \\ &= \frac{1}{\lambda^2} [b\Lambda \partial_s P_{B_1} + b_s \Lambda P_{B_1} + \Psi_B]_\lambda + \partial_t \left[\frac{1}{\lambda} (\partial_s P_{B_1})_\lambda \right] \end{aligned}$$

and rewrite the equation for w :

$$\partial_t^2 w + \tilde{H}_{B_1} w = -\frac{1}{\lambda^2} [b\Lambda \partial_s P_{B_1} + b_s \Lambda P_{B_1} + \Psi_B]_\lambda - \partial_t \left[\frac{1}{\lambda} (\partial_s P_{B_1})_\lambda \right] + N_\lambda(w). \quad (3.16)$$

For most of our arguments we prefer to view the linear operator H_{B_1} acting on w in (3.16) as a perturbation of the linear operator H_λ associated to Q_λ . Then

$$\begin{aligned} \partial_t^2 w + H_\lambda w &= F_{B_1} \\ &= -\frac{1}{\lambda^2} [b\Lambda \partial_s P_{B_1} + b_s \Lambda P_{B_1} + \Psi_{B_1}]_\lambda - \partial_t \left[\frac{1}{\lambda} (\partial_s P_{B_1})_\lambda \right] \\ &\quad - [f'(Q_\lambda) - f'((P_{B_1})_\lambda)] w + N_\lambda(w) \end{aligned} \quad (3.17)$$

with

$$H_\lambda w = -\Delta w + f'(Q_\lambda) w. \quad (3.18)$$

3.3. The set of bootstrap estimates. At first, we fix some notations. We introduce the energy $\mathcal{E}(t)$ associated to the Hamiltonian H_λ :

$$\mathcal{E}(t) = \lambda^2 \int [(H_\lambda \partial_t w, \partial_t w) + (H_\lambda w)^2]. \quad (3.19)$$

Given $\zeta \in (0, \infty)$ the unstable eigenvalue, we set:

$$V_+ = \begin{vmatrix} 1 \\ \sqrt{\zeta} \end{vmatrix} \quad V_- = \begin{vmatrix} 1 \\ -\sqrt{\zeta} \end{vmatrix} \quad (3.20)$$

and, we introduce the decomposition of the unstable direction

$$\begin{vmatrix} (\varepsilon, \psi) \\ (\partial_s \varepsilon, \psi) \end{vmatrix} = \tilde{a}_+(s) V_+ + \tilde{a}_-(s) V_- \quad (3.21)$$

Let us denote:

$$\kappa_+(s) = \tilde{a}_+(s) + \frac{b_s}{2\sqrt{\zeta}} (\partial_b P_{B_1}, \psi), \quad \kappa_-(s) = \tilde{a}_-(s) - \frac{b_s}{2\sqrt{\zeta}} (\partial_b P_{B_1}, \psi). \quad (3.22)$$

We note that the vectors V_+, V_- given by (3.20) yield an eigenbasis of

$$\begin{pmatrix} 0 & 1 \\ \zeta & 0 \end{pmatrix}$$

and hence correspond respectively to the unstable and stable mode of the two dimensional dynamical system

$$\frac{dY}{ds} = \begin{pmatrix} 0 & 1 \\ \zeta & 0 \end{pmatrix} Y$$

which to first order in b is verified by the projection onto the unstable mode (ε, ψ) , see (4.57). The deformation term $b_s(\partial_b P_{B_1}, \psi)$ in (3.22) is present to handle some possible time oscillations induced by the $\partial_s^2 P_{B_1}$ term in the RHS of (3.11) which cannot be estimated in absolute value but will be proved to be lower order.

With these conventions, we may now parameterize the set of initial data described by Lemma 3.1 by $a_+ = \kappa_+(0)$, and then reformulate the initial smallness properties in terms of suitable initial bounds for ε , see Appendix A for the proof which is standard.

Lemma 3.3 (Initial parametrization of the unstable mode and initial bounds). *Let M and b_0 be given as in Lemma 3.1 and denote by $C(M)$ a sufficiently large constant. Then, given (η_0, η_1, a_+) satisfying*

$$|a_+| + \|\eta_0, \nabla \eta_0, \eta_1 + b_0(1 - \chi_{B_1(b_0)})\Lambda Q, \nabla \eta_1\|_{\dot{H}^1 \times \dot{H}^1 \times L^2 \times L^2} \leq \frac{b_0^2}{|\log b_0|}, \quad (3.23)$$

there exists a unique d_+ with $|d_+| \lesssim b_0^2/|\log(b_0)|$ and $T_0 > 0$ such that the unique decomposition

$$u(t) = (P_{B_1(b(t))} + \varepsilon)\lambda(t) = (P_{B_1(b(t))})\lambda(t) + w(t),$$

of the unique smooth solution u to (1.1) on $[0, T_0]$ with initial data (3.2) satisfies the initialization

$$\kappa_+(0) = a_+, \quad (3.24)$$

and the following smallness condition on $[0, T_0]$:

- *Smallness and positivity of b :*

$$0 < b(t) < 5b_0; \quad (3.25)$$

- *Pointwise bound on b_s :*

$$|b_s(t)|^2 \leq C(M) \frac{[b(t)]^4}{|\log b(t)|^2}; \quad (3.26)$$

- *Smallness of the energy norm:*

$$\|(\nabla w(t), \partial_t w(t) + \frac{b(t)}{\lambda(t)}((1 - \chi_{B_1(b(t))})\Lambda Q))\lambda(t)\|_{L^2 \times L^2} \leq \sqrt{b_0}; \quad (3.27)$$

- *Global \dot{H}^2 bound:*

$$|\mathcal{E}(t)| \leq C(M) \frac{[b(t)]^4}{|\log b(t)|^2}; \quad (3.28)$$

- *A priori bound on the stable mode:*

$$|\kappa_-(t)| \leq (C(M))^{\frac{1}{8}} \frac{[b(t)]^2}{|\log b(t)|}. \quad (3.29)$$

- *A priori bound of the unstable mode:*

$$|\kappa_+(t)| \leq 2 \frac{[b(t)]^2}{|\log b(t)|}. \quad (3.30)$$

We may now describe the bootstrap regime as follows:

Definition 3.4 (Exit time). *Let $K(M)$ denote some large enough constant.*

Given $a_+ \in [-\frac{b_0^2}{|\log b_0|}, \frac{b_0^2}{|\log b_0|}]$, we let $T(a_+)$ be the life time of the solution to (1.1) with initial data (3.2), and $T_1(a_+) > 0$ be the supremum of $T \in (0, T(a_+))$ such that for all $t \in [0, T]$, the following estimates hold:

- *Smallness and positivity of b :*

$$0 < b(t) < 5b_0; \quad (3.31)$$

- *Pointwise bound on b_s :*

$$|b_s|^2 \leq K(M) \frac{[b(t)]^4}{|\log b(t)|^2}; \quad (3.32)$$

- *Smallness of the energy norm:*

$$\|(\nabla w(t), \partial_t w(t) + \frac{b(t)}{\lambda(t)}((1 - \chi_{B_1(b(t))})\Lambda Q))_{\lambda(t)}\|_{L^2 \times L^2} \leq \sqrt{b_0}; \quad (3.33)$$

- *Global \dot{H}^2 bound:*

$$|\mathcal{E}(t)| \leq K(M) \frac{[b(t)]^4}{|\log b(t)|^2}; \quad (3.34)$$

- *A priori bound on the stable and unstable modes:*

$$|\kappa_+(t)| \leq 2 \frac{[b(t)]^2}{|\log b(t)|}, \quad |\kappa_-(t)| \leq (K(M))^{\frac{1}{8}} \frac{[b(t)]^2}{|\log b(t)|}. \quad (3.35)$$

The existence of blow up solutions in the regime described by Theorem 1.1 now follows from the following:

Proposition 3.5. *There exists $a_+ \in [-\frac{b_0^2}{|\log b_0|}, \frac{b_0^2}{|\log b_0|}]$ such that*

$$T_1(a_+) = T(a_+)$$

and then corresponding solution to (1.1) blows up in finite time in the regime described by Theorem 1.1.

The proof of Proposition 3.5 relies on a monotonicity argument on the energy \mathcal{E} which is the core of the analysis, see Proposition 4.6, and the strictly outgoing behavior of the unstable mode induced by the non trivial eigenvalue $-\zeta < 0$ of H , see Lemma 4.10. The fact that the regime described by the bootstrap bounds (3.31), (3.32), (3.33), (3.34), (3.35) corresponds to a finite blow up solution with a specific blow up speed will then follow from the modulation equations and the sharp derivation of the blow speed as in [31].

4. Improved bounds

This section is devoted to the derivation of the main dynamical properties of the flow in the bootstrap regime described by Definition 3.4. The three main steps are first the derivation of a monotonicity property on \mathcal{E} which allows us to improve the bounds (3.31), (3.32), (3.33), (3.34) in $[0, T_1(a_+)]$, second the derivation of the dynamics of the eigenmode and the outgoing behavior of the unstable direction, and eventually the derivation of the sharp law for the parameter b which allows to bootstrap its smallness (3.31) and will eventually allow us to derive the sharp blow up speed.

Remark 4.1. All along the proof, we will introduce various constants $C(M), \delta(M) > 0$ which do not depend on the bootstrap constant $K(M)$. An important feature of all these constants is that, up to a smaller choice of $b^*(M)$ or a larger choice of $K(M)$, we assume that any product of the form $C(M) f(b)$ where $\lim_{b \rightarrow 0} f(b) = 0$ or any ratio $\delta(M)/K(M)$ is small in the trapped regime. This will be used implicitly in this section.

4.1. Coercitivity of \mathcal{E} . Let us start with showing that the linearized energy \mathcal{E} yields a control of suitable weighted norms of (w, ε) in the regime $t \in [0, T_1(a_+)]$.

Lemma 4.2 (Coercitivity of \mathcal{E}). *There exists $M_0 \geq 1$ such that for all $M \geq M_0$, there exists⁵ $\delta(M) > 0$ and $C(M) < \infty$ such that in the interval $[0, T_1(a_+)]$, there holds:*

$$\mathcal{E} \geq \frac{1}{2} \lambda^2 \int (H_\lambda w)^2 + \delta(M) \lambda^2 \left[\int (\nabla \partial_t w)^2 + \int \frac{(\partial_r w)^2}{r^2} \right] - C(M) [K(M)]^{\frac{1}{4}} \frac{b^4}{|\log b|^2}. \quad (4.1)$$

Proof of Lemma 4.2.

This is a consequence of the explicit distribution of the negative eigenvalues of H and the a priori bound on the unstable mode (3.35). Indeed, let $t \in [0, T_1(a_+)]$, then first observe from (3.21), (3.22), (3.35) that

$$\begin{aligned} |(\varepsilon, \psi)|^2 + |(\partial_s \varepsilon, \psi)|^2 &\lesssim |\kappa_+|^2 + |\kappa_-|^2 + |b_s|^2 (\partial_b P_{B_1}, \psi)^2 \\ &\lesssim [K(M)]^{\frac{1}{4}} \frac{b^4}{|\log b|^2} + C(M) b^2 |b_s|^2 \lesssim [K(M)]^{\frac{1}{4}} \frac{b^4}{|\log b|^2} \end{aligned} \quad (4.2)$$

where we used the estimates of Proposition 2.1 and the well localization of ψ . This yields

$$\begin{aligned} \frac{1}{\lambda^4} (w, \psi_\lambda)^2 + \frac{1}{\lambda^2} (\partial_t w, \psi_\lambda)^2 &= (\varepsilon, \psi)^2 + (\partial_s \varepsilon + b \Lambda \varepsilon, \psi)^2 \\ &\lesssim [K(M)]^{\frac{1}{4}} \frac{b^4}{|\log b|^2} + b^2 \left[\int \frac{\varepsilon^2}{y^4 (1 + |\log(y)|)^2} + \int \frac{|\nabla \varepsilon|^2}{y^2} \right] \end{aligned} \quad (4.3)$$

and similarly using the orthogonality condition (3.4):

$$\begin{aligned} \frac{1}{\lambda^4} (w, (\chi_M \Phi)_\lambda)^2 + \frac{1}{\lambda^2} (\partial_t w, (\chi_M \Phi)_\lambda)^2 &= (b \Lambda \varepsilon, \chi_M \Phi)^2 \\ &\lesssim b^2 M^C \left[\int \frac{\varepsilon^2}{y^4 (1 + |\log(y)|)^2} + \int \frac{|\nabla \varepsilon|^2}{y^2} \right]. \end{aligned} \quad (4.4)$$

Moreover, applying Lemma C.3 yields:

$$\begin{aligned} \lambda^2 \int |H_\lambda w|^2 &= \int |H \varepsilon|^2 \\ &\geq \delta(M) \left[\int \frac{|\nabla \varepsilon|^2}{y^2} + \frac{\varepsilon^2}{y^4 (1 + |\log(y)|)^2} \right] \end{aligned}$$

Introducing the rescaled version (C.13) of Lemma C.3, we then conclude:

$$\begin{aligned} \mathcal{E} &\geq \frac{1}{2} \int \lambda^2 (H_\lambda w)^2 + \delta_1(M) \left[\lambda^2 \int (\nabla \partial_t w)^2 + \int \frac{|\nabla \varepsilon|^2}{y^2} + \int \frac{\varepsilon^2}{y^4 (1 + |\log(y)|)^2} \right] \\ &\quad - b^2 M^C \left[\int \frac{\varepsilon^2}{y^4 (1 + |\log(y)|)^2} + \int \frac{|\nabla \varepsilon|^2}{y^2} \right] - C(M) [K(M)]^{\frac{1}{4}} \frac{b^4}{|\log b|^2} \\ &\geq \frac{1}{2} \int \lambda^2 (H_\lambda w)^2 + \delta(M) \lambda^2 \left[\int (\nabla \partial_t w)^2 + \int \frac{(\partial_r w)^2}{r^2} \right] - C(M) [K(M)]^{\frac{1}{4}} \frac{b^4}{|\log b|^2} \end{aligned}$$

where we used the Hardy bound (C.3), and (4.1) is proved. This concludes the proof of Lemma 4.2.

Remark 4.3. Note that (4.1) together with the Hardy estimate (C.1), the coercitivity estimate (C.9) and (4.4) yield the following weighted bound on ε which will be extensively used in the paper: let

$$\eta(s, y) = \lambda^{\frac{N-2}{2}+1} \partial_t w(t, \lambda y) = \partial_s \varepsilon(s, y) + b \Lambda \varepsilon(s, y), \quad (4.5)$$

⁵recall remark 4.1

then:

$$\int \frac{\varepsilon^2}{y^4(1+|\log y|^2)} + \int \frac{\eta^2}{y^2} + \int \frac{|\nabla \varepsilon|^2}{y^2} + \int |\nabla \eta|^2 \lesssim c(M) \left[|\mathcal{E}| + [K(M)]^{\frac{1}{4}} \frac{b^4}{|\log b|^2} \right], \quad (4.6)$$

$$\lesssim c(M) |\mathcal{E}| + \sqrt{K(M)} \frac{b^4}{|\log b|^2}. \quad (4.7)$$

4.2. First bound on b_s . We now derive a crude bound on b_s which appears as an *order one* forcing term in the RHS of the equation for ε (3.11). This bound is a simple consequence of the construction of the profile Q_b and the choice of the orthogonality condition (3.4).

Lemma 4.4 (Rough pointwise bound on b_s). *There holds the rough pointwise bound⁶:*

$$\left(b_s + \frac{(\varepsilon, H\Phi)}{(\Lambda Q, \Phi)} \right)^2 \lesssim \frac{1}{M} |\mathcal{E}| + \sqrt{K(M)} \frac{b^4}{|\log b|^2}. \quad (4.8)$$

Remark 4.5. This is in contrast with [31] where the b_s term could be treated as degenerate with respect to ε thanks to a specific choice of orthogonality conditions and the factorization of the operator H in the wave map case. This difficulty in our case will be treated using a specific algebra generated by our choice of orthogonality condition (3.4) which gives the *right sign* to the leading order terms involving b_s in the energy identity (4.6), see (4.24), (4.38).

Proof of Lemma 4.4.

Let us recall that the equation for ε in rescaled variables is given by (3.11), (3.12), (3.13). Observe also that from (1.19), the adjoint of H_B with respect to the $L^2(\mathbb{R}^N)$ inner product is given by:

$$H_{B_1}^* = H_{B_1} + 2b^2 D. \quad (4.9)$$

To compute b_s we take the scalar product of (3.11) with $\chi_M \Phi$. Using the orthogonality relations

$$(\partial_s^m \varepsilon, \chi_M \Phi) = (\partial_s^m (P_{B_1} - Q), \chi_M \Phi) = 0, \quad \forall m \geq 0$$

we integrate by parts to get the algebraic identity:

$$\begin{aligned} b_s [(\Lambda P_{B_1}, \chi_M \Phi) + 2b(\Lambda \partial_b P_{B_1}, \chi_M \Phi) + (\Lambda \varepsilon, \chi_M \Phi)] \\ = -(\Psi_{B_1}, \chi_M \Phi) - (\varepsilon, H_{B_1}^* (\chi_M \Phi)) + 2b(\partial_s \varepsilon, \Lambda (\chi_M \Phi)) + (N(\varepsilon), \chi_M \Phi). \end{aligned} \quad (4.10)$$

We first derive from the estimates of Proposition 2.1:

$$(\Psi_{B_1}, \chi_M \Phi)^2 \lesssim \frac{b^4}{|\log b|^2}. \quad (4.11)$$

Similarly, using (4.6) yields:

$$(\partial_s \varepsilon, \Lambda (\chi_M \Phi))^2 \lesssim C(M) \left[c(M) |\mathcal{E}| + \sqrt{K(M)} \frac{b^4}{|\log b|^2} \right] \quad (4.12)$$

and

$$(\varepsilon, H_{B_1}^* (\chi_M \Phi)) = (\varepsilon, H\Phi) - (H\varepsilon, (1 - \chi_M)\Phi) + O \left(M^C b^2 \sqrt{c(M) |\mathcal{E}| + \sqrt{K(M)} \frac{b^4}{|\log b|^2}} \right).$$

We then use the improved decay (2.2) and (4.7) to estimate:

$$(H\varepsilon, (1 - \chi_M)\Phi)^2 \lesssim \left(\int_{y \geq M} \frac{|H\varepsilon|}{1 + y^N} \right)^2 \lesssim \frac{|\mathcal{E}|}{M} + \sqrt{K(M)} \frac{b^4}{|\log b|^2}$$

⁶recall remark 4.1

Thus:

$$|(\varepsilon, H_{B_1}^*(\chi_M \Phi)) - (\varepsilon, H\Phi)|^2 \lesssim \frac{1}{M} |\mathcal{E}| + \sqrt{K(M)} \frac{b^4}{|\log b|^2}. \quad (4.13)$$

similarly,

$$\begin{aligned} & (\Lambda P_{B_1}, \chi_M \Phi) + 2b(\Lambda \partial_b P_{B_1}, \chi_M \Phi) + (\Lambda \varepsilon, \chi_M \Phi) \\ &= (\Lambda Q, \Phi) + O\left(\frac{b}{\log(b)} + M^C \sqrt{|\mathcal{E}| + \sqrt{K(M)} \frac{b^4}{|\log b|^2}}\right) \\ &= (\Lambda Q, \Phi) + O\left(\frac{b}{\log(b)}\right) \end{aligned} \quad (4.14)$$

where we have used that in the trapped regime $\mathcal{E} \leq K(M)b^4/[\log(b)]^2$. Finally, on the support of χ_M and for $b < b_0^*(M)$ small enough, the term Q dominates in $Q_b = Q + b^2 T_1$. Hence, for the nonlinear term, we have from Sobolev and (4.7):

$$\begin{aligned} |(N(\varepsilon), \chi_M \Phi)| &\lesssim \int \left(\frac{\varepsilon^2}{1+y^6} + \frac{\varepsilon^3}{1+y^4} \right) \lesssim \int \frac{|\varepsilon|^2}{(1+y^5)} [1 + \|y\varepsilon\|_{L^\infty}] \\ &\lesssim C(M) \left[\mathcal{E} + \sqrt{K(M)} \frac{b^4}{|\log b|^2} \right]. \end{aligned}$$

Injecting this together with (4.11), (4.12), (4.13), (4.14) into (4.10) yields (4.8)⁷ and concludes the proof of Lemma 4.4.

4.3. Global \dot{H}^2 bound. We derive in the section a monotonicity statement for the energy \mathcal{E} which provides a global \dot{H}^2 estimate for the solution. The monotonicity statement involves suitable repulsivity properties of the rescaled Hamiltonian H_λ in the focusing regime under the orthogonality condition (3.11) and the a priori control of the unstable mode (3.35), which themselves rely on the positivity of an explicit quadratic form, see Lemma 4.7.

Proposition 4.6 (H^2 control of the radiation). *In the trapped regime, there exists a function \mathcal{F} satisfying*

$$\mathcal{F} \lesssim \frac{\mathcal{E}}{M} + \sqrt{K(M)} \frac{b^4}{|\log b|^2} \quad (4.15)$$

and such that, for some $0 < \alpha < 1$ close enough to 1, there holds:

$$\frac{d}{dt} \left\{ \frac{\mathcal{E} + \mathcal{F}}{\lambda^{2(1-\alpha)}} \right\} \leq \frac{b}{\lambda^{3-2\alpha}} \left[\sqrt{K(M)} \frac{b^4}{|\log b|^2} \right]. \quad (4.16)$$

Proof of Proposition 4.6

step 1 Energy identity.

Let

$$\tilde{V}(t, r) = \frac{N+2}{N-2} Q_\lambda^{\frac{4}{N-2}}(r) = \frac{1}{\lambda^2} V\left(\frac{r}{\lambda}\right), \quad V(y) = \frac{N+2}{N-2} Q^{\frac{4}{N-2}}(y).$$

We first have the following algebraic energy identity which follows by integrating by parts from (3.17):

$$\frac{1}{2} \frac{d}{dt} \left\{ \int (\partial_{tr} w)^2 - \int \tilde{V} (\partial_t w)^2 + \int (H_\lambda w)^2 \right\} = - \int \partial_t \tilde{V} \left[\frac{(\partial_t w)^2}{2} + w H_\lambda w \right] + \int \partial_t w H_\lambda F_{B_1}. \quad (4.17)$$

⁷recall remark 4.1

We now use the w equation and integration by parts to compute:

$$-\int \partial_t \tilde{V} w H_\lambda w = -\int \partial_t \tilde{V} w (F_{B_1} - \partial_{tt} w) \quad (4.18)$$

$$= \frac{d}{dt} \left\{ \int \partial_t \tilde{V} w \partial_t w \right\} - \int \partial_t \tilde{V} w F_{B_1} - \int \partial_t \tilde{V} (\partial_t w)^2 - \int \partial_{tt} \tilde{V} w \partial_t w \quad (4.19)$$

We next pick $0 < \alpha < 1$ close enough to 1 and combine the above identities to get:

$$\begin{aligned} \frac{1}{2\lambda^{2\alpha}} \frac{d}{dt} \left\{ \lambda^{2\alpha} \left[\int (\partial_{tr} w)^2 - \int \tilde{V} (\partial_t w)^2 + \int (H_\lambda w)^2 - 2 \int \partial_t \tilde{V} w \partial_t w \right] \right\} \\ = -R_1 + R_2 + \frac{2\alpha b}{\lambda} \int \partial_t \tilde{V} w \partial_t w - \int \partial_{tt} \tilde{V} w \partial_t w \end{aligned} \quad (4.20)$$

where R_1 collects the quadratic terms:

$$\begin{aligned} R_1 &= \frac{\alpha b}{\lambda} \left[\int (\partial_{tr} w)^2 - \int \tilde{V} (\partial_t w)^2 + \int (H_\lambda w)^2 \right] + \frac{3}{2} \int \partial_t \tilde{V} (\partial_t w)^2 - \frac{b_s}{\lambda^2} \int \partial_t \tilde{V} (\Lambda Q)_\lambda w \\ &= \frac{b}{\lambda^3} \left[\alpha \int (\partial_y \eta)^2 - \alpha \int V \eta^2 + \alpha \int (H\varepsilon)^2 + \frac{3}{2} \int (2V + y \cdot \nabla V) \eta^2 \right. \\ &\quad \left. - b_s \int \varepsilon (2V + y \cdot \nabla V) \Lambda Q \right] \end{aligned} \quad (4.21)$$

and R_2 collects the nonlinear higher order terms:

$$R_2 = \int \partial_t w H_\lambda F_{B_1} - \int \partial_t \tilde{V} w \left[F_{B_1} + \frac{b_s}{\lambda^2} (\Lambda Q)_\lambda \right] \quad (4.22)$$

step 2 Derivation of the quadratic terms and treatment of the b_s term.

Let us now obtain a suitable *lower bound* for the quadratic term R_1 . The main enemy is the b_s term which is order one in ε and will be treated using a specific algebra generated by the choice of orthogonality condition (3.4).

Observe from $H(\Lambda Q) = 0$ that $(\Lambda Q/\lambda)_\lambda(y) = (1/\lambda)^{\frac{N}{2}} (\Lambda Q)(y/\lambda)$ satisfies:

$$-\Delta(\Lambda Q/\lambda)_\lambda(y) - (1/\lambda)^2 V(y/\lambda) (\Lambda Q/\lambda)_\lambda(y) = 0.$$

Differentiating this relation at $\lambda = 1$ yields:

$$H\Phi = H(D\Lambda Q) = (2V + y \cdot \nabla V) \Lambda Q.$$

We inject this into the modulation equation (4.8) to get:

$$-b_s \int \varepsilon (2V + y \cdot \nabla V) \Lambda Q = b_s^2 (\Phi, \Lambda Q) + |b_s| O \left(\frac{|\mathcal{E}|}{M} + \sqrt{K(M)} \frac{b^4}{|\log b|^2} \right)^{\frac{1}{2}}. \quad (4.23)$$

We thus conclude using the sign

$$(\Phi, \Lambda Q) > 0$$

and (4.21), (4.8) that:

$$\begin{aligned} R_1 \geq \frac{b}{\lambda^3} \left[\alpha \int (\partial_y \eta)^2 + \int [(3 - \alpha)V + \frac{3}{2} y \cdot \nabla V] \eta^2 + \alpha \int (H\varepsilon)^2 + c_1 (b_s)^2 \right. \\ \left. + O \left(\frac{|\mathcal{E}|}{M} + \sqrt{K(M)} \frac{b^4}{|\log b|^2} \right) \right] \end{aligned} \quad (4.24)$$

for some universal constant $c_1 > 0$ independent of M .

step 3 Coercitivity of the quadratic form.

We now claim the following coercitivity property of the quadratic form in η appearing in the RHS of (4.24) in the limit case $\alpha = 1$, see Appendix B:

Lemma 4.7 (Coercitivity of the quadratic form). *There exists a universal constant $c_0 > 0$ such that for all $\eta \in \dot{H}_{rad}^1$, there holds:*

$$\int (\partial_y \eta)^2 + \int \left[2V + \frac{3}{2} y \cdot \nabla V \right] \eta^2 \geq c_0 \int (\partial_y \eta)^2 - \frac{1}{c_0} [(\eta, \psi)^2 + (\eta, \Phi)^2].$$

From a simple continuity argument, there exists $0 < \alpha^* < 1$ such that given $0 < \alpha^* < \alpha \leq 1$, for all $\eta \in \dot{H}_{rad}^1$, there holds:

$$\alpha \int (\partial_y \eta)^2 + \int \left[(3 - \alpha)V + \frac{3}{2} y \cdot \nabla V \right] \eta^2 \geq \frac{c_0}{2} \int (\partial_y \eta)^2 - \frac{2}{c_0} [(\eta, \psi)^2 + (\eta, \Phi)^2].$$

We now pick once and for all such an $\alpha < 1$ and control the negative directions.

Using (4.3) and (4.7), it yields:

$$(\eta, \psi)^2 \lesssim b|\mathcal{E}| + \sqrt{K(M)} \frac{b^4}{|\log b|^2}$$

Similarly, we compute $(\eta, \Phi) = (\eta, \chi_M \Phi) + (\eta, (1 - \chi_M) \Phi)$ for which (4.4) and (4.7) yield

$$(\eta, \chi_M \Phi)^2 \lesssim b|\mathcal{E}| + \sqrt{K(M)} \frac{b^4}{|\log b|^2}$$

and we have, applying (C.1):

$$(\eta, (1 - \chi_M) \Phi)^2 \leq \|y\eta\|_{L^\infty}^2 \left[\int_{y \geq M/2} \frac{|\Phi|}{y} \right]^2 \lesssim \frac{1}{M} \int |\partial_y \eta|^2$$

This together with (4.24) yields the lower bound on quadratic terms:

$$R_1 \geq \frac{b}{\lambda^3} \left[c_1 ((b_s)^2 + |\mathcal{E}|) + O \left(\sqrt{K(M)} \frac{b^4}{|\log b|^2} \right) \right] \quad (4.25)$$

for some universal constant $c_1 > 0$. Indeed, a straightforward integration by parts in (3.19) yields:

$$\mathcal{E} \lesssim \int |\partial_y \eta|^2 + \int |H\varepsilon|^2.$$

step 4 Control of lower order quadratic terms.

The lower order quadratic terms in (4.20) are controlled similarly:

$$\begin{aligned} \left| \int \partial_t \tilde{V} w \partial_t w \right| &\lesssim \frac{b}{\lambda^2} \left[\int \frac{\varepsilon^2}{1 + y^6} + \int \frac{\eta^2}{y^2} \right] \lesssim \frac{1}{\lambda^2} \left(bC(M)|\mathcal{E}| + \sqrt{K(M)} \frac{b^4}{|\log b|^2} \right), \\ &\lesssim \frac{1}{\lambda^2} \left(\frac{|\mathcal{E}|}{M} + \sqrt{K(M)} \frac{b^4}{|\log b|^2} \right), \end{aligned} \quad (4.26)$$

and, with the help of (3.32),

$$\begin{aligned} \left| \int \partial_{tt} \tilde{V} w \partial_t w \right| &\lesssim \left(\frac{b^2}{\lambda^3} + \frac{|b_s|}{\lambda^3} \right) \left[\int \frac{\varepsilon^2}{1 + y^6} + \int \frac{\eta^2}{y^2} \right], \\ &\lesssim \frac{b}{\lambda^3} \left(bC(M)|\mathcal{E}| + \sqrt{K(M)} \frac{b^4}{|\log b|^2} \right). \end{aligned}$$

Remark 4.8. We note here that (4.26) is sufficient for the proof of our theorem. Indeed, the estimated term $\int \partial_t \tilde{V} w \partial_t w$ has been integrated by parts with respect to time, so that it becomes a part of \mathcal{F} . Furthermore, we note that to compute (4.16), we multiply \mathcal{F} by $\lambda^{2\alpha}$. Consequently, the commutator $b\alpha/\lambda \int \partial_t \tilde{V} w \partial_t w$ appears on the right-hand side. However, (4.26) yields that, in the trapped regime, this supplementary term is controlled by $b/\lambda^3 \sqrt{K(M)} b^4 / |\log b|^2$. Similar arguments will be repeated implicitly below for the terms which require an integration by parts with respect to time.

step 5 Rewriting of the nonlinear R_2 terms.

It remains to control the nonlinear R_2 terms in (4.20) given by (4.22). According to (3.17), this term contains b_{ss} type of terms which cannot be estimated in absolute value and require a further integration by parts in time. Let

$$F_{B_1} = F_1 - \partial_t F_2 \quad \text{with} \quad F_2 = \frac{1}{\lambda} (\partial_s P_{B_1})_\lambda \quad (4.27)$$

and rewrite:

$$R_2 = \int \partial_t w H_\lambda F_1 - \int \partial_t \tilde{V} w \left[F_1 + \frac{b_s}{\lambda^2} (\Lambda Q)_\lambda \right] - \int \partial_t w H_\lambda \partial_t F_2 + \int \partial_t \tilde{V} w \partial_t F_2$$

We now integrate by parts in time to treat the F_2 term:

$$\begin{aligned} & - \int \partial_t w H_\lambda \partial_t F_2 + \int \partial_t \tilde{V} w \partial_t F_2 \\ &= - \frac{d}{dt} \left\{ \int \partial_t w H_\lambda F_2 - \int \partial_t \tilde{V} w F_2 \right\} - \int (\partial_{tt} \tilde{V} w + 2\partial_t \tilde{V} \partial_t w) F_2 + \int \partial_{tt} w H_\lambda F_2. \end{aligned}$$

The last term is rewritten using (3.17) and integration by parts:

$$\begin{aligned} \int \partial_{tt} w H_\lambda F_2 &= \int [F_1 - \partial_t F_2 - H_\lambda w] H_\lambda F_2 \\ &= - \frac{1}{2} \frac{d}{dt} \left\{ \int |\nabla F_2|^2 - \int \tilde{V} F_2^2 \right\} - \frac{1}{2} \int \partial_t \tilde{V} F_2^2 + \int [F_1 - H_\lambda w] H_\lambda F_2. \end{aligned}$$

eventually arrive at a manageable expression for R_2 :

$$\begin{aligned} R_2 &= - \frac{d}{dt} \left\{ \int \partial_t w H_\lambda F_2 - \int \partial_t \tilde{V} w F_2 + \frac{1}{2} \int |\nabla F_2|^2 - \frac{1}{2} \int \tilde{V} F_2^2 \right\} \\ &\quad - \int \partial_t \tilde{V} w \left[F_1 + \frac{b_s}{\lambda^2} (\Lambda Q)_\lambda \right] + \int \partial_t w H_\lambda F_1 - \int (\partial_{tt} \tilde{V} w + 2\partial_t \tilde{V} \partial_t w) F_2 \\ &\quad - \frac{1}{2} \int \partial_t \tilde{V} F_2^2 + \int [F_1 - H_\lambda w] H_\lambda F_2. \end{aligned} \quad (4.28)$$

We now aim at estimating all the terms in the RHS of (4.28). According to (3.17), we split F_1 into four terms:

$$F_1 + \frac{b_s}{\lambda^2} (\Lambda Q)_\lambda = - \frac{1}{\lambda^2} [\Psi_{B_1} + F_{1,1} + F_{1,2} + N(\varepsilon)]_\lambda \quad (4.29)$$

with

$$F_{1,1} = b\Lambda \partial_s P_{B_1} + b_s (\Lambda P_{B_1} - \Lambda Q), \quad F_{1,2} = [f'(Q) - f'(P_{B_1})] \varepsilon. \quad (4.30)$$

step 6 F_1 terms.

The F_1 terms are the leading order terms.

Ψ_{B_1} terms: We first extract from (2.9) the rough bound:

$$|\Psi_{B_1}| \lesssim \frac{b^2}{|\log b|(1+y^2)} + C(M)b^4 \mathbf{1}_{y \leq 2B_1} \quad (4.31)$$

which yields:

$$\int \frac{1 + |\log y|^2}{1 + y^4} |\Psi_{B_1}|^2 \lesssim \frac{b^4}{|\log b|^2}$$

and thus from (4.7):

$$\begin{aligned} \left| \int \partial_t \tilde{V} w \frac{1}{\lambda^2} (\Psi_{B_1})_\lambda \right| &\lesssim \frac{b}{\lambda^3} \int \frac{|\varepsilon| |\Psi_{B_1}|}{(1+y^4)} \\ &\lesssim \frac{b}{\lambda^3} \frac{b^2}{|\log b|} C(M) \sqrt{|\mathcal{E}| + \sqrt{K(M)}} \frac{b^4}{|\log b|^2} \\ &\lesssim \frac{b}{\lambda^3} \sqrt{K(M)} \frac{b^4}{|\log b|^2}. \end{aligned}$$

Next, we use the fundamental cancellation $H(\Lambda Q) = 0$ and (2.9) to estimate:

$$\begin{aligned} |H\Psi_{B_1}| &\lesssim \frac{b^4}{1+y^2} \left[\frac{1 + |\log(by)|}{|\log b|} \mathbf{1}_{2 \leq y \leq 2B_0} + \frac{1}{b^2 y^2 |\log b|} \mathbf{1}_{\frac{B_0}{2} \leq y \leq 2B_1} + \frac{\log(M) + |\log(1+y)|}{1+y^2} \mathbf{1}_{y \leq 2B_1} \right] \\ &+ \frac{b^2}{(1+y^4)|\log b|} \mathbf{1}_{y \geq B_1/2}, \end{aligned}$$

and thus

$$\int (1+y^2) |H(\Psi_{B_1})|^2 \lesssim \frac{b^6}{|\log b|^2}. \quad (4.32)$$

Hence:

$$\begin{aligned} \left| \int \partial_t w H_\lambda \left(\frac{1}{\lambda^2} (\Psi_{B_1})_\lambda \right) \right| &\lesssim \frac{b}{\lambda^3} \|\eta/y\|_{L^2} \left[\int \frac{1}{b^2} (1+y)^2 |H(\Psi_{B_1})|^2 \right]^{\frac{1}{2}} \\ &\lesssim \frac{b}{\lambda^3} \sqrt{K(M)} \frac{b^4}{|\log b|^2}. \end{aligned}$$

$F_{1,1}$ terms: From (2.7), (2.8), there holds :

$$|F_{1,1}| \lesssim |b_s| b^2 \left[\frac{1 + |\log(by)|}{|\log b|} \mathbf{1}_{2 \leq y \leq \frac{B_0}{2}} + \frac{1}{b^2 y^2 |\log b|} \mathbf{1}_{\frac{B_0}{2} \leq y \leq 2B_1} + \frac{\log(M) + |\log y|}{1+y^2} \right]$$

and, recalling that differentiation w.r.t. y acts as a multiplication by $1/(1+y)$:

$$|HF_{1,1}| \lesssim C(M) \frac{|b_s| b^2}{1+y^2} \left[\frac{1 + |\log(by)|}{|\log b|} \mathbf{1}_{2 \leq y \leq \frac{B_0}{2}} + \frac{1}{b^2 y^2 |\log b|} \mathbf{1}_{\frac{B_0}{2} \leq y \leq 2B_1} + \frac{\log(M) + |\log y|}{1+y^2} \right]$$

from which

$$\int (1+y^2) |H(F_{1,1})|^2 \lesssim |b_s|^2 \frac{b^2}{|\log b|^2}, \quad \int \frac{(1 + |\log y|^2)}{(1+y^4)} |F_{1,1}|^2 \lesssim |b_s|^2 b^2. \quad (4.33)$$

Hence similar arguments as with the Ψ_{B_1} terms yield:

$$\begin{aligned} \left| \int \partial_t \tilde{w} F_{1,1} \right| &\lesssim \frac{b}{\lambda^3} b |b_s| C(M) \sqrt{|\mathcal{E}| + \sqrt{K(M)} \frac{b^4}{|\log b|^2}} \\ &\lesssim \frac{b}{\lambda^3} \sqrt{K(M)} \frac{b^4}{|\log b|^2}, \end{aligned}$$

and

$$\begin{aligned} \left| \int \partial_t w H_\lambda F_{1,1} \right| &\lesssim \frac{C(M)b}{\lambda^3} \frac{|b_s|}{|\log b|} \sqrt{|\mathcal{E}| + \sqrt{K(M)} \frac{b^4}{|\log b|^2}} \lesssim \frac{b}{\lambda^3} \left[\frac{|b_s|^2}{|\log b|} + \frac{\mathcal{E}}{|\log b|} + \sqrt{K(M)} \frac{b^4}{|\log b|^2} \right] \\ &\lesssim \frac{b}{\lambda^3} \sqrt{K(M)} \frac{b^4}{|\log b|^2}. \end{aligned}$$

$F_{1,2}$ terms: The explicit expansion of the cubic nonlinearity and the bound (2.7) yield:

$$|F_{1,2}| \lesssim \frac{C(M)b^2}{1+y^2} |\varepsilon|, \quad |\nabla F_{1,2}| \lesssim \frac{C(M)b^2}{1+y^3} |\varepsilon| + \frac{C(M)b^2}{1+y^2} |\nabla \varepsilon| \quad (4.34)$$

from which:

$$\frac{1}{\lambda^2} \left| \int \partial_t \tilde{w} (F_{1,2})_\lambda \right| \lesssim \frac{C(M)b^3}{\lambda^3} \int \frac{\varepsilon^2}{1+y^6} \lesssim \frac{b}{\lambda^3} \left(b |\mathcal{E}| + \sqrt{K(M)} \frac{b^4}{|\log b|^2} \right),$$

and, after integration by parts of the laplacian term:

$$\begin{aligned} \frac{1}{\lambda^2} \left| \int \partial_t w H_\lambda (F_{1,2})_\lambda \right| &\lesssim \frac{C(M)}{\lambda^3} \left[\int \frac{|\eta|}{1+y^4} \frac{b^2}{1+y^2} |\varepsilon| + \int |\nabla \eta| \left(\frac{b^2}{1+y^3} |\varepsilon| + \frac{b^2}{1+y^2} |\nabla \varepsilon| \right) \right] \\ &\lesssim \frac{b}{\lambda^3} \left[\frac{|\mathcal{E}|}{M} + \sqrt{K(M)} \frac{b^4}{|\log b|^2} \right]. \end{aligned}$$

Nonlinear term $N(\varepsilon)$: We expand the nonlinearity:

$$N(\varepsilon) = 3P_{B_1} \varepsilon^2 + \varepsilon^3.$$

This yields using (3.27), (C.1) the rough bound:

$$|N(\varepsilon)| \lesssim \frac{\varepsilon^2}{1+y}.$$

In what follows, we will use the following bound on η which follows from (4.6), (C.1):

$$|y\eta|_{L^\infty} \lesssim |\nabla \eta|_{L^2} \lesssim \left(c(M) |\mathcal{E}| + \sqrt{K(M)} \frac{b^4}{|\log b|^2} \right)^{\frac{1}{2}}.$$

We then estimate:

$$\begin{aligned} \left| \frac{1}{\lambda^2} \int \partial_t \tilde{w} (N(\varepsilon))_\lambda \right| &\lesssim \frac{b}{\lambda^3} \int \frac{|\varepsilon|^3}{1+y^5} \lesssim \frac{b}{\lambda^3} |\nabla \varepsilon|_{L^2} \left(c(M) |\mathcal{E}| + \sqrt{K(M)} \frac{b^4}{|\log b|^2} \right) \\ &\lesssim \frac{b}{\lambda^3} \sqrt{K(M)} \frac{b^4}{|\log b|^2} \end{aligned}$$

for $b_0 < b^*(M)$ small enough. We split the second term:

$$\int \partial_t w H_\lambda \left(\frac{(N(\varepsilon))_\lambda}{\lambda^2} \right) = \int \nabla \partial_t w \cdot \nabla \left(\frac{(N(\varepsilon))_\lambda}{\lambda^2} \right) - \int \tilde{V} \partial_t w \left(\frac{(N(\varepsilon))_\lambda}{\lambda^2} \right). \quad (4.35)$$

The second term is estimated in brute force:

$$\begin{aligned}
\left| \int \tilde{V} \partial_t w \left(\frac{(N(\varepsilon))_\lambda}{\lambda^2} \right) \right| &\lesssim \frac{1}{\lambda^3} \int \frac{|\eta| |\varepsilon|^2}{1+y^5} \lesssim \frac{1}{\lambda^3} |y\eta|_{L^\infty} \int \frac{|\varepsilon|^2}{1+y^6} \\
&\lesssim \frac{1}{\lambda^3} \left(c(M) |\mathcal{E}| + \sqrt{K(M)} \frac{b^4}{|\log b|^2} \right)^{\frac{3}{2}} \\
&\lesssim \frac{b}{\lambda^3} \frac{b^4}{|\log b|^2}.
\end{aligned}$$

The first term in (4.35) is split into two parts:

$$\int \nabla \partial_t w \cdot \nabla \left(\frac{(N(\varepsilon))_\lambda}{\lambda^2} \right) = \int \nabla \partial_t w \cdot [\nabla(w^3) + 3(P_{B_1})_\lambda \nabla(w^2)] + \frac{3}{\lambda^3} \int \varepsilon^2 \nabla \eta \cdot \nabla P_{B_1}.$$

The last term is integrated by parts in space and then estimated in brute force:

$$\begin{aligned}
\left| \frac{3}{\lambda^3} \int \varepsilon^2 \nabla \eta \cdot \nabla P_{B_1} \right| &= \frac{3}{\lambda^3} \left| \int \eta [\varepsilon^2 \Delta P_{B_1} + 2\varepsilon \nabla P_{B_1} \cdot \nabla \varepsilon] \right| \\
&\lesssim \frac{1}{\lambda^3} \int |\eta| \left[\frac{\varepsilon^2}{1+y^4} + \frac{|\varepsilon| |\nabla \varepsilon|}{1+y^3} \right] \lesssim \frac{1}{\lambda^3} |y\eta|_{L^\infty} \left[\int \frac{\varepsilon^2}{1+y^5} + \int \frac{|\nabla \varepsilon|^2}{y^2} \right] \\
&\lesssim \frac{1}{\lambda^3} \left(c(M) |\mathcal{E}| + \sqrt{K(M)} \frac{b^4}{|\log b|^2} \right)^{\frac{3}{2}} \lesssim \frac{b}{\lambda^3} \frac{b^4}{|\log b|^2}.
\end{aligned}$$

The first term is the most delicate one and requires first a time integration by parts:

$$\begin{aligned}
\int \nabla \partial_t w \cdot [\nabla(w^3) + 3(P_{B_1})_\lambda \nabla(w^2)] &= \frac{d}{dt} \left\{ \int |\nabla w|^2 \left[\frac{3}{2} w^2 + 3(P_{B_1})_\lambda w \right] \right\} \\
- 3 \int w \partial_t w |\nabla w|^2 - 3 \int |\nabla w|^2 [w \partial_t (P_{B_1})_\lambda + (P_{B_1})_\lambda \partial_t w].
\end{aligned}$$

We may now estimate all terms in brute force:

$$\left| \int |\nabla w|^2 \left[\frac{3}{2} w^2 + 3(P_{B_1})_\lambda w \right] \right| \lesssim \frac{1}{\lambda^2} [|y\varepsilon|_{L^\infty} + |yP_{B_1}|_{L^\infty}] |y\varepsilon|_{L^\infty} \int \frac{|\nabla \varepsilon|^2}{y^2} \lesssim \frac{1}{\lambda^2} \frac{b^4}{|\log b|^2},$$

$$\begin{aligned}
\left| \int w \partial_t w |\nabla w|^2 \right| &\lesssim \frac{1}{\lambda^2} |y\varepsilon|_{L^\infty} |y\eta|_{L^\infty} \int \frac{|\nabla \varepsilon|^2}{y^2} \lesssim \left(c(M) |\mathcal{E}| + \sqrt{K(M)} \frac{b^4}{|\log b|^2} \right)^{\frac{3}{2}} \\
&\lesssim \frac{b}{\lambda^3} \frac{b^4}{|\log b|^2},
\end{aligned}$$

$$\begin{aligned}
\left| \int |\nabla w|^2 w \partial_t (P_{B_1})_\lambda \right| &\lesssim \frac{|yw|_{L^\infty}}{\lambda^3} \int \frac{|\nabla w|^2}{y} \left[\frac{b}{1+y^2} + C(M) b |b_s| \mathbf{1}_{y \leq B_1} \right] \\
&\lesssim \frac{b}{\lambda^3} |\nabla \varepsilon|_{L^2} \left(1 + C(M) |b_s| \frac{|\log b|}{b} \right) \int \frac{|\nabla \varepsilon|^2}{y^2} \\
&\lesssim \frac{b}{\lambda^3} \frac{b^4}{|\log b|^2}
\end{aligned}$$

where we used the rough bound extracted from (2.8): $|\partial_b P_{B_1}| \lesssim C(M)b\mathbf{1}_{y \leq B_1}$, and finally:

$$\begin{aligned} \left| \int |\nabla w|^2 (P_{B_1})_\lambda \partial_t w \right| &\lesssim \frac{1}{\lambda^3} |y\eta|_{L^\infty} \int \frac{|\nabla \varepsilon|^2}{1+y^3} \lesssim \left(C(M)\mathcal{E} + \sqrt{K(M)} \frac{b^4}{|\log b|^2} \right)^{\frac{3}{2}} \\ &\lesssim \frac{b}{\lambda^3} \frac{b^4}{|\log b|^2}, \end{aligned}$$

for $b_0 < b^*(M)$ small enough. The above chain of estimates together with remark 4.8 closes the control of the nonlinear term $N(\varepsilon)$.

step 9 F_2 terms.

We estimate from (2.8):

$$\int \left| \frac{\partial_b P_{B_1}}{(1+y)} \right|^2 + \int |\nabla \partial_b P_{B_1}|^2 \lesssim \frac{1}{|\log b|^2}, \quad \int \frac{1}{1+y^3} |\partial_b P_{B_1}|^2 \lesssim \frac{b}{|\log b|^2} \quad (4.36)$$

and hence:

$$\begin{aligned} \left| \int \partial_t w H_\lambda F_2 \right| &\lesssim \frac{|b_s|}{\lambda^2} \left[\int \frac{|\eta| |\partial_b P_{B_1}|}{(1+y^4)} + \int |\nabla \eta| |\nabla \partial_b P_{B_1}| \right] \\ &\lesssim \frac{1}{\lambda^2} \frac{|b_s|}{|\log b|} \sqrt{c(M)|\mathcal{E}| + \sqrt{K(M)} \frac{b^4}{|\log b|^2}} \\ &\lesssim \frac{1}{\lambda^2} \left[\frac{|\mathcal{E}|}{M} + \sqrt{K(M)} \frac{b^4}{|\log b|^2} \right], \\ \left| \int \partial_t \tilde{V} w F_2 \right| &\lesssim \frac{|b_s|b}{\lambda^2} \int \frac{|\partial_b P_{B_1}| |\varepsilon|}{(1+y^4)} \lesssim \frac{|b_s|b}{\lambda^2 |\log b|} \left[c(M)|\mathcal{E}| + \sqrt{K(M)} \frac{b^4}{|\log b|^2} \right]^{\frac{1}{2}} \\ &\lesssim \frac{1}{\lambda^2} \left[\frac{|\mathcal{E}|}{M} + \sqrt{K(M)} \frac{b^4}{|\log b|^2} \right], \\ \int |\nabla F_2|^2 + \int V F_2^2 &\lesssim \frac{|b_s|^2}{\lambda^2} \left[\int \frac{|\partial_b P_{B_1}|^2}{(1+y^4)} + \int |\nabla \partial_b P_{B_1}|^2 \right] \\ &\lesssim \frac{1}{\lambda^2} \frac{(b_s)^2}{|\log b|^2} \lesssim \frac{1}{\lambda^2} \frac{b^4}{|\log b|^2}. \end{aligned}$$

similarly:

$$\begin{aligned} \left| \int (\partial_{tt} \tilde{V} w + 2\partial_t \tilde{V} \partial_t w) F_2 \right| + \left| \int \partial_t \tilde{V} F_2^2 \right| &\lesssim \frac{|b_s|}{\lambda^3} \left[\int \frac{((|b_s| + b^2)|\varepsilon| + b|\eta|) |\partial_b P_{B_1}|}{(1+y^4)} + |b_s|b \int \frac{|\partial_b P_{B_1}|^2}{1+y^4} \right] \\ &\lesssim \frac{|b_s|}{\lambda^3} \left[\frac{(|b_s| + b)}{|\log b|} \sqrt{c(M)|\mathcal{E}| + \sqrt{K(M)} \frac{b^4}{|\log b|^2}} + \frac{b^2}{|\log b|^2} |b_s| \right] \\ &\lesssim \frac{b}{\lambda^3} \left[\frac{|\mathcal{E}|}{M} + \sqrt{K(M)} \frac{b^4}{|\log b|^2} \right]. \end{aligned}$$

Eventually, (4.32), (4.33) ensure:

$$\int (1+y^2) |H(\Psi_{B_1} + F_{1,1})|^2 \lesssim \left[\frac{b^6}{|\log b|^2} + \frac{b^2 |b_s|^2}{|\log b|^2} \right] \lesssim \frac{b^6}{|\log b|^2}$$

which together with (4.36) yields:

$$\left| \int \frac{1}{\lambda^2} (\Psi_{B_1} + F_{1,1})_\lambda H_\lambda F_2 \right| \lesssim \frac{1}{\lambda^3} \frac{b^3 |b_s|}{|\log b|^2} \lesssim \frac{b}{\lambda^3} \frac{b^4}{|\log b|^2}.$$

We similarly estimate from (4.34) and after integration by parts:

$$\begin{aligned} \left| \int \frac{1}{\lambda^2} (F_{1,2})_\lambda H_\lambda F_2 \right| &\lesssim \frac{|b_s|}{\lambda^3} \left[\int \frac{b^2 |\varepsilon| |\partial_b P_{B_1}|}{1+y^6} + \int |\nabla \partial_b P_{B_1}| \left(\frac{b^2 |\varepsilon|}{1+y^3} + \frac{b^2 |\nabla \varepsilon|}{1+y^2} \right) \right] \\ &\lesssim C(M) \frac{b^4}{\lambda^3 |\log b|} \left(\int \frac{\varepsilon^2}{1+y^6} + \int \frac{|\nabla \varepsilon|^2}{1+y^4} \right)^{\frac{1}{2}} \lesssim \frac{b}{\lambda^3} \frac{b^4}{|\log b|^2}. \end{aligned}$$

For the nonlinear term, we extract from (2.8) the rough bound

$$|H(\partial_b P_{B_1})| \lesssim [C(M) + \log(b)] \frac{b}{1+y^2} \mathbf{1}_{y \leq B_1},$$

which together with (C.1) ensures:

$$\begin{aligned} \left| \int \frac{1}{\lambda^2} (N(\varepsilon))_\lambda H_\lambda F_2 \right| &\lesssim \frac{[C(M) + \log(b)]}{\lambda^3} |b_s| \int \frac{b}{1+y^2} \frac{\varepsilon^2}{1+y} \mathbf{1}_{y \leq B_1} \\ &\lesssim C(M) \frac{|b_s| |\log b|^4}{\lambda^3} \int \frac{\varepsilon^2}{(1+y^4) |\log y|^2} \\ &\lesssim \frac{b}{\lambda^3} \sqrt{b} \left(c(M) |\mathcal{E}| + \sqrt{K(M)} \frac{b^4}{|\log b|^2} \right) \lesssim \frac{b}{\lambda^3} \frac{b^4}{|\log b|^2}. \end{aligned}$$

step 10 The remaining F_2 term has the right sign.

It remains to estimate the term

$$- \int H_\lambda w H_\lambda F_2$$

in the RHS of (4.28). Let us stress onto the fact that this term is a priori no better $O(\frac{1}{\lambda^3} \mathcal{E})$ due to the b_s contribution and the bound (4.8), recall remark 4.5.

We now claim that the *main contribution has the right sign again*.

Indeed, we first compute from the T_1 equation (2.16):

$$HT_1 = -\Phi + c_b \chi_{\frac{B_0}{4}} \Lambda Q, \quad H \partial_b T_1 = O \left(\frac{1}{b |\log b|} \frac{\mathbf{1}_{2 \leq y \leq \frac{B_0}{2}}}{(1+y^2)} \right) \quad (4.37)$$

We then apply the decomposition (2.22):

$$H(\partial_b P_{B_1}) = H \left(2bT_1 + 2b(\chi_{B_1} - 1)T_1 + b^2 \partial_b \log(B_1) \rho_{B_1} T_1 + b^2 \chi_{B_1} \partial_b T_1 \right) = -2b\Phi + \Sigma$$

and estimate using (2.8), (2.21), (4.37):

$$|\Sigma| \lesssim \frac{b}{1+y^2} \left[\frac{1}{|\log b|} \mathbf{1}_{2 \leq y \leq \frac{B_0}{2}} + \frac{1}{b^2 y^2 |\log b|} \mathbf{1}_{\frac{B_0}{2} \leq y} \right].$$

In particular,

$$\int \Sigma^2 \lesssim \frac{b^2}{|\log b|}$$

and thus using the modulation equation (4.8) :

$$\begin{aligned}
& - \int H_{\lambda} w H_{\lambda} F_2 = - \frac{b_s}{\lambda^3} \int (H\varepsilon) H(\partial_b P_{B_1}) = - \frac{b_s}{\lambda^3} \int H\varepsilon (-2b\Phi + \Sigma) \\
& = 2 \frac{b}{\lambda^3} b_s (\varepsilon, H\Phi) + \frac{b}{\lambda^3} O \left(\frac{|b_s|}{\sqrt{|\log b|}} \sqrt{|\mathcal{E}| + \sqrt{K(M)} \frac{b^4}{|\log b|^2}} \right) \\
& = 2 \frac{b}{\lambda^3} \left[- \frac{(\varepsilon, H\Phi)}{(\Lambda Q, \Phi)} + O \left(\sqrt{\frac{|\mathcal{E}|}{M} + \sqrt{K(M)} \frac{b^4}{|\log b|^2}} \right) \right] (\varepsilon, H\Phi) + \frac{b}{\lambda^3} O \left(\frac{b^4}{|\log b|^2} \right) \\
& = - \frac{2b}{\lambda^3} \frac{(\varepsilon, H\Phi)^2}{(\Lambda Q, \Phi)} + O \left(\frac{|\mathcal{E}|}{M} + \sqrt{K(M)} \frac{b^4}{|\log b|^2} \right) + \frac{b}{\lambda^3} O \left(\frac{b^4}{|\log b|^2} \right) \tag{4.38} \\
& \leq O \left(\frac{|\mathcal{E}|}{M} + \sqrt{K(M)} \frac{b^4}{|\log b|^2} \right). \tag{4.39}
\end{aligned}$$

The recollection of all above estimates yields (4.16) and concludes the proof of Proposition 4.6.

4.4. Improved bound. We now claim that the a priori bound on the unstable direction (3.35) coupled with the monotonicity property of Proposition 4.6 imply the following improved bounds:

Lemma 4.9 (Improved bounds under the a priori control (3.35)). *There holds in $[0, T_1(a_+)]$:*

$$\|(\nabla w(t), \partial_t w(t) + \frac{b(t)}{\lambda(t)}((1 - \chi_{B_1(b(t))})\Lambda Q))_{\lambda(t)}\|_{L^2 \times L^2} \lesssim b_0 |\log b_0|, \tag{4.40}$$

$$\frac{b^4(t)}{|\log b(t)|^2 \lambda^{2(1-\alpha)}(t)} \geq \frac{b^4(0)}{|\log b(0)|^2 \lambda^{2(1-\alpha)}(0)}, \tag{4.41}$$

$$|b_s|^2 \leq \frac{K(M)}{2} \frac{b^4}{|\log b|^2}, \tag{4.42}$$

$$|\mathcal{E}(t)| \leq \frac{K(M)}{2} \frac{b^4}{(\log b)^2}. \tag{4.43}$$

Proof of Lemma 4.9

step 1 Energy bound.

The energy bound (4.40) is a consequence of the conservation of the energy. Indeed, the conservation of the energy and the initial bounds of Lemma 3.1 ensure

$$E(u, \partial_t u) = E(u_0, u_1) = E(Q) + O(b_0 \sqrt{|\log b_0|}),$$

(see Appendix 3.1) and thus:

$$\begin{aligned}
& E(Q) + O(b_0 |\log b_0|) \\
& = \frac{1}{2} \int [\partial_t (P_{B_1})_{\lambda} + \partial_t w]^2 + \frac{1}{2} \int |\nabla (P_{B_1})_{\lambda} + \nabla w|^2 - \frac{1}{4} \int [(P_{B_1})_{\lambda} + w]^4. \tag{4.44}
\end{aligned}$$

We lower bound the first term by expanding,

$$\begin{aligned}
\partial_t (P_{B_1})_{\lambda} + \partial_t w & = \partial_t w + \frac{b}{\lambda} ((1 - \chi_{B_1})\Lambda Q)_{\lambda} + \frac{b}{\lambda} (\chi_{B_1} \Lambda Q)_{\lambda} + \frac{b^3}{\lambda} (\Lambda[\chi_{B_1} T_1])_{\lambda} + \frac{b_s}{\lambda} (\partial_b P_{B_1})_{\lambda} \\
& = \partial_t w + \frac{b}{\lambda} ((1 - \chi_{B_1})\Lambda Q)_{\lambda} + \Sigma,
\end{aligned}$$

with

$$\int \Sigma^2 \lesssim b_0^2 |\log b_0|,$$

where we used the bootstrap bounds (3.31), (3.32). Finally :

$$\int [\partial_t(P_{B_1})_\lambda + \partial_t w]^2 \geq \frac{1}{2} \int \left[\frac{b}{\lambda} ((1 - \chi_{B_1}) \Lambda Q)_\lambda + \partial_t w \right]^2 - O(b_0^2 |\log b_0|). \quad (4.45)$$

We then expand the second term:

$$\begin{aligned} & \frac{1}{2} \int |\nabla(P_{B_1})_\lambda + \nabla w|^2 - \frac{1}{4} \int [(P_{B_1})_\lambda + w]^4 = \frac{1}{2} \int |\nabla P_{B_1} + \nabla \varepsilon|^2 - \frac{1}{4} \int [P_{B_1} + \varepsilon]^4 \\ &= \frac{1}{2} \int |\nabla P_{B_1}|^2 - \frac{1}{4} \int |P_{B_1}|^4 - (\varepsilon, \Delta P_{B_1} + P_{B_1}^3) + \frac{1}{2} \left(\int |\nabla \varepsilon|^2 - 3 \int P_{B_1}^2 \varepsilon^2 \right) \\ & - \frac{1}{4} (4P_{B_1} \varepsilon^3 + \varepsilon^4) \end{aligned}$$

From the construction of P_{B_1} ,

$$\frac{1}{2} \int |\nabla P_{B_1}|^2 - \frac{1}{4} \int |P_{B_1}|^4 = E(Q) + O(b^2 |\log b|). \quad (4.46)$$

The linear term is treated using (2.9), the improved decay (2.2) and (4.31):

$$\begin{aligned} |(\varepsilon, \Delta P_{B_1} + P_{B_1}^3)| &= |(\varepsilon, b^2 D \Lambda P_{B_1} - \Psi_{B_1})| \\ &\lesssim \|\varepsilon/y\|_{L^2} \|y(b^2 D \Lambda P_{B_1} - \Psi_{B_1})\|_{L^2} \lesssim b \|\nabla \varepsilon\|_{L^2} \end{aligned} \quad (4.47)$$

We now rewrite the quadratic term as a small deformation of H and use the coercivity bound (C.8) to ensure:

$$\int |\nabla \varepsilon|^2 - 3 \int P_{B_1}^2 \varepsilon^2 \geq c_0 \int |\nabla \varepsilon|^2 + Def, \quad (4.48)$$

with

$$Def := 3 \int (Q^2 - P_{B_1}^2) \varepsilon^2 - \frac{(\varepsilon, \psi)^2}{c_0}.$$

Collecting (2.7) and (C.1), on the one hand, and (4.2) on the other hand, we compute:

$$\left| \int (Q^2 - P_{B_1}^2) \varepsilon^2 \right| \leq \|y^2(Q^2 - P_{B_1}^2)\|_{L^\infty} \|\nabla \varepsilon\|_{L^2}^2 \lesssim b \|\nabla \varepsilon\|_{L^2}^2, \quad |(\varepsilon, \psi)|^2 \lesssim b^2 |\log b|. \quad (4.49)$$

The nonlinear term is easily estimated from Sobolev:

$$\int |(3P_{B_1} + \varepsilon) \varepsilon^3| \leq \|y P_{B_1}\|_{L^\infty} \|y \varepsilon\|_{L^\infty} \|\nabla \varepsilon\|_{L^2}^2 \lesssim \sqrt{b_0} \|\nabla \varepsilon\|_{L^2}^2 \quad (4.50)$$

Injecting (4.45), (4.47), (4.46), (4.49), (4.48), (4.50) into (4.44) yields (4.40).

step 2 Lower bound on b .

We now turn to the proof of (4.41). First observe from the bootstrap estimate (3.32) that

$$|b_s| \leq \sqrt{K(M)} \frac{b^2}{|\log b|} \leq \frac{1 - \alpha}{10} b^2 \quad (4.51)$$

This implies:

$$\frac{d}{ds} \left(\frac{b^4}{(\log b)^2 \lambda^{2(1-\alpha)}} \right) = \frac{4b^3}{\lambda^{2(1-\alpha)} (\log b)^2} \left[b_s \left(1 - \frac{1}{2 \log b} \right) + \frac{1 - \alpha}{2} b^2 \right] > 0$$

and (4.41) follows.

step 3 Improved \dot{H}^2 bound.

We now turn to the proof of (4.43). We integrate (4.16) in time and conclude from (4.1), (4.15):

$$\begin{aligned} |\mathcal{E}(t)| &\lesssim \left(\frac{\lambda(t)}{\lambda(0)} \right)^{2(1-\alpha)} |\mathcal{E}(0)| \\ &+ (K(M))^{\frac{1}{2}} \left[\frac{b^4(t)}{|\log b(t)|^2} + [\lambda(t)]^{2(1-\alpha)} \int_0^t \frac{b(\tau)}{[\lambda(\tau)]^{3-2\alpha}} \frac{b^4(\tau)}{|\log b(\tau)|^2} d\tau \right] \end{aligned} \quad (4.52)$$

We then derive from (4.51):

$$\begin{aligned} &\int_0^t \frac{b(\tau)}{[\lambda(\tau)]^{3-2\alpha}} \frac{b^4(\tau)}{|\log b(\tau)|^2} d\tau = - \int_0^t \frac{\lambda_t}{\lambda^{3-2\alpha}} \frac{b^4}{|\log b|^2} d\tau \\ &\leq \frac{1}{2(1-\alpha)} \frac{b^4(t)}{\lambda^{2(1-\alpha)}(t) |\log b(t)|^2} - \frac{1}{2(1-\alpha)} \int_0^t \frac{b_s}{\lambda^{3-2\alpha}} \frac{b^3}{|\log b|^2} \left[1 - \frac{2}{|\log b|^2} \right] \\ &\lesssim \frac{b^4(t)}{\lambda^{2(1-\alpha)}(t) |\log b(t)|^2} + \sqrt{K(M)} \int_0^t \frac{b(\tau)}{[\lambda(\tau)]^{3-2\alpha}} \frac{b^4(\tau)}{|\log b(\tau)|^2} \frac{1}{|\log b(\tau)|} d\tau \end{aligned}$$

and hence the bound:

$$\lambda^{2(1-\alpha)}(t) \int_0^t \frac{b(\tau)}{[\lambda(\tau)]^{3-2\alpha}} \frac{b^4(\tau)}{|\log b(\tau)|^2} d\tau \lesssim \frac{b^4(t)}{|\log b(t)|^2}.$$

Injecting this into (4.52) and using the initial bound (A.12),(A.17) and the monotonicity (4.41) yields:

$$\begin{aligned} \mathcal{E}(t) &\lesssim \left(\frac{\lambda(t)}{\lambda(0)} \right)^{2(1-\alpha)} \frac{b^4(0)}{|\log b(0)|^2} + (K(M))^{\frac{1}{2}} \frac{b^4(t)}{|\log b(t)|^2} \\ &\lesssim \sqrt{K(M)} \frac{b^4(t)}{|\log b(t)|^2} \end{aligned} \quad (4.53)$$

and (4.43) follows. (4.42) now follows from (4.4) and (4.53).

This concludes the proof of Lemma 4.9.

4.5. Dynamic of the unstable mode. We now focus onto the dynamic of the unstable mode. We recall the decomposition

$$Y(t) = \begin{pmatrix} (\varepsilon, \psi) \\ (\partial_s \varepsilon, \psi) \end{pmatrix} = \tilde{a}_+(t) V_+ + \tilde{a}_-(t) V_-, \quad (4.54)$$

and the variables given by (3.22):

$$\kappa_+(s) = \tilde{a}_+(s) + \frac{b_s}{2\sqrt{\zeta}} (\partial_b P_{B_1}, \psi), \quad \kappa_-(s) = \tilde{a}_-(s) - \frac{b_s}{2\sqrt{\zeta}} (\partial_b P_{B_1}, \psi).$$

Lemma 4.10 (Control of the unstable mode). *There holds: for all $t \in [0, T_1(a_+)]$,*

$$|\kappa_-(t)| \leq \frac{1}{2} (K(M))^{\frac{1}{8}} \frac{b^2}{|\log b|} \quad (4.55)$$

and κ_+ is strictly outgoing:

$$\left| \frac{d\kappa_+}{ds} - \sqrt{\zeta} \kappa_+ \right| \leq \sqrt{b} \frac{b^2}{|\log b|}. \quad (4.56)$$

Proof of Lemma 4.10

We compute the equation satisfied by the unstable direction (ε, ψ) by taking the inner product of (3.11) with the well localized direction ψ to get:

$$\frac{d^2}{ds^2}(\varepsilon, \psi) - \zeta(\varepsilon, \psi) = E(\varepsilon) - (\partial_s^2 P_{B_1}, \psi) \quad (4.57)$$

with

$$\begin{aligned} E(\varepsilon) &= -(\Psi_{B_1}, \psi) - b_s(\Lambda P_{B_1}, \psi) - b(\partial_s P_{B_1} + 2\Lambda \partial_s P_{B_1}, \psi) - b(\partial_s \varepsilon + 2\Lambda \partial_s \varepsilon, \psi) \\ &\quad - b_s(\Lambda \varepsilon, \psi) + (N(\varepsilon), \psi) + b^2(\Lambda \varepsilon, D\psi) + ((f'(P_{B_1}) - f'(Q))\varepsilon, \psi). \end{aligned} \quad (4.58)$$

Simple algebraic manipulations using (4.54), (3.22) and the initial condition yield the equivalent system:

$$\frac{d}{ds}\kappa_+ = \sqrt{\zeta}\kappa_+(s) + \frac{E_+(s)}{2\sqrt{\zeta}}, \quad \frac{d}{ds}\kappa_- = -\sqrt{\zeta}\kappa_-(s) - \frac{E_-(s)}{2\sqrt{\zeta}}\kappa_-(0) \quad (4.59)$$

with

$$E_+(s) = E(s) - \frac{b_s}{2}(\partial_b P_{B_1}, \psi) \quad E_-(s) = E(s) + \frac{b_s}{2}(\partial_b P_{B_1}, \psi) \quad (4.60)$$

We now have from the explicit formula (4.58), (4.60), the exponential localization of ψ , the orthogonality

$$(\psi, \Lambda Q) = 0,$$

the estimates of Proposition 2.1 and the bootstrap estimate (3.32) the bound:

$$\frac{1}{\sqrt{\zeta}}|E_{\pm}| \lesssim |b|(|b_s| + \sqrt{|\mathcal{E}|} + \sqrt{K(M)}) \frac{b^2}{|\log b|} \leq \sqrt{b} \frac{b^2}{|\log b|}, \quad (4.61)$$

which together with (4.59) yields (4.56). Let then

$$\mathcal{G} = \kappa_-^2 \frac{|\log b|^2}{b^4},$$

then from (4.59), (4.61), (3.32), we estimate:

$$\begin{aligned} \frac{d\mathcal{G}}{ds} &= 2\kappa_- \frac{d\kappa_-}{ds} \frac{|\log b|^2}{b^4} + \kappa_-^2 b_s \left[-\frac{4|\log b|^2}{b^5} + \frac{2\log b}{b^5} \right] \\ &= 2 \frac{|\log b|^2}{b^4} \left[\kappa_- \left(-\sqrt{\zeta}\kappa_- - \frac{E_-}{\sqrt{\zeta}} \right) \right] + \kappa_-^2 \frac{|\log b|^2}{b^4} O\left(\frac{|b_s|}{b}\right) \\ &\leq -\frac{\sqrt{\zeta}}{2} \frac{|\log b|^2}{b^4} \kappa_-^2 + \frac{|\log b|^2}{b^4} \kappa_- \sqrt{b} \frac{b^2}{|\log b|} \lesssim -\frac{\sqrt{\zeta}}{2} \mathcal{G} + 1. \end{aligned}$$

We integrate this in time

$$\mathcal{G}(s) \leq \mathcal{G}(0)e^{-\frac{\sqrt{\zeta}}{2}s} + \int_0^s e^{-\frac{\sqrt{\zeta}}{2}(s-\sigma)} d\sigma \lesssim 1$$

where we used the initial inequality (A.18) yielding that $\mathcal{G}(0) \lesssim 1$. This concludes the proof of (4.55) and of Lemma 4.10.

4.6. Derivation of the sharp law for b . We now turn to the derivation of the sharp law for b which will yield the required monotonicity statement on b to close the smallness bootstrap estimate (3.31), and will eventually lead to the derivation of the sharp blow up speed (1.10).

Lemma 4.11 (Sharp derivation of the b law). *Let*

$$\tilde{P}_{B_0} = \chi_{\frac{B_0}{4}} Q, \quad (4.62)$$

$$G(b) = b|\Lambda\tilde{P}_{B_0}|_{L^2}^2 + \int_0^b \tilde{b}(\partial_b \tilde{P}_{B_0}, \Lambda\tilde{P}_{B_0}) d\tilde{b}, \quad (4.63)$$

$$\mathcal{I}(s) = (\partial_s \varepsilon, \Lambda\tilde{P}_{B_0}) + b(\varepsilon + 2\Lambda\varepsilon, \Lambda\tilde{P}_{B_0}) + b_s(\partial_b \tilde{P}_{B_0}, \Lambda\tilde{P}_{B_0}) - b_s \left(\partial_b (P_{B_1} - \tilde{P}_{B_0}), \Lambda\tilde{P}_{B_0} \right), \quad (4.64)$$

then there holds:

$$G(b) = 64b|\log b| + O(b), \quad |\mathcal{I}| \lesssim K(M)b, \quad (4.65)$$

$$\left| \frac{d}{ds} \{G(b) + \mathcal{I}(s)\} + 32b^2 \right| \lesssim K(M) \frac{b^2}{\sqrt{|\log b|}}. \quad (4.66)$$

Remark 4.12. Observe that (4.65), (4.66) essentially yield a pointwise differential equation

$$b_s \sim -\frac{b^2}{2|\log b|}$$

which will allow us to derive the sharp scaling law via the relationship $-\frac{\lambda_s}{\lambda} = b$.

Proof of Lemma 4.11

The proof is inspired by the one in [31]. We multiply (3.11) with $\Lambda\tilde{P}_{B_0}$ and compute:

$$\begin{aligned} & (b_s \Lambda P_{B_1} + b(\partial_s P_{B_1} + 2\Lambda \partial_s P_{B_1}) + \partial_s^2 P_{B_1}, \Lambda\tilde{P}_{B_0}) = -(\Psi_{B_1}, \Lambda\tilde{P}_{B_0}) - (H_{B_1} \varepsilon, \Lambda\tilde{P}_{B_0}) \\ & - \left(\partial_s^2 \varepsilon + b(\partial_s \varepsilon + 2\Lambda \partial_s \varepsilon) + b_s \Lambda \varepsilon, \Lambda\tilde{P}_{B_0} \right) + (N(\varepsilon), \Lambda\tilde{P}_{B_0}) \end{aligned}$$

We further rewrite this as follows:

$$\begin{aligned} & (b_s \Lambda \tilde{P}_{B_0} + b(\partial_s \tilde{P}_{B_0} + 2\Lambda \partial_s \tilde{P}_{B_0}) + \partial_s^2 \tilde{P}_{B_0}, \Lambda\tilde{P}_{B_0}) = -(\Psi_{B_1}, \Lambda\tilde{P}_{B_0}) \\ & - (b_s \Lambda (P_{B_1} - \tilde{P}_{B_0}) + b(\partial_s (P_{B_1} - \tilde{P}_{B_0}) + 2\Lambda \partial_s (P_{B_1} - \tilde{P}_{B_0})) + \partial_s^2 (P_{B_1} - \tilde{P}_{B_0}), \Lambda\tilde{P}_{B_0}) \\ & - (H_{B_1} \varepsilon, \Lambda\tilde{P}_{B_0}) - \left(\partial_s^2 \varepsilon + b(\partial_s \varepsilon + 2\Lambda \partial_s \varepsilon) + b_s \Lambda \varepsilon, \Lambda\tilde{P}_{B_0} \right) + (N(\varepsilon), \Lambda\tilde{P}_{B_0}). \end{aligned} \quad (4.67)$$

We now estimate all terms in the above identity.

step 1 b terms.

An integration by parts in time allows us to rewrite the left-hand side of (4.67) as follows:

$$(b_s \Lambda \tilde{P}_{B_0} + b(\partial_s \tilde{P}_{B_0} + 2\Lambda \partial_s \tilde{P}_{B_0}) + \partial_s^2 \tilde{P}_{B_0}, \Lambda\tilde{P}_{B_0}) = \frac{d}{ds} \left[G(b) + b_s(\partial_b \tilde{P}_{B_0}, \Lambda\tilde{P}_{B_0}) \right] + |b_s|^2 |\partial_b \tilde{P}_{B_0}|_{L^2}^2 \quad (4.68)$$

with G given by (4.63). Observe from (3.32) the bound

$$|b_s|^2 |\partial_b \tilde{P}_{B_0}|_{L^2}^2 \lesssim \frac{|b_s|^2}{b^2} \lesssim (K(M))^2 \frac{b^2}{|\log b|^2} \lesssim \frac{b^2}{\sqrt{|\log b|}}.$$

We now turn to the key step in the derivation of the sharp b law which corresponds to the following outgoing flux computation⁸:

$$(\Psi_{B_1}, \Lambda \tilde{P}_{B_0}) = 32b^2 \left(1 + O\left(\frac{1}{|\log b|}\right) \right) \quad \text{as } b \rightarrow 0. \quad (4.69)$$

Indeed, we first estimate from (2.9):

$$\begin{aligned} \left| (\Psi_{B_1} - c_b b^2 \chi_{\frac{B_0}{4}} \Lambda Q, \Lambda \tilde{P}_{B_0}) \right| &\lesssim b^4 \int_{y \leq \frac{B_0}{2}} \left[\frac{1 + |\log(by)|}{|\log b|(1+y^2)} + \frac{1 + |\log(1+y)|}{(1+y^2)^2} \right] \\ &\lesssim \frac{b^2}{|\log b|}. \end{aligned}$$

The remainder term is computed from (2.10) and the explicit formula for Q (1.3):

$$\begin{aligned} (c_b b^2 \chi_{\frac{B_0}{4}} \Lambda Q, \Lambda \tilde{P}_{B_0}) &= \frac{b^2}{2|\log b|} \left(1 + O\left(\frac{1}{|\log b|}\right) \right) \left[\int_{y \leq \frac{1}{2b}} (\Lambda Q)^2 + O(1) \right] \\ &= 32b^2 \left(1 + O\left(\frac{1}{|\log b|}\right) \right), \end{aligned}$$

and (4.69) follows.

We now estimate the lower order terms in b which correspond to the second line of (4.67). One term is reintegrated by parts in time:

$$-(\partial_s^2(P_{B_1} - \tilde{P}_{B_0}), \Lambda \tilde{P}_{B_0}) = -\frac{d}{ds} \left\{ b_s (\partial_b(P_{B_1} - \tilde{P}_{B_0}), \Lambda \tilde{P}_{B_0}) \right\} + b_s^2 (\partial_b(P_{B_1} - \tilde{P}_{B_0}), \partial_b \Lambda \tilde{P}_{B_0}).$$

The remaining terms are estimated in brute force using (2.8) and (3.32) which yield:

$$\begin{aligned} &\left| (b_s \Lambda(P_{B_1} - \tilde{P}_{B_0}) + b(\partial_s(P_{B_1} - \tilde{P}_{B_0}) + 2\Lambda \partial_s(P_{B_1} - \tilde{P}_{B_0}))), \Lambda \tilde{P}_{B_0} \right| \\ &+ b_s^2 \left| (\partial_b(P_{B_1} - \tilde{P}_{B_0}), \partial_b \Lambda \tilde{P}_{B_0}) \right| \lesssim |b_s| + \frac{|b_s|^2}{b^2} \lesssim K(M) \frac{b^2}{|\log b|}. \end{aligned}$$

step 2 ε terms .

We are left with estimating the third line on the RHS of (4.67). We first treat the linear term from (4.1), (4.7), (3.34):

$$\left| (H_{B_1} \varepsilon, \Lambda \tilde{P}_{B_0}) \right| \lesssim |(H \varepsilon, \Lambda \tilde{P}_{B_0})| + \int |\varepsilon| |P_{B_1}^2 - Q^2| |\Lambda \tilde{P}_{B_0}| + b^2 \left| (D \Lambda \varepsilon, \Lambda \tilde{P}_{B_0}) \right| \quad (4.70)$$

On the one hand, (4.7) together with bootstrap estimates yield:

$$\begin{aligned} \int |\varepsilon| |P_{B_1}^2 - Q^2| |\Lambda \tilde{P}_{B_0}| &\lesssim b^2 \int_{y \leq B_0} \frac{|\varepsilon|}{(1+y^2)^2} \leq b^{\frac{3}{2}} \left(\int \frac{|\varepsilon|^2}{(1+y)^5} \right)^{\frac{1}{2}} \\ &\lesssim \frac{b^2}{|\log(b)|} \end{aligned}$$

⁸see again [31] for more details about the flux computation statement and its connection to the Pohozaev integration by parts formula

On the other hand, after integration by parts, we repeat the same arguments and apply (C.4). This yield:

$$\begin{aligned}
b^2 \left| (D\Lambda\varepsilon, \Lambda\tilde{P}_{B_0}) \right| &\leq b^2 \int_{y \leq B_0} \frac{|\varepsilon|}{(1+y^4)} + b^2 \int_{B_0/4 \leq y \leq B_0/2} \frac{|\varepsilon|}{(1+y^2)} + b^2 \int_{y \leq B_0} |\nabla\varepsilon| \frac{y}{1+y^2} \\
&\lesssim b^{\frac{3}{2}} \left(\int \frac{|\varepsilon|^2}{(1+y^5)} \right)^{\frac{1}{2}} + \left(\int_{B_0/4 \leq y \leq B_0/2} \frac{|\varepsilon|^2}{(1+y^4)} \right)^{\frac{1}{2}} + \left(\int_{y \leq B_0} \frac{|\nabla\varepsilon|^2}{1+y^2} \right)^{\frac{1}{2}} \\
&\lesssim \sqrt{|\log(b)|} \left(c(M)|\mathcal{E}| + \sqrt{K(M)} \frac{b^4}{|\log b|^2} \right)^{\frac{1}{2}} \\
&\lesssim \sqrt{K(M)} \frac{b^2}{\sqrt{|\log(b)|}}.
\end{aligned}$$

Finally:

$$\begin{aligned}
|(H\varepsilon, \Lambda\tilde{P}_{B_0})| &\lesssim |H\varepsilon|_{L^2} \sqrt{|\log b|} + \sqrt{K(M)} \frac{b^2}{\sqrt{|\log(b)|}} \\
&\lesssim \sqrt{|\log b|} \sqrt{|\mathcal{E}| + \sqrt{K(M)} \frac{b^4}{|\log b|^2}} \lesssim \sqrt{K(M)} \frac{b^2}{\sqrt{|\log b|}}
\end{aligned}$$

We further integrate by parts in time to obtain:

$$\begin{aligned}
\left(\partial_s^2 \varepsilon + b(\partial_s \varepsilon + 2\Lambda \partial_s \varepsilon) + b_s \Lambda \varepsilon, \Lambda \tilde{P}_{B_0} \right) &= \frac{d}{ds} \left[(\partial_s \varepsilon, \Lambda \tilde{P}_{B_0}) + b(\varepsilon + 2\Lambda \varepsilon, \Lambda \tilde{P}_{B_0}) \right] \\
&\quad - b_s \left[(\partial_s \varepsilon + b\Lambda \varepsilon, \Lambda \partial_b \tilde{P}_{B_0}) + (\varepsilon, \Phi_b) \right]
\end{aligned}$$

with

$$\Phi_b = -\Lambda \tilde{P}_{B_0} - \Lambda^2 \tilde{P}_{B_0} - b\Lambda \partial_b \tilde{P}_{B_0} - b\Lambda^2 \partial_b \tilde{P}_{B_0}.$$

We thus estimate from (4.1), (4.5), (4.7), (3.32), (3.34):

$$\begin{aligned}
|b_s| \left| (\partial_s \varepsilon + b\Lambda \varepsilon, \Lambda \partial_b \tilde{P}_{B_0}) + (\varepsilon, \Phi_b) \right| &\lesssim |b_s| \left[\int_{\frac{B_0}{4} \leq y \leq B_0} \frac{|\eta|}{y} + \int_{y \leq B_0} \frac{|\varepsilon|}{1+y^2} \right] \\
&\lesssim \frac{|b_s| |\log b|}{b^2} C(M) \sqrt{|\mathcal{E}| + \sqrt{K(M)} \frac{b^4}{|\log b|^2}} \lesssim K(M) \frac{b^2}{\sqrt{|\log b|}}
\end{aligned}$$

The non linear term is estimated as previously. Indeed, we have:

$$\begin{aligned}
\left| (N(\varepsilon), \Lambda \tilde{P}_{B_0}) \right| &\lesssim \int (|P_{B_1}| + |\varepsilon|) \varepsilon^2 |\Lambda \tilde{P}_{B_0}| \\
&\lesssim \frac{1}{b^2} \|y(|P_{B_1}| + |\varepsilon|)\|_{L^\infty} \|(1+y^2)\Lambda \tilde{P}_{B_0}\|_{L^\infty} \int_0^{B_0} \frac{|\varepsilon|^2}{y(1+y^4)} \\
&\lesssim \frac{C(M)}{b^2} \left[\mathcal{E} + K(M) \frac{b^4}{|\log b|^2} \right] \lesssim K(M) \frac{b^2}{\sqrt{|\log b|}}.
\end{aligned}$$

step 5 Control of $G(b)$ and \mathcal{I} .

Injecting the estimates of step 1 and step 2 into (4.67) yields (4.66). It remains to prove (4.65). The estimate for $G(b)$ is a straightforward consequence of the choice (4.62) and the explicit formula

(1.3). It remains to control \mathcal{I} . We integrate by parts in space in (4.64) to rewrite:

$$\mathcal{I}(s) = (\partial_s \varepsilon + b\Lambda \varepsilon, \Lambda \tilde{P}_{B_0}) - b(\varepsilon, \Lambda \tilde{P}_{B_0} + \Lambda^2 \tilde{P}_{B_0}) + b_s(\partial_b \tilde{P}_{B_0}, \Lambda \tilde{P}_{B_0}) - b_s \left(\partial_b (P_{B_1} - \tilde{P}_{B_0}), \Lambda \tilde{P}_{B_0} \right).$$

The b terms are estimated as in step 1:

$$|b_s| \left| (\partial_b \tilde{P}_{B_0}, \Lambda \tilde{P}_{B_0}) - (\partial_b (P_{B_1} - \tilde{P}_{B_0}), \Lambda \tilde{P}_{B_0}) \right| \lesssim \frac{|b_s|}{b} \lesssim b.$$

The linear term is estimated using (4.1), (4.5), (4.7), (3.32), (3.34):

$$\begin{aligned} & \left| (\partial_s \varepsilon + b\Lambda \varepsilon, \Lambda \tilde{P}_{B_0}) - b(\varepsilon, \Lambda \tilde{P}_{B_0} + \Lambda^2 \tilde{P}_{B_0}) \right| \lesssim \int_{y \leq B_0} \frac{|\eta|}{y^2} + b \int_{y \leq B_0} \frac{|\varepsilon|}{y^2} \\ & \lesssim \frac{1}{b} \left(\int \frac{|\eta|^2}{y^2} \right)^{\frac{1}{2}} + \frac{|\log b|}{b^2} \left(\int_{y \leq B_0} \frac{|\varepsilon|^2}{y^4(1+|\log y|^2)} \right)^{\frac{1}{2}} \lesssim K(M)b \end{aligned}$$

and (4.65) is proved.

This concludes the proof of Lemma 4.11.

5. Sharp description of the singularity formation

We are now in position to conclude the proof of Proposition 3.5 and Theorem 1.1 as a simple consequence of the a priori bounds obtained in the previous section. The proof relies on a topological argument which closes the bootstrap argument, and then the sharp description of the blow up dynamic is a consequence of the a priori bounds obtained on the solution and in particular the modulation equation (4.66).

Proof of Proposition 3.5

We argue by contradiction and assume that for all $a_+ \in \left[-\frac{b_0^2}{|\log b_0|}, \frac{b_0^2}{|\log b_0|} \right]$,

$$T_1(a_+) < T(a_+).$$

In view of the Definition 4.9 of the bootstrap regime and the improved bounds of Lemma 4.9 and Lemma 4.10, a simple continuity argument ensures that $T_1(a_+)$ is attained at the first time t where

$$|\kappa_+(t)| = \frac{|b(t)|^2}{2|\log(b(t))|}. \quad (5.1)$$

The fundamental fact now is the outgoing behaviour (4.56) which together with (5.1) ensures

$$\left| \frac{d\kappa_+}{dt}(T_1(a_+)) \right| > 0.$$

Thus from standard argument⁹, the map

$$\begin{aligned} \left[-\frac{b_0^2}{|\log b_0|}, \frac{b_0^2}{|\log b_0|} \right] & \rightarrow \mathbb{R}^*_{+} \\ a_+ & \mapsto T_1(a_+) \end{aligned} \quad \text{is continuous.}$$

We may thus consider the continuous map:

$$\begin{aligned} \Phi : \left[-\frac{b_0^2}{|\log b_0|}, \frac{b_0^2}{|\log b_0|} \right] & \rightarrow \mathbb{R} \\ a_+ & \rightarrow \kappa_+(T_1(a_+)) \frac{2|\log b(T_1(a_+))|}{b^2(T_1(a_+))} \end{aligned}$$

⁹see [5, Lemma 6] for a complete exposition

On the one hand, (5.1) implies:

$$\Phi \left(\left[-\frac{b_0^2}{|\log b_0|}, \frac{b_0^2}{|\log b_0|} \right] \right) \subset \{-1, 1\}.$$

On the other hand, the outgoing behavior (4.56) together with the initialization $\kappa_+(0) = a_+$ ensures:

$$\Phi \left(-\frac{b_0^2}{|\log b_0|} \right) = -1, \quad \Phi \left(\frac{b_0^2}{|\log b_0|} \right) = 1$$

and a contradiction follows.¹⁰This concludes the proof of Proposition 3.5.

Proof of Theorem 1.1

step 1 Finite time blow up and derivation of the blow up speed.

Let from Proposition 3.5 an initial data with $T_1(a_+) = T(a_+)$. We first claim that u blows up in finite time

$$T = T(a_+) < +\infty. \quad (5.2)$$

Indeed, from (4.41),

$$\lambda^{2(1-\alpha)} \lesssim b^3 \quad \text{and thus} \quad \lambda^{\frac{2}{3}} \lesssim \lambda^{\frac{2(1-\alpha)}{3}} \lesssim b = -\lambda_t.$$

Integrating this differential inequation yields

$$t \lesssim \lambda^{\frac{1}{3}}(0) - \lambda^{\frac{1}{3}}(t) \lesssim 1$$

and (5.2) follows. The $(\dot{H}^1 \cap \dot{H}^2) \times (L^2 \cap \dot{H}^1)$ bounds (3.33), (3.34) on $(\varepsilon, \partial_t \varepsilon)$ and hence on $(u, \partial_t u)$ in the bootstrap regime and standard H^2 local well posedness theory ensure that blow up corresponds to

$$\lambda(t) \rightarrow 0 \quad \text{as} \quad t \rightarrow T(a_+).$$

We now derive the blow up speed by reintegrating the ODE (4.66) and briefly sketch the proof which follows as in [31].

First recall the standard scaling lower bound

$$\lambda(t) \leq C(u_0)(T - t)$$

which implies that the rescaled time is global:

$$s(t) = \int_0^t \frac{d\tau}{\lambda(\tau)} \rightarrow +\infty \quad \text{as} \quad t \rightarrow T.$$

Let

$$\mathcal{J} = G + \mathcal{I}$$

so that from (4.65):

$$\mathcal{J} = 64b|\log b| \left(1 + O \left(\frac{1}{|\log b|} \right) \right) \quad \text{ie} \quad b = \frac{\mathcal{J}}{64|\log \mathcal{J}|} \left(1 + O \left(\frac{1}{\sqrt{|\log \mathcal{J}|}} \right) \right) \quad (5.3)$$

and \mathcal{J} satisfies from (4.66) the ODE:

$$\mathcal{J}_s + \frac{\mathcal{J}^2}{128|\log \mathcal{J}|^2} \left(1 + O \left(\frac{1}{\sqrt{|\log \mathcal{J}|}} \right) \right) = 0.$$

¹⁰This topological argument is of course the one dimensional version of Brouwer's fixed point argument used in [5].

We multiply the above by $\frac{|\log \mathcal{J}|^2}{\mathcal{J}^2}$, integrate in time and obtain to leading order:

$$\mathcal{J} = \frac{128(\log s)^2}{s} \left(1 + O \left(\frac{1}{\sqrt{|\log s|}} \right) \right) \text{ ie } -\frac{\lambda_s}{\lambda} = b = \frac{2\log s}{s} \left(1 + O \left(\frac{1}{\sqrt{|\log s|}} \right) \right).$$

where we used (5.3). Integrating this once more in time yields:

$$-\log \lambda = (\log s)^2 \left(1 + O \left(\frac{1}{\sqrt{|\log s|}} \right) \right)$$

and thus

$$b = -\lambda_t = \exp \left(-\sqrt{|\log \lambda|} \left(1 + O \left(\frac{1}{|\log \lambda|^{\frac{1}{4}}}} \right) \right) \right).$$

Integrating this from t to T where $\lambda(T) = 0$ yields the asymptotic

$$\lambda(t) = (T - t) \exp \left(-\sqrt{|\log \lambda(t)|} \left(1 + O \left(\frac{1}{|\log \lambda(t)|^{\frac{1}{4}}}} \right) \right) \right)$$

which yields (1.10).

step 2 Energy quantization.

It remains to prove (1.9) which can be derived exactly as in [31], this is left to the reader. This concludes the proof of Theorem 1.1.

Appendix A. Modulation theory

This appendix is devoted to the proof of Lemmas 3.1 and 3.3. The arguments are standard in the framework of modulation theory and we briefly sketch the main computations.

A.1. Proof of Lemma 3.1. First note that the bounds

$$|\nabla(P_{B_1} - Q)|_{L^2} + b|\Lambda P_{B_1} - b(1 - \chi_{B_1})\Lambda Q|_{L^2} \lesssim b|\log b|$$

ensure that our initial data are of the form

$$u_0 = Q + \tilde{\eta}_0, \quad u_1 = \tilde{\eta}_1$$

for a small excess of energy in the sense that:

$$\|\nabla \tilde{\eta}_0, \tilde{\eta}_1\|_{L^2 \times L^2} \lesssim b_0 |\log b_0|, \quad \|\nabla^2 \tilde{\eta}_0, \nabla \tilde{\eta}_1\|_{L^2 \times L^2} \lesssim b_0, \quad (\text{A.1})$$

Hence the continuity of the flow associated to (1.1) ensures the existence of a time $T_0 > 0$ (uniform in $\tilde{\eta}_0, \tilde{\eta}_1$) for which the solution u to (1.1) (u_0, u_1) satisfies on $[0, T_0]$:

$$\sup_{[0, T_0]} \|\nabla(u - Q), \partial_t u\|_{L^2 \times L^2} \lesssim b_0 |\log b_0|, \quad (\text{A.2})$$

Step 1 : Modulation near Q .

The non degeneracy $(\Lambda Q, \Phi) \neq 0$ ensures¹¹ that u admits on $[0, T_0]$ a decomposition

$$u(t) = (Q + \tilde{\varepsilon}(t))_{\lambda(t)} \quad (\text{A.3})$$

with:

$$(\tilde{\varepsilon}(t), \chi_M \Phi) = 0. \quad (\text{A.4})$$

¹¹as a direct consequence of the implicit function theorem and the smoothness of the flow (1.1)

Moreover, $\lambda \in \mathcal{C}^2([0, T_0]; \mathbb{R}_+^*)$ and noting that $\tilde{\eta}_0$ satisfies

$$|(\tilde{\eta}_0, \chi_M \Phi)| \lesssim \frac{b_0^2}{|\log b_0|},$$

we obtain the bound:

$$|\lambda(0) - 1| \lesssim \frac{b_0^2}{|\log b_0|}. \quad (\text{A.5})$$

We then let $b(t) = -\lambda_t(t)$ on $[0, T_0]$.

Step 2 : Positivity of b .

Straightforward computations yield:

$$\partial_t \tilde{\varepsilon}(t) = \left(\partial_t u - \frac{b(t)}{\lambda(t)} \Lambda u \right)_{\frac{1}{\lambda(t)}}.$$

Taking the scalar product with $\chi_M \Phi$, we obtain at the initial time:

$$b(0) = \lambda(0) \frac{((u_1)_{\frac{1}{\lambda(0)}}), \chi_M \Phi}{((\Lambda u_0)_{\frac{1}{\lambda(0)}}), \chi_M \Phi}, \quad (\text{A.6})$$

where (2.5) together with (A.5) imply:

$$((u_1)_{\frac{1}{\lambda(0)}}), \chi_M \Phi = b_0 (\Lambda Q, \chi_M \Phi) + O\left(\frac{b_0^2}{|\log(b_0)|}\right), \quad (\text{A.7})$$

$$((\Lambda u_0)_{\frac{1}{\lambda(0)}}), \chi_M \Phi = (\Lambda Q, \chi_M \Phi) + O(b_0^2 |\log(b_0)|). \quad (\text{A.8})$$

This yields the positivity of $b(0)$ and moreover, the positivity of $b(t)$ for small time together with:

$$b(t) = b_0 + O\left(\frac{b_0^2}{|\log(b_0)|}\right) \quad (\text{A.9})$$

As $b > 0$, we may introduce the decomposition:

$$u(t) = (Q + \tilde{\varepsilon})_{\lambda(t)} = (P_{B_1(b(t))} + \varepsilon)_{\lambda(t)} \quad \text{ie} \quad \varepsilon(t) = \tilde{\varepsilon}(t) - (P_{B_1(b(t))} - Q). \quad (\text{A.10})$$

Observe from (2.4), (A.4) that

$$\forall t \in [0, T_0], \quad (\varepsilon(t), \chi_M \Phi) = 0. \quad (\text{A.11})$$

The uniqueness of such a decomposition is guaranteed by the (local) uniqueness of $(\lambda, \tilde{\varepsilon})$.

Step 3 : Smallness of ε .

To complete the proof, we obtain the smallness of ε in \dot{H}^1 and \dot{H}^2 . To this end, we note that:

$$\varepsilon(0) = (u_0)_{\frac{1}{\lambda(0)}} - P_{B_1(b(0))} = \left[(P_{B_1(b_0)})_{\frac{1}{\lambda(0)}} - P_{B_1(b(0))} \right] + (\eta_0 + d_+ \psi)_{\frac{1}{\lambda(0)}}.$$

Simple computations based on the estimates of Proposition 2.1 yield the expected result :

$$\|\nabla \varepsilon(0)\|_{L^2} \lesssim b_0 |\log(b_0)| \quad \left\| \frac{\varepsilon(0)}{1+y^4} \right\|_{L^2} + \|\nabla^2 \varepsilon(0)\|_{L^2} \lesssim \frac{b_0^2}{|\log(b_0)|}. \quad (\text{A.12})$$

A.2. Proof of Lemma 3.3. The proof of this lemma is divided into two steps. First, given (η_0, η_1, d_+) satisfying smallness condition (3.1) for small b_0 , we prove that b, b_s and w satisfy (3.31)–(3.34). Then, we show that, given (b_0, η_0, η_1) , we can apply the inverse mapping theorem to $d_+ \mapsto \kappa_+(0)$ close to 0. The arguments are standard and we refer to [5] for a detailed proof in a similar setting.

Step 1: Smallness of initial modulation given (η_0, η_1, d_+) .

Given (η_0, η_1, d_+) satisfying smallness condition (3.1) we can apply Lemma 3.1 this yields T_0 and b, ε, w such that (3.31) holds and

$$\|\nabla w(t)\|_{L^2} \lesssim b_0 |\log(b_0)| \quad \|\nabla^2 w(t)\|_{L^2} \lesssim \frac{b_0^2}{|\log(b_0)|^2} \quad (\text{A.13})$$

We emphasize in particular that Lemma 3.1 implies $b_0/2 < b(0) < 2b_0$ for sufficiently small b_0 .

As previously, we focus now on bounds satisfied initially. We first compute $b_s(0)$ using (1.1) and the orthogonality condition (A.11). Recalling that $(\partial_b^k P_{B_1}, \chi_M \Phi) = (\partial_s^{k-1} \varepsilon, \chi_M \Phi) = 0$ for any integer k , we get like for (4.10):

$$\begin{aligned} b_s [(\Lambda P_{B_1}, \chi_M \Phi) + 2b(\Lambda \partial_b P_{B_1}, \chi_M \Phi) + (\Lambda \varepsilon, \chi_M \Phi)] \\ = -(\Psi_{B_1}, \chi_M \Phi) - (\varepsilon, H_{B_1}^*(\chi_M \Phi)) + b(\partial_s \varepsilon, \Lambda(\chi_M \Phi)) + (N(\varepsilon), \chi_M \Phi) \end{aligned}$$

where, denoting LHS and RHS the left-hand and right-hand side at initial time, we compute, for sufficiently small b_0 w.r.t. M :

$$|RHS| \leq C(M) \left(\frac{b_0^2}{|\log(b_0)|} + \|\partial_s \varepsilon\|_{L^2(y < M)} \right), \quad \frac{|b_s(0)|}{2} (\Lambda Q, \chi_M \Phi) \leq |LHS|. \quad (\text{A.14})$$

On the other hand, after time-differentiation, we obtain :

$$\partial_s \varepsilon(0) = \lambda(0) \partial_t \varepsilon(0) = -b_s(0) \partial_b P_{B_1(b(0))} - b(0) \Lambda u_0 + \lambda(0) (b_0 \Lambda P_{B_1(b_0)})_{\frac{1}{\lambda(0)}}. \quad (\text{A.15})$$

Observe now from (2.8) that

$$\|\partial_b P_{B_1(b_0)}\|_{L^2(y \leq 2M)} \lesssim C(M) b_0 \leq \sqrt{b_0}$$

which together with (A.5), (A.9) and (3.1) yields:

$$\|\partial_s \varepsilon(0)\|_{L^2(y \leq 2M)} = \lambda(0) \|\partial_t \varepsilon(0)\|_{L^2(y \leq 2M)} \lesssim \frac{b_0^2}{|\log b_0|} + |b_s(0)| \sqrt{b_0}, \quad (\text{A.16})$$

which together with (A.14) concludes the proof of the initial bound (3.26) on b_s .

Then, we compute :

$$\partial_t w(0) = u_1 - \left(\frac{b_s(0)}{\lambda(0)} \partial_b P_{B_1(b(0))} + \frac{b(0)}{\lambda(0)} \Lambda P_{B_1(b(0))} \right)_{\lambda(0)}$$

so that, introducing (A.15) and previous estimates on $b(0)$, we compute :

$$\|\partial_t w(0) + \frac{b(0)}{\lambda(0)} ((1 - \chi_{B_1(b(0))}) \Lambda Q)_{\lambda(0)}\|_{L^2} \lesssim b_0 |\ln(b_0)| \leq \sqrt{b_0}.$$

and

$$\|\nabla \partial_t w(0)\|_{L^2} \lesssim \frac{b_0^2}{|\log b_0|}, \quad (\text{A.17})$$

Together with (A.13), this yields (3.27) and (3.28).

Finally, straightforward computations yield:

$$\kappa_- = \frac{1}{2}(\varepsilon, \psi) - \frac{1}{\zeta}(\partial_s \varepsilon, \psi) - \frac{b_s}{2\zeta}(\partial_b P_{B_1}, \psi),$$

Consequently, we apply (3.28), noting that $w(t) = (\varepsilon(t))_{\lambda(t)}$, and (A.15) because of the exponential decay of ψ to compute

$$|\kappa_-(0)| \lesssim \frac{b_0^2}{|\log b_0|} \quad (\text{A.18})$$

Step 2: Computation of d_+ .

We now claim from an explicit computation that given a_+ , the initialization (3.24) can be reformulated in the form

$$F(d_+) = a_+ \quad \text{with} \quad \frac{\partial F}{\partial d_+}|_{d_+=0} = \frac{\|\psi\|_{L^2}^2}{2} + O(b_0) \quad (\text{A.19})$$

which from the implicit function theorem concludes the proof of Lemma 3.3.

Let us briefly justify (A.19). We want to study the mapping

$$\begin{aligned} \mathcal{V} &\rightarrow \mathbb{R}^4 \\ d_+ &\mapsto [b(t), b_s(t), (\varepsilon(0), \psi), (\partial_s \varepsilon(t), \psi)] \end{aligned}$$

where \mathcal{V} is a neighborhood of 0. To this end, it is necessary to study the dependencies of all initial parameters on d_+ . For conciseness, we denote by d differentiation w.r.t. d_+ in what follows

Computation of $(\lambda(0), \tilde{\varepsilon}(0))$. As a first step in the modulation theory, we proved that $(\lambda(0), \tilde{\varepsilon}(0)) = \Phi(u_0)$ where Φ is a smooth mapping $\dot{H}^1(\mathbb{R}^N) \rightarrow \mathbb{R} \times \dot{H}^1(\mathbb{R}^N)$ defined in a neighborhood of Q . Due to the exponential decay of $\psi \in C^\infty(\mathbb{R}^N)$ we thus have that $\lambda(0)$ is a smooth function of d_+ with differential $d\lambda(0) = d\lambda \in \mathbb{R}$. We have the same result for ε with differential $d\tilde{\varepsilon}(0) = d\tilde{\varepsilon} \in \dot{H}^1(\mathbb{R}^N)$. By definition, we have

$$\tilde{\varepsilon}(0) = u_0 - Q_{\frac{1}{\lambda}}$$

so that:

$$d\tilde{\varepsilon} = \psi + \frac{d\lambda}{\lambda(0)}(\Lambda Q)_{\frac{1}{\lambda(0)}}.$$

Computation of $b(0)$: From (A.6), $b(0)$ is a C^1 mapping with:

$$\begin{aligned} db(0) &= d\lambda \left[\frac{((u_1)_{\frac{1}{\lambda(0)}}, \chi_M \Phi)}{((\Lambda u_0)_{\frac{1}{\lambda(0)}}, \chi_M \Phi)} + \frac{((\Lambda^2 u_0)_{\frac{1}{\lambda(0)}}, \chi_M \Phi) - ((\Lambda u_1)_{\frac{1}{\lambda(0)}}, \chi_M \Phi)}{((\Lambda u_0)_{\frac{1}{\lambda(0)}}, \chi_M \Phi)^2} \right] \\ &\quad - \lambda(0) \frac{((u_1)_{\frac{1}{\lambda(0)}}, \chi_M \Phi)((\Lambda \psi)_{\frac{1}{\lambda(0)}}, \chi_M \Phi)}{((\Lambda u_0)_{\frac{1}{\lambda(0)}}, \chi_M \Phi)^2} \end{aligned}$$

where (A.6) and (A.7) ensure that, for some $db \in \mathbb{R}$, there holds :

$$db(0) = db + O(b_0).$$

Computation of $\varepsilon(0)$: Next,

$$\varepsilon(0) = \tilde{\varepsilon}(0) - (P_{B_1(b(0))} - Q)$$

Consequently, $(\varepsilon(0), \psi)$ is also a smooth function of d_+ with derivative $dps_1(0)$ satisfying

$$dps_1(0) = (d\tilde{\varepsilon}, \psi) - db(0)(\partial_b P_{B_1(b(0))}, \psi)$$

Replacing $d\tilde{\varepsilon}$ by its values, and applying that $(\Lambda Q, \psi) = 0$ together with $|\lambda(0) - 1| \lesssim b_0^2/|\log(b_0)|$, we get:

$$(d\tilde{\varepsilon}, \psi) = \|\psi\|_{L^2}^2 + O(b_0)$$

so that:

$$dps_1(0) = \|\psi\|_{L^2}^2 + O(b_0).$$

Computation of $\partial_s \varepsilon(0) + b_s(0) \partial_b P_{B_1(b(0))}$: From (A.15),

$$\partial_s \varepsilon(0) = -b_s(0) \partial_b P_{B_1(b(0))} - b(0) \Lambda u_0 + \lambda(0) (b_0 \Lambda P_{B_1(b_0)})_{\frac{1}{\lambda(0)}}$$

so that $(\partial_s \varepsilon(0) + b_s(0) \partial_b P_{B_1(b(0))}, \psi)$ is a smooth function of d_+ with derivative :

$$dps_2(0) = -db(0)(\Lambda u_0, \psi) + d\lambda \left(\left[(b_0 \Lambda P_{B_1(b_0)})_{\frac{1}{\lambda(0)}} + (b_0 \Lambda^2 P_{B_1(b_0)})_{\frac{1}{\lambda(0)}} \right], \psi \right) - b(0)(\Lambda \psi, \psi),$$

where, for the same orthogonality reason $(\Lambda Q, \psi) = 0$, we have:

$$(\Lambda u_0, \psi) = (\Lambda Q, \psi) + O(b_0) = O(b_0)$$

Consequently $dps_2(0) = O(b_0)$.

Conclusion: Finally, there holds

$$\kappa_+(0) = \frac{1}{2} \left[(\varepsilon(0), \psi) + \frac{1}{\sqrt{\zeta}} (\partial_s \varepsilon(0) + b_s(0) \partial_b P_{B_1(b(0))}, \psi) \right].$$

and $\kappa_+(0) = a_+$ reduces to a simple 1D equation $F(d_+) = a_+$ with F computed as combination of the above functions so that it is smooth in a neighborhood of 0. Moreover, there holds:

$$dF = \frac{1}{2} \left[dps_1(0) + \frac{1}{\sqrt{\zeta}} dps_2(0) \right] = \frac{\|\psi\|_{L^2}^2}{2} + O(b_0),$$

and (A.19) is proved. This concludes the proof of Lemma 3.3.

Appendix B. Coercivity estimates

The aim of this section is a proof of the coercivity properties of the quadratic form:

$$B(\eta, \eta) = (\mathcal{B}v, v) = \int_{\mathbb{R}^4} |\partial_r \eta|^2 + \int_{\mathbb{R}^4} W \eta^2$$

where

$$W(r) = 2V + \frac{3}{2} rV' = \frac{6}{(1+r^2/8)^2} - \frac{9}{4} \frac{r^2}{(1+r^2/8)^3}$$

We use the elementary method developed in [8]. The coercivity property of Lemma 4.7 is a consequence of the two following facts. First the index of B on $\dot{H}_r^1 = \{u \text{ radial with } \int |\nabla u|^2 + \int \frac{u^2}{r^2} < +\infty\}$ is at most 2. From standard Sturm Liouville oscillation theorems, see Theorem XIII.8 [33], this is equivalent to counting the number of zeroes of

$$\begin{cases} \mathcal{B}U = 0 & \text{on } (0, \infty), \\ U(0) = 1 \quad U'(0) = 0, \end{cases} \quad (\text{B.1})$$

and this can be analytically reduced to counting the number of zeroes of a Bessel function. Then we need to show that the orthogonality conditions $(\eta, \psi) = (\eta, \Phi) = 0$ are enough to treat the two negative directions. Arguing exactly as in [8], see also [13], this is equivalent to first invert the operator \mathcal{B} on \dot{H}_{rad}^1 , and then show that B restricted to $Span\{\mathcal{B}^{-1}\psi, \mathcal{B}^{-1}\Phi\}$ is definite negative, which is an elementary numerical check. We shall check these two facts below and refer to [8] for the proofs that this implies the claimed coercivity property. Note that the proofs in [8] are given for exponentially decaying functions and potentials, but one checks easily that the decay of the




FIGURE 1. Solution to (B.1) computed by MAPLE

potential $|W(r)| \sim \frac{1}{r^4}$ at infinity and $|\Phi(r)| \sim \frac{1}{r^4}$ are more than enough to have all proofs go through.

B.1. Computation of the index of B . We claim:

Lemma B.1 (Derivation of the index). *The index of \mathcal{B} on \dot{H}_r^1 is at most 2..*

Proof. First, we note that $W(r) \geq \hat{W}(r)$ where:

$$\hat{W}(r) = -\frac{3}{2} \frac{r^2}{(1 + r^2/8)^3}.$$

Hence, classical Sturm-Liouville theory ensures that U has less zeros than \hat{U} the unique solution to :

$$\begin{cases} -\frac{1}{r^3} \frac{d}{dr} \left[r^3 \frac{d}{dr} \hat{U} \right] + \hat{W} \hat{U} = 0 & \text{on } (0, \infty), \\ \hat{U}(0) = 1 \quad \hat{U}'(0) = 0, \end{cases} \quad (\text{B.2})$$

Second, we look for \hat{U} of the form:

$$\hat{U}(r) = \frac{2}{r^2} \bar{U}(r^2/2),$$

with \bar{U} a sufficiently smooth function. Denoting by s the new variable $r^2/2$, straightforward calculations yield that \bar{U} is a solution to :

$$\begin{cases} -\frac{d^2}{ds^2} \bar{U} + \bar{W} \bar{U} = 0 & \text{on } (0, \infty), \\ \bar{U}(0) = 0 \quad \bar{U}'(0) = 1, \end{cases} \quad (\text{B.3})$$

where:

$$\bar{W}(s) = -\frac{3}{2} \frac{1}{(1+s/4)^3}.$$

Setting then $\bar{U}(s) = \sqrt{1+s/4} \tilde{U}(1/\sqrt{1+s/4})$, we obtain that \bar{U} is a solution to (B.3) if and only if \tilde{U} is a solution to

$$\begin{cases} \tau^2 \frac{d^2}{d\tau^2} \tilde{U} + \tau \frac{d}{d\tau} \tilde{U} + (96\tau^2 - 1)\tilde{U} = 0 & \text{on } (0, 1), \\ \tilde{U}(1) = 0 & \tilde{U}'(1) = -8, \end{cases}$$

Hence, \tilde{U} is a combination of Bessel functions:

$$\tilde{U}(\tau) = C_1 J(1, 4\sqrt{6}\tau) + C_2 Y(1, 4\sqrt{6}\tau)$$

We compute (C_1, C_2) and draw the explicit combination with MAPLE. We obtain Figure 2. The computed solution \tilde{U} has two zeros on $(0, 1)$. Moreover, it diverges in 0 so that $\tilde{U}(\tau) \sim K/\tau$ close to 0 with $K \neq 0$. As a consequence

$$\hat{U}(r) \sim \frac{K}{4} \neq 0 \quad \text{when } r \rightarrow \infty,$$

and thus the index of $-\Delta + \hat{W}$ on \dot{H}_{rad}^1 is exactly two. Hence the index of \mathcal{B} is at most 2. This completes the proof of Lemma B.1. \square




FIGURE 2. Solution to (B.3) computed by MAPLE

B.2. Choice for the orthogonality conditions. We now invert \mathcal{B} . We first check numerically that the solution U does not vanish at infinity ie

$$\lim_{r \rightarrow +\infty} U(r) > 0,$$

see Figure 1.

Hence U is not a resonance -note that if U had been a resonance, we could have removed the resonance by diminishing a bit the potential and getting a potential with index 2 and no resonance-, and thus from standard ODE arguments, [8], there exists unique smooth solution in \dot{H}^1_{rad} of:

$$\begin{cases} \mathcal{B}U = -\frac{1}{r^3} \frac{d}{dr} \left[r^3 \frac{d}{dr} U \right] + WU = \psi & \text{on } (0, \infty), \\ U'(0) = 0, \end{cases} \quad \text{with } (1+r^2)U \in L^\infty \quad (\text{B.4})$$

and

$$\begin{cases} \mathcal{B}U = -\frac{1}{r^3} \frac{d}{dr} \left[r^3 \frac{d}{dr} U \right] + WU = \Phi & \text{on } (0, \infty), \\ U'(0) = 0, \end{cases} \quad \text{with } \left(1 + \frac{r^2}{\log r}\right)U \in L^\infty \quad (\text{B.5})$$

We denote $\mathcal{B}^{-1}\psi$ and $\mathcal{B}^{-1}\Phi$ the respective solutions to these systems. We recall the explicit formula

$$\Phi(r) = D\Lambda Q(r) = \frac{2 - 3r^2/4}{(1 + r^2/8)^3}.$$

In the remainder of this section we, check numerically that the restriction of B to $\text{Span}(\mathcal{B}^{-1}\psi, \mathcal{B}^{-1}\Phi)$ is definite negative, or equivalently:

Lemma B.2 (Numerical check of the orthogonality conditions). *The symmetric matrix*

$$\mathbb{B} = \begin{bmatrix} (\mathcal{B}^{-1}\psi, \psi) & (\mathcal{B}^{-1}\Phi, \psi) \\ (\mathcal{B}^{-1}\Phi, \psi) & (\mathcal{B}^{-1}\Phi, \Phi) \end{bmatrix}$$


satisfies:

$$(\mathcal{B}^{-1}\psi, \psi) < 0 \quad \text{and} \quad \det \mathbb{B} > 0, \quad (\text{B.6})$$


and is thus definite-negative.

Numerical proof of Lemma B.2 We use standard MATLAB routines for the computation of solutions to (B.4,B.5). We note that we only fixed the initial value for $U'(0)$. The value $U(0)$ is left open in order to achieve the expected decay at infinity which characterizes the inverse. In order to obtain $\mathcal{B}^{-1}\psi$, we first compute ψ . We obtain that the corresponding eigenvalue is approximatively $l = -0.5860808922$. Because ψ decays exponentially, we only need to obtain an approximation on a short time-range. We computed our solutions until $T_{\psi, max} = 30$. We emphasize here that we use an explicit scheme. As a drawback, the accumulation of errors tends to make the numerical solution to become negative when the exact solution is exponential small. Hence, our scheme becomes unstable after time $\tilde{T}_{\psi, max} = 18$. Nevertheless, we extend our numerical solution with 0 after this time. This induces an exponentially small error. The pictures in Figure 3 illustrate this computation. On the left-hand side is drawn the obtained solution. On the right-hand side, we draw $\psi_{test}(r) = \psi(r)\exp(\sqrt{-l}r)$. We observe here that our solution enters the exponential asymptotic regime before the instability comes into play.

The solution $\mathcal{B}^{-1}\psi$ is computed with the extension of ψ . Straightforward ode analysis shows that the unique solution decaying fast at infinity behaves like $1/r^2$ asymptotically. The choice of $U(0)$ is made with respect to this criterion. Figure 4 illustrates that we obtained a solution with the suitable decay. As previously, on the left-hand side is a picture of the numerical solution. On the right-hand side we plot $\mathcal{B}^{-1}\psi_{test}(r) = r^2\mathcal{B}^{-1}\psi(r)$. In the latter computations, this solution is involved in scalar

FIGURE 3. Numerical simulations for ψ

products with ψ . Hence even if drawn until $T_{max} = 300$, we only need a precise computation of this solution until $T_{\mathcal{B}^{-1}\psi, max} = 18$.

FIGURE 4. Numerical simulations for $\mathcal{B}^{-1}\psi$


The last solution $\mathcal{B}^{-1}\Phi$ is computed with the same method. In this second case, the expected decay of the solution is $\log(r)/r^2$. Figure 5 illustrates that we obtained a solution with the suitable decay. The picture on the right-hand side restricts to the time-interval $r = 0..100$ because this is the significant region. In the latter computations, this solution is involved in integrals which converge slowly. Hence, we compute this solution until $T_{\mathcal{B}^{-1}\Phi, max} = 1000$.

We now compute numerically the entries of the matrix \mathbb{B} . We first compute $(\mathcal{B}^{-1}\Phi, \psi) = (\mathcal{B}^{-1}\psi, \Phi)$. The exponential decay of ψ implies that we need to compute the first integral $(\mathcal{B}^{-1}\Phi, \psi)$ on a shorter time-interval. Hence, we prefer this computation to the second one. We compute the L^2 -scalar products with a standard trapezoidal method. Changing the time-interval and the time-step, the computations are stable up to an error of 10^{-2} . We get the following approximations for the integrals involving ψ .

$$(\mathcal{B}^{-1}\psi, \psi) = -4.63 \pm 10^{-2} \quad (\mathcal{B}^{-1}\Phi, \psi) = 32.65 \pm 10^{-2}.$$

The last integral is a more involved computation. Indeed, standard real analysis implies that there holds :

$$I(M) := \int_0^M \mathcal{B}^{-1}\Phi Q(r)\Phi(r)r^3 dr = (\mathcal{B}^{-1}\Phi, \Phi) + err(M)$$

FIGURE 5. Numerical simulations for $\mathcal{B}^{-1}\Phi$

with a remainder satisfying $err(M) = (K + o(1)) \ln(M)/M^2$ for some constant K . This remainder going slowly to 0, we see numerically that our computations has not converged even when integrating until $T_{\mathcal{B}^{-1}\Phi, max} = 1000$ (see Figure 6, red crosses). In order to improve the rate of convergence we compute an approximation of coefficient K and substract the estimated error term of our computations. This yields Figure 6, blue circles. On this second computation we obtain a very good rate of convergence. Hence, we provide the approximation

$$(\mathcal{B}^{-1}\Phi, \Phi) = -574.25 \pm 10^{-2}$$

Hence

$$\det(\mathbb{B}) = 1591 \pm 10$$

which concludes the numerical proof of Lemma B.2.

Appendix C. Some linear estimates

We start by recalling some obvious integration-by-part results :

Lemma C.1. *For any $N \geq 3$, there exists a constant C for which there holds, for any $v \in H_{rad}^1(\mathbb{R}^N)$*

$$\left[\int_{\mathbb{R}^N} \frac{|v(y)|^2}{|y|^2} \right]^{\frac{1}{2}} + \sup_{y \in \mathbb{R}^N} \left(|y|^{\frac{N-2}{2}} |v(y)| \right) \leq C \left[\int_{\mathbb{R}^n} |\nabla v(y)|^2 \right]^{\frac{1}{2}}. \quad (\text{C.1})$$

Looking for control on further derivatives, we prove:

Lemma C.2 (Hardy inequalities). *Let $N = 4$. Then $\forall R > 2$, $\forall v \in H_{rad}^2(\mathbb{R}^N)$, there holds the following controls:*

$$\int \frac{|\partial_y v|^2}{y^2} \lesssim \int (\Delta v)^2, \quad (\text{C.2})$$

$$\int_{y \leq R} \frac{|v|^2}{y^4 (1 + |\log y|)^2} \lesssim \int_{y \leq R} \frac{|\partial_y v|^2}{y^2} + \int_{y \leq 2} |v|^2. \quad (\text{C.3})$$

$$\int_{R \leq y \leq 2R} \frac{|v|^2}{y^4} \lesssim \log R \int_{y \leq R} \frac{|\partial_y v|^2}{y^2} + \int_{y \leq 2} |v|^2. \quad (\text{C.4})$$




FIGURE 6. Computations for $(\mathcal{B}^{-1}\Phi, \Phi)$

Proof. Let v smooth. (C.2) follows from the explicit formula after integration by parts

$$\int (\Delta v)^2 = \int (\partial_{yy}v + \frac{N-1}{y}\partial_y v)^2 = \int (\partial_{yy}v)^2 + (N-1) \int \frac{|\partial_y v|^2}{y^2}.$$

To prove (C.3), let

$$a \in [1, 2] \text{ such that } |v(a)|^2 \leq \int_{1 \leq y \leq 2} |v|^2. \quad (\text{C.5})$$

Let $f(y) = -\frac{e_y}{y^3(1+\log(y))}$ so that $\nabla \cdot f = \frac{1}{y^4(1+|\log y|^2)}$, and integrate by parts to get:

$$\begin{aligned} \int_{a \leq y \leq R} \frac{|v|^2}{y^4(1+\log y)^2} &= \int_{a \leq y \leq R} |v|^2 \nabla \cdot f \\ &= - \left[\frac{|v|^2}{1+\log(y)} \right]_a^R + 2 \int_{y \leq R} \frac{v \partial_y v}{y^3(1+\log y)} \\ &\lesssim |v(a)|^2 + \left(\int_{y \leq R} \frac{|v|^2}{y^4(1+|\log y|^2)} \right)^{\frac{1}{2}} \left(\int_{y \leq R} \frac{|\partial_y v|^2}{y^2} \right)^{\frac{1}{2}}. \end{aligned} \quad (\text{C.6})$$

similarly, using $\tilde{f}(y) = \frac{e_y}{y^3(1-\log(y))}$, we get:

$$\begin{aligned} \int_{\varepsilon \leq y \leq a} \frac{|v|^2}{y^4(1-\log y)^2} &= \int_{a \leq y \leq R} |v|^2 \nabla \cdot \tilde{f} \\ &= \left[\frac{|v|^2}{1-\log(y)} \right]_{\varepsilon}^a + 2 \int_{y \leq a} v \partial_y v \frac{1}{y^3(1-\log y)} \\ &\lesssim |v(a)|^2 + \left(\int_{y \leq R} \frac{|v|^2}{y^4(1+|\log y|)^2} \right)^{\frac{1}{2}} \left(\int_{y \leq R} \frac{|\partial_y v|^2}{y^2} \right)^{\frac{1}{2}}. \end{aligned} \quad (\text{C.7})$$

(C.5), (C.6) and (C.7) now yield (C.3). The last inequality (C.4) is a straightforward variant of [31, Lemma B.1, (B.4)] and is left to the reader. \square

Lemma C.3 (Coercitivity estimates with H). *Let ψ be the first eigenvector of H . Then there exists $c > 0$ and $M_0 \geq 1$ such that for $M \geq M_0$, there exists $\delta(M) > 0$ such that given $u \in H_{rad}^1(\mathbb{R}^N)$, there holds*

$$(Hu, u) \geq c \int (\partial_y u)^2 - \frac{1}{c} [(u, \psi)^2 + (u, \chi_M \Phi)^2] \quad (\text{C.8})$$

$$\int (Hu)^2 \geq \delta(M) \left[\int \frac{(\partial_y u)^2}{y^2} + \int \frac{u^2}{y^4(1+|\log y|)^2} \right] - \frac{1}{\delta(M)} (u, \chi_M \Phi)^2. \quad (\text{C.9})$$

Proof. (C.8) is a standard consequence of the coercitivity of the linearized energy which admits exactly ψ as bound state and ΛQ as resonance at the origin, the good enough localization of Φ (2.1) and the nondegeneracy (2.2). The detailed proof is left to the reader.

To prove (C.9), we first observe the key subcoercivity property:

$$\begin{aligned} \int (Hu)^2 &= \int (\Delta u + Vu)^2 = \int (\Delta u)^2 - 2 \int V(\partial_y u)^2 + \int (\Delta V + V^2)u^2 \\ &\geq c \left[\int (\Delta u)^2 + \int \frac{u^2}{1+y^6} \right] - \frac{1}{c} \left[\int \frac{(\partial_y u)^2}{1+y^4} + \int \frac{u^2}{1+y^8} \right]. \end{aligned} \quad (\text{C.10})$$

where we used the asymptotic value

$$V(y) = \frac{N(N+2)(N-2)}{y^4} \left[1 + O\left(\frac{1}{y^2}\right) \right] \quad \text{as } y \rightarrow +\infty.$$

(C.9) now follows by contradiction. Let $M > 0$ fixed and consider a sequence u_n such that

$$\int \frac{(\partial_y u_n)^2}{y^2} + \int \frac{u_n^2}{y^4(1+|\log y|)^2} = 1 \quad (\text{C.11})$$

and

$$\int (Hu_n)^2 \leq \frac{1}{n}, \quad (u_n, \chi_M \Phi) = 0. \quad (\text{C.12})$$

Then by semicontinuity of the norm, u_n weakly converges on a subsequence to $u_\infty \in H_{loc}^1$ solution to $Hu_\infty = 0$. u_∞ is smooth away from the origin and hence the explicit integration of the ODE and the regularity assumption at the origin $u_\infty \in H_{loc}^1$ implies

$$u_\infty = \alpha \Lambda Q.$$

On the one hand, the uniform bound (C.11) together with the local compactness of Sobolev embeddings ensure up to a subsequence:

$$\int \frac{(\partial_y u_n)^2}{1+y^4} + \int \frac{|u_n|^2}{1+y^8} \rightarrow \int \frac{(\partial_y u_\infty)^2}{1+y^4} + \int \frac{|u_\infty|^2}{1+y^8} \quad \text{and} \quad (u_n, \chi_M \Phi) \rightarrow (u_\infty, \chi_M \Phi)$$

thanks to the χ_M localization. We thus conclude that

$$\alpha(\Lambda Q, \chi_M \Phi) = (u_\infty, \chi_M \Phi) = 0 \quad \text{and thus} \quad \alpha = 0.$$

On the other hand, the subcoercivity property (C.10), the Hardy control (C.2), (C.3) and (C.11), (C.12) ensure

$$\int \frac{(\partial_y u_n)^2}{1+y^4} + \int \frac{u_n^2}{1+y^8} \geq C > 0$$

from which

$$\alpha^2 \left[\int \frac{(\partial_y \Lambda Q)^2}{1+y^4} + \int \frac{|\Lambda Q|^2}{1+y^8} \right] = \int \frac{(\partial_y u_\infty)^2}{1+y^4} + \int \frac{|u_\infty|^2}{1+y^8} \geq C > 0 \quad \text{and thus} \quad \alpha \neq 0.$$

A contradiction follows. This concludes the proof of (C.9) and Lemma C.3. \square

Straightforward computations show that the coercitivity estimates with H can be adapted to any of the operator H_λ yielding, for any $\lambda > 0$ and $u \in H_{rad}^1(\mathbb{R}^N)$,

$$(H_\lambda u, u) \geq c \int (\partial_y u)^2 - \frac{1}{c\lambda^4} [(u, (\psi)_\lambda)^2 + (u, (\chi_M \Phi)_\lambda)^2] \quad (\text{C.13})$$

for the same c and $\delta(M)$ as in Lemma C.3.

References

- [1] Blow up for nonlinear hyperbolic equations, volume 17 of Progress in Nonlinear Differential Equations and their Applications, Birkhäuser Boston Inc., Boston, MA, 1995.
- [2] van den Berg, J. B.; Hulshof, J.; King, J.R., Formal asymptotics of bubbling in the harmonic map heat flow, SIAM J. Appl. Math. 63 (2003), no. 5, 1682–1717.
- [3] Bizoń, P.; Chmaj, T.; Tabor, Z., Formation of singularities for equivariant $(2+1)$ -dimensional wave maps into the 2-sphere, Nonlinearity 14 (2001), no. 5, 1041–1053.
- [4] Christodoulou, D.; Tahvildar-Zadeh, A. S., On the regularity of spherically symmetric wave maps, Comm. Pure Appl. Math. 46 (1993), no. 7, 1041–1091.
- [5] Cote, R.; Martel, Y.; Merle, F., Construction of multisolitons solutions for the L^2 -supercritical gKdV and NLS equations, arXiv:0905.0470 (2010).
- [6] Duyckaerts, T.; Kenig, C.E.; Merle, F., Universality of blow up profile for small radial type II blow up solutions of energy critical wave equation, arXiv:0910.2594 (2009)
- [7] Duyckaerts, T.; Kenig, C.E.; Merle, F., Universality of blow up profile for small type II blow up solutions of energy critical wave equation: the non radial case, arXiv:1003.0625 (2010)
- [8] Fibich, G.; Merle, F.; Raphael, P., Numerical proof of a spectral property related to the singularity formation for the L^2 critical nonlinear Schrödinger equation, Phys. D 220 (2006), no. 1, 1–13.
- [9] Grillakis, M.G., Regularity and asymptotic behaviour of the wave equation with a critical nonlinearity. Ann. of Math. (2) 132 (1990), no. 3, 485–509.
- [10] Jörgens, K., Das Anfangswertproblem im Grossen für eine Klasse nichtlinearer Wellengleichungen. (German) Math. Z. 77 1961 295–308.
- [11] Karageorgis, P.; Strauss, W. A., Instability of steady states for nonlinear wave and heat equations, J. Differential Equations 241 (2007), no. 1, 184–205.
- [12] Kavian, O.; Weissler, F.B., Finite energy self-similar solutions of a nonlinear wave equation, Comm. Partial Differential Equations 15 (1990), no. 10, 1381–1420.
- [13] Marzuola, J.; Simpson, G., Spectral Analysis for matrix Hamiltonian operators, arXiv:1003.2474 (2010)
- [14] Kenig, C.E.; Merle, F., Global well-posedness, scattering and blow-up for the energy-critical focusing non-linear wave equation. Acta Math. 201 (2008), no. 2, 147–212.
- [15] Krieger, J.; Schlag, W., On the focusing critical semi-linear wave equation, Amer. J. Math. 129 (2007), no. 3, 843–913.MK
- [16] Krieger, J.; Schlag, W., Non-generic blow-up solutions for the critical focusing NLS in 1-D. J. Eur. Math. Soc. (JEMS) 11 (2009), no. 1, 1–125.
- [17] Krieger, J.; Schlag, W.; Tataru, D., Renormalization and blow up for charge one equivariant critical wave maps, Invent. Math. 171 (2008), no. 3, 543–615.

- [18] Krieger, J.; Schlag, W.; Tataru, D., Slow blow-up solutions for the $H^1(\mathbb{R}^3)$ critical focusing semilinear wave equation, *Duke Math. J.* 147 (2009), no. 1, 1–53.
- [19] Krieger, J.; Schlag, W.; Tataru, D., Renormalization and blow up for the critical Yang-Mills problem, *Adv. Math.* 221 (2009), no. 5, 1445–1521.
- [20] Levine, H.A., Instability and nonexistence of global solutions to nonlinear wave equations of the form $Pu_{tt} = -Au + F(u)$, *Trans. Amer. Math. Soc.* 192 (1974), 1–21.
- [21] Martel, Y., Asymptotic N -soliton-like solutions of the subcritical and critical generalized Korteweg-de Vries equations, *Amer. J. Math.* 127 (2005), no. 5, 1103–1140.
- [22] Martel, Y.; Merle, F., Stability of blow-up profile and lower bounds for blow-up rate for the critical generalized KdV equation, *Ann. of Math. (2)* 155 (2002), no. 1, 235–280.
- [23] Merle, F.; Raphaël, P., Blow up dynamic and upper bound on the blow up rate for critical nonlinear Schrödinger equation, *Ann. Math.* 161 (2005), no. 1, 157–222.
- [24] Merle, F.; Raphaël, P., Sharp upper bound on the blow up rate for critical nonlinear Schrödinger equation, *Geom. Funct. Anal.* 13 (2003), 591–642.
- [25] Merle, F.; Raphaël, P., On universality of blow up profile for L^2 critical nonlinear Schrödinger equation, *Invent. Math.* 156, 565–672 (2004).
- [26] Merle, F.; Raphaël, P., Sharp lower bound on the blow up rate for critical nonlinear Schrödinger equation, *J. Amer. Math. Soc.* 19 (2006), no. 1, 37–90.
- [27] Merle, F.; Raphaël, P., Profiles and quantization of the blow up mass for critical nonlinear Schrödinger equation, *Comm. Math. Phys.* 253 (2005), no. 3, 675–704.
- [28] Merle, F.; Zaag, H., Determination of the blow-up rate for the semilinear wave equation, *Amer. J. Math.* 125 (2003), no. 5, 1147–1164.
- [29] Merle, F.; Zaag, H., Openness of the set of non-characteristic points and regularity of the blow-up curve for the 1 D semilinear wave equation, *Comm. Math. Phys.* 282 (2008), no. 1, 55–86.
- [30] Raphaël, P., Stability of the log-log bound for blow up solution to the critical non linear Schrödinger equation, *Math. Ann.* 331 (2005), no. 3, 577–609.
- [31] Raphaël, P.; Rodnianski, I., Stable blow up dynamics for the critical corotational wave maps and equivariant Yang Mills problems, submitted.
- [32] Raphaël, P.; Szeftel, J. Existence and uniqueness of minimal blow up solutions to an inhomogeneous mass critical NLS, arXiv:1001.1627 (2009).
- [33] Reed, M.; Simon, B., *Methods of modern mathematical physics, vol I-IV*, Academic Press, New York, 1972-1979.
- [34] Rodnianski, I.; Sterbenz, J., On the formation of singularities in the critical $O(3)$ σ -model, to appear *Ann. Math.*
- [35] Sogge, C.D., *Lectures on nonlinear wave equations*, Monographds in Analysis, II, International Press, Boston, MA, 1995.
- [36] Shatah, J.; Tahvildar-Zadeh, A. S., On the Cauchy problem for equivariant wave maps, *Comm. Pure Appl. Math.* 47 (1994), no. 5, 719–754.
- [37] Strauss, W. A., *Nonlinear wave equations*, CBMS Regional Conference Series in Mathematics, 73, AMS, Providence, RI, 1989.
- [38] Struwe, M., Globally regular solutions to the u^5 Klein-Gordon equation. *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)* 15 (1988), no. 3, 495–513.
- [39] Struwe, M., Equivariant wave maps in two space dimensions. Dedicated to the memory of Jürgen K. Moser, *Comm. Pure Appl. Math.* 56 (2003), no. 7, 815–823

CEREMADE, UNIVERSITÉ PARIS DAUPHINE, FRANCE
E-mail address: hillairet@ceremade.dauphine.fr

INSTITUT DE MATHÉMATIQUES DE TOULOUSE, UNIVERSITÉ TOULOUSE III, FRANCE
E-mail address: pierre.raphael@math.univ-toulouse.fr