



Non parametric estimation of the structural expectation of a stochastic increasing function

Jean-François Dupuy, Jean-Michel Loubes, Elie Maza

► To cite this version:

Jean-François Dupuy, Jean-Michel Loubes, Elie Maza. Non parametric estimation of the structural expectation of a stochastic increasing function. 2008. <hal-00347829>

HAL Id: hal-00347829

<https://hal.archives-ouvertes.fr/hal-00347829>

Submitted on 16 Dec 2008

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Non parametric estimation of the structural expectation of a stochastic increasing function

J-F. DUPUY & J-M. LOUBES & E. MAZA

Abstract

This article introduces a non parametric warping model for functional data. When the outcome of an experiment is a sample of curves, data can be seen as realizations of a stochastic process, which takes into account the small variations between the different observed curves. The aim of this work is to define a mean pattern which represents the main behaviour of the set of all the realizations. So we define the structural expectation of the underlying stochastic function. Then we provide empirical estimators of this structural expectation and of each individual warping function. Consistency and asymptotic normality for such estimators are proved.

Keywords: Functional data analysis; Non parametric warping model; Structural expectation; Curve registration; .

Subject Class. MSC-2000: 62G05, 62G20 .

1 Statistical model for functional phase variations

Functional data analysis deals with the analysis of experiments where one or several quantities are recorded during a time period for different individuals, resulting in a sample of observed curves. One of the main difficulties is given by the fact that curves usually not only present amplitude variability (a variation in the y -axis) but also time or phase variability (a variation in the x -axis). Hence the classical cross-sectional mean does not make sense and the definition of an appropriate population mean is even not obvious. Giving a sense to the common behaviour of a sample of curves, and finding a meaningful mean curve in this context is thus an important issue, called curve registration, or time warping problem, which first appeared in the engineering literature in [11].

Several methods have been proposed over the years to estimate the mean pattern of a sample of curves. A popular method consists in i) first aligning the curves to a given template by warping the time axis, ii) then taking the mean of all the dewarped curves. Such methods are increasingly common in statistics, see [9] for a review. A landmark registration methodology is proposed by [5] and further developed by [1]. A non parametric method is investigated in [8] and in [6], using local regressions. Dynamic time warping methodology is developed by [14]. An alternative approach is provided in [4], where semi-parametric estimation of shifted curves is studied. But these methods imply choosing a starting curve as a fixed point for the alignment process. This initial choice may either bias the estimation procedure, or imply strong and restrictive identifiability conditions.

In this work, we consider a second point of view. We define an archetype representing the common behaviour of the sample curves directly from the data, without stressing a particular curve. Such a method has the advantage of not assuming any technical restrictions on the data and so, enables to handle a large variety of cases. However, the registration procedure and the common pattern have to be clearly defined.

We observe $i = 1, \dots, m$ curves $f_i : [a, b] \rightarrow \mathbb{R}$ at equidistant discrete times $t_{ij} \in [a, b]$, $j = 1, \dots, n$. So the data can be written as

$$Y_{ij} = f_i(t_{ij}), \quad i = 1, \dots, m, \quad j = 1, \dots, n. \quad (1)$$

So, the registration problem aims at finding a mean pattern f and warping functions h_i which align the observed curves, i.e such that $\forall i = 1, \dots, m, f = f_i \circ h_i$. Hence each curve is obtained by warping the original curve f using the warping functions h_i .

Defining the registration operator is a difficult task. In this paper we propose a random warping procedure which takes into account the variability of the deformation as a random effect. So, we assume that there exists a random process H such that the data are i.i.d realizations of this process. Let H be this warping stochastic process defined as

$$\begin{aligned} H : \Omega &\rightarrow \mathcal{C}([a, b]) \\ w &\mapsto H(w, \cdot), \end{aligned} \quad (2)$$

where $(\Omega, \mathcal{A}, \mathbf{P})$ is an unknown probability space, and $(\mathcal{C}([a, b]), \|\cdot\|_\infty, \mathcal{B})$ is the set of all real continuous functions defined on the interval $[a, b]$, equipped with the uniform norm and with its Borel algebra. Consider h_1, \dots, h_m i.i.d realizations of the process $H(t)$. h_i warps a mean pattern f onto the i -th observation curve f_i . Hence, model (1) can be modeled by

$$Y_{ij} = f_i(t_{ij}) = f \circ h_i^{-1}(t_{ij}). \quad (3)$$

For sake of simplicity, we will write $f_i(t) = f \circ h_i^{-1}(t)$, for all $i \in \{1, \dots, m\}$. We point out that h_i^{-1} is well defined since the warping processes are assumed to be continuous increasing functions.

Under general assumptions (2), model (3) is not identifiable. More precisely, the unknown function f and the unknown warping process H can not be estimated. Indeed, if $\tilde{h} : [a, b] \rightarrow [a, b]$ is an increasing continuous function, with $\tilde{h}(a) = a$ and $\tilde{h}(b) = b$, then, for all $i \in \{1, \dots, m\}$ and all $j \in \{1, \dots, n\}$, we have that $Y_{ij} = f \circ \tilde{h}^{-1} \circ \tilde{h} \circ h_i^{-1}(\cdot, t_{ij})$. Hence, the function $f \circ \tilde{h}^{-1}$, associated with the warping process $H \circ \tilde{h}^{-1}$, is also a solution of model (3).

The aim of this paper is to build a new kind of pattern which represents the common functional feature of the data but still that can be estimated together with the warping procedure. For this, let $\phi(\cdot)$ be the expectation of the warping process and define the *structural expectation* f_{ES} as

$$f_{ES} := f \circ \phi^{-1}.$$

The structural expectation is obviously not the function f , but the function f composed with ϕ^{-1} , the inverse of the expectation of H . Hence it can be seen as the mean warping of the function f by the stochastic process H . In this article, we aim at studying the properties of the structural expectation, and finding an estimator of the structural expectation and of the warping paths h_i .

The article is organized as follows. In Section 1, we introduce a warping functional model with a stochastic phase variation, and we introduce the structural expectation. In Section 2, we define empirical estimators of the structural expectation and of the individual warping functions. Asymptotic properties of these estimators are investigated. Proofs are postponed to Section A. Section 3 investigates some extensions of the proposed methodology, in particular to the case of noisy non increasing functions. The results of a simulation study are reported in Section 4. There, we also apply the proposed estimators to a real data set.

2 Theoretical study of a particular case: warping of strictly increasing functions

First, consider the case where f is a strictly increasing. Hence the inverse function f^{-1} exists and is also strictly increasing. Moreover, a phase warping of function f (i.e. on x -axis) corresponds to an amplitude warping of function f^{-1} (i.e. on y -axis). We propose estimators of both the inverse of the structural expectation $f_{ES}^{-1} = \phi \circ f^{-1}$, and also of each individual warping function $\phi \circ h_i^{-1}$, for all $i \in \{1, \dots, m\}$, and finally an of the structural expectation $f_{ES} = f \circ \phi^{-1}$.

Note that all the asymptotic results are taken with respect to m and n , so we recall that $u(m, n) \xrightarrow{m, n \rightarrow \infty} c$ if, and only if, we have

$$\forall \epsilon > 0, \exists(m_0, n_0) \in \mathbb{N}^2, \forall(m, n) \in \mathbb{N}^2, m > m_0 \text{ and } n > n_0 \Rightarrow |u(m, n) - c| < \epsilon.$$

We will assume the following conditions on the warping process in order to define a good registration procedure.

The warping process does not change the timeline (not time inversion) and leaves fixed the two extreme points, so for almost all $w \in \Omega$, assume that

- i) $H(w, \cdot)$ is an increasing function,
- ii) $H(w, a) = a$ and $H(w, b) = b$.

The following proposition introduces respectively the expectation, the second order moment and the covariance function of H .

Proposition 2.1. *Under assumption (2), the expectation $\phi(\cdot)$, the second order moment $\gamma(\cdot)$ and the covariance function $\mathfrak{r}(\cdot, \cdot)$ of the stochastic process H are well defined. ϕ and γ are also continuous increasing functions. Moreover, we have $\phi(a) = a$, $\phi(b) = b$, $\gamma(a) = a^2$ and $\gamma(b) = b^2$. As a consequence, we have that $\text{var}H(\cdot, a) = \text{var}H(\cdot, b) = 0$.*

Proof. The process H is bounded and increasing. As a consequence, ϕ and γ exist. Moreover, H is a continuous increasing process, which leads to continuous and increasing first and second order moments. \square

In order to prove asymptotic results, the following technical assumptions on the warping process H and on the function f are needed:

Assumptions

1. There exists a constant $C_1 > 0$ such that, for all $(s, t) \in [f(a), f(b)]^2$, we have

$$\mathbf{E} |H(s) - \mathbf{E}H(s) - (H(t) - \mathbf{E}H(t))|^2 \leq C_1 |s - t|^2. \quad (4)$$

2. There exists a constant $C_2 > 0$ such that, for all $(s, t) \in [f(a), f(b)]^2$, we have

$$|f^{-1}(s) - f^{-1}(t)|^2 \leq C_2 |s - t|^2. \quad (5)$$

3. There exists a constant $C_3 > 0$ such that, for all $\omega \in \Omega$, for all $(s, t) \in [a, b]^2$, we have

$$|H^{-1}(\omega, s) - H^{-1}(\omega, t)|^2 \leq C_3 |s - t|^2. \quad (6)$$

2.1 Estimator of the structural expectation f_{ES}

Since $f_i = f \circ h_i^{-1}$ we have $f_i^{-1} = h_i \circ f^{-1}$. Hence $\mathbf{E}(f_i) = (\mathbf{E}(H)) \circ f^{-1}$. Hence it seems natural to consider the mean of functions $(f_i^{-1})_{i \in \{1, \dots, m\}}$ in order to estimate the inverse of the structural expectation. For all $y \in [f(a), f(b)]$, and for all $i \in \{1, \dots, m\}$, define

$$j_i(y) = \arg \min_{j \in \{1, \dots, n\}} |Y_{ij} - y| \quad \text{and} \quad T_i(y) := t_{ij_i(y)}. \quad (7)$$

Then, the empirical estimator of the inverse of the structural expectation is defined by

$$\widehat{f_{ES}^{-1}}(y) = \frac{1}{m} \sum_{i=1}^m T_i(y). \quad (8)$$

The following theorem provides consistency and asymptotic normality of estimator (8).

Theorem 2.1 (Consistency of the inverse of the structural expectation). Under Assumption (2),

$$\left\| \widehat{f_{ES}^{-1}} - f_{ES}^{-1} \right\|_{\infty} \xrightarrow[m, n \rightarrow \infty]{as} 0.$$

Moreover, let $n = m^{\frac{1}{2} + \alpha}$ with $\alpha > 0$, and assume Conditions (4) and (5). Then,

$$\sqrt{m} \left(\widehat{f_{ES}^{-1}} - f_{ES}^{-1} \right) \xrightarrow[m \rightarrow \infty]{\mathbf{D}} G,$$

where G is a centered Gaussian process with covariance given by: for all $(s, t) \in [f(a), f(b)]^2$,

$$\text{cov}(G(s), G(t)) = r(f^{-1}(s), f^{-1}(t)).$$

From (8) and (7), $\widehat{f_{ES}^{-1}}$ is an increasing step function with jumps occurring at say, $K(m, n)$ points $v_1, \dots, v_{K(m, n)}$ in $[f(a), f(b)]$, such that $f(a) = v_0 < v_1 < \dots < v_{K(m, n)} < v_{K(m, n)+1} = f(b)$. Hence for all $y \in [f(a), f(b)] \setminus (v_k)_{k \in \mathcal{K}}$ ($\mathcal{K} = \{0, \dots, K(m, n) + 1\}$), $\widehat{f_{ES}^{-1}}(y)$ can be expressed as

$$\widehat{f_{ES}^{-1}}(y) = \sum_{k=0}^{K(m, n)} u_k \mathbf{1}_{(v_k, v_{k+1})}(y)$$

with $a = u_0 < u_1 < \dots < u_{K(m, n)-1} < u_{K(m, n)} = b$. A natural estimator of the structural expectation f_{ES} is then obtained by linear interpolation between the points (u_k, v_k) . For all $t \in [a, b]$, let define

$$\widehat{f_{ES}}(t) = \sum_{k=0}^{K(m, n)-1} \left(v_k + \frac{v_{k+1} - v_k}{u_{k+1} - u_k} (t - u_k) \right) \mathbf{1}_{[u_k, u_{k+1})}(t) + v_{K(m, n)} \mathbf{1}_{\{b\}}(t).$$

Note that by construction, this estimator is strictly increasing and continuous on $[a, b]$. The following theorem states its consistency.

Theorem 2.2 (Consistency of the estimator of the Structural Expectation). Under Assumption (2), we have

$$\left\| \widehat{f_{ES}} - f_{ES} \right\|_{\infty} \xrightarrow[m, n \rightarrow \infty]{as} 0.$$

Obtaining confidence bands for f_{ES}^{-1} is useful for describing and visualizing the uncertainty in the estimate of f_{ES}^{-1} . This requires finding an estimator of $\text{var}(G(y)) = \gamma \circ f^{-1}(y) - \{f_{ES}^{-1}(y)\}^2$, $y \in [f(a), f(b)]$.

Lemma 2.3. Let $y \in [f(a), f(b)]$. Let $\widehat{\gamma \circ f^{-1}}(y) = \frac{1}{m} \sum_{i=1}^m T_i^2(y)$, with $T_i(\cdot)$ defined as in (7). Then,

$$\widehat{\gamma \circ f^{-1}}(y) - \left\{ \widehat{f_{ES}^{-1}}(y) \right\}^2 \xrightarrow[m, n \rightarrow \infty]{as} \text{var}(G(y)).$$

Proof of this lemma is given in Section 4.2. Combining this lemma with the asymptotic normality result stated by Theorem 2.1 yields a pointwise asymptotic confidence band for f_{ES}^{-1} .

Corollary 2.4. An asymptotic $(1 - \alpha)$ -level pointwise confidence band for f_{ES}^{-1} is given by

$$\left[\widehat{f_{ES}^{-1}}(y) - u_{1-\frac{\alpha}{2}} \sqrt{\frac{\widehat{\text{var}}(G(y))}{m}}, \widehat{f_{ES}^{-1}}(y) + u_{1-\frac{\alpha}{2}} \sqrt{\frac{\widehat{\text{var}}(G(y))}{m}} \right],$$

where $\widehat{\text{var}}(G(y)) = \widehat{\gamma \circ f^{-1}}(y) - \left\{ \widehat{f_{ES}^{-1}}(y) \right\}^2$ and $u_{1-\frac{\alpha}{2}}$ is the quantile of order $1 - \frac{\alpha}{2}$ of the standard normal distribution.

Note that the construction of a simultaneous asymptotic confidence band for f_{ES}^{-1} would require determination of the distribution of $\sup_{f(a) \leq y \leq f(b)} |G(y)|$. This, however, falls beyond the scope of this paper.

2.2 Estimator of an individual warping function

In a warping framework, it is necessary to estimate the mean pattern but also the individual warping functions, i.e. $(h_i^{-1})_{i \in \{1, \dots, m\}}$. As previously, we can not directly estimate the functions h_i^{-1} only the functions $\phi \circ h_i^{-1}$.

Let $i_0 \in \{1, \dots, m\}$. Now we want to compute for all $i \neq i_0$, $T_i^*(t) = f_i^{-1} \circ f_{i_0}(t)$. For this, define

$$j_0(t) = \arg \min_{j \in \{1, \dots, n\}} |t_{i_0 j} - t|. \quad (9)$$

This point is the observation time for the i_0 curve which is the closest to t . Note that the index $j_0(t)$ depends on i_0 but, for sake of simplicity we drop this index in the notations. Then $\forall t \in [a, b]$, and $\forall i \in \{1, \dots, m\} \setminus i_0$, compute

$$T_i(t) = \arg \min_{t_j \in \{t_{i_0 1}, \dots, t_{i_0 n}\}} |Y_{ij} - Y_{i_0 j_0(t)}|, \quad (10)$$

an estimate of T_i^* . Then for a fixed i_0 , noting that $T_i^* = h_i \circ h_{i_0}^{-1}$, we can see that an empirical estimator of each individual warping function $\phi \circ h_{i_0}^{-1}$ is given by

$$\widehat{\phi \circ h_{i_0}^{-1}}(t) := \frac{1}{m-1} \sum_{\substack{i=1 \\ i \neq i_0}}^m T_i(t). \quad (11)$$

The following theorem asserts consistency and asymptotic normality of this estimator.

Theorem 2.5. Under assumption (2),

$$\left\| \widehat{\phi \circ h_{i_0}^{-1}} - \phi \circ h_{i_0}^{-1} \right\|_{\infty} \xrightarrow[m, n \rightarrow \infty]{a.s.} 0.$$

Let $n = m^{\frac{1}{2} + \alpha}$ (with $\alpha > 0$) and assume that (4) and (6) hold. Then $\sqrt{m}(\widehat{\phi \circ h_{i_0}^{-1}} - \phi \circ h_{i_0}^{-1})$ converges weakly to a centered Gaussian process Z ,

$$\sqrt{m}(\widehat{\phi \circ h_{i_0}^{-1}} - \phi \circ h_{i_0}^{-1}) \xrightarrow[m \rightarrow \infty]{\mathbf{D}} Z,$$

with covariance function defined for all $(s, t) \in [a, b]^2$ by

$$\text{cov}(Z(s), Z(t)) = r(h_{i_0}^{-1}(s), h_{i_0}^{-1}(t)).$$

We may also compute confidence bands for $\phi \circ h_{i_0}^{-1}$, based on a consistent estimator of $\text{var}(Z(t)) = \gamma \circ h_{i_0}^{-1}(t) - \{\phi \circ h_{i_0}^{-1}(t)\}^2$.

Lemma 2.6. Let $t \in [a, b]$. Let $\widehat{\gamma \circ h_{i_0}^{-1}}(t) = \frac{1}{m} \sum_{i=1}^m T_i^2(t)$, with $T_i(\cdot)$ defined by (10) and (9). Then

$$\widehat{\gamma \circ h_{i_0}^{-1}}(t) - \left\{ \widehat{\phi \circ h_{i_0}^{-1}}(t) \right\}^2 \xrightarrow[m, n \rightarrow \infty]{a.s.} \text{var}(Z(t)).$$

Proof of this lemma relies on the same arguments as proof of Lemma 2.3 and is outlined in Section 4.2. A pointwise asymptotic confidence band for $\phi \circ h_{i_0}^{-1}$ is now given by

Corollary 2.7. An asymptotic $(1 - \alpha)$ -level pointwise confidence band for $\phi \circ h_{i_0}^{-1}$ is given by

$$\left[\widehat{\phi \circ h_{i_0}^{-1}}(t) - u_{1-\frac{\alpha}{2}} \sqrt{\frac{\widehat{\text{var}}(Z(t))}{m}}, \widehat{\phi \circ h_{i_0}^{-1}}(t) + u_{1-\frac{\alpha}{2}} \sqrt{\frac{\widehat{\text{var}}(Z(t))}{m}} \right],$$

where $\widehat{\text{var}}(Z(t)) = \widehat{\gamma \circ h_{i_0}^{-1}}(t) - \left\{ \widehat{\phi \circ h_{i_0}^{-1}}(t) \right\}^2$.

3 Extensions to the general case

In the preceding part, we studied the asymptotic behaviour of a new warping methodology. However, drastic restrictions over the class of functions are needed, mostly dealing with monotonicity of the observed functions and on a non noisy model. In this part, we get rid of such assumptions and provide a practical way of handling more realistic observations.

First, note that the assumptions $H(a) \stackrel{a.s.}{=} a$ et $H(b) \stackrel{a.s.}{=} b$ can be weakened into the following assumptions

ii') $H^{-1}(\cdot, a)$ and $H^{-1}(\cdot, b)$ random variables compactly supported such that

$$\sup_{w \in \Omega} H^{-1}(w, a) \stackrel{a.s.}{<} \inf_{w \in \Omega} H^{-1}(w, b).$$

Then, we focus on the other assumptions.

3.1 Breaking monotonicity

The main idea is to build a transformation \mathcal{G} which turns a nonmonotone function into a monotone function while preserving the warping functions. For sake of simplicity, the observation times will be taken equal for all the curves, hence t_{ij} will be denoted t_j . Hence, the observations

$$Y_{ij} = f \circ f_i^{-1}(t_j), \quad i = 1, \dots, m, \quad j = 0, \dots, n,$$

are transformed into

$$Z_{ij} = \mathcal{G}(f) \circ h_i^{-1}(t_j) := g \circ h_i^{-1}(t_j), \quad i = 1, \dots, m, \quad j = 0, \dots, n, \quad (12)$$

where g is a monotone function. So, estimating the warping process of the monotonized model can be used to estimate the real warping functions, and then align the original observations Y_{ij} to their structural mean.

For this, consider a nonmonotone function $f : [a, b] \rightarrow \{1, \dots, m\}$ and let $a = s_0 < s_1 < \dots < s_r < s_{r+1} = b$ be the different variational change points, in the sense that $\forall k \in \{0, \dots, r-1\}$,

$$\begin{aligned} & \forall (t_1, t_2) \in]s_k, s_{k+1}[, \forall (t_3, t_4) \in]s_{k+1}, s_{k+2}[, \\ & t_1 < t_2 \text{ et } t_3 < t_4 \Rightarrow (f(t_1) - f(t_2))(f(t_3) - f(t_4)) < 0. \end{aligned} \quad (13)$$

So, consider functional warping over the subset

$$\mathcal{F} = \{f : [a, b] \rightarrow \{1, \dots, m\} \subset \mathbb{R} \text{ such that (13) holds} \}.$$

Let $\pi :]a, b[\setminus \{s_1, \dots, s_r\} \rightarrow \{-1, 1\}$ be a tool function which indicates whether, around a given point t , the function is f increasing or decreasing, defined by

$$\pi : s_{l(t)} < t < s_{l(t)+1} \mapsto \pi(t) = \begin{cases} -1 & \text{si } s_{l(t)} - s_{l(t)+1} > 0, \\ 1 & \text{si } s_{l(t)} - s_{l(t)+1} < 0, \end{cases}$$

with $l(t) \in \{0, \dots, r\}$.

Monotonizing Operator

Define for all $f \in \mathcal{F}$ the operator $\mathcal{G}(\cdot, f) : t \in]a, b[\setminus \{s_0, \dots, s_{r+1}\} \rightarrow \mathcal{G}(t, f)$ by

$$\mathcal{G}(t, f) = f(t)\pi(t) - \sum_{k=0}^r \pi(t)f(s_k)\mathbf{1}_{]s_k, s_{k+1}[}(t) + f(s_0) + \sum_{k=1}^r |f(s_{k-1}) - f(s_k)|\mathbf{1}_{]s_k, b[}(t),$$

and, for all $k \in \{0, \dots, r+1\}$,

$$\mathcal{G}(s_k, f) = f(a) + \sum_{l=1}^k |f(s_{l-1}) - f(s_l)|,$$

with the notation $\sum_{l=1}^0 |f(s_{l-1}) - f(s_l)| = 0$.

By construction, it is obvious that $t \rightarrow \mathcal{G}(t, f)$ is strictly increasing. Moreover, the following proposition proves that the warping functions remain unchanged

Proposition 3.1. Consider $f \in \mathcal{F}$ and warping functions h_i , $i = 1, \dots, m$. Set $\forall i = 1, \dots, m$, $\mathcal{G}(\cdot, f_i) = g_i(\cdot)$. We have $g_i = g \circ h_i^{-1}$.

Proof. For $i \in \{1, \dots, m\}$, let us prove that $f_i \in \mathcal{F}$. Using (13), consider the change points $(s_k)_{k=1, \dots, r}$ and set $s_k^i = h_i(s_k)$. For $k \in \{0, \dots, r-1\}$, consider

$$(t_1, t_2) \in]s_k^i, s_{k+1}^i[, (t_3, t_4) \in]s_{k+1}^i, s_{k+2}^i[/ t_1 < t_2 \text{ et } t_3 < t_4,$$

then,

$$(f_i(t_1) - f_i(t_2))(f_i(t_3) - f_i(t_4)) = (f \circ h_i^{-1}(t_1) - f \circ h_i^{-1}(t_2))(f \circ h_i^{-1}(t_3) - f \circ h_i^{-1}(t_4)).$$

Since h_i^{-1} is strictly increasing, we get $s_k < h_i^{-1}(t_1) < h_i^{-1}(t_2) < s_{k+1}$ and $s_{k+1} < h_i^{-1}(t_3) < h_i^{-1}(t_4) < s_{k+2}$. Hence

$$(f_i(t_1) - f_i(t_2))(f_i(t_3) - f_i(t_4)) < 0,$$

which yields that $f_i \in \mathcal{F}$. So define $g_i = \mathcal{G}(f_i)$. For all $t \in]a, b[\setminus \{s_1^i, \dots, s_r^i\}$, we get

$$g_i(t) = f_i(t)\Pi(t, f_i) - \sum_{k=0}^r \Pi(t, f_i) f_i(s_k^i) \mathbf{1}_{]s_k^i, s_{k+1}^i[}(t) + f_i(s_0) + \sum_{k=1}^r |f_i(s_{k-1}^i) - f_i(s_k^i)| \mathbf{1}_{]s_k^i, b[}(t),$$

But

$$\begin{aligned} f_i(t) &= f \circ h_i^{-1}(t), \\ \Pi(t, f_i) &= \Pi(t, f \circ h_i^{-1}) = \Pi(h_i^{-1}(t), f), \\ \mathbf{1}_{]s_k^i, s_{k+1}^i[}(t) &= \mathbf{1}_{]h_i(s_k), h_i(s_{k+1})[}(t) = \mathbf{1}_{]s_k, s_{k+1}[}(h_i^{-1}(t)), \end{aligned}$$

which implies that

$$g_i = g \circ h_i^{-1}.$$

□

The discretization implies however that the Z_{ij} can not be computed directly since the functions f_i , $i = 1, \dots, m$, are known on the grid t_j , $j = 0, \dots, n$, while the values, s_k and s_k^i are unknown. So consider estimates of Z_{ij} defined as follows

$$\tilde{Z}_{i0} = Y_{i0}$$

$$\forall j \in \{1, \dots, n\}, \tilde{Z}_{ij} = \tilde{Z}_{ij-1} + |Y_{ij} - Y_{ij-1}|. \quad (14)$$

The following proposition proves the consistency of such estimation procedure.

Proposition 3.2. For $f \in \mathcal{F}$, $i \in \{1, \dots, m\}$ and $t \in [a, b]$, define a sequence $j(n)$ such that $\frac{j(n)}{n} \xrightarrow{n \rightarrow +\infty} t$. Then,

$$\tilde{Z}_{ij(n)} - Z_{ij(n)} \xrightarrow[n \rightarrow +\infty]{a.s.} 0.$$

Proof. Set $t \in]a, b[\setminus \{s_1^i, \dots, s_r^i\}$, and $l \in \{0, 1, \dots, r\}$ such that $t \in]s_l^i, s_{l+1}^i[$. Consider $j(n)$, $\frac{j(n)}{n} \xrightarrow{n \rightarrow +\infty} t$, then $\exists n_0 \in \mathbb{N} / \forall n \in \mathbb{N}, n \geq n_0 \Rightarrow \frac{j(n)}{n} \in]s_l^i, s_{l+1}^i[$. $\forall n \geq n_0$, we have

$$\begin{aligned} \tilde{Z}_{ij(n)} &= \tilde{Z}_{ij(n)-1} + |Y_{ij(n)} - Y_{ij(n)-1}| \\ &= Y_{i0} + \sum_{k=1}^{j(n)} |Y_{ik} - Y_{ik-1}|. \end{aligned}$$

Moreover,

$$\begin{aligned} Z_{ij(n)} &= g \circ h_i^{-1} (t_{j(n)}) \\ &= f_i (t_{j(n)}) \Pi (t_{j(n)}, f_i) + \Pi (t_{j(n)}, f_i) f_i (s_l^i) + f_i(s_0) + \sum_{k=1}^l |f_i (s_{k-1}^i) - f_i (s_k^i)|, \end{aligned}$$

which yields that

$$\begin{aligned} Z_{ij(n)} &= \Pi (t_{j(n)}, f_i) (Y_{ij(n)} - f_i (s_l^i)) + Y_{i0} + \sum_{k=1}^l |f_i (s_{k-1}^i) - f_i (s_k^i)| \\ &= |Y_{ij(n)} - f_i (s_l^i)| + Y_{i0} + \sum_{k=1}^l |f_i (s_{k-1}^i) - f_i (s_k^i)| \\ &= A + Y_{i0} + B. \end{aligned}$$

Write $\forall k = 1, \dots, l+1$, $s_{k-1}^i \leq t_{j_k - p_k} < \dots < t_{j_k} \leq s_k^i$, we get that

$$\begin{aligned} &|f_i (s_{k-1}^i) - f_i (s_k^i)| \\ &= \left| f_i (s_{k-1}^i) - f_i (t_{j_k - p_k}) + \sum_{q=1}^{p_k} (f_i (t_{j_k - q}) - f_i (t_{j_k - q + 1})) + f_i (t_{j_k}) - f_i (s_k^i) \right| \\ &= |f_i (s_{k-1}^i) - f_i (t_{j_k - p_k})| + \sum_{q=1}^{p_k} |f_i (t_{j_k - q}) - f_i (t_{j_k - q + 1})| + |f_i (t_{j_k}) - f_i (s_k^i)| \\ &= |f_i (s_{k-1}^i) - Y_{ij_k - p_k}| + \sum_{q=1}^{p_k} |Y_{ij_k - q + 1} - Y_{ij_k - q}| + |Y_{ij_k} - f_i (s_k^i)|, \end{aligned}$$

Hence

$$\begin{aligned} B &= \sum_{k=1}^l |f_i (s_{k-1}^i) - f_i (s_k^i)| \\ &= \sum_{k=1}^l |f_i (s_{k-1}^i) - Y_{ij_k - p_k}| + \sum_{k=1}^l |Y_{ij_k} - f_i (s_k^i)| + \sum_{k=1}^l \sum_{q=1}^{p_k} |Y_{ij_k - q + 1} - Y_{ij_k - q}| \\ &= \sum_{k=1}^l |f_i (s_{k-1}^i) - Y_{ij_k - p_k}| + \sum_{k=1}^l |Y_{ij_k} - f_i (s_k^i)| + \sum_{k=1}^{j_l} |Y_{ik} - Y_{ik-1}| \\ &\quad - \sum_{k=1}^{l-1} |Y_{ij_{k+1} - p_{k+1}} - Y_{ij_{k+1} - p_{k+1} - 1}| - |Y_{i1} - Y_{i0}|. \end{aligned}$$

With the same ideas, we can write

$$\begin{aligned} A &= |Y_{ij(n)} - f_i (s_l^i)| = \\ &\quad \sum_{q=j_{l+1} - j(n) + 1}^{p_{l+1} + 1} |Y_{ij_{l+1} - q + 1} - Y_{ij_{l+1} - q}| + |Y_{ij_{l+1} - p_{l+1}} - f_i (s_l^i)| - |Y_{ij_{l+1} - p_{l+1}} - Y_{ij_{l+1} - p_{l+1} - 1}|. \end{aligned}$$

As a result,

$$\begin{aligned} Z_{ij(n)} - \tilde{Z}_{ij(n)} &= \sum_{k=1}^{l+1} |f_i(s_{k-1}^i) - Y_{ij_{k-p_k}}| + \sum_{k=1}^l |Y_{ij_k} - f_i(s_k^i)| \\ &\quad - \sum_{k=1}^l |Y_{ij_{k+1-p_{k+1}}} - Y_{ij_{k+1-p_{k+1}-1}}| - |Y_{i1} - Y_{i0}|. \end{aligned}$$

By continuity of f , f_i is also continuous, hence

$$Z_{ij(n)} - \tilde{Z}_{ij(n)} \xrightarrow[n \rightarrow +\infty]{a.s.} 0.$$

For $t \in \{s_0^i, \dots, s_{r+1}^i\}$, we get similar results, leading to the conclusion. \square

As a conclusion, we can extend our results to the case of nonmonotone functions since we transform the problem into a monotone warping problem with the same warping functions. These functions h_i $i = 1, \dots, m$ can be estimated by our methodology using the new observations $\tilde{Z}_{ij} \forall i = 1, \dots, m, \forall j = 0, \dots, n$. The estimator can be written in the following form

$$\widetilde{\phi \circ h_{i_0}^{-1}}_{mn}(t) = \frac{1}{m-1} \sum_{\substack{i=1 \\ i \neq i_0}}^m T_{j_i}, \quad (15)$$

with

$$T_{j_i} = \arg \min_{t_j \in \{t_0, \dots, t_n\}} |\tilde{Z}_{ij} - \tilde{Z}_{i_0 j_0}|,$$

and

$$t_{j_0} = \arg \min_{t_j \in \{t_0, \dots, t_n\}} |t_j - t|.$$

3.2 Dealing with noisy data

If theoretical asymptotical results are only given in a non noisy framework, we can still handle the case where the data are observed in the standard regression model

$$Y_{ij} = f \circ h_i^{-1}(t_j) + \varepsilon_{ij}, \quad i = 1, \dots, m, \quad j = 0, \dots, n, \quad (16)$$

with $\varepsilon_{ij} \stackrel{\text{i.i.d.}}{\sim} \mathcal{N}(0, \sigma^2)$. To be able to apply our algorithm, we first denoise the data. For this, we estimate separately each function f_i , $i = 1, \dots, m$, by a Kernel estimator. On a practical point of view, we describe the estimation procedure used in the simulations.

1. $\forall i \in \{1, \dots, m\}$, f_i is estimated

$$\hat{f}_i(t_0) = \frac{1}{m} \sum_{i=1}^m Y_{i0} \xrightarrow[m \rightarrow +\infty]{a.s.} f(t_0),$$

$$\hat{f}_i(t_n) = \frac{1}{m} \sum_{i=1}^m Y_{in} \xrightarrow[m \rightarrow +\infty]{a.s.} f(t_n).$$

Given a Gaussian kernel Φ , $\forall j \in \{1, \dots, n-1\}$, $f_i(t_j)$ is estimated by

$$\hat{f}_i(t_j) = \frac{\sum_{k=0}^n Y_{ik} \Phi\left(\frac{t_k - t_j}{\nu_i}\right)}{\sum_{k=0}^n \Phi\left(\frac{t_k - t_j}{\nu_i}\right)}. \quad (17)$$

The bandwidths ν_i are to be properly chosen.

2. The estimation procedure can be conducted using the denoised observations $\hat{f}_i(t_j)$, leading to new estimates $\hat{f}(t)$ of the structural expectation $f \circ \phi^{-1}$.

We point out that the efficiency of the procedure heavily relies on a proper choice of the bandwidths ν_i , $i = 1, \dots, m$. Cross-validation technics do not provide good results since the aim is not to get a good estimation of the function but only a good separation of the different functions. Hence oversmoothing the data is not a drawback in this settings. So, the smoothing parameters $\nu = \nu_i, \forall i = 1, \dots, m$ are obtained by minimizing the following matching criterion

$$\hat{\nu} = \arg \min_{\nu \in L} \sum_{i=1}^m \sum_{j=0}^n \left| \hat{f}^i(t_j) - \hat{f}(t_j) \right|,$$

over a grid L . Practical applications of this algorithm are given in Figure

4 Numerical Study

In this part, we estimate the structural expectation, using both the proposed method and the analytic registration approach developed in [9]. Note that an alternative to analytic registration is provided by the so-called landmarks registration approach in [5], but this approach requires the determination of landmarks (such as local extrema) which can be difficult in our simulations. Hence, this method is not implemented here. First, results on simulated data are given in order to compare these two methods. Then, an application of our methodology is given for a real data set.

4.1 Simulations

Two simulation studies are carried out in this section. The first one involves a strictly increasing function and the second one involves a non monotone function.

Warped functions. Let f and g (see Figures 1 and 2) be defined by

$$\forall t \in [0, 1], f(t) = \sin(3\pi t) + 3\pi t \text{ and } g(t) = \frac{\sin(6\pi t)}{6\pi t}.$$

These two functions will be warped by the following random warping process.

Warping processes. The Stochastic warping functions H_i , $i = 1, \dots, m$, are simulated by the iterative process described below.

Let $\epsilon > 0$. First, for all $i = 1, \dots, m$, let $H_i^{(0)}$ be the identity function. Then, the warping functions $H_i^{(k+1)}$, $i = 1, \dots, m$, are successively carried out from functions $H_i^{(k)}$, $i = 1, \dots, m$, by iterating N times the following process :

1. Let U be a uniformly distributed random variable on $[10\epsilon, 1 - 10\epsilon]$.
2. Let V_i , $i = 1, \dots, m$, be independant and identically uniformly distributed variables on $[U - \epsilon, U + \epsilon]$.
3. For all $i = 1, \dots, m$, the warping function $H_i^{(k)}$ is warped as follows :

$$H_i^{(k+1)} = W_i \circ H_i^{(k)}$$

where W_i is defined by

$$W_i(t) = \begin{cases} \frac{V_i}{U}t & \text{if } 0 \leq t \leq U, \\ \frac{1-V_i}{1-U}t + \frac{V_i-U}{1-U} & \text{if } U < t \leq 1. \end{cases}$$

We have thus defined strictly increasing stochastic functions $H_i : [0, 1] \rightarrow [0, 1]$, such that

$$\forall t \in [0, 1], \mathbf{E}(H_i(t)) = \phi(t) = t.$$

Hence, the warping process is centered, in the sense that, the structural expectation $f \circ \phi^{-1}$ is equal to f .

Our simulated warping functions H_i , $i = 1, \dots, m$, have been carried out by using the above iterating process with $m = 30$, $N = 3000$ and $\epsilon = 0.005$. These processes are shown on Figure 1 and on Figure 2. In these figure, we can see the very large phase variations. For example, point 0.2 is warped between approximately 0.05 and 0.35 for the first case, and between approximately 0.00 and 0.40 for the second case.

Simulated data. Finally, simulated data are carried out on an equally spaced grid as follows :

$$Y_{ij} = f(H_i^{-1}(t_j))$$

and

$$Y_{ij} = g(H_i^{-1}(t_j)) + \epsilon_{ij}$$

with $t_j = \frac{j}{n}$, $j = 0, 1, \dots, n$, $n = 100$ points. Simulated warped functions are respectively shown on Figures 1 and 2.

Figures 1 and 2 show the functions f and g and the mean functions of the warped functions. We can easily see that the classical mean is not adapted to describe the data. In particular, the mean function of the first simulated data does not reflect the flat component of the function f (in the range $[0.2, 0.4]$) which yet appears in each individual warped function. In the same way, the mean function of the second simulated data set does not reflect the structure of the individual curves. The classical mean attenuated curve variations.

The estimated structural expectations with both the analytic registration approach and the proposed method are shown on Figures 1 and 2 (bottom right figures). We

can easily see that estimations with our proposed method are closer to the structural expectations f and g . These results can be explained as follows :

- For the first simulated data set, the analytic registration approach does not directly work on the strictly increasing functions but on first derivatives. However, theoretically registering a given function data set is not the same issue as the registration of first derivatives of these functions.
- The analytic registration approach uses the mean curve to register all the functions. Due to the drawbacks of the mean curve when dealing with large deformations (for instance in the second data set, where the result is a very flat mean curve), the structural mean approach provides better results.
- For both first and second simulated data sets, the analytic registration approach works on estimated functions and not directly with given data, which implies an additional source of error.




Figure 1: Function f is shown on top left figure (solid line). Simulated warping processes are shown on top right figure. Simulated warped functions are shown on bottom left figure. The classical mean of these functions is draw on top left figure (dotted line). Finally, estimations of f with the Analytic registration procedure (dashed line) and with our method (dotted-dashed line) are shown on bottom right figure.

4.2 A concrete application : multiple referees and equity

The field of application of the results presented in this paper is large. Here we consider an example in the academic field : how to guarantee equality in an exam with several different referees?

Consider an examination with a large number of candidates, such that it is impossible to evaluate the candidates one after another. So the students are divided into m groups




Figure 2: Function g is shown on top left figure (solid line). Simulated warping processes are shown on top right figure. Simulated warped functions are shown on bottom left figure. The classical mean of these functions is draw on top left figure (dotted line). Finally, estimations of g with the Analytic registration procedure (dashed line) and with our method (dotted-dashed line) are shown on bottom right figure.

and a board of examiners is charged to grade the students of one group. The evaluation is performed by assigning a score from 0 to 20.

The m different boards of examiners are supposed to behave the same way in order to respect the equality among the candidates. Moreover it is assumed that the sampling of the candidates is perfect in the sense that it is done in such a way that each board of examiners evaluates candidates with the same global level. Hence, if all the examiners had the same requirement levels, the distribution of the ranks would be the same for all the boards of examiners. Here, we aim at balancing the effects of the differences between the examiners, and gaining equity for the candidates, and at studying real data provided by the french competitive teacher exam *Agrégation de mathématiques*.

This situation can be modeled as follows. Define for each group $i = 1, \dots, m$, the score obtained by students as $\mathbf{X}^i = \{X_j^i \in (0, 20), j = 1, \dots, n_i\}$. Let $f_i, i = 1, \dots, m$ be the repartition function of the group i defined by

$$f_i(t) = \frac{1}{n} \sum_{j=1}^n \mathbf{1}_{X_j^i \leq t}.$$

Under the assumption that all the examiners give the same ranking to two candidates with the same level, which will be denoted H_0 , it implies that the random variables $\mathbf{X}^1, \dots, \mathbf{X}^m$ are equally distributed. Figure 3 shows the distinct empirical functions for all the $m = 13$ groups.

First, we test for every couple of sets (i, j) the assumption $\mathbf{X}^i \sim \mathbf{X}^j$. For this we perform the following homogeneity test, see for instance [3]. Define for all $k = 1, \dots, 20$,

$n_k^i = \sum_{l=1}^m \mathbf{1}_{X_l^i=k}$ and $n_k^j = \sum_{l=1}^m \mathbf{1}_{X_l^j=k}$. Define $\hat{\mu}_k = \frac{n_k^i + n_k^j}{2n}$. Finally set $d_k = n\hat{\mu}_k$ and

$$D_n^i = \sum_{k=1}^{20} \frac{(d_k - n_k^i)^2}{d_k}, \quad D_n^j = \sum_{k=1}^{20} \frac{(d_k - n_k^j)^2}{d_k}.$$

Under H_0 , $D_n = D_n^i + D_n^j$ converges in distribution to a χ^2 distribution and almost surely to $+\infty$ under the assumption that the two laws are different.

We have $n = 4000$ candidates and $m = 13$ examiners. We test the assumption H_0 for all the different possibilities. In 60% of cases and for a level equal to 5%, we reject the assumption that the rankings follow the same distribution.

As a consequence the following procedure is proposed. We aim at finding the average way of ranking, with respect to the ranks that were given within the 13 bunches of candidates. For this, assume that there is such average empirical distribution function and that the empirical distribution function of each group is warped by a random process from this reference distribution function. Hence a good choice is given by the structural expectation, since the functions f_j , $j = 1, \dots, 13$ are increasing so that Theorem 2.1 may apply.

Figure 3 shows the different empirical distributions and the structural empirical distribution function. At each dot on the empirical distribution function, corresponding to the rank given within one group, is associated its correspondent structural rank. It is obtained by simply projecting onto the structural empirical distribution function. As a result we obtain rescaled structural ranks corresponding to the rank obtained by each candidate if they could have been judged by an average board of examiner, leading to a more fair rankings. Indeed the difference of judgments between each subgroup are revised according to the average judgment of all the groups of examiners.

In conclusion, structural expectation provides a data-driven pattern, which plays the role of a reference pattern to which all the different curves can be compared. We applied successfully this method to rescale the ranks obtained by candidates evaluated by different boards of examiners. This use is not restrictive since it can be used to provide mean patterns for other types of functional data in various fields such as econometry, biology or data traffic for instance.

A Proofs and technical lemmas

In practice, functions $(f_i)_{i \in \{1, \dots, m\}}$ are evaluated on a discrete interval of \mathbb{R} as described in Section 1. In order to prove Theorem 2.1 and Theorem 2.5, we first study asymptotic results on the theoretical continuous model, i.e.

$$f_i(t) = f \circ h_i^{-1}(t), \quad i \in \{1, \dots, m\}, \quad t \in [a, b], \quad (18)$$

where H is defined in the same way as in model (3). In a second step, we will extend the proofs to the discretized model and prove results of Section 3.

So consider that all functions are measured on the entire interval $[a, b]$. After asymptotic results are proved for this continuous model in Section A.1, we use these results to prove Theorem 2.1 and Theorem 2.5 (and subsequently Theorem 2.2) in Section A.2.




Figure 3: Empirical distribution functions (left figure) and structural expectation estimation (right figure).

A.1 Asymptotic results for the continuous model

For the continuous model (18), we provide asymptotic results (analogous to Theorem 2.1 and Theorem 2.5) and their proofs.

A.1.1 Estimator and asymptotic results of the inverse of the structural expectation

Considering the continuous model (18), we define an empirical estimator of the inverse of the structural expectation in the following way. Set

$$\overline{f_{ES}^{-1}} = \frac{1}{m} \sum_{i=1}^m f_i^{-1}. \quad (19)$$

The following theorem gives us consistency and asymptotic normality of this estimator.

Theorem A.1. Under assumption (2), we have that $\overline{f_{ES}^{-1}}$ converges almost surely to f_{ES}^{-1}

$$\left\| \overline{f_{ES}^{-1}} - f_{ES}^{-1} \right\|_{\infty} \xrightarrow[m \rightarrow \infty]{as} 0.$$

Moreover, let assume that assumptions (4) and (5) are allowed. Then, we have that $\sqrt{m}(\overline{f_{ES}^{-1}} - f_{ES}^{-1})$ converges weakly toward a zero-mean Gaussian process G :

$$\sqrt{m} \left(\overline{f_{ES}^{-1}} - f_{ES}^{-1} \right) \xrightarrow[m \rightarrow \infty]{\mathbf{D}} G,$$

where the covariance function of G is defined, for all $(s, t) \in [f(a), f(b)]^2$, by

$$\text{Cov}(G(s), G(t)) = r \left(f^{-1}(s), f^{-1}(t) \right).$$

Proof. Almost sure convergence of estimator $\overline{f_{ES}^{-1}}$ is directly deduced from corollary 7.10 (p. 189) in [7]. This corollary is an extension of the Strong Law of Large Numbers to Banach spaces.

For all $i \in \{1, \dots, m\}$, the functions $(f_i^{-1})_{i \in \{1, \dots, m\}}$ are obviously strictly increasing, hence $\overline{f_{ES}^{-1}}$ is strictly increasing, and we have

$$\overline{f_{ES}^{-1}} = \frac{1}{m} \sum_{i=1}^m f_i^{-1} = \frac{1}{m} \sum_{i=1}^m (f \circ h_i^{-1})^{-1} = \frac{1}{m} \sum_{i=1}^m h_i \circ f^{-1}.$$

For all $i \in \mathbb{N}^*$, let

$$X_i = h_i \circ f^{-1} - f_{ES}^{-1},$$

and, for all $m \in \mathbb{N}^*$, let

$$S_m = \sum_{i=1}^m X_i.$$

The $(X_i)_{i \in \{1, \dots, m\}}$ are B -valued random variables, where $B = \mathcal{C}([f(a), f(b)])$ is a separable Banach space. Moreover, the dual space of B is the set of bounded measures on $[f(a), f(b)]$ ([10]). Hence, our framework corresponds to Chapter 7 of [7] and thus we can apply Corollary 7.10. Indeed, we have

$$\mathbf{E}(\|X_1\|_{\infty}) < +\infty \text{ et } \mathbf{E}(X_1) = 0,$$

then

$$\frac{S_m}{m} \xrightarrow[m \rightarrow +\infty]{a.s.} 0,$$

which proves almost sure convergence.

We now turn to weak convergence. From the multivariate CLT, for any $k \in \mathbb{N}^*$ and fixed $(y_1, \dots, y_k) \in [f(a), f(b)]^k$,

$$\sqrt{m} \left(\begin{pmatrix} \overline{f_{ES}^{-1}}(y_1) \\ \vdots \\ \overline{f_{ES}^{-1}}(y_k) \end{pmatrix} - \begin{pmatrix} f_{ES}^{-1}(y_1) \\ \vdots \\ f_{ES}^{-1}(y_k) \end{pmatrix} \right) \xrightarrow[m \rightarrow \infty]{\mathcal{D}} \mathcal{N}_k(0, \Gamma),$$

where the covariance matrix $\Gamma = (\Gamma_{ij})_{i,j}$ is given by $\Gamma_{ij} = \text{cov}(F^{-1}(y_i), F^{-1}(y_j)) = \text{cov}(H(f^{-1}(y_i)), H(f^{-1}(y_j)))$, $i, j = 1, \dots, k$. It remains to show that $\{\sqrt{m}(\overline{f_{ES}^{-1}} - f_{ES}^{-1})\}$ is tight. We verify the moment condition stated by [13] [Example 2.2.12].

$$\begin{aligned} & \mathbf{E} \left[\left| \sqrt{m}(\overline{f_{ES}^{-1}}(s) - f_{ES}^{-1}(s)) - \sqrt{m}(\overline{f_{ES}^{-1}}(t) - f_{ES}^{-1}(t)) \right|^2 \right] \\ &= \mathbf{E} \left[m \left| \frac{1}{m} \sum_{i=1}^m f_i^{-1}(s) - \mathbf{E}F^{-1}(s) - \left(\frac{1}{m} \sum_{i=1}^m f_i^{-1}(t) - \mathbf{E}F^{-1}(t) \right) \right|^2 \right] \\ &= \mathbf{E} \left[\left| F^{-1}(s) - \mathbf{E}F^{-1}(s) - (F^{-1}(t) - \mathbf{E}F^{-1}(t)) \right|^2 \right], \end{aligned}$$

where the last equality follows from the fact that the h_i 's are i.i.d. Then, from (4) and (5), we get that

$$\mathbf{E} \left[\left| \sqrt{m}(\overline{f_{ES}^{-1}}(s) - f_{ES}^{-1}(s)) - \sqrt{m}(\overline{f_{ES}^{-1}}(t) - f_{ES}^{-1}(t)) \right|^2 \right] \leq C_1 C_2 |s - t|^2,$$

which completes the proof. \square

A.1.2 Estimator and asymptotic results of an individual warping function

For the continuous model, we define an empirical estimator of the individual warping function $\phi \circ h_{i_0}^{-1}$ ($i_0 \in \{1, \dots, m\}$) as follows. Conditionally to $F_{i_0} = f_{i_0}$, for all $t \in [a, b]$, let

$$\overline{\phi \circ h_{i_0}^{-1}}(t) = \frac{1}{m-1} \sum_{\substack{i=1 \\ i \neq i_0}}^m f_i^{-1} \circ F_{i_0}(t). \quad (20)$$

The following theorem gives us consistency and asymptotic normality of this estimator.

Theorem A.2. Under assumption (2), we have that $\overline{\phi \circ h_{i_0}^{-1}}$ converges almost surely to $\phi \circ h_{i_0}^{-1}$:

$$\left\| \overline{\phi \circ h_{i_0}^{-1}} - \phi \circ h_{i_0}^{-1} \right\|_{\infty} \xrightarrow[m \rightarrow \infty]{as} 0.$$

Let $n = m^{\frac{1}{2} + \alpha}$ (with $\alpha > 0$) and assume that (4) and (6) hold. Then, we have that $\sqrt{m}(\overline{\phi \circ h_{i_0}^{-1}} - \phi \circ h_{i_0}^{-1})$ converges weakly to a zero-mean Gaussian process Z ,

$$\sqrt{m}(\overline{\phi \circ h_{i_0}^{-1}} - \phi \circ h_{i_0}^{-1}) \xrightarrow[m \rightarrow \infty]{\mathbf{D}} Z,$$

with covariance function defined for all $(s, t) \in [a, b]^2$ by

$$\mathbf{Cov}_{\langle Z, (\cdot) s \rangle, Z(t)} = r(h_{i_0}^{-1}(s), h_{i_0}^{-1}(t)).$$

Proof. Let $i_0 \in \{1, \dots, m\}$. Given $F_{i_0} = f_{i_0}$,

$$\overline{\phi \circ h_{i_0}^{-1}} = \frac{1}{m-1} \sum_{\substack{i=1 \\ i \neq i_0}}^m f_i^{-1} \circ f_{i_0}.$$

Noting also that $\phi \circ h_{i_0}^{-1} = f_{ES}^{-1} \circ f_{i_0}$, consistency of $\overline{\phi \circ h_{i_0}^{-1}}$ follows by the same arguments as in proof of Theorem A.1.

We now turn to weak convergence. From the multivariate CLT, for any $k \in \mathbb{N}^*$ and fixed $(t_1, \dots, t_k) \in [a, b]^k$,

$$\sqrt{m} \left(\begin{pmatrix} \overline{\phi \circ h_{i_0}^{-1}}(t_1) \\ \vdots \\ \overline{\phi \circ h_{i_0}^{-1}}(t_k) \end{pmatrix} - \begin{pmatrix} \phi \circ h_{i_0}^{-1}(t_1) \\ \vdots \\ \phi \circ h_{i_0}^{-1}(t_k) \end{pmatrix} \right) \xrightarrow{\mathcal{D}} \mathcal{N}_k(0, \Gamma_0),$$

where the covariance matrix $\Gamma_0 = (\Gamma_{0,ij})_{i,j}$ is given by $\Gamma_{0,ij} = \text{cov}(H(h_{i_0}^{-1}(t_i)), H(h_{i_0}^{-1}(t_j))) = r(h_{i_0}^{-1}(t_i), h_{i_0}^{-1}(t_j))$, $i, j = 1, \dots, k$. It remains to show that $\{\sqrt{m}(\overline{\phi \circ h_{i_0}^{-1}} - \phi \circ h_{i_0}^{-1})\}$ is tight. Again, we verify the moment condition stated by [13] [Example 2.2.12].

$$\begin{aligned} & \mathbf{E} \left[\left| \sqrt{m}(\overline{\phi \circ h_{i_0}^{-1}}(s) - \phi \circ h_{i_0}^{-1}(s)) - \sqrt{m}(\overline{\phi \circ h_{i_0}^{-1}}(t) - \phi \circ h_{i_0}^{-1}(t)) \right|^2 \right] \\ &= \frac{m}{(m-1)^2} \mathbf{E} \left[\left| \sum_{\substack{i=1 \\ i \neq i_0}}^m (f_i^{-1}(f_{i_0}(s)) - \mathbf{E}f_i^{-1}(f_{i_0}(s)) - (f_i^{-1}(f_{i_0}(t)) - \mathbf{E}f_i^{-1}(f_{i_0}(t)))) \right|^2 \right] \\ &= \frac{m}{m-1} \mathbf{E} \left[\left| H(h_{i_0}^{-1}(s)) - \mathbf{E}H(h_{i_0}^{-1}(s)) - (H(h_{i_0}^{-1}(t)) - \mathbf{E}H(h_{i_0}^{-1}(t))) \right|^2 \right]. \end{aligned}$$

Now, from assumptions (4) and (6), we get that

$$\mathbf{E} \left[\left| \sqrt{m}(\overline{\phi \circ h_{i_0}^{-1}}(s) - \phi \circ h_{i_0}^{-1}(s)) - \sqrt{m}(\overline{\phi \circ h_{i_0}^{-1}}(t) - \phi \circ h_{i_0}^{-1}(t)) \right|^2 \right] \leq 2C_1 C_3 |s - t|^2,$$

which completes the proof. \square

A.2 Proofs of asymptotic results

We now use Theorem A.1 (given for the continuous model) to prove Theorem 2.1.

PROOF OF THEOREM 2.1 Let $y \in [f(a), f(b)]$. The n observation times are equidistant and for each $i = 1, \dots, m$, f_i is almost surely increasing, hence $f_i^{-1}(y) - \frac{1}{n} \leq T_i(y) \leq f_i^{-1}(y) + \frac{1}{n}$. This implies that almost surely,

$$\overline{f_{ES}^{-1}}(y) - \frac{1}{n} \leq \widehat{f_{ES}^{-1}}(y) \leq \overline{f_{ES}^{-1}}(y) + \frac{1}{n}. \quad (21)$$

Since

$$\left\| \overline{f_{ES}^{-1}} + \frac{1}{n} - f_{ES}^{-1} \right\|_{\infty} \leq \left\| \overline{f_{ES}^{-1}} - f_{ES}^{-1} \right\|_{\infty} + \frac{1}{n},$$

we get that $\left\| \overline{f_{ES}^{-1}} + \frac{1}{n} - f_{ES}^{-1} \right\|_{\infty} \xrightarrow[m, n \rightarrow \infty]{as} 0$, by Theorem A.1. A similar argument holds for

LHS(21) and finally, $\left\| \widehat{f_{ES}^{-1}} - f_{ES}^{-1} \right\|_{\infty} \xrightarrow[m, n \rightarrow \infty]{as} 0$.

Let $n = m^{\frac{1}{2} + \alpha}$ ($\alpha > 0$). From (21), we get that almost surely,

$$\left\| \sqrt{m}(\widehat{f_{ES}^{-1}} - \overline{f_{ES}^{-1}}) \right\|_{\infty} \leq \frac{1}{m^{\alpha}}.$$

Since $\left\| \sqrt{m}(\widehat{f_{ES}^{-1}} - \overline{f_{ES}^{-1}}) \right\|_{\infty} = \left\| \sqrt{m}(\widehat{f_{ES}^{-1}} - f_{ES}^{-1}) - \sqrt{m}(\overline{f_{ES}^{-1}} - f_{ES}^{-1}) \right\|_{\infty}$, we get that $\left\| \sqrt{m}(\widehat{f_{ES}^{-1}} - f_{ES}^{-1}) - \sqrt{m}(\overline{f_{ES}^{-1}} - f_{ES}^{-1}) \right\|_{\infty}$ converges almost surely to 0 as m tends to infinity. Combining Theorem A.1 and Theorem 4.1 in [2], it follows that $\sqrt{m}(\widehat{f_{ES}^{-1}} - f_{ES}^{-1}) \xrightarrow[m \rightarrow \infty]{\mathbf{D}} G$. We now turn to proof of Lemma 2.3.

PROOF OF LEMMA 2.3 Consider first the continuous model (18), and define

$$\overline{\gamma \circ f^{-1}} = \frac{1}{m} \sum_{i=1}^m (f_i^{-1})^2.$$

Using similar arguments as in proof of Theorem A.1, we get that $\left\| \overline{\gamma \circ f^{-1}} - \gamma \circ f^{-1} \right\|_{\infty} \xrightarrow[m \rightarrow \infty]{as} 0$.

Now, since $|T_i(y) - f_i^{-1}(y)| \leq \frac{1}{n}$, we obtain by straightforward calculations that almost surely,

$$-\frac{1}{n} \cdot \frac{1}{m} \sum_{i=1}^m |f_i^{-1}(y)| \leq \frac{1}{m} \sum_{i=1}^m T_i \cdot f_i^{-1}(y) - \overline{\gamma \circ f^{-1}}(y) \leq \frac{1}{n} \cdot \frac{1}{m} \sum_{i=1}^m |f_i^{-1}(y)|,$$

which implies that $\frac{1}{m} \sum_{i=1}^m T_i \cdot f_i^{-1}(y) - \overline{\gamma \circ f^{-1}}(y) \xrightarrow[m, n \rightarrow \infty]{as} 0$, from which we deduce that $\frac{1}{m} \sum_{i=1}^m T_i \cdot f_i^{-1}(y) \xrightarrow[m, n \rightarrow \infty]{as} \gamma \circ f^{-1}(y)$.

From $|T_i(y) - f_i^{-1}(y)| \leq \frac{1}{n}$, we also get that

$$0 \leq \frac{1}{m} \sum_{i=1}^m T_i^2 - \frac{2}{m} \sum_{i=1}^m T_i \cdot f_i^{-1}(y) + \frac{1}{m} \sum_{i=1}^m (f_i^{-1}(y))^2 \leq \frac{1}{n^2},$$

that is, $0 \leq \widehat{\gamma \circ f^{-1}}(y) - \frac{2}{m} \sum_{i=1}^m T_i \cdot f_i^{-1}(y) + \overline{\gamma \circ f^{-1}}(y) \leq \frac{1}{n^2}$, from which we deduce that $\widehat{\gamma \circ f^{-1}}(y) \xrightarrow[m, n \rightarrow \infty]{as} \gamma \circ f^{-1}(y)$. Combining this with Theorem 2.1 completes the proof.

Proof of Theorem 2.2. For F an arbitrary function, let F^{-1} denote its generalized inverse, defined by $F^{-1}(t) = \inf\{y : F(y) \geq t\}$. By Theorem 2.1, for all $y \in [f(a), f(b)]$,

$$\widehat{f_{ES}^{-1}}(y) \xrightarrow[m, n \rightarrow \infty]{as} f_{ES}^{-1}(y).$$

By Lemma 21.2 in [12],

$$(\widehat{f_{ES}^{-1}})^{-1}(t) \xrightarrow[m, n \rightarrow \infty]{as} (f_{ES}^{-1})^{-1}(t)$$

at every t where $(f_{ES}^{-1})^{-1}$ is continuous. Since f_{ES}^{-1} is continuous and strictly increasing, $(f_{ES}^{-1})^{-1}$ is a proper inverse and is equal to $f \circ \phi^{-1}$, hence for all $t \in [a, b]$,

$$(\widehat{f_{ES}^{-1}})^{-1}(t) \xrightarrow[m, n \rightarrow \infty]{as} f \circ \phi^{-1}(t).$$

Now, for all $t \in [a, b]$, $(\widehat{f_{ES}^{-1}})^{-1}(t)$ can be rewritten as

$$(\widehat{f_{ES}^{-1}})^{-1}(t) = v_0 \mathbf{1}_{\{a\}}(t) + \sum_{k=0}^{K(m,n)-1} v_{k+1} \mathbf{1}_{(u_k, u_{k+1}]}(t),$$

and by construction, letting $t \in (u_{k(m,n)}, u_{k(m,n)+1}]$ ($k(m,n) \in \mathcal{K}$), we have $v_{k(m,n)} \leq \widehat{f_{ES}}(t) \leq v_{k(m,n)+1}$. Combining this and the above equality yields that for all $t \in (u_{k(m,n)}, u_{k(m,n)+1}]$, $|\widehat{f_{ES}}(t) - (\widehat{f_{ES}^{-1}})^{-1}(t)| \leq v_{k(m,n)+1} - v_{k(m,n)}$. Since f_{ES} is continuous, $v_{k(m,n)+1} - v_{k(m,n)} \xrightarrow{m,n \rightarrow \infty} 0$, and $|\widehat{f_{ES}}(t) - (\widehat{f_{ES}^{-1}})^{-1}(t)| \xrightarrow{m,n \rightarrow \infty} 0$. It follows that almost surely, $\widehat{f_{ES}}(t)$ converges to $f_{ES}(t)$. Uniform convergence finally follows from Dini's theorem. \square

We now use Theorem A.2 (given for the continuous model) to prove Theorem 2.5. A preliminary definition and a lemma are needed.

Let define the modulus of continuity of H as

$$K_H(\delta) = \sup_{\substack{(s,t) \in [a,b]^2 \\ |s-t| \leq \delta}} |H_1 \circ H_2^{-1}(s) - H_1 \circ H_2^{-1}(t)| \quad (\delta \geq 0),$$

where H_1 and H_2 are two independent copies of H . Since for all $\delta \geq 0$, $0 \leq K_H(\delta) \leq b-a$ almost surely, we can define $L_H^1(H_2, \delta)$ and $L_H^2(H_2, \delta)$ as $L_H^1(H_2, \delta) = \mathbf{E}[K_H(\delta)|H_2]$ and $L_H^2(H_2, \delta) = \mathbf{E}[K_H^2(\delta)|H_2]$ ($\delta \geq 0$), and we have $L_H^1(H_2, 0) = L_H^2(H_2, 0) = 0$. From continuity of H , it holds that $L_H^1(H_2, \cdot)$ and $L_H^2(H_2, \cdot)$ are right-continuous at 0. Given $H_2 = h$, this implies that for $\epsilon > 0$, there exists $\eta_\epsilon > 0$ such that $L_H^1(h, \eta_\epsilon) < \epsilon$. We shall use this result in proving the following lemma:

Lemma A.3. Let $t \in [a, b]$. Under assumption (2), the following holds:

$$\sup_{t \in [a,b]} \left| \overline{\phi \circ h_{i_0}^{-1}} \left(t + \frac{1}{n} \right) - \overline{\phi \circ h_{i_0}^{-1}}(t) \right| \xrightarrow{m,n \rightarrow \infty} 0.$$

Proof. Letting $t \in [a, b]$, one easily shows that

$$\left| \overline{\phi \circ h_{i_0}^{-1}} \left(t + \frac{1}{n} \right) - \overline{\phi \circ h_{i_0}^{-1}}(t) \right| \leq \frac{1}{m-1} \sum_{\substack{i=1 \\ i \neq i_0}}^m \left| h_i \circ h_{i_0}^{-1} \left(t + \frac{1}{n} \right) - h_i \circ h_{i_0}^{-1}(t) \right|.$$

This in turn implies that

$$\begin{aligned} \sup_{t \in [a,b]} \left| \overline{\phi \circ h_{i_0}^{-1}} \left(t + \frac{1}{n} \right) - \overline{\phi \circ h_{i_0}^{-1}}(t) \right| &\leq \frac{1}{m-1} \sum_{\substack{i=1 \\ i \neq i_0}}^m \sup_{\substack{(s,t) \in [a,b]^2 \\ |s-t| \leq \frac{1}{n}}} |h_i \circ h_{i_0}^{-1}(s) - h_i \circ h_{i_0}^{-1}(t)| \\ &= \frac{1}{m-1} \sum_{\substack{i=1 \\ i \neq i_0}}^m \tilde{K}_H^i \left(\frac{1}{n} \right), \end{aligned}$$

where the $\tilde{K}_H^i \left(\frac{1}{n} \right)$ ($i \in \{1, \dots, m\} \setminus i_0$) are independent random variables distributed as $K_H \left(\frac{1}{n} \right) |H_2 = h_{i_0}$. Since $K_H(\cdot)$ is increasing, if $n \in \mathbb{N}$ is sufficiently large so that $\frac{1}{n} < \eta_\epsilon$, we get that

$$\sup_{t \in [a,b]} \left| \overline{\phi \circ h_{i_0}^{-1}} \left(t + \frac{1}{n} \right) - \overline{\phi \circ h_{i_0}^{-1}}(t) \right| \leq \frac{1}{m-1} \sum_{\substack{i=1 \\ i \neq i_0}}^m \tilde{K}_H^i(\eta_\epsilon).$$

By the law of large numbers, we get that almost surely,

$$0 \leq \limsup_{m,n} \sup_{t \in [a,b]} \left| \overline{\phi \circ h_{i_0}^{-1}} \left(t + \frac{1}{n} \right) - \overline{\phi \circ h_{i_0}^{-1}}(t) \right| \leq L_H^1(h_{i_0}, \eta_\epsilon) < \epsilon.$$

This holds for any $\epsilon > 0$, hence

$$\sup_{t \in [a,b]} \left| \overline{\phi \circ h_{i_0}^{-1}} \left(t + \frac{1}{n} \right) - \overline{\phi \circ h_{i_0}^{-1}}(t) \right| \xrightarrow[m,n \rightarrow \infty]{as} 0.$$

□

PROOF OF THEOREM 2.5 Let $t \in [a, b]$. For each $i \in \{1, \dots, m\}$, f_i is almost surely increasing, hence almost surely $f_i^{-1} \circ f_{i_0}(t_{j_0}) - \frac{1}{n} \leq T_i \leq f_i^{-1} \circ f_{i_0}(t_{j_0}) + \frac{1}{n}$. Recall that $t_{j_0} = \arg \min_{j \in \{1, \dots, n\}} |t_j - t|$, hence $t - \frac{1}{n} \leq t_{j_0} \leq t + \frac{1}{n}$. Thus, combining the above two inequalities, we get that almost surely $f_i^{-1} \circ f_{i_0}(t - \frac{1}{n}) - \frac{1}{n} \leq T_i \leq f_i^{-1} \circ f_{i_0}(t + \frac{1}{n}) + \frac{1}{n}$, from which we deduce:

$$\overline{\phi \circ h_{i_0}^{-1}} \left(t - \frac{1}{n} \right) - \frac{1}{n} \leq \widehat{\phi \circ h_{i_0}^{-1}}(t) \leq \overline{\phi \circ h_{i_0}^{-1}} \left(t + \frac{1}{n} \right) + \frac{1}{n}. \quad (22)$$

We shall focus on the upper bound in this inequality, since the same kind of argument holds for the lower bound. Straightforward calculation yields

$$\begin{aligned} & \sup_{t \in [a,b]} \left| \overline{\phi \circ h_{i_0}^{-1}} \left(t + \frac{1}{n} \right) + \frac{1}{n} - \phi \circ h_{i_0}^{-1}(t) \right| \\ & \leq \sup_{t \in [a,b]} \left| \overline{\phi \circ h_{i_0}^{-1}} \left(t + \frac{1}{n} \right) - \overline{\phi \circ h_{i_0}^{-1}}(t) \right| + \frac{1}{n} + \left\| \overline{\phi \circ h_{i_0}^{-1}} - \phi \circ h_{i_0}^{-1} \right\|_\infty, \end{aligned}$$

where the RHS of this inequality tends to 0 as m and n tend to infinity, by Lemma A.3 and Theorem A.2 in [2]. It follows that $\left\| \widehat{\phi \circ h_{i_0}^{-1}} - \phi \circ h_{i_0}^{-1} \right\|_\infty \xrightarrow[m,n \rightarrow \infty]{as} 0$.

We now turn to weak convergence. Assume that $n = m^{\frac{1}{2} + \alpha}$ ($\alpha > 0$). From (22), it holds

$$\begin{aligned} \left\| \sqrt{m} \left(\widehat{\phi \circ h_{i_0}^{-1}} - \overline{\phi \circ h_{i_0}^{-1}} \right) \right\|_\infty & \leq \sup_{t \in [a,b]} \left| \sqrt{m} \left(\overline{\phi \circ h_{i_0}^{-1}} \left(t + \frac{1}{n} \right) - \overline{\phi \circ h_{i_0}^{-1}}(t) \right) \right| \\ & \quad + \sup_{t \in [a,b]} \left| \sqrt{m} \left(\overline{\phi \circ h_{i_0}^{-1}} \left(t - \frac{1}{n} \right) - \overline{\phi \circ h_{i_0}^{-1}}(t) \right) \right| + \frac{2}{m^\alpha} \\ & \leq 2Z_m + \frac{2}{m^\alpha}, \end{aligned}$$

where $Z_m = \sqrt{m} \frac{1}{m-1} \sum_{\substack{i=1 \\ i \neq i_0}}^m \tilde{K}_H^i \left(\frac{1}{n} \right)$. Since

$$\text{var}(Z_m) = \frac{m}{m-1} \left(L_H^2 \left(h_{i_0}, \frac{1}{n} \right) - \left\{ L_H^1 \left(h_{i_0}, \frac{1}{n} \right) \right\}^2 \right) \xrightarrow{m \rightarrow \infty} 0$$

(recall that $n = m^{\frac{1}{2} + \alpha}$ and that $L_H^1(h_{i_0}, \cdot)$ and $L_H^2(h_{i_0}, \cdot)$ are right-continuous at 0), we get that

$$\left\| \sqrt{m} \left(\widehat{\phi \circ h_{i_0}^{-1}} - \overline{\phi \circ h_{i_0}^{-1}} \right) \right\|_\infty \xrightarrow[m \rightarrow \infty]{P} 0,$$

hence by Theorem A.2 and Theorem 4.1 in [2], $\sqrt{m}(\widehat{\phi \circ h_{i_0}^{-1}} - \phi \circ h_{i_0}^{-1}) \xrightarrow[m \rightarrow \infty]{\mathbf{D}} Z$.

PROOF OF LEMMA 2.6 Consider first the continuous model (18), and define conditionally to $F_{i_0} = f_{i_0}$

$$\overline{\gamma \circ h_{i_0}^{-1}} = \frac{1}{m-1} \sum_{\substack{i=1 \\ i \neq i_0}}^m (f_i^{-1} \circ f_{i_0})^2.$$

Using similar arguments as in proof of Theorem A.2, we get that

$$\left\| \overline{\gamma \circ h_{i_0}^{-1}} - \gamma \circ h_{i_0}^{-1} \right\|_{\infty} \xrightarrow[m \rightarrow \infty]{as} 0.$$

Now, from $f_i^{-1} \circ f_{i_0}(t - \frac{1}{n}) - \frac{1}{n} \leq T_i \leq f_i^{-1} \circ f_{i_0}(t + \frac{1}{n}) + \frac{1}{n}$, we get the following inequality

$$|T_i - f_i^{-1} \circ f_{i_0}(t)| \leq \left| f_i^{-1} \circ f_{i_0} \left(t - \frac{1}{n} \right) - f_i^{-1} \circ f_{i_0}(t) \right| + \left| f_i^{-1} \circ f_{i_0} \left(t + \frac{1}{n} \right) - f_i^{-1} \circ f_{i_0}(t) \right| + \frac{2}{n}.$$

Acting as in proof of Lemma 2.3, it is fairly straightforward to show that for $t \in [a, b]$, $\frac{1}{m-1} \sum_{\substack{i=1 \\ i \neq i_0}}^m T_i \cdot f_i^{-1} \circ f_{i_0}(t) \xrightarrow[m, n \rightarrow \infty]{as} \gamma \circ h_{i_0}^{-1}(t)$.

From the above inequality, we also obtain that

$$|T_i - f_i^{-1} \circ f_{i_0}(t)| \leq 2\tilde{K}_H^i \left(\frac{1}{n} \right) + \frac{2}{n}.$$

Summing over $i \in \{1, \dots, m\} \setminus i_0$, and then using convergence of $\frac{1}{m-1} \sum_{i \neq i_0} T_i \cdot f_i^{-1} \circ f_{i_0}(t)$, of $\overline{\gamma \circ h_{i_0}^{-1}}(t)$, together with right-continuity of $L_H^1(h_{i_0}, \cdot)$ and $L_H^2(h_{i_0}, \cdot)$ at 0, yield that $\widehat{\gamma \circ h_{i_0}^{-1}}(t) \xrightarrow[m, n \rightarrow \infty]{as} \gamma \circ h_{i_0}^{-1}(t)$. This and convergence of $\widehat{\phi \circ h_{i_0}^{-1}}(t)$ complete the proof of Lemma 2.6.

References

- [1] Jérémie Bigot. A scale-space approach with wavelets to singularity estimation. *ESAIM Probab. Stat.*, 9:143–164 (electronic), 2005.
- [2] P. Billingsley. *Convergence of probability measures*. John Wiley & Sons, New York, 1968.
- [3] J-M. Bouché and G. Saporta. *L'analyse des données. Que Sais-Je ?* Presses Universitaires de France, Paris, 1980.
- [4] Fabrice Gamboa, Jean-Michel Loubes, and Elie Maza. Semi-parametric estimation of shifts. *Electron. J. Stat.*, 1:616–640, 2007.
- [5] A. Kneip and T. Gasser. Statistical tools to analyze data representing a sample of curves. *Ann. Statist.*, 20(3):1266–1305, 1992.
- [6] A. Kneip, X. Li, K.B. MacGibbon, and J.O. Ramsay. Curve registration by local regression. *Canad. J. Statist.*, 28(1):19–29, 2000.
- [7] M. Ledoux and M. Talagrand. *Probability in Banach spaces*, volume 23 of *Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)]*. Springer-Verlag, Berlin, 1991. Isoperimetry and processes.

- [8] J.O. Ramsay and X. Li. Curve registration. *J. R. Stat. Soc. Ser. B Stat. Methodol.*, 60(2):351–363, 1998.
- [9] J.O. Ramsay and B.W. Silverman. *Applied functional data analysis*. Springer Series in Statistics. Springer-Verlag, New York, 2002. Methods and case studies.
- [10] W. Rudin. *Real and complex analysis*. McGraw-Hill Book Co., New York, third edition, 1987.
- [11] H. Sakoe and S. Chiba. Dynamic programming algorithm optimization for spoken word recognition. *IEEE Transactions, ASSP-26*, 1:43–49, 1978.
- [12] A.W. van der Vaart. *Asymptotic statistics*. Cambridge Series in Statistical and Probabilistic Mathematics. Cambridge University Press, Cambridge, 1998.
- [13] A.W. van der Vaart and J.A. Wellner. *Weak convergence and empirical processes*. Springer Series in Statistics. Springer-Verlag, New York, 1996.
- [14] K. Wang and T. Gasser. Synchronizing sample curves nonparametrically. *Ann. Statist.*, 27(2):439–460, 1999.