



## Boutet de Monvel's Calculus and Groupoids I

Johannes Aastrup, Severino T. Melo, Bertrand Monthubert, Elmar Schrohe

► **To cite this version:**

Johannes Aastrup, Severino T. Melo, Bertrand Monthubert, Elmar Schrohe. Boutet de Monvel's Calculus and Groupoids I. *J. Noncommut. Geom.*, 2010, 4 (3), pp.313-329. <hal-00113161>

**HAL Id: hal-00113161**

**<https://hal.archives-ouvertes.fr/hal-00113161>**

Submitted on 11 Nov 2006

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Boutet de Monvel's Calculus and Groupoids I

J. Aastrup, S. T. Melo\*, B. Monthubert & E. Schrohe

## Abstract

Can Boutet de Monvel's algebra on a compact manifold with boundary be obtained as the algebra  $\Psi^0(G)$  of pseudodifferential operators on some Lie groupoid  $G$ ? If it could, the kernel  $\mathcal{G}$  of the principal symbol homomorphism would be isomorphic to the groupoid  $C^*$ -algebra  $C^*(G)$ . While the answer to the above question remains open, we exhibit in this paper a groupoid  $G$  such that  $C^*(G)$  possesses an ideal  $\mathcal{I}$  isomorphic to  $\mathcal{G}$ . In fact, we prove first that  $\mathcal{G} \simeq \Psi \otimes \mathcal{K}$  with the  $C^*$ -algebra  $\Psi$  generated by the zero order pseudodifferential operators on the boundary and the algebra  $\mathcal{K}$  of compact operators. As both  $\Psi \otimes \mathcal{K}$  and  $\mathcal{I}$  are extensions of  $C(S^*Y) \otimes \mathcal{K}$  by  $\mathcal{K}$  ( $S^*Y$  is the co-sphere bundle over the boundary) we infer from a theorem by Voiculescu that both are isomorphic.

## Introduction

Boutet de Monvel's calculus [2, 3, 4, 10, 24, 28] is a pseudodifferential calculus on manifolds with boundary. It includes the classical differential boundary value problems as well as the parametrices to elliptic elements. Many operator-algebraic aspects of this algebra (spectral invariance, noncommutative residues and traces, composition sequence,  $K$ -theory) have been studied recently [9, 11, 18, 19, 22, 27]. The problem of identifying this algebra as the pseudodifferential algebra (or as an ideal of one) of a Lie groupoid may be the key to an effective application of the methods of noncommutative geometry. If that is accomplished, one could then seek for extensions of the calculus, and for a better understanding of its index theory [4, 8, 24]. Basic definitions and certain facts about Boutet de Monvel's algebra are recalled in Section 1.

The groupoid approach to pseudodifferential calculus was developed in noncommutative geometry, following the seminal work of A. Connes for foliations [7]. The guiding principle in that approach is that the central object in global analysis is a groupoid. To study a particular situation, for a class of singular manifolds for instance, one needs to define a groupoid adapted to the context and then use the general pseudodifferential tools for groupoids, as developed in [21, 23, 20, 16, 17]. This has been applied to the context of manifolds with corners, with the goal of studying Melrose's  $b$ -calculus (see [21, 23, 20]). Groupoids

---

\*S.T. Melo and B. Monthubert were partly funded by a cooperation agreement CAPES-COFECUB.

were defined whose pseudodifferential calculi recover the  $b$ -calculus and the cusp-calculus. Basic definitions and certain facts about pseudodifferential calculus on groupoids are recalled in Section 3.

The starting point of this work is the following result (see [18]): The kernel of the principal symbol map for Boutet de Monvel's calculus is equal to the norm closure  $\mathcal{G}$  of the ideal of singular Green operators. Since in the pseudodifferential calculus on a groupoid, the  $C^*$ -algebra of the groupoid is the kernel of the principal symbol map, this gives a hint that finding a groupoid whose  $C^*$ -algebra is  $\mathcal{G}$  could give some insight about the relationship between the Boutet de Monvel algebra and groupoid pseudodifferential algebras.

Besides,  $\mathcal{G}$  fits into a short exact sequence (see [19, Section 7]):

$$0 \rightarrow \mathcal{K} \rightarrow \mathcal{G} \rightarrow C(S^*Y) \otimes \mathcal{K} \rightarrow 0,$$

which is similar to that for pseudodifferential operators on smooth manifolds:

$$0 \rightarrow \mathcal{K} \rightarrow \Psi \rightarrow C(S^*Y) \rightarrow 0.$$

In Section 2, we show that  $\mathcal{G}$  is actually Morita-equivalent to the norm-closure  $\Psi$  of the algebra of pseudodifferential operators on the boundary. Since  $\mathcal{G}$  is a stable  $C^*$ -algebra, it is thus isomorphic to  $\Psi \otimes \mathcal{K}$ .

On the other hand, we define in Section 4 a groupoid whose  $C^*$ -algebra contains an ideal  $\mathcal{I}$  which fits in an extension analogous as that of  $\Psi \otimes \mathcal{K}$ . By showing in Section 5 that the  $KK$ -theory elements induced by these extensions coincide, we infer from a theorem by Voiculescu that  $\mathcal{G}$  and  $\mathcal{I}$  are isomorphic.

## 1 Boutet de Monvel's Calculus

Let  $X$  denote a compact manifold of dimension  $n$  with boundary  $Y$  and interior  $\dot{X}$ . Given a pseudodifferential operator  $P$ , defined on an open neighborhood  $\tilde{X}$  of  $X$ , and  $u \in C^\infty(X)$ , one defines  $P_+u$  as equal to the restriction to  $\dot{X}$  of  $PEu$ , where  $Eu$  is the extension of  $u$  to  $\tilde{X}$  which vanishes outside  $X$ . In general, singularities may develop at the boundary, and one gets only a mapping  $P_+ : C^\infty(X) \rightarrow C^\infty(\dot{X})$ . One says that  $P$  has the *transmission property* if the image of the truncated operator  $P_+$  is contained in  $C^\infty(X)$  (a subspace of  $C^\infty(\dot{X})$ ). This was defined by Boutet de Monvel in [2, 3], where he proved that the transmission property for a classical (polyhomogeneous) pseudodifferential operator is equivalent to certain symmetry conditions for the homogeneous components of the asymptotic expansion of its symbol at the boundary. Later [4], he constructed an algebra whose elements are operators of the form

$$\begin{pmatrix} P_+ + G & K \\ T & S \end{pmatrix} : \begin{array}{c} C^\infty(X) \\ \oplus \\ C^\infty(Y) \end{array} \longrightarrow \begin{array}{c} C^\infty(X) \\ \oplus \\ C^\infty(Y) \end{array}, \quad (1)$$

where  $P$  is a pseudodifferential operator on  $X$  satisfying a condition that ensures the transmission property,  $S$  is a pseudodifferential operator on  $Y$ , while  $G, K$

and  $T$  belong to classes of operators he then defined and named, respectively, singular Green, Poisson and trace operators. We call an operator as in (1) a Boutet de Monvel operator. For detailed expositions of his calculus, we refer to [10, 24].

A Boutet de Monvel operator has an *order*, roughly the usual order of pseudodifferential operators. The entries  $T$  and  $G$  have, moreover, an integer *class* assigned to them. The class of a trace operator is related to the order of the derivatives that appear in the boundary-value conditions it prescribes. One must assign a class also to singular Green operators due to the fact that the composition  $KT$  is a singular Green operator. There exist isomorphisms between suitable Sobolev spaces such that the composition of a given operator of arbitrary order and class with one of them has order and class zero. For index theory purposes it is therefore sufficient to consider operators of order and class zero. These form an adjoint invariant subalgebra of the algebra  $\mathcal{L}(\mathcal{H})$  of all bounded operators on a suitable Hilbert space  $\mathcal{H}$ . Adopting the definition of order in [24, 28] for  $K$  and  $T$ , we here choose  $\mathcal{H} = L^2(X) \oplus L^2(Y)$ . If, as does Grubb [10], one keeps the original definition (which makes more sense if one needs general  $L^p$  estimates) then one must take a Sobolev space of order  $-1/2$  over the boundary.

Boutet de Monvel operators can also be defined as mappings between smooth sections of vector bundles. If  $E$  is a bundle of positive rank over  $X$ , and  $F$  is an arbitrary bundle over  $Y$ , then the algebra of all Boutet de Monvel operators of order and class zero acting between sections of  $E$  and  $F$  is Morita equivalent [18, Section 1.5] to the algebra obtained by taking a rank-one trivial bundle over  $X$  and the zero-bundle over  $Y$ . This partly justifies, again if one is interested in index theory, to consider only the operators appearing in the upper-left corner of the matrix in (1) and to assume, as we did at the beginning, that the bundle over  $X$  is  $X \times \mathbb{C}$ .

The problem of computing the Fredholm index of a Boutet de Monvel operator acting between sections of different bundles over each side can be reduced to the case of equal bundles on both sides by a device developed by Boutet de Monvel [2, 3], recalled in [19, Section 1.1].

Let us now explain what a singular Green operator  $G$  is, in the case of order and class zero and of a rank-one trivial bundle over  $X$ . Its distribution kernel is smooth outside the boundary diagonal; i.e. if  $\varphi \in C_c^\infty(\dot{X})$ , and if we denote by  $M_\varphi$  the operator of multiplication by  $\varphi$ , then  $GM_\varphi$  and  $M_\varphi G$  are integral operators with smooth kernels. The push-forward of  $G$  by a boundary chart is an operator-valued-symbol pseudodifferential operator on the variables tangential to the boundary, as we describe below. It is perhaps worth stressing, however, that it is in general not a pseudodifferential operator on all variables, because of its particular way of acting on the normal variable.

Given  $u \in C_c^\infty(\mathbb{R}_+^n)$ ,  $\mathbb{R}_+^n = \{(x', x_n) \in \mathbb{R}^{n-1} \times \mathbb{R}; x_n \geq 0\}$ , let  $\hat{u}$  denote the vector-valued Fourier transform of  $u$  with respect to the  $n - 1$  first variables,

$$\hat{u}(\xi') = \int e^{ix' \cdot \xi'} u(x', \cdot) dx' \in C_c^\infty(\mathbb{R}_+). \quad (2)$$

In local coordinates for which the boundary corresponds to  $x_n = 0$  and the interior to  $x_n > 0$ ,  $G$  is given by

$$Gu(x', \cdot) = (2\pi)^{1-n} \int e^{ix' \cdot \xi'} g(x', \xi', D_n) \hat{u}(\xi') d\xi' \quad (3)$$

The integrals in (2) or in (3) should be regarded, for fixed  $\xi'$  or  $x'$ , respectively, as  $L^2(\mathbb{R}_+)$ -valued integrals. For each  $(x', \xi')$ ,  $g(x', \xi', D_n)$  in (3) is an integral operator with kernel  $\tilde{g}(x', \cdot, \cdot, \xi')$  equal to the restriction to  $\mathbb{R}_+ \times \mathbb{R}_+$  of a function belonging to the Schwartz space of rapidly decreasing functions on  $\mathbb{R}^2$ . The function  $\tilde{g}(x', x_n, y_n, \xi')$  (called by Grubb the *symbol-kernel* of  $G$ ) is smooth and satisfies the estimates [10, (1.2.38)]. This is invariantly defined [10, Theorem 2.4.11] with respect to coordinate changes that preserve the set  $\{x_n = 0\}$ .

We denote by  $\mathcal{A}_0$  the set of all polyhomogeneous operators  $P_+ + G$  of order and class zero on  $X$ , and by  $\mathcal{G}_0$  its subset of all singular Green operators. It follows from the rules of Boutet de Monvel's calculus that  $\mathcal{A}_0$  is an algebra and that  $\mathcal{G}_0$  is an ideal in  $\mathcal{A}_0$ .

In the sequel, we shall restrict ourselves to coordinate changes which preserve the variable  $x_n$ , i.e., we choose a normal coordinate. Then two \*-homomorphisms are defined on  $\mathcal{A}_0$ , the principal symbol and the boundary principal symbol:

$$\sigma : \mathcal{A}_0 \rightarrow C^\infty(S^*X) \quad \text{and} \quad \gamma : \mathcal{A}_0 \rightarrow C^\infty(S^*Y, \mathcal{L}(L^2(\mathbb{R}_+))).$$

The principal symbol of a given  $P_+ + G$  is, by definition, the usual principal symbol of  $P$

$$\sigma(P_+ + G) = p_0,$$

where  $p_0$  is the leading term in the asymptotic expansion of the symbol of  $P$ .

At a point  $(x', \xi')$  in  $S^*Y$ , the boundary principal symbol of  $P_+$  is defined to be the truncated Fourier multiplier

$$\gamma_{P_+}(x', \xi') = p_0(x', 0, \xi', D_n)_+$$

of symbol  $\xi_n \mapsto p_0(x', 0, \xi', \xi_n)$ . The boundary principal symbol of  $G \in \mathcal{G}_0$  is the integral operator

$$\gamma_G(x', \xi') = g_0(x', \xi', D_n) \quad (4)$$

with the rapidly decreasing kernel  $\tilde{g}_0(x', \cdot, \cdot, \xi')$ , where  $\tilde{g}_0$  denotes the leading term in the asymptotic expansion of  $\tilde{g}$ , cf. [10, (1.2.39)]. Then  $\gamma$  maps  $\mathcal{G}_0$  into  $C^\infty(S^*Y, \mathcal{K}_{\mathbb{R}_+})$ , with the ideal  $\mathcal{K}_{\mathbb{R}_+}$  of compact operators on  $L^2(\mathbb{R}_+)$ .

Let  $\mathcal{A}$  and  $\mathcal{G}$  denote the norm closures of  $\mathcal{A}_0$  and  $\mathcal{G}_0$ , respectively; and let  $\mathcal{K}_X$  denote the set of all compact operators on  $L^2(X)$ . It follows from Theorem 1 in [24, 2.3.4.4] that  $\sigma$  and  $\gamma$  can be extended to  $C^*$ -algebra homomorphisms

$$\bar{\sigma} : \mathcal{A} \rightarrow C(S^*X) \quad \text{and} \quad \bar{\gamma} : \mathcal{A} \rightarrow C(S^*Y, \mathcal{L}(L^2(\mathbb{R}_+))).$$

Moreover, by Corollary 2 in [24, 2.3.4.4] and [18, Theorems 5 and 6], we have that:

$$\ker \bar{\gamma} \cap \ker \bar{\sigma} = \mathcal{K}_X, \quad \ker \bar{\sigma} = \mathcal{G}, \quad (5)$$

and  $\bar{\gamma}$  maps  $\mathcal{G}$  onto  $C(S^*Y, \mathcal{K}_{\mathbb{R}_+})$ . In other words, the restriction of the boundary principal symbol to  $\mathcal{G}$  gives rise to the exact sequence of  $C^*$ -algebras

$$0 \longrightarrow \mathcal{K}_X \longrightarrow \mathcal{G} \xrightarrow{\bar{\gamma}} C(S^*Y, \mathcal{K}_{\mathbb{R}_+}) \longrightarrow 0. \quad (6)$$

In Section 2 we use (6) to prove that  $\mathcal{G}$  is isomorphic to the tensor product  $\Psi \otimes \mathcal{K}$  of the  $C^*$ -closure  $\Psi$  of the pseudodifferential operators of order zero on  $Y$  by the compacts. For that we need to use trace and Poisson operators.

Similarly as for the singular Green operators, the trace operators and the Poisson operators ( $T$  and  $K$  in (1)) are, locally, operator-valued-symbol pseudodifferential operators on the variables tangential to the boundary, given by

$$Tu(x') = (2\pi)^{1-n} \int e^{ix' \cdot \xi'} t(x', \xi', D_n) \hat{u}(\xi') \mathcal{d}\xi', \quad u \in C_c^\infty(\mathbb{R}_+^n), \quad (7)$$

and

$$Ku(x', \cdot) = (2\pi)^{1-n} \int e^{ix' \cdot \xi'} k(x', \xi', D_n) \hat{u}(\xi') \mathcal{d}\xi', \quad u \in C_c^\infty(\mathbb{R}^{n-1}). \quad (8)$$

The mappings  $t(x', \xi', D_n) : L^2(\mathbb{R}_+) \rightarrow \mathbb{C}$  and  $k(x', \xi', D_n) : \mathbb{C} \rightarrow L^2(\mathbb{R}_+)$  are defined, for each  $(x', \xi') \in \mathbb{R}^{n-1} \times \mathbb{R}^{n-1}$ , each  $v \in L^2(\mathbb{R}_+)$  and each  $\alpha \in \mathbb{C}$ , by

$$t(x', \xi', D_n)v = \int \tilde{t}(x', y_n, \xi') v(y_n) dy_n \quad (9)$$

and

$$[k(x', \xi', D_n)\alpha](x_n) = \alpha \tilde{k}(x', x_n, \xi'). \quad (10)$$

For each  $(x', \xi')$ ,  $\tilde{t}(x', \cdot, \xi')$  and  $\tilde{k}(x', \cdot, \xi')$  are restrictions to  $\mathbb{R}_+$  of functions in the Schwartz class on  $\mathbb{R}$ . The functions  $\tilde{t}(x', y_n, \xi')$  and  $\tilde{k}(x', x_n, \xi')$ , called the symbol-kernels of  $T$  and  $K$ , are smooth and satisfy certain estimates. In the polyhomogenous case, they have asymptotic expansions in homogeneous components, whose leading terms we denote by  $\tilde{t}_0$  and  $\tilde{k}_0$ , respectively. The estimates and expansions for  $\tilde{t}$  and  $\tilde{k}$  listed or explained in [10, Section 1.2] are not the right ones for our definition of order (and consequent choice of Hilbert space): we need to shift some of the exponents there by  $\pm 1/2$ .

The boundary-principal symbols of  $T$  and  $K$  are

$$\gamma_T(x', \xi') = t_0(x', \xi', D_n) \quad \text{and} \quad \gamma_K(x', \xi') = k_0(x', \xi', D_n),$$

defined as in (9) and (10), except that  $\tilde{t}_0$  and  $\tilde{k}_0$  replace  $\tilde{t}$  and  $\tilde{k}$ . Lastly, the boundary principal symbol of a polyhomogeneous pseudodifferential operator on  $Y$  is simply its usual principal symbol, and we get a \*-homomorphism

$$\gamma : \mathcal{B}_0 \longrightarrow C^\infty(S^*Y, \mathcal{L}(L^2(\mathbb{R}_+) \oplus \mathbb{C})),$$

where  $\mathcal{B}_0$  denotes the set of all polyhomogeneous Boutet de Monvel operators of order and class zero on  $X$ . It has a continuous extension to the norm-closure of  $\mathcal{B}_0$ , but we will not use this fact.

## 2 A Product Description of the Singular Green Operators

**Lemma 1.** *There exists a zero-order Poisson operator  $K$  such that  $K^*K$  is a strictly positive operator on  $L^2(Y)$ .*

*Proof.* It is well-known that the Dirichlet problem

$$\begin{pmatrix} \Delta \\ \gamma_0 \end{pmatrix} : H^2(X) \longrightarrow \begin{matrix} L^2(X) \\ \oplus \\ H^{3/2}(Y) \end{matrix}$$

defines a bounded invertible operator. We denote by  $\lambda^{3/2}$  an order reduction of order  $3/2$  on  $Y$  and by  $\Lambda^{-2}$  and order reduction of order  $-2$  on  $X$ . This gives us an isomorphism

$$\begin{pmatrix} \Delta \Lambda^{-2} \\ \lambda^{3/2} \gamma_0 \Lambda^{-2} \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & \lambda^{3/2} \end{pmatrix} \begin{pmatrix} \Delta \\ \gamma_0 \end{pmatrix} \Lambda^{-2} : L^2(X) \longrightarrow \begin{matrix} L^2(X) \\ \oplus \\ L^2(Y) \end{matrix}$$

which is an element of order and class 0 in Boutet de Monvel's calculus. Its inverse therefore also is in Boutet de Monvel's calculus; it is of the form

$$(P_+ + G \ K) : \begin{matrix} L^2(X) \\ \oplus \\ L^2(Y) \end{matrix} \longrightarrow L^2(X)$$

with suitable  $P, G$ , and  $K$  of order and class zero. In particular,  $K$  is a right inverse for the trace operator  $T = \lambda^{3/2} \gamma_0 \Lambda^{-2} : L^2(X) \rightarrow L^2(Y)$ . For  $v \in L^2(Y)$  we thus have

$$\|v\|_{L^2(Y)} = \|TKv\|_{L^2(Y)} \leq \|T\|_{\mathcal{L}(L^2(X), L^2(Y))} \|Kv\|_{L^2(X)}.$$

We then get  $\|Kv\| \geq c\|v\|$  for some  $c > 0$ , so that  $K^*K$  is strictly positive.  $\square$

**Lemma 2.** *There exist a trace operator of order and class zero  $T : L^2(X) \rightarrow L^2(Y)$  and a Poisson operator of order zero  $K : L^2(Y) \rightarrow L^2(X)$  such that  $TK$  is equal to the identity operator on  $L^2(Y)$ ,  $K^* = T$  and  $T^* = K$ .*

*Proof.* Let  $K_0$  be a zero-order Poisson operator such that  $K_0^*K_0$  is a strictly positive operator on  $L^2(Y)$ , and let  $Q = (K_0^*K_0)^{-1/2}$ .  $Q$  is a zero-order pseudodifferential operator on  $Y$ . Take  $K = K_0Q$  and  $T = QK_0^*$ .  $\square$

We denote by  $\Psi$  the norm closure of the algebra of all polyhomogeneous pseudodifferential operators of order zero on  $Y$ , and by  $\bar{\sigma} : \Psi \rightarrow C(S^*Y)$  the continuous extension of the principal-symbol homomorphism. It is well-known (this is mentioned in [1] and follows from [14, Theorem A.4], or from [12, Theorem 3.3]) that  $\bar{\sigma}$  induces the short exact sequence of  $C^*$ -algebras

$$0 \longrightarrow \mathcal{K}_Y \longrightarrow \Psi \xrightarrow{\bar{\sigma}} C(S^*Y) \longrightarrow 0, \quad (11)$$

where  $\mathcal{K}_Y$  denotes the ideal of compact operators on  $L^2(Y)$ .

By Lemma 2 a  $C^*$ -homomorphism  $\Xi : \Psi \rightarrow \mathcal{G}$  can be defined by

$$\Xi(A) = KAT.$$

Since  $\Xi(A)$  is compact if  $A$  is compact, we can use  $\Xi$  to couple the sequences (6) and (11). Together they yield the commutative diagram of exact sequences of  $C^*$ -algebras

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathcal{K}_X & \longrightarrow & \mathcal{G} & \xrightarrow{\bar{\gamma}} & C(S^*Y, \mathcal{K}_{\mathbb{R}_+}) & \longrightarrow & 0 \\ & & \uparrow & & \uparrow \Xi & & \uparrow h & & \\ 0 & \longrightarrow & \mathcal{K}_Y & \longrightarrow & \Psi & \xrightarrow{\bar{\sigma}} & C(S^*Y) & \longrightarrow & 0 \end{array} \quad (12)$$

**Lemma 3.** *The homomorphism  $\Xi$  imbeds  $\Psi$  as a hereditary subalgebra of  $\mathcal{G}$ .*

*Proof:* We have to prove, that if  $0 \leq G \leq KAT$  then  $G$  is again of the form  $KA_1T$  with  $A_1 \in \Psi$ . Since  $KT$  acts as the identity on  $KAT$  it also acts as the identity on  $G$  and we therefore get  $G = KTGKT = K(TGK)T$ .  $\square$

**Lemma 4.** *Let*

$$\begin{array}{ccccccc} 0 & \longrightarrow & I_1 & \longrightarrow & A_1 & \xrightarrow{q_1} & B_1 & \longrightarrow & 0 \\ & & \uparrow \phi_1 & & \uparrow \phi_2 & & \uparrow \phi_3 & & \\ 0 & \longrightarrow & I_2 & \longrightarrow & A_2 & \xrightarrow{q_2} & B_2 & \longrightarrow & 0 \end{array}$$

*be a commutative diagram of short exact sequences, where  $\phi_1, \phi_2$  and  $\phi_3$  are embeddings. Then  $\phi_2$  is full provided that  $\phi_1$  and  $\phi_3$  are full.*

*Proof:* We have to prove that the two-sided ideal generated by  $\phi_2(A_2)$  is dense in  $A_1$ . We thus have to prove that to a given  $a \in A_1$  and a given  $\varepsilon > 0$  we can find an element  $b$  in the twosided ideal generated by the image of  $\phi_2$  such that  $\|a - b\| < \varepsilon$ . Since  $\phi_3$  is full we can find an element  $c$  in the twosided ideal generated by  $\phi_3(B_2)$  such that  $\|q_1(a) - c\| < \frac{\varepsilon}{2}$ . The element  $c$  can be lifted to an element  $b_1$  in the twosided ideal generated by  $\phi_2(A_2)$  and we can therefore find an element  $d_1 \in I_1$  such that  $\|a - b_1 - d_1\| < \frac{\varepsilon}{2}$ . Since  $\phi_1$  is full there is an element  $d_2$  in the twosided ideal generated by  $\phi_1(I_2)$  with  $\|d_1 - d_2\| < \frac{\varepsilon}{2}$ . As the desired  $b$  we can therefore choose  $b = b_1 + d_2$ .  $\square$

**Theorem 1.** *The algebras  $\mathcal{G}$  and  $\Psi \otimes \mathcal{K}$  are isomorphic.*

*Proof:* By Lemma 3, the diagram 12 and Lemma 4 the imbedding  $\Xi$  is full and hereditary. It follows from the remark below Theorem 8 on page 155 in [7] that  $\mathcal{G}$  and  $\Psi$  are strongly Morita equivalent. By the results in [5] and [6] we have  $\mathcal{G} \otimes \mathcal{K}$  is isomorphic to  $\Psi \otimes \mathcal{K}$ . However  $\mathcal{G}$  is stable since it is the extension of  $\mathcal{K}$  with a stable algebra, namely  $C(S^*Y, \mathcal{K}_{\mathbb{R}_+})$ , (see Proposition 6.12 in [26]). This gives the isomorphism.  $\square$




### 3 Pseudodifferential Operators and Groupoids

Groupoids were introduced in the context of global analysis when A. Connes showed that in the case of foliations the index takes values in a  $C^*$ -algebra which is defined as the  $C^*$ -algebra of the holonomy groupoid of the foliation. He defined a pseudodifferential calculus on a foliation using the groupoid structure.

In several papers ([21, 23, 16, 20]), generalizations of this approach to a larger class of groupoids were achieved. One particular aspect of this theory is that, as A. Connes showed in [7] for smooth manifolds, it is possible to define the analytic index using a groupoid, the *tangent groupoid*.

A groupoid is a small category in which all morphisms are invertible. This means that a groupoid  $G$  has a set of units, denoted by  $G^{(0)}$ , and two maps called *range* and *source*,  $G \begin{smallmatrix} \xrightarrow{r} \\ \xleftarrow{s} \end{smallmatrix} G^{(0)}$ .

Two elements  $\gamma, \gamma' \in G$  are composable if and only if  $r(\gamma) = s(\gamma')$ :



We recall briefly the main aspects of this theory. Let  $G$  be a Lie groupoid, which means that it has a smooth structure. Then one can define an algebra of pseudodifferential operators  $\Psi^\infty(G)$ : A pseudodifferential operator on  $G$  is a  $G$ -equivariant continuous family of pseudodifferential operators on the fibers of  $G$ .

For example, if  $M$  is a manifold without boundary, and  $G = M \times M$ , with set of units  $G^{(0)} = M$ , and range and source maps  $r(x, y) = x$ ,  $s(x, y) = y$ , and composition  $(x, y)(y, z) = (x, z)$ , then  $\Psi^\infty(G)$  is the algebra of pseudodifferential operators on  $M$ .

If  $G$  is a Lie group,  $\Psi^\infty(G)$  is the algebra of  $G$ -equivariant pseudodifferential operators on  $G$ .

In order to work with singular manifolds, the framework of Lie groupoids needs to be extended. That was done in [16], where the algebras of pseudodifferential operators on continuous family groupoids, which are groupoids whose fibers are smooth manifolds, were defined.

On the algebra of pseudodifferential operators one can define a symbol map,  $\sigma$ . The algebra of order 0 operators can be completed as a  $C^*$ -algebra, denoted by  $\overline{\Psi^0(G)}$ , and the symbol map extends to this algebra. The “regularizing operators” of the calculus, which are the operators with trivial symbol, are the elements of the  $C^*$ -algebra of the groupoid, and we have the following Atiyah-

Singer exact sequence:

$$0 \rightarrow C^*(G) \rightarrow \overline{\Psi^0}(G) \rightarrow C(S^*(G)) \rightarrow 0,$$

where  $S^*(G)$  is the cosphere bundle of the Lie algebroid  $A(G)$ , which can be thought of as a tangent space.

We next recall in more detail the construction of the adiabatic groupoid  $^{ad}(Y \times Y)$  associated with a smooth manifold  $Y$ :

$$^{ad}(Y \times Y) = (TY \times \{0\}) \cup (Y \times Y \times \mathbb{R}_+^*)$$

with the tangent bundle  $TY$  of  $Y$ . The groupoid structure is given as follows:

$$r(x, \xi, 0) = s(x, \xi, 0) = x, \quad (x, \xi, 0) \circ (x, \xi', 0) = (x, \xi + \xi', 0),$$

$$r(x, y, \lambda) = (x, \lambda), \quad s(x, y, \lambda) = (y, \lambda), \quad (x, y, \lambda) \circ (y, z, \lambda) = (x, z, \lambda), \quad \lambda > 0.$$

This groupoid is endowed with a differential structure, through an exponential, in the following way:

- On  $Y \times Y \times \mathbb{R}_+^*$ , the structure is that of a product of manifolds.
- Define a map on an open neighborhood  $U$  of  $TY \times \{0\}$  in  $TY \times \mathbb{R}_+$ , with values in  $^{ad}(Y \times Y)$ , by

$$\begin{cases} \psi(x, \xi, \lambda) = (x, \exp_x(-\lambda\xi), \lambda) & \text{if } \lambda > 0 \\ \psi(x, \xi, 0) = (x, \xi, 0). \end{cases}$$

In other terms, the topology is such that a sequence of terms  $(x_n, y_n, \lambda_n)$  of  $Y \times Y \times \mathbb{R}_+^*$  converges to  $(x, \xi, 0) \in TY \times \{0\}$ , if and only if we have locally

$$x_n \rightarrow x, \quad y_n \rightarrow x, \quad \frac{x_n - y_n}{\lambda_n} \rightarrow \xi.$$

Note that A. Connes' tangent groupoid is just the restriction of  $^{ad}(Y \times Y)$  to  $\lambda \in [0, 1]$ .

The main interest of this groupoid is that it provides a way to define the analytic index. Consider indeed the decomposition of the groupoid as an open and a closed subgroupoid, which gives rise to the exact sequence:

$$0 \rightarrow C^*(Y \times Y \times \mathbb{R}_+^*) \rightarrow C^*(^{ad}(Y \times Y)) \rightarrow C^*(TY) \rightarrow 0. \quad (13)$$

This simplifies since  $C^*(Y \times Y) \simeq \mathcal{K}$ , and  $C^*(TY) \simeq C_0(T^*Y)$ . A. Connes proved that the boundary map of the 6-terms exact sequence induced by this extension is nothing but the analytic index

$$ind_a : K_c^0(T^*Y) \rightarrow K_1(C_0(\mathbb{R}) \otimes \mathcal{K}) = \mathbb{Z}.$$

## 4 A Groupoid Associated to the Singular Green Operators

Suppose we could identify the  $C^*$ -closure of Boutet de Monvel's algebra with the  $C^*$ -algebra  $\Psi^0(G)$  of pseudodifferential operators on a Lie groupoid  $G$ . Then, as pointed out above, the kernel of the principal symbol map would be isomorphic to  $C^*(G)$ . As the kernel of the principal symbol map in Boutet de Monvel's calculus consists of the singular Green operators, we thus wish to identify these with the  $C^*$ -algebra of a groupoid.

We will actually not identify them with a groupoid  $C^*$ -algebra, but with an ideal in a groupoid  $C^*$ -algebra.

Let us consider the following action of the group  $\mathbb{R}_+^*$  on  $ad(Y \times Y)$ :

- On  $TY$ ,  $\mathbb{R}_+^*$  acts by dilations:  $\lambda.(x, \xi) = (x, \lambda\xi)$
- On  $Y \times Y \times \mathbb{R}_+^*$ ,  $\mathbb{R}_+^*$  acts by  $\lambda.(x, y, t) = (x, y, \frac{t}{\lambda})$ .

This is a continuous action: If  $(x_n, y_n, t_n)$  converges to  $(x, \xi)$  (which means that  $x_n, y_n \rightarrow x, t_n \rightarrow 0, \frac{x_n - y_n}{t_n} \rightarrow \xi$ ), then  $\lambda.(x_n, y_n, t_n) = (x_n, y_n, \frac{t_n}{\lambda}) \rightarrow (x, \lambda\xi, 0)$ , since

$$\frac{x_n - y_n}{\frac{t_n}{\lambda}} \rightarrow \lambda\xi.$$

It is thus possible to construct the semi-direct product  $G = ad(Y \times Y) \rtimes \mathbb{R}_+^*$  of the adiabatic groupoid by  $\mathbb{R}_+^*$ : As a set, it is  $ad(Y \times Y) \times \mathbb{R}_+^*$ , with set of units  $Y \times \mathbb{R}_+^*$ , such that:

- $r(x, y, t, \lambda) = (x, t), s(x, y, t, \lambda) = (y, \frac{t}{\lambda})$ , for  $t > 0$ ;
- $r(x, \xi, \lambda) = (x, 0), s(x, \xi, \lambda) = (x, 0)$ , for  $t = 0$ ;
- $(x, y, t, \lambda)(y, z, \frac{t}{\lambda}, \mu) = (x, z, t, \lambda\mu)$ ;
- $(x, \xi, \lambda)(x, \eta, \mu) = (x, \xi + \lambda\eta, \lambda\mu)$ .

Note that the action of  $\mathbb{R}_+^*$  on the adiabatic groupoid induces an action on its  $C^*$ -algebra and that J. Renault proved in [25] that for any locally compact groupoid  $\mathcal{G}$  one has

$$C^*(\mathcal{G} \rtimes \mathbb{R}_+^*) \simeq C^*(\mathcal{G}) \rtimes \mathbb{R}_+^*.$$

The evaluation at  $t = 0$  provides a map  $e_0 : C^*(G) \rightarrow C_0(T^*Y) \rtimes \mathbb{R}_+^*$ . Also, the evaluation at the zero-section  $\xi = 0$  induces a map  $r_0 : C_0(T^*Y) \rtimes \mathbb{R}_+^* \rightarrow C(Y) \rtimes \mathbb{R}_+^*$ .

But since the action of  $\mathbb{R}_+^*$  on  $Y$  is trivial, the latter algebra is just the algebra of the (regular) product:

$$C(Y) \rtimes \mathbb{R}_+^* = C_0(Y \times \mathbb{R}_+^*).$$

Let  $C = \ker r_0$  and  $\mathcal{I} = \ker r_0 \circ e_0$ .

The kernel of  $e_0$  is  $C^*(Y \times Y \times \mathbb{R}_+^* \rtimes \mathbb{R}_+^*)$ . But  $\mathbb{R}_+^* \rtimes \mathbb{R}_+^*$  is directly isomorphic to the pair groupoid  $\mathbb{R}_+^* \times \mathbb{R}_+^*$ : To clarify the proof, let us denote  $G_1 = \mathbb{R}_+^* \rtimes \mathbb{R}_+^*$  and  $G_2 = \mathbb{R}_+^* \times \mathbb{R}_+^*$ . Then let  $\phi : G_1 \rightarrow G_2$  be defined by

$$\phi(t, \lambda) = \left( t, \frac{t}{\lambda} \right).$$

This a morphism of groupoids: The composition of  $(t, \lambda)$  with  $(\frac{t}{\lambda}, \mu)$  gives  $(t, \lambda\mu)$ , and

$$\phi(t, \lambda\mu) = \left( t, \frac{t}{\lambda\mu} \right)$$

while

$$\phi(t, \lambda) \circ \phi\left(\frac{t}{\lambda}, \mu\right) = \left( t, \frac{t}{\lambda} \right) \circ \left( \frac{t}{\lambda}, \frac{t}{\lambda\mu} \right) = \left( t, \frac{t}{\lambda\mu} \right).$$

Hence the kernel of  $e_0$  is just the algebra of compact operators,  $\mathcal{K}$ .

To make this clear, here is the commutative diagram describing this:

$$\begin{array}{ccccccc}
 & & & 0 & & 0 & \\
 & & & \downarrow & & \downarrow & \\
 0 & \longrightarrow & \mathcal{K} & \longrightarrow & \mathcal{I} & \longrightarrow & C & \longrightarrow & 0 \\
 & & \parallel & & \downarrow & & \downarrow j & & \\
 0 & \longrightarrow & \mathcal{K} & \longrightarrow & C^*(G) & \xrightarrow{e_0} & C_0(T^*Y) \rtimes \mathbb{R}_+^* & \longrightarrow & 0 \\
 & & & & \searrow & & \downarrow r_0 & & \\
 & & & & & & C_0(Y \times \mathbb{R}_+^*) & & 
 \end{array} \tag{14}$$

We will use this diagram and extension theory to prove that  $\mathcal{I}$  is isomorphic to the algebra of singular Green operators.

**Proposition 1.**  $C$  is isomorphic to  $C(S^*Y) \otimes \mathcal{K}$ , where  $S^*Y$  is the sphere bundle in  $T^*Y$ .

*Proof.* First of all notice that  $C$  is isomorphic to  $C_0(T^*Y \setminus Y) \rtimes \mathbb{R}_+^*$ : Indeed, the exact sequence

$$0 \rightarrow C_0(T^*Y \setminus Y) \rightarrow C_0(T^*Y) \rightarrow C(Y) \rightarrow 0$$

induces the exact sequence

$$0 \rightarrow C_0(T^*Y \setminus Y) \rtimes \mathbb{R}_+^* \rightarrow C_0(T^*Y) \rtimes \mathbb{R}_+^* \rightarrow C(Y) \rtimes \mathbb{R}_+^* = C_0(Y \times \mathbb{R}_+^*) \rightarrow 0.$$

But  $T^*Y \setminus Y \simeq S^*Y \times \mathbb{R}_+^*$ , so that  $C_0(T^*Y \setminus Y) \simeq C(S^*Y) \otimes C_0(\mathbb{R}_+^*)$  and

$$C_0(T^*Y \setminus Y) \rtimes \mathbb{R}_+^* \simeq C(S^*Y) \otimes C_0(\mathbb{R}_+^*) \rtimes \mathbb{R}_+^*.$$

Now

$$C_0(\mathbb{R}_+^*) \rtimes \mathbb{R}_+^* \simeq C^*(\mathbb{R}_+^*) \rtimes \mathbb{R}_+^* \simeq C^*(\mathbb{R}_+^* \times \mathbb{R}_+^*) \simeq C^*(\mathbb{R}_+^* \times \mathbb{R}_+^*) \simeq \mathcal{K}$$

where we again used Renault's result for the second isomorphism and the isomorphism of  $\mathbb{R}_+^* \times \mathbb{R}_+^*$  with the pair groupoid  $\mathbb{R}_+^* \times \mathbb{R}_+^*$  for the third. This ends the proof.  $\square$

## 5 Identification of the Ideal with the Singular Green Operators

We have just shown that  $\mathcal{I}$  is an extension of  $C_0(S^*Y) \otimes \mathcal{K}$  by  $\mathcal{K}$ , and this is also the case for the algebra of singular Green operators. The main result is the following:

**Theorem 2.** *The  $C^*$ -algebra  $\mathcal{I}$  is isomorphic to  $\Psi \otimes \mathcal{K}$ .*

*Proof.* We shall prove that the extensions

$$0 \rightarrow \mathcal{K} \rightarrow \mathcal{I} \rightarrow C(S^*Y) \otimes \mathcal{K} \rightarrow 0$$

and

$$0 \rightarrow \mathcal{K} \rightarrow \Psi \otimes \mathcal{K} \rightarrow C(S^*Y) \otimes \mathcal{K} \rightarrow 0$$

satisfy the conditions of a theorem by Voiculescu which we recall now.

D. Voiculescu proved in [30] (look also at the survey [29], Theorem 10.9) that if two extensions  $0 \rightarrow \mathcal{K} \rightarrow D_1 \rightarrow A \rightarrow 0$  and  $0 \rightarrow \mathcal{K} \rightarrow D_2 \rightarrow A \rightarrow 0$ , are such that:

- $D_1$  and  $D_2$  define the same element in  $Ext(A)$ ,
- $D_1$  and  $D_2$  are not unital,
- $\mathcal{K}$  is essential in  $D_1$  and in  $D_2$ ,

then  $D_1$  and  $D_2$  are isomorphic. Recall that an ideal  $J$  of  $D$  is essential if and only if for every  $x \in D$ ,  $x \neq 0 \Rightarrow \exists y \in J, xy \neq 0$ .

Let us apply this important result in our context.

First of all,  $\mathcal{I}$  and  $\Psi \otimes \mathcal{K}$  are non-unital since their quotients by the compacts are isomorphic to  $C(S^*Y) \otimes \mathcal{K}$ , which is non-unital.

Since the algebra of compact operators on a Hilbert space  $\mathcal{H}$  is essential in any  $C^*$ -algebra included in  $\mathcal{L}(\mathcal{H})$ , we obtain that  $\mathcal{K}_X$  is essential in  $\Psi \otimes \mathcal{K}_{\mathbb{R}_+^*}$ .

For the other algebra, notice that  $C^*(Y \times Y \times \mathbb{R}_+^*)$  is an essential ideal of  $C^*(ad(Y \times Y))$ , thus its crossed product by  $\mathbb{R}_+^*$  is also an essential ideal of  $C^*(ad(Y \times Y)) \rtimes \mathbb{R}_+^*$ , hence of  $\mathcal{I}$  (see [15]).

It remains to show that the extensions give rise to the same element of  $Ext(C(S^*Y) \otimes \mathcal{K})$ . But since  $C(S^*Y) \otimes \mathcal{K}$  is separable,  $KK_1(C(S^*Y) \otimes \mathcal{K}, \mathbb{C})$  is

isomorphic to the group of invertibles of  $Ext(C(S^*Y) \otimes \mathcal{K})$ , thanks to a result of Kasparov ([13]). The  $C^*$ -algebra  $C(S^*Y) \otimes \mathcal{K}$  being nuclear,  $Ext(C(S^*Y) \otimes \mathcal{K})$  is actually a group, thus it is isomorphic to  $KK_1(C(S^*Y) \otimes \mathcal{K}, \mathbb{C})$ .

The element  $i_S \in KK_1(C(S^*Y) \otimes \mathcal{K}, \mathbb{C}) \simeq Ext(C(S^*Y) \otimes \mathcal{K})$  associated to the extension

$$0 \rightarrow \mathcal{K}_X \rightarrow \Psi \otimes \mathcal{K}_{\mathbb{R}_+} \rightarrow C(S^*Y) \otimes \mathcal{K}_{\mathbb{R}_+} \rightarrow 0$$

provides a map  $KK_1(C(S^*Y) \otimes \mathcal{K}_{\mathbb{R}_+}) \simeq KK_1(C(S^*Y)) \rightarrow K_0(\mathbb{C}) = \mathbb{Z}$ , which is the analytic index.

For the class of the extension

$$0 \rightarrow \mathcal{K} \rightarrow \mathcal{I} \rightarrow C(S^*Y) \otimes \mathcal{K} \rightarrow 0,$$

let us consider first the extension

$$0 \rightarrow C^*(Y \times Y \times \mathbb{R}_+^*) \rightarrow C^*(ad(Y \times Y)) \rightarrow C_0(T^*Y) \rightarrow 0,$$

whose class is denoted by  $i_T \in KK_1(C_0(T^*Y), \mathcal{K} \otimes C_0(\mathbb{R}_+^*)) \simeq KK_0(C_0(T^*Y), \mathcal{K})$ .

It induces the extension

$$0 \rightarrow C^*(Y \times Y \times \mathbb{R}_+^*) \rtimes \mathbb{R}_+^* \simeq \mathcal{K} \rightarrow C^*(ad(Y \times Y)) \rtimes \mathbb{R}_+^* \simeq C^*(G) \rightarrow C_0(T^*Y) \rtimes \mathbb{R}_+^* \rightarrow 0$$

whose class is denoted by  $\alpha \in KK_1(C_0(T^*Y) \rtimes \mathbb{R}_+^*, \mathcal{K})$ .

The relation between  $i_T$  and  $i_S$  is made clear by considering the following exact sequence

$$0 \rightarrow C_0(T^*Y) \rightarrow C(B^*Y) \rightarrow C(S^*Y) \rightarrow 0$$

where  $B^*Y$  is the ball bundle over  $Y$ . Its class is denoted by  $\psi \in KK_1(C(S^*Y), C_0(T^*Y))$ , and one has the well-known equality:

$$i_S = \psi i_T.$$

For the convenience of the reader, we now recall the diagram (14):

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathcal{K} & \longrightarrow & \mathcal{I} & \longrightarrow & C & \longrightarrow & 0 & (\partial) \\ & & \parallel & & \downarrow & & \downarrow j & & & \\ 0 & \longrightarrow & \mathcal{K} & \longrightarrow & C^*(G) & \xrightarrow{e_0} & C_0(T^*Y) \rtimes \mathbb{R}_+^* & \longrightarrow & 0 & (\alpha) \\ & & & & \searrow & & \downarrow r_0 & & & \\ & & & & & & C_0(Y \times \mathbb{R}_+^*) & & & \end{array}$$

Let us denote the class of the first sequence by  $\partial \in KK_1(C, \mathcal{K})$ ; it is thus given by the Kasparov product:

$$\partial = j_* \alpha.$$

Let us make the relation between  $j_*$  and  $\varphi$  precise.  
 Consider the following commutative diagram:

$$\begin{array}{ccccccc}
 0 & \longrightarrow & C_0(T^*Y \setminus Y) & \longrightarrow & C_0(B^*Y \setminus Y) & \longrightarrow & C(S^*Y) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \parallel \\
 0 & \longrightarrow & C_0(T^*Y) & \longrightarrow & C(B^*Y) & \longrightarrow & C(S^*Y) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \\
 & & C(Y) & \xlongequal{\quad} & C(Y) & & 
 \end{array}$$

The first exact sequence actually decomposes as

$$0 \rightarrow C(S^*Y) \otimes C_0(\mathbb{R}_+^*) \rightarrow C(S^*Y) \otimes C_0(\mathbb{R}_+^* \cup \{\infty\}) \rightarrow C(S^*Y) \rightarrow 0$$

so that its  $KK_1$ -class is the identity of  $KK_1(C(S^*Y), C(S^*Y) \otimes C_0(\mathbb{R}_+^*))$ .

There is an action of  $\mathbb{R}_+^*$  on each algebra of the previous diagram, which is trivial on  $C(S^*Y)$  and  $C(Y)$ . This gives the following:

$$\begin{array}{ccccccc}
 0 & \longrightarrow & C_0(T^*Y \setminus Y) \rtimes \mathbb{R}_+^* & \longrightarrow & C_0(B^*Y \setminus Y) \rtimes \mathbb{R}_+^* & \longrightarrow & C(S^*Y \times \mathbb{R}_+^*) \longrightarrow 0 & (\partial_1) \\
 & & \downarrow j & & \downarrow & & \parallel \\
 0 & \longrightarrow & C_0(T^*Y) \rtimes \mathbb{R}_+^* & \longrightarrow & C(B^*Y) \rtimes \mathbb{R}_+^* & \longrightarrow & C(S^*Y \times \mathbb{R}_+^*) \longrightarrow 0 & (\varphi) \\
 & & \downarrow & & \downarrow & & \\
 & & C(Y \times \mathbb{R}_+^*) & \xlongequal{\quad} & C(Y \times \mathbb{R}_+^*) & & 
 \end{array}$$

Denote by  $\partial_1$  (resp.  $\varphi$ ) the class of the first (resp. second) exact sequence of this diagram, and by  $j_* \in KK(C_0(T^*Y \setminus Y), C_0(T^*Y))$  the element induced by  $C_0(T^*Y \setminus Y) \rightarrow C_0(T^*Y)$ . One has thus the equality

$$\varphi = \partial_1 j_*,$$

so that

$$\partial = j_* \alpha = \partial_1^{-1} \varphi \alpha.$$

But  $\varphi \alpha$  is the image of  $\psi i_T$  under the Thom-Connes isomorphism, and  $\partial_1$  is also a Thom-Connes element in  $KK$ -theory. Hence the classes in  $KK_1$  of the extensions of  $\mathcal{I}$  and of  $\Psi \otimes \mathcal{K}$  are the same. Voiculescu's theorem implies that these algebras are isomorphic.  $\square$

**Corollary 1.** *The algebra of singular Green operators is isomorphic to  $\mathcal{I}$ , an ideal of  $C^*({}^{ad}(Y \times Y) \rtimes \mathbb{R}_+^*)$ .*

*Proof.* This is a direct consequence of Theorem 2, since the algebra of singular Green operators is isomorphic to  $\Psi \otimes \mathcal{K}$ .  $\square$

## References

- [1] M. F. Atiyah and I. M. Singer. The index of elliptic operators, I. *Annals of Math.*, 87:484–530, 1968.
- [2] Louis Boutet de Monvel. Comportement d’un opérateur pseudo-différentiel sur une variété à bord. I. La propriété de transmission. *J. Analyse Math.*, 17:241–253, 1966.
- [3] Louis Boutet de Monvel. Comportement d’un opérateur pseudo-différentiel sur une variété à bord. II. Pseudo-noyaux de Poisson. *J. Analyse Math.*, 17:255–304, 1966.
- [4] Louis Boutet de Monvel. Boundary problems for pseudo-differential operators. *Acta Math.*, 126(1-2):11–51, 1971.
- [5] Lawrence G. Brown. Stable isomorphism of hereditary subalgebras of  $C^*$ -algebras. *Pacific J. Math.*, 71(2):335–348, 1977.
- [6] Lawrence G. Brown, Philip Green, and Marc A. Rieffel. Stable isomorphism and strong Morita equivalence of  $C^*$ -algebras. *Pacific J. Math.*, 71(2):349–363, 1977.
- [7] Alain Connes. *Noncommutative Geometry*. Academic Press, Inc., 1994.
- [8] B. V. Fedosov. Index theorems. In *Partial differential equations, VIII*, volume 65 of *Encyclopaedia Math. Sci.*, pages 155–251. Springer, Berlin, 1996.
- [9] Boris V. Fedosov, François Golse, Eric Leichtnam, and Elmar Schrohe. The noncommutative residue for manifolds with boundary. *J. Funct. Anal.*, 142(1):1–31, 1996.
- [10] Gerd Grubb. *Functional calculus of pseudodifferential boundary problems*, volume 65 of *Progress in Mathematics*. Birkhäuser Boston Inc., Boston, MA, second edition, 1996.
- [11] Gerd Grubb and Elmar Schrohe. Trace expansions and the noncommutative residue for manifolds with boundary. *J. Reine Angew. Math.*, 536:167–207, 2001.
- [12] Lars Hörmander. Pseudo-differential operators and hypoelliptic equations. In *Singular integrals (Proc. Sympos. Pure Math., Vol. X, Chicago, Ill., 1966)*, pages 138–183. Amer. Math. Soc., Providence, R.I., 1967.



- [13] G. G. Kasparov. The operator  $K$ -functor and extensions of  $C^*$ -algebras. *Izv. Akad. Nauk SSSR Ser. Mat.*, 44(3):571–636, 719, 1980.
- [14] J. J. Kohn and L. Nirenberg. An algebra of pseudo-differential operators. *Comm. Pure Appl. Math.*, 18:269–305, 1965.
- [15] Masaharu Kusuda. Characterizations of certain classes of hereditary  $C^*$ -subalgebras. *Proc. Amer. Math. Soc.*, 116(4):999–1005, 1992.
- [16] Robert Lauter, Bertrand Monthubert, and Victor Nistor. Pseudodifferential analysis on continuous family groupoids. *Doc. Math.*, 5:625–655 (electronic), 2000.
- [17] Robert Lauter, Bertrand Monthubert, and Victor Nistor. Spectral invariance of certain algebras of pseudodifferential operators. *J. Inst. Math. Jussieu*, 3(4), 2005.
- [18] S. T. Melo, R. Nest, and E. Schrohe.  $C^*$ -structure and  $K$ -theory of Boutet de Monvel’s algebra. *J. Reine Angew. Math.*, 561:145–175, 2003.
- [19] S.T. Melo, T. Schick, and E. Schrohe. A  $K$ -theoretic proof of Boutet de Monvel’s index theorem. Preprint math.KT/0403059, to appear in *J. Reine Angew. Math.*.
- [20] Bertrand Monthubert. Groupoids and pseudodifferential calculus on manifolds with corners. *J. Funct. Anal.*, 199(1):243–286, 2003.
- [21] Bertrand Monthubert and François Pierrot. Indice analytique et groupoïdes de Lie. *C. R. Acad. Sci. Paris Sér. I Math.*, 325(2):193–198, 1997.
- [22] Ryszard Nest and Elmar Schrohe. Dixmier’s trace for boundary value problems. *Manuscripta Math.*, 96(2):203–218, 1998.
- [23] Victor Nistor, Alan Weinstein, and Ping Xu. Pseudodifferential operators on differential groupoids. *Pacific J. Math.*, 189(1):117–152, 1999.
- [24] Stephan Rempel and Bert-Wolfgang Schulze. *Index theory of elliptic boundary problems*. North Oxford Academic Publishing Co. Ltd., London, 1985. Reprint of the 1982 edition.
- [25] Jean Renault. *A Groupoid Approach to  $C^*$ -Algebras*, volume 793 of *Lecture Notes in Mathematics*. Springer-Verlag, 1980.
- [26] Mikael Rørdam. Stable  $C^*$ -algebras. In *Operator algebras and applications*, volume 38 of *Adv. Stud. Pure Math.*, pages 177–199. Math. Soc. Japan, Tokyo, 2004.
- [27] Elmar Schrohe. Fréchet algebra techniques for boundary value problems on noncompact manifolds: Fredholm criteria and functional calculus via spectral invariance. *Math. Nachr.*, 199:145–185, 1999.

- [28] Elmar Schrohe. A short introduction to Boutet de Monvel's calculus. In *Approaches to singular analysis (Berlin, 1999)*, volume 125 of *Oper. Theory Adv. Appl.*, pages 85–116. Birkhäuser, Basel, 2001.
- [29] Georges Skandalis. Kasparov's bivariant  $K$ -theory and applications. *Exposition. Math.*, 9(3):193–250, 1991.
- [30] Dan Voiculescu. A non-commutative Weyl-von Neumann theorem. *Rev. Roumaine Math. Pures Appl.*, 21(1):97–113, 1976.