



Nova Southeastern University  
NSUWorks

Biology Faculty Articles

Department of Biological Sciences

2-2008

# Genomic Organization, Sequence Divergence, and Recombination of Feline Immunodeficiency Virus from Lions in the Wild

Jill Pecon-Slattery

*National Cancer Institute at Frederick*

Carrie L. McCracken

*National Cancer Institute at Frederick*

Jennifer L. Troyer

*National Cancer Institute at Frederick*

Sue VandeWoude


*Colorado State University - Fort Collins*

Melody E. Roelke

*National Cancer Institute at Frederick*

*See next page for additional authors*

Follow this and additional works at: [http://nsuworks.nova.edu/cnso\\_bio\\_facarticles](http://nsuworks.nova.edu/cnso_bio_facarticles)

 Part of the [Genetics and Genomics Commons](#), [Immunology and Infectious Disease Commons](#), and the [Veterinary Medicine Commons](#)

## NSUWorks Citation

Pecon-Slattery, Jill; Carrie L. McCracken; Jennifer L. Troyer; Sue VandeWoude; Melody E. Roelke; Kerry Sondgeroth; Christiaan Winterbach; and Stephen J. O'Brien. 2008. "Genomic Organization, Sequence Divergence, and Recombination of Feline Immunodeficiency Virus from Lions in the Wild." *BMC Genomics* 9, (66): 1-13. [http://nsuworks.nova.edu/cnso\\_bio\\_facarticles/776](http://nsuworks.nova.edu/cnso_bio_facarticles/776)

This Article is brought to you for free and open access by the Department of Biological Sciences at NSUWorks. It has been accepted for inclusion in Biology Faculty Articles by an authorized administrator of NSUWorks. For more information, please contact [nsuworks@nova.edu](mailto:nsuworks@nova.edu).

---

**Authors**

Jill Pecon-Slattery, Carrie L. McCracken, Jennifer L. Troyer, Sue VandeWoude, Melody E. Roelke, Kerry Sondgeroth, Christiaan Winterbach, and Stephen J. O'Brien

Research article

Open Access

## Genomic organization, sequence divergence, and recombination of feline immunodeficiency virus from lions in the wild

Jill Pecon-Slattery\*<sup>1</sup>, Carrie L McCracken<sup>2</sup>, Jennifer L Troyer<sup>2</sup>,  
Sue VandeWoude<sup>3</sup>, Melody Roelke<sup>2</sup>, Kerry Sondgeroth<sup>4</sup>,  
Christiaan Winterbach<sup>5</sup>, Hanlie Winterbach<sup>5</sup> and Stephen J O'Brien<sup>1</sup>

Address: <sup>1</sup>Laboratory of Genomic Diversity, National Cancer Institute-Frederick, Frederick MD 21702 USA, <sup>2</sup>Laboratory of Genomic Diversity, Basic Research Program-SAIC Frederick – National Cancer Institute, Frederick, MD 21702 USA, <sup>3</sup>Department of Microbiology, Immunology and Pathology, Colorado State University, Fort Collins CO 80532 USA, <sup>4</sup>Department of Veterinary Microbiology and Pathology, Washington State University, Pullman WA 99164 USA and <sup>5</sup>Tau Consultants, Private Bag 83, Maun, Botswana

Email: Jill Pecon-Slattery\* - slattery@ncifcrf.gov; Carrie L McCracken - cmccracken@ncifcrf.gov; Jennifer L Troyer - jtroyer@ncifcrf.gov; Sue VandeWoude - suev@lamar.colostate.edu; Melody Roelke - roelke@ncifcrf.gov; Kerry Sondgeroth - kss@vetmed.wsu.edu; Christiaan Winterbach - tau@dynabyte.bw; Hanlie Winterbach - tau@dynabyte.bw; Stephen J O'Brien - obrien@ncifcrf.gov

\* Corresponding author

Published: 5 February 2008

Received: 31 August 2007

BMC Genomics 2008, 9:66 doi:10.1186/1471-2164-9-66

Accepted: 5 February 2008

This article is available from: <http://www.biomedcentral.com/1471-2164/9/66>

© 2008 Pecon-Slattery et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

### Abstract

**Background:** Feline immunodeficiency virus (FIV) naturally infects multiple species of cat and is related to human immunodeficiency virus in humans. FIV infection causes AIDS-like disease and mortality in the domestic cat (*Felis catus*) and serves as a natural model for HIV infection in humans. In African lions (*Panthera leo*) and other exotic felid species, disease etiology introduced by FIV infection are less clear, but recent studies indicate that FIV causes moderate to severe CD4 depletion.

**Results:** In this study, comparative genomic methods are used to evaluate the full proviral genome of two geographically distinct FIV subtypes isolated from free-ranging lions. Genome organization of FIV<sub>ple</sub> subtype B (9891 bp) from lions in the Serengeti National Park in Tanzania and FIV<sub>ple</sub> subtype E (9899 bp) isolated from lions in the Okavango Delta in Botswana, both resemble FIV genome sequence from puma, Pallas cat and domestic cat across 5' LTR, *gag*, *pol*, *vif*, *orfA*, *env*, *rev* and 3'LTR regions. Comparative analyses of available full-length FIV consisting of subtypes A, B and C from FIV<sub>Fca</sub>, Pallas cat FIV<sub>Oma</sub> and two puma FIV<sub>Pco</sub> subtypes A and B recapitulate the species-specific monophyly of FIV marked by high levels of genetic diversity both within and between species. Across all FIV<sub>ple</sub> gene regions except *env*, lion subtypes B and E are monophyletic, and marginally more similar to Pallas cat FIV<sub>Oma</sub> than to other FIV. Sequence analyses indicate the SU and TM regions of *env* vary substantially between subtypes, with FIV<sub>ple</sub> subtype E more related to domestic cat FIV<sub>Fca</sub> than to FIV<sub>ple</sub> subtype B and FIV<sub>Oma</sub> likely reflecting recombination between strains in the wild.

**Conclusion:** This study demonstrates the necessity of whole-genome analysis to complement population/gene-based studies, which are of limited utility in uncovering complex events such as recombination that may lead to functional differences in virulence and pathogenicity. These full-length lion lentiviruses are integral to the advancement of comparative genomics of human pathogens, as well as emerging disease in wild populations of endangered species.

## Background

Feline immunodeficiency viruses (FIV) naturally infect cat species in the wild and are related to other lentiviruses known to infect primates (human and simian immunodeficiency viruses, HIV and SIV), sheep and goats (caprine arthritis encephalitis virus -CAEV), horse (equine infectious anemia virus-EIAV), and cattle (bovine immunodeficiency virus-BIV). FIV is endemic in Felidae species [1-12], many of which are considered endangered or threatened with extinction [13]. A recent comprehensive survey of serum and lymphocyte specimens from 3055 individuals affirm that at least 11 free-ranging and, if captive animals are included, as many as 31 species of cat are infected with FIV [4]. Phylogenetic analyses of the *pol-RT* region sequenced from six of these felid species, plus spotted hyaena, *Crocuta crocuta*, affirm the high level of species-specificity worldwide [4,11,14-17]. Each species specific FIV forms a distinct monophyletic lineage, separated by substantial genetic divergence that suggests virus-host adaptation and rare episodes of interspecies transmission in the wild [4,18].

The effects of FIV infection and disease are well described in domestic cat (*Felis catus*) but less so in exotic felids. FIV<sub>Fca</sub> infection in domestic cat is analogous to HIV infection of humans causing early flu-like symptoms, followed by severe weight loss, chronic wasting disease, and increased susceptibility to rare cancers and opportunistic disease, neurologic disease and death [19,20]. Captive and wild populations of two species, the African lion (*Panthera leo*) infected with FIV<sub>ple</sub> and the puma (*Puma concolor*), infected with FIV<sub>pco</sub> exhibit less severe disease associations. However, infected lions show a dramatic decline in CD4+ subsets, a reduction of the CD4+/CD8+ ratio, reduction of CD8+β<sup>high</sup> cells, and expansion of the CD8+β<sup>low</sup> subset relative to uninfected lions [21-23]. Further, FIV<sub>pco</sub> infected puma display a more generalized response of lymphopenia expressed as a significant decline in total lymphocytes, CD5+ T-cells, and CD5-lymphocytes as well as a significant reduction in CD4+ T-cells [23]. Like lions, seropositive pumas have a significant decline in CD8+β<sup>high</sup> cells but differ by not showing compensatory expansion of CD8+β<sup>low</sup> cells relative to controls [23]. The results observed with FIV-infected lion and puma parallels human (HIV) and Asian monkey (SIV) CD4+ diminution, and suggests there may be an immunological cost of FIV infection in these two species of large cats.

Identification of genetic correlates of FIV virulence, infectivity, and pathogenicity in different cat species is limited due to a paucity of complete genome sequence. Only subtypes A, B and C from domestic cat FIV<sub>Fca</sub> [24-26], subtypes A and B from puma FIV<sub>pco</sub> [14,27] and a single strain (FIV<sub>Oma</sub>) from Pallas cat (*Otocolobus manul*) [16] have

been sequenced in entirety. Here we present full-length provirus sequenced from FIV<sub>ple</sub> subtype B isolated from lions in the Serengeti National Park in Tanzania and FIV<sub>ple</sub> subtype E from lions dwelling in the Okavango Delta in Botswana. These two FIV<sub>ple</sub> subtypes exhibit a range in sequence divergence throughout the genome, share motifs unique to this lion-specific lentivirus, yet also exhibit unusual and significant differences in the *env* gene.

## Results and Discussion

### Genomic Organization and Sequence Divergence of FIV<sub>ple</sub> Subtypes


FIV<sub>ple</sub> subtypes B (accession number EU117991) and E (accession number EU117992) share a similar genome organization with other FIV which consists of LTR, *gag*, *pol*, *vif*, *orfA*, *env*, and additional small ORFs that may represent accessory genes including *rev* (Table 1). The total proviral genome size was conserved between FIV<sub>ple</sub> subtype B (9899 bp) and subtype E (9891 bp) (Table 1). FIV<sub>ple</sub> *gag* encodes three putative structural proteins of matrix, capsid and nucleocapsid. *Pol* is conserved and encodes key viral enzymes of protease, reverse transcriptase, RNAase, dUTPase and integrase. FIV<sub>ple</sub> *vif*, an accessory protein essential for viral replication, resembles that of FIV<sub>Fca</sub>. *OrfA* in FIV<sub>ple</sub> is similar to FIV<sub>Fca</sub> and likely corresponds to HIV *tat*, which targets transcription factors in the LTR. FIV<sub>ple</sub> *env* encodes the putative leader, surface (SU), and transmembrane (TM) regions of the envelope glycoprotein, essential components for viral binding to and entry into the host cell. FIV<sub>ple</sub> *rev* is similar to HIV/FIV *rev*, and is thought to be critical in viral replication. FIV<sub>ple</sub> *rev* appears to be encoded by splicing two exons: the first in the leader region of *env*, the second located near the 3' region adjacent to *env* (Table 1).

The LTR of FIV<sub>ple</sub> contains transcription and regulatory elements common to other FIV. These include the direct 2 bp repeat (IR) defining the 5' and 3' termini of LTR, AP-4, Aml-1 (EPB20), AP-1, TATA box, Poly A, and the cap transcription initiation site (Figure 1). FIV<sub>ple</sub> subtypes have additional transcription factors characteristic of FIV, but placed in alternate locations within the LTR U3 including NF-AT and CREBP-1/c-Jun. These and other motifs were determined by homology search with a threshold value of 85% with the Motif Search database [28] [see Additional file 1]. Overall, lion LTRs are not identical between subtypes B and E, differing by 15% in nucleotide substitutions, comparable to that observed between FIV<sub>Fca</sub> subtypes A, B and C (Figure 1, Figure 2A).


Deep genetic divergence between FIV strains from different cat species made alignments problematic. For coding regions, we first translated each gene into amino acid residues, which are less divergent as changes occur at a lower rate of substitution, to serve as a "scaffold" for alignment

**Table 1: Gene size and location within FIV<sub>Ple</sub> Subtypes B and E compared with previously published FIV<sub>Fca</sub>, FIV<sub>Oma</sub> and FIV<sub>Pco</sub>.**

	5'LTR	5'UTR	Gag	Pol	Vif	OrfA	Env	PPT	3'LTR
FIV <sub>Ple</sub> Subtype B (Serengeti)									
Gene position	1-398	399-704	705-2213	2018-5464	5461-6171	6288-6542	6601-9213	9484-9498	9501-9899
Gene length (bp)	398	306	1509	3447	711	255	2613	15	398
Translated Protein Size (# aa)			503	1149	237	85	871		
FIV <sub>Ple</sub> Subtype E (Botswana)									
Gene Position	1-397	398-702	703-2199	2004-5450	5447-6211	6198-6452	6532-9222	9478-9492	9495-9891
Gene length (bp)	397	306	1497	3447	765	255	2691	15	397
Translated Protein Size (# aa)			498	1149	255	85	897		
FIV <sub>Fca</sub> Petaluma (Subtype A)									
Gene Position	1-355	356-627	628-1980	1868-5243	5236-5991	5992-6228	6266-8836	9098-9117	9120-9474
Gene length (bp)	355	272	1353	3375	756	237	2571	19	
Translated Protein Size (# aa)			451	1125	252	79	857		
FIV <sub>Fca</sub> USIL (Subtype B)									
Gene position	1-361	362-633	634-1983	1875-5248	5239-5994	5995-6231	6269-8830	9092-9110	9102-9462
Gene length (bp)	361	272	1350	3374	756	237	2562	17	361
Translated Protein Size (# aa)			451	1124	252	79	854		
FIV <sub>Fca</sub> Subtype C									
Gene Position	1-354	355-632	633-1985	1874-5248	5241-5996	5997-6233	6271-8835	9092-9100	9113-9466
Gene length (bp)	354	278	1353	3375	756	237	2565	19	354
Translated Protein Size (# aa)			451	1125	252	79	855		
FIV <sub>Oma</sub>									
Gene Position	1-376	377-684	685-2181	1980-5432	5429-6187	6188-6448	6512-9103	9360-9375	9378-9751
Gene length (bp)	376	308	1497	3453	759	261	2592	16	374
Translated Protein Size (# aa)			499	1161	253	87	864		
FIV <sub>Pco</sub> PLV-14 Subtype A (Florida)									
Gene Position	1-311	312-615	616-2055	2199-5459	5419-6249	5759-5938	6250-8772	8771-8787	8790-9100
Gene length (bp)	311	304	1440	3261	831	180	2523	17	311
Translated Protein Size (# aa)			480	1087	277	59	841		
FIV <sub>Pco</sub> PLV-1695 Subtype B (British Columbia)									
Gene Position	1-306	307-638	639-2024	1886-5323	5298-6008	5972-6310	6283-8715	8772-8784	8787-9092
Gene length (bp)	306	332	1386	3438	711	339	2433	13	305
Translated Protein Size (# aa)			462	1146	237	113	811		



**Figure 1**  
Alignment of FIV<sub>Ple</sub> subtype B and E LTR showing the U3, R and U5 regions. Grey shadow indicates inverted repeat, boxed regions indicate putative transcription elements common in FIV.



**Figure 2**

Phylogenetic reconstruction based on nucleotide sequence of LTR and coding genes from full-length FIV nucleotide sequences excluding *env*. (A-E) Shown are the maximum likelihood trees (ML) which are identical to tree topologies using maximum parsimony (MP) and minimum evolution (ME) for each gene region. See methods and Additional file 3 for specific parameters as implemented in PAUP ver 4.10b. (E) *OrfA* phylogeny does not include FIV<sub>Pco</sub> subtype A due to lack of sufficient homology for proper gene identification. (F) Phylogenetic tree of concatenated combined data of coding genes *gag*, *pol*, *vif*, and *orfA*. All nodes supported by 100% bootstrap proportions in ME, MP and ML analyses except for relative positions of FIV<sub>Fca</sub> subtypes which were supported by bootstraps >50% but less than 100% within the FIV<sub>Fca</sub> clade.

of nucleotides using the program RevTrans [29]. Our results indicate that *pol* (3657 bp) is the most conserved gene across FIV, albeit exhibiting substantial average pairwise genetic distances of 60% and 54% for nucleotide and amino acid data, respectively (Table 2). *Gag* sequences (1551 bp) differed by an average pairwise genetic distance of 65.8% for nucleotides, a 53.2% amino acids (Table 2). However, *vif* (870 bp), *orfA* (351 bp), and *env* (2958 bp) were highly divergent. For these genes, sufficient homology existed to both identify the gene, and to create a multiple sequence alignment across all FIV yet, phylogenetic models for patterns of substitution at variable sites were saturated resulting in an average genetic distance of 100% for both nucleotide and amino acid data (Table 2). Such differences in rates of evolution between viral genes corroborate previous findings describing functional con-

straints for *gag* and *pol* [7,8,17], while also demonstrating that *vif*, *orfA*, and *env* rapidly evolve in each host species.

**Phylogenetic Analyses of FIV<sub>Ple</sub> Subtypes**

The evolution of FIV<sub>Ple</sub> subtypes is defined by separate phylogenetic analyses of each viral gene as well as combined data of concatenated sequences representing the entire coding region of FIV. LTR, *gag*, *pol*, *vif* and *orfA* affirm the species-specificity of FIV both in individual gene analyses (Figure 2A-E) and in the combined concatenated data phylogeny excluding *env* (Figure 2F). The three subtypes of FIV<sub>Fca</sub> from the domestic cat exhibit the least amount of genetic divergence within each viral gene phylogeny. Sharing a monophyletic lineage with distantly related FIV<sub>Oma</sub>, the FIV<sub>Ple</sub> subtypes B and E have intermediate levels of genetic distance with each viral gene exam-

**Table 2: Estimates of genetic divergence of FIV genes.**

	FIV GENE				
	Gag	Pol	Vif	OrfA <sup>1</sup>	Env
Genetic distance	Nucleotide % Genetic Distance (GTR)				
Average Pairwise (N = 8)	65.8	60.3	100*	100*	100*
Selected comparisons					
FIV <sub>Ple</sub> B vs FIV <sub>Ple</sub> E	20.3	20.4	33.1	27.9	100*
FIV <sub>Ple</sub> B vs FIV <sub>Oma</sub>	28.8	29.3	44.3	55.8	42.7
FIV <sub>Ple</sub> B vs FIV <sub>Fca</sub> C	55.6	53.5	100*	100*	100*
FIV <sub>Ple</sub> E vs FIV <sub>Oma</sub>	32.9	28.1	36.7	61.3	100*
FIV <sub>Ple</sub> E vs FIV <sub>Fca</sub> C	61.4	51.4	79.3	100*	64.4
	Amino Acid % Genetic Distance (Pam-Dayhoff <sup>2</sup> )				
Average Pairwise (N = 8)	53.2	44.1	100*	100*	100*
FIV <sub>Ple</sub> B vs FIV <sub>Ple</sub> E	9.4	10.5	36.2	24.4	100*
FIV <sub>Ple</sub> B vs FIV <sub>Oma</sub>	24.8	20.1	59.1	58.4	42.8
FIV <sub>Ple</sub> B vs FIV <sub>Fca</sub> C	47.6	38.2	100*	100*	100*
FIV <sub>Ple</sub> E vs FIV <sub>Oma</sub>	25.3	19.9	42.1	68.3	100*
FIV <sub>Ple</sub> E vs FIV <sub>Fca</sub> C	46.2	37.9	91.9	100*	79.7

\* 100% genetic distance means sufficient homology present to create alignment, but no meaningful phylogenetic associations are detected.

<sup>1</sup> FIV<sub>Pco</sub> A not included as no homologous OrfA identified.

<sup>2</sup> See Methods

ined. Subtypes A and B of FIV<sub>Pco</sub> are the most divergent and have substantial differences across the viral genome. Thus, the hierarchical pattern of genetic divergence among full-length genomic analyses of FIV<sub>Fca</sub>, FIV<sub>Ple</sub> and FIV<sub>Pco</sub> recapitulates earlier evolutionary studies based on portions of *pol-RT* and *gag* [4,7,8,10-12,17,30,31].


The relative differences in genetic diversity among FIV strains may be correlated with the amount of time since the virus entered modern felids and therefore, can be interpreted in the context of the evolutionary and phylogeographic history of each host species. The domestic cat evolved as a unique felid lineage only around 10,000 year ago [32] from subspecies of wildcat *Felis silvestris* inhabiting Near East Asia [33]. Preliminary results from limited seroprevalence studies, indicate that FIV appears to be absent from nearly all of the close relatives of domestic cat [(genus *Felis* after [34]) except for French European wildcat *F. silvestris* [4,35]. Thus, the pattern of FIV<sub>Fca</sub> divergence may represent recent emergence combined with rapid viral diversification within the domestic cat world-wide. In contrast, the puma is one of the oldest species within Felidae, sharing an evolutionary lineage with the African cheetah (*Acinonyx jubatus*) and the New World jaguarundi (*Puma yagouaroundi*) and arose approximately 4.5 MYA [34]. The extreme divergence between subtypes A and B

within the FIV<sub>Pco</sub> lineage suggests an ancient origin of FIV infection of puma, a result consistent with the published *pol-RT* phylogeny marked by high levels of intra-subtype divergence of FIV<sub>Pco</sub> subtypes from throughout the host species range [4,8,11]. Lastly, the African lion species arose approximately 2 MYA and spread throughout Africa, Asia and the Americas [34]. However, due to episodes of population reduction followed by expansion from East Africa and recolonization, genomic diversity in modern lion populations coalesces to approximately 325,000 years ago and is confined the African continent [36]. FIV<sub>Oma</sub> is found in wild populations of the Eurasian Pallas cat [4], a species that arose during the late Pleistocene [34]. The monophyletic lineage of Pallas cat FIV<sub>Oma</sub> and African lion FIV<sub>Ple</sub> observed here suggest more ancient inter-species transmission as the last time lions and Pallas cats were in geographic contact was during the Pleistocene when lion ranges spread throughout Asia, providing a possible opportunity for FIV transmission between these species [37].


#### **Discordant env phylogeny between FIV<sub>Ple</sub> subtypes reveals ancestral FIV recombination events in the wild**

The patterns of phylogenetic divergence between FIV strains from different cat species are concordant between all viral gene regions with one notable exception, the *env*

A. Combined All Genes (9391 bp)



B. Env (2958 bp)



**Figure 3**


Phylogenetic reconstruction based on nucleotide sequence of fulllength proviral FIV including *env* and separate analysis of *env*. A. Phylogenetic tree of concatenated combined data of coding genes *gag*, *pol*, *vif*, *orfA* and *env*. B. Phylogenetic tree of *env* sequences only. Shown is the maximum likelihood tree (ML) identical to tree topology using maximum parsimony (MP) and minimum evolution (ME) for each gene region. See methods and Additional file 3 for specific parameters as implemented in PAUP ver 4.10b. All nodes supported by 100% bootstrap proportions in ME, MP and ML analyses except for relative positions of FIV<sub>Fca</sub> subtypes which were supported by bootstraps >50% but less than 100% within the FIV<sub>Fca</sub> clade.

gene. Phylogenetic analyses of the entire concatenated coding region (9391 bp) and separate analysis of the *env* gene (2958 bp) show the two FIV<sub>Ple</sub> subtypes are no longer monophyletic (Figure 3A and 3B). A closer examination of the *env* gene shows only two shared regions of homology between FIV<sub>Ple</sub> subtypes. The first spans the sites 1–519 of *env*, containing exon 1 of *rev* (Table 1), within the leader region exhibiting 80% nucleotide and 68% amino acid homology between FIV<sub>Ple</sub> subtypes. The second region occurs at the terminal 3' region of *env* (sites 2506–2958) with 87% and 71% genetic identity for nucleotides and amino acid, respectively. Based on comparison with FIV<sub>Fca</sub>, this region of FIV<sub>Ple</sub> may be the *rev* responsive element (RRE), which is critical for targeting *rev* to the nucleolus of the cell [38]. As *rev* is conserved between lion subtypes, it is likely that RRE must remain conserved as well.

By contrast, the SU and TM regions of *env* differ substantially between FIV<sub>Ple</sub> subtypes (Figure 4). A contiguous region of *env*, from amino acid sites 181 through 931 (green in Figure 4), shows that FIV<sub>Ple</sub> subtype E is more similar to FIV<sub>Fca</sub> than to FIV<sub>Ple</sub> subtype B. Further, *env* of FIV<sub>Ple</sub> subtype B, concordant with results from other gene trees (Figure 2A–E), shares more homology with FIV<sub>Oma</sub> (blue in Figure 4). Moreover, the lack of monophyly between FIV<sub>Ple</sub> subtype B and FIV<sub>Oma</sub> (Figure 3) is a consequence of the recombinant *env* of FIV<sub>Ple</sub> subtype E, as exclusion of this subtype from the analyses (data not shown) recovered the monophyletic relationship observed with other genome regions (Figure 2A–E).

The predicted *env* protein from both FIV<sub>Ple</sub> strains were compared to other published FIV strains with respect to inferred structural elements, with particular focus on regions known to be important for receptor binding. Con-





**Figure 4**

Multiple sequence alignment of amino acids of *env* from FIV<sub>ple</sub> subtypes B and E compared with FIV<sub>Fca</sub> subtypes A, B and C, FIV<sub>Pco</sub> subtypes A and B, and FIV<sub>Oma</sub>. Significant structures within the *env* gene inferred from Smirnova et al. 2005 are indicated in colored boxes. Putative regions include: conserved amino acids (white box), variable regions VI-V9 (red box); epitope binding sites (orange box); conserved cysteine (red highlight); N glycosylation sites (yellow highlight). Homologous region shared between FIV<sub>ple</sub> subtypes B and E are highlighted in grey. Amino acid sites 1-176 contain the first exon of *rev* (see Table 1) in lion FIV<sub>ple</sub> subtypes B and E. The portion of *env* proposed to be a result of recombination in FIV<sub>ple</sub> subtype E is highlighted in green. The corresponding region of *env* thought to represent FIV<sub>ple</sub> without recombination, as it is more homologous to FIV<sub>Oma</sub>, is highlighted in blue. Amino acids sites 931-978 (grey) likely contain the RRE element shared by lion FIV<sub>ple</sub> subtypes B and E.

served (white in Figure 4) and variable regions (red in Figure 4) and epitope binding sites (orange in Figure 4) were identified based on their locations in the domestic cat FIV sequences [39]. The V3-V5 regions shared least homology between the two strains. In FIV<sub>Fca</sub>, this region has been

shown to contain the CXCR4 binding site [40], neutralizing antibody binding sites [41-43] and several epitopes important for cell tropism and cell line adaptation [44-46]. Within the V3-V5 region, several biochemical differences have been noted between domestic and non-

domestic cat lentiviruses [39]. *FIV<sub>ple</sub>* subtype B demonstrated properties more similar to other non-domestic cat lentiviruses including a negative charge and fewer cysteine residues within this region. Conversely, *FIV<sub>ple</sub>* subtype E had a positive charge and more cysteines in V3-V5, more similar to the domestic cat lentiviruses. Both lion FIVs had similar numbers of predicted N-glycosylation sites (10 and 11 for B and E, respectively) and these numbers are intermediate to the domestic cat FIVs (8–10) and the other non-domestic FIVs (13–14). A similar trend of lower charge and more cysteine residues in B than E was noted in V3, the region implicated as receptor binding domain for FIV [44,46,47]. In contrast, the more conserved regions flanking V3-V5 were more positively charged in *FIV<sub>ple</sub>* subtype B than in *FIV<sub>ple</sub>* subtype E, but contained similar numbers of cysteine residues and putative N-glycosylation sites. Such differences suggest that substantial divergence may occur in secondary and tertiary structures at the receptor-binding region of these two lion lentiviruses.

Recombination in lentiviruses is not uncommon. In the ongoing global HIV pandemic, at least 34 circulating recombinant forms from HIV-1 subtypes have been so far described in patients world-wide [48]. SIV full genome sequence comparisons increasingly depict extant primate lentiviruses with mosaic structures indicative of multiple recombination events over time [49-54]. In *FIV<sub>Fca'</sub>*, recombination in the V3-V5 region of *env* was detected between subtypes A and B in feral cats [7], and different recombination frequencies occur between large regions of *FIV<sub>pco</sub>* subtype B in domestic cat experimentally infected with *FIV<sub>pco</sub>* B [31]. Whereas the frequency of *FIV<sub>ple</sub>* recombination is not yet known, our studies show that over 40% of Serengeti lions in Tanzania are multiply infected with *FIV<sub>ple</sub>* subtypes A, B and C, which circulate freely within this large population [6] and thus offer opportunities for recombination.

The recombination of *env* in FIV infected lions has interesting evolutionary significance because the divergence in this region is extensive between the two subtypes. Therefore, subtype E recombination may represent an ancient event of recombination followed by a long period of divergence, or a more recent recombination with a highly divergent but as yet unsequenced strain either from lions or another African felid species. Although *FIV<sub>ple</sub>* subtype E *env* is more similar to *FIV<sub>Fca</sub>* than to any other known FIV the extent of genetic divergence is still quite substantial, i.e. 64.4% nucleotide relative to *FIV<sub>Fca</sub>* subtype C (Table 2), suggesting that if recombination has occurred recently, it is likely to have been with strain that has not yet been sequenced for the *env* gene. This recombination event may also have functional implications, as *FIV<sub>ple</sub>* subtype E *env* has structural features more similar to pathogenic *FIV<sub>Fca</sub>*.

Further investigation into complete genome analyses of *FIV<sub>ple</sub>* subtypes A, C, D and F as well as FIV from other seropositive African felids, will likely provide new insights into the role of recombination in *env* in the wild. Clinical studies will help to clarify the significance of these recombination events.

## Conclusion

Ongoing efforts to sequence full genome FIV from all seropositive exotic cat species will be essential to understanding the evolutionary trajectory of these viruses including the origin and frequency of recombination within FIV. This study demonstrates the necessity of whole-genome analysis to compliment population/gene-based studies, which are of limited utility in uncovering complex events such as recombination that may lead to functional differences in virulence and pathogenicity. The changes observed in the *env* gene as a consequence of recombination in *FIV<sub>ple</sub>* will provide important clues to the natural history of these viruses and their hosts, and may lead to insights into genetic determinants of pathogenicity and virulence differences between domestic cat and lion FIV; findings with important implications for HIV pathogenesis in humans and virus attenuation in wild populations of endangered species.

## Methods

### Cell Culture of *FIV<sub>ple</sub>* Subtype E Botswana lion Ple-1027

*FIV<sub>ple</sub>* subtype E was isolated from PBMCs (whole blood with EDTA) collected from wild lions in the Okavango Delta in Botswana, viably frozen under field conditions [23] and stored in liquid nitrogen. In preparation for cell culture, viably frozen PBMCs from Ple-1027 were thawed at 37°C, washed twice in LBT media (RPMI 1640 (Invitrogen Life Sciences, Carlsbad, Calif.) containing 20% fetal bovine serum (Atlanta Biologicals, Norcross, Ga.), 1% Glutamax I, 1 mM sodium pyruvate, 0.1 mM nonessential amino acids,  $5 \times 10^{-5}$  M  $\beta$ -2-mercaptoethanol, 100 U of penicillin/ml, 100  $\mu$ g of streptomycin/ml, (all from Invitrogen Life Sciences), and 9 g of glucose (Sigma)/liter), and resuspended at a final concentration of  $1-1.5 \times 10^6$  cells/ml in LBT + interleukin-2 at 100 U/ml (Invitrogen Life Sciences).

Domestic cat Mya-1 naïve feeder cells [55] were prepared for co-culture by cultivation in LBT + IL-2, plated in MEM containing 10% fetal bovine serum, 100 U of penicillin, 100  $\mu$ g of streptomycin/ml, and 1% glutamax, and dispensed at  $2 \times 10^6$  cells/ml in appropriate media in a 24 well plate. Reconstituted lion PBMCs were then added to Mya-1 cells at a volume of 400  $\mu$ l ( $4-6 \times 10^5$  cells).

Media was collected biweekly and subjected to microtiter reverse transcriptase assay as follows. Briefly, 15  $\mu$ l of culture supernatant in triplicate was incubated with 50  $\mu$ l of

0.05 M Tris (pH 7.8) with 75 mM KCl, 5 mM MgCl<sub>2</sub>, 0.5 mM EGTA, 2 mM dithiothreitol, 5 nM oligo(dT), 0.05% NP-40, poly(A) at 50 µg/ml, and <sup>32</sup>P at 20 µCi/ml for 90 to 120 minutes at 37°C. Aliquots of 2.5 µl of each reaction mixture were spotted onto a nylon filter (Wallac, Turku, Finland) and allowed to dry. Un-incorporated label was washed away with five 10 to 60 minute washes with 0.03 M sodium citrate, pH 7.0, in 0.3 M sodium chloride (SSC) buffer, and the membrane was then fixed in 100% ethanol. Counts per minute were measured using a Microbeta Counter (Wallac).

Starting on day 34 post co-culture, supernatant from lion PBMC cocultures with Mya-1 cells had reverse transcriptase (RT) values approximately 3 to 10 times naïve supernatant levels, indicating productive lentiviral replication. RT activity was not detected in any other control supernatants through 49 days of culture.

RNA was extracted from 200 µl of supernatant from positive cultures using QIAamp viral RNA mini kit (QIAGEN) and reversed transcribed to cDNA with Superscript II (Invitrogen) according to manufacturer's instructions. PCR was then performed to amplify a diagnostic region of *pol* as previously described [4]. Amplicons were sequenced to confirm the presence of a Botswana strain of FIV (*FIV<sub>ple</sub>* subtype E). One ml aliquots of supernatant were frozen at -70°C. Aliquots were then thawed and used to inoculate 3 × 10<sup>6</sup> Mya-1 cells, which were grown 14 days to achieve positive RT values as above. Cells were supplemented with fresh media weekly and grown to 1 × 10<sup>7</sup> cells at which point cells were harvested by centrifugation and cell pellets were frozen at -70°C.

#### **Cell Culture of *FIV<sub>ple</sub>* Subtype B: Serengeti lion Ple-458**

Isolation and culture methods for *FIV<sub>ple</sub>* Subtype B are similar to the methods described for Subtype E (above) with the following exceptions. *FIV<sub>ple</sub>* Subtype B was isolated from PBMCs from a wild, sero-positive lion (Ple-458, Serengeti National Park), separated from heparinized whole blood by sucrose gradient centrifugation using Histopaque (Sigma). Cells were mixed with 10% DMSO with 90% fetal calf serum and viably frozen in nitrogen vapor in aliquots of ~10<sup>6</sup> cells per ml. Post-freezing, thawed PBMCs (10<sup>6</sup> cells) from the wild lion were co-cultivated with an equal number of lion donor cells (Ple-73, captive, National Zoological Park, Wash., D.C.; this lion was sero-positive but had repeatedly tested negative for virus isolation). All PBMCs were mitogen stimulated with concanavalin A (5 µg/ml) for 72 hrs. Co-cultures were propagated in RPMI 1640 with 10% bovine serum and 10% human interleukin-2 (Gibco-BRL). Fresh media was added every 72 hours and new donor cells (10<sup>6</sup> cells) were added every 14 days. Replicating virus was confirmed in the supernatant by demonstrating both positive

Mg<sub>2</sub><sup>+</sup> – dependent reverse transcriptase (RT) and the presence of typical lentiviral particles seen by electron microscopy [10]. Virus rich supernatants were clarified by slow speed centrifugation and stored in liquid nitrogen freezers.

In order to expand the culture sufficiently to harvest viral supernatant for Western blot assays and to conduct the genetic analysis, 1 ml RT positive supernatant (LLV-2, SV lab) was used to inoculate 3201 cells (5 ml at 2 × 10<sup>6</sup>/ml), FeLV negative lymphosarcoma cells [56]. Cells were maintained in equal parts Leibovitz's L-15 media and RPMI 1640 with 20% fetal calf serum with glutamine (2x) and penicillin/streptomycin (1x). Initially, this culture was difficult to maintain in 3201 cells because it caused rapid cell death thus, in order to keep the culture alive, fresh media and naïve 3201 cells had to be added every 3–4 days. After 21 days post infection (dPI), fresh media continued to be added to the culture every 3–4 days, but the addition of naïve 3201 cells was stopped and the % viability was allowed to decline (in the hope that a cell adapted virus could emerge that would enhance our ability to grow up viral stocks for use in Western blot assays). From dPI 28 to 49 the culture viability hovered between 18–24%, but after dPI 52 it was clear that both the viability and cell numbers began to improve (viability from 46 to 86%). By dPI 71 the cell viability was holding at >90% and the culture was growing at 40–50% per day. Infected cells for DNA extraction and genetic analysis of subtype B virus were harvested on dPI 88, centrifuged, and the pellets frozen at -70°C.

#### **DNA extraction, Cloning and Sequencing of *FIV<sub>ple</sub>* Subtypes B and E**

DNA was extracted and purified from frozen cell culture pellets following the manufacturer's protocols established for blood products (Quiagen). Following extraction, DNA quality was checked by gel electrophoresis, and quantified by spectrophotometer (NanoDrop).

*FIV<sub>ple</sub>* proviral DNA was amplified using long PCR to generate overlapping proviral genome regions of approximately 5 kb (Roche's Expand PCR kit). For *FIV<sub>ple</sub>* subtype B, three over-lapping regions were amplified using the following primer pairs: FSHltr2F and FIVpol6R (LTR-pol); FIVgag2aF and FIVpol5R (gag-pol); and FIVpol5F and FIVltr4R (pol-LTR) (see Additional file 2). For subtype E, two over-lapping regions were amplified using the two primer pairs FSHltr2F and FIVpol5R (LTR-pol), and FIVgag2aF and FIVltr4R (gag-LTR) (see Additional file 2). PCR reactions used 0.2–2.0 µg DNA with the following thermocycling conditions: 94°C for 2 minutes; ten cycles of 94°C for 10 seconds, 52°C for 30 seconds and 68°C for 4 minutes; 25 cycles of 94°C for 10 seconds, 52°C for 30 seconds and 68°C for 4 minutes and 20 seconds, with

each having an extension time 20 seconds longer than the one before it; followed by 68°C for 7 minutes and 4°C hold. Additional "internal" primers were developed to fill in sequence gaps within each subtype (see Additional file 2) using the same PCR conditions listed above. Biometra T1 thermocyclers were used for all PCR reactions and amplicons were visualized on a 1% agarose gel.

The PCR products were cloned using TOPO TA XL cloning kits (Invitrogen). The resultant colonies were grown on LB agar plates with kanamycin and mini-prepped using Qiagen's REAL Prep 96. A restriction digest with EcoRI was performed to confirm successful cloning. The ends of the inserted PCR product were sequenced using the primers provided with the cloning kit for additional verification of *FIV<sub>ple</sub>* cloned products.

The final full-length sequence for each over-lapping long PCR product generated for each *FIV<sub>ple</sub>* subtype was obtained using transposon bombing [GPS-1 kit (New England BioLabs)]. In this method, transposons were randomly inserted into one of the successfully transformed plasmids for each primer combination for each sample. The results were used to transform OneShot Chemically Competent *E. coli* cells (Invitrogen) and grown on LB agar plates with kanamycin and chloramphenicol and were mini-prepped for DNA extraction using REAL Prep 96 (Quiagen). Restriction digest with EcoRI was performed to confirm successful insertion of the transposon. Using sequencing primers provided in the GPS-1 kit, 48 transposon fragments/PCR reaction were sequenced using an automated sequencer model ABI 3730.

Sequences (average read was approximately 600 bp) randomly generated by transposon bombing of individual clones defined multiple overlapping regions and were assembled into the full-length viral genome using Sequencher version 4.1 (Gene Codes Corporation) and submitted to GenBank [accession number [EU117991](#) (Subtype B) and [EU117992](#) (Subtype E)].

#### **Genetic and Phylogenetic Analyses of *FIV<sub>ple</sub>* Subtypes B and E**

Gene annotation of *gag*, *pol*, *env*, *orfA*, *vif* and *rev* from open reading frames in lion subtypes B and E used translation into amino acids and comparison with existing full-length FIV strains. The following sequences were used from GenBank: for *FIV<sub>Pco</sub>* Subtype B in pumas strain PLV-1695 accession number [DQ192583](#)[27]; for *FIV<sub>Pco</sub>* Subtype A in pumas strain PLV-14 accession number [U03982](#)[14]; for *FIV<sub>Oma</sub>* in Pallas cat accession number [AY713445](#)[16]; for *FIV<sub>Fca</sub>* subtype C in domestic cat strain C36 accession number [AY600517](#)[24]; for *FIV<sub>Fca</sub>* subtype B, strain usil2489, accession number [U11820](#); for *FIV<sub>Fca</sub>* [57]; Subtype A, strain PPR accession number [M36968](#)

and strain Petaluma accession number [M25381](#)[15,25,58]. Open reading frames were determined and regions of homology between *FIV<sub>ple</sub>* with other FIV strains using pair-wise comparisons implemented by BLAST of two sequences [59]. The boundaries of both the 5'LTR and 3'LTR regions were identified by the conserved polypurine tract (PPT) shared by all FIV [60] and the primer binding site (PBS) which mark the boundary between the 3'LTR and the 5'LTR, respectively.

The genome of lion *FIV<sub>ple</sub>* was compared with existing full-length FIV by multiple sequence alignments of each viral gene. LTR regions were aligned using Clustal X [61] and verified and edited by eye using Se-AL ver 2.0 [62]. Due to large genetic divergence between FIV from different species, alignment for coding regions of FIV used the program REVTRANS ver 1.4 [29] which takes a multiple sequence file, translates that file into amino acid residues, aligns the amino acids, and uses this alignment as the scaffold for nucleotide alignment. Aligned multiple sequence files were imported into Modeltest ver 3.7 [63] and the optimal model of nucleotide substitution was selected using the AIC criterion (see Additional file 3).

Viral genes were analyzed separately, as well as combined, for genome comparison and phylogenetic reconstruction. Phylogenetic trees based on nucleotide data were obtained using a heuristic search with three different optimality criteria of maximum likelihood (ML), minimum evolution (ME) and maximum parsimony (MP) as implemented in PAUP\* ver 4.0b10 [64]. Conditions for the ML analysis included starting trees obtained by stepwise addition, and branch swapping using the tree-bisection-reconnection (TBR) algorithm. Specific conditions for the ME search included starting trees obtained by neighbor – joining, TBR branch-swapping algorithm, and no collapsing of zero-length branches. The MP analyses coded gaps as "missing", with step-wise addition of taxa and TBR branch swapping. Support for nodes within the phylogeny used bootstrap analysis with identical settings established for each method of phylogenetic reconstruction and retention of node bootstrap values greater than 50%. The number of bootstrap iterations consisted of 1000 for ME and MP methods and 100 for ML. Additional analyses were conducted on FIV coding sequences after translation into amino acids. Genetic distances between strains were derived using the Pam-Dayhoff model of amino acid substitution as implemented in MEGA version 3.1 [65] with gamma-correction (alpha = 2.5) and pairwise deletion of missing data.

#### **Authors' contributions**

JPS conceived the experiments, conducted genetic and phylogenetic analyses, and wrote the paper. CLMcC conducted all PCR and sequencing experiments, analyzed

data, and contributed to writing the manuscript. JLT assisted in experimental design and helped write the manuscript. SVW provided cell culture expertise, reagents, and helped write the manuscript. MR collected blood samples from animals in the wild, conducted cell culture of subtype B and helped write the manuscript. KS conducted cell culture of subtype E and helped in writing the manuscript. CW and HW contributed expertise and essential logistic support in obtaining lion blood samples. SJ O'B contributed expertise, reagents, and helped write the manuscript. All authors have read and approved the final version of the manuscript.

## Additional material

### Additional file 1

Transcription factors present within FIV<sub>PLV</sub> LTR of lion subtypes B and E. These motifs were identified by the setting a threshold similarity score of 85% for screening against the TRANSFAC database at the website <http://motif.genome.jp/>.

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1471-2164-9-66-S1.doc>]

### Additional file 2

Primers for lion FIV amplification. The primers span the entire proviral genome of subtype E (Ple1027) and subtype B (Ple458). Shown are the primer sequence, relative position and orientation.

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1471-2164-9-66-S2.XLS>]

### Additional file 3

Parameters used in PAUP\* analyses for LTR, each viral gene and combined analyses. These parameters were determined using the program Modeltest (see main text) and were implemented for the maximum likelihood and minimum evolution analyses in PAUP.

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1471-2164-9-66-S3.XLS>]

## Acknowledgements

All tissue samples were collected in full compliance with specific Federal Fish and Wildlife permits from the Conservation of International Trade in Endangered Species of Wild flora and Fauna: Endangered and Threatened Species. Captive Bred issued to the National Cancer Institute (NCI)-National Institutes of Health (NIH) (S.J.O. principal officer) by the U.S. Fish and Wildlife Services of the Department of the Interior. This project has supported in whole and in part with federal funds from the National Cancer Institute, National Institutes of Health, under contract N01-CO-12400. This research has been supported (in part) by the Intramural Research Program of the NIH, NCI, Center for Cancer Research. The content of this publication does not necessarily reflect the views or policies of the Department of Health and Human Services, nor does mention of trade names, commercial products, or organizations imply endorsement by the U.S. Government.

## References

- Olmsted RA, Langley R, Roelke ME, Goeken RM, Adger-Johnson D, Goff JP, Albert JP, Packer C, Laurenson MK, Caro TM, Scheepers I, Wildt D, Bush M, Martenson JS, O'Brien SJ: **Worldwide prevalence of lentivirus infection in wild feline species: epidemiologic and phylogenetic aspects.** *J Virol* 1992, **66(10)**:6008-6018.
- Driciru M, Siefert L, Prager KC, Dubovi E, Sande R, Princee F, Friday T, Munson L: **A serosurvey of viral infections in lions (Panthera leo), from Queen Elizabeth National Park, Uganda.** *J Wildl Dis* 2006, **42(3)**:667-671.
- Biek R, Ruth TK, Murphy KM, Anderson CR Jr, Johnson M, DeSimone R, Gray R, Hornocker MG, Gillin CM, Poss M: **Factors associated with pathogen seroprevalence and infection in Rocky Mountain cougars.** *J Wildl Dis* 2006, **42(3)**:606-615.
- Troyer JL, Pecon-Slattery J, Roelke ME, Johnson W, VandeWoude S, Vazquez-Salat N, Brown M, Frank L, Woodroffe R, Winterbach C, Hemson G, Bush M, Alexander KA, Revilla E, O'Brien SJ: **Seroprevalence and genomic divergence of circulating strains of feline immunodeficiency virus among Felidae and Hyaenidae species.** *J Virol* 2005, **79(13)**:8282-8294.
- Munson L, Marker L, Dubovi E, Spencer JA, Evermann JF, O'Brien SJ: **Serosurvey of viral infections in free-ranging Namibian cheetahs (Acinonyx jubatus).** *J Wildl Dis* 2004, **40(1)**:23-31.
- Troyer JL, Pecon-Slattery J, Roelke ME, Black L, Packer C, O'Brien SJ: **Patterns of feline immunodeficiency virus multiple infection and genome divergence in a free-ranging population of African lions.** *J Virol* 2004, **78(7)**:3777-3791.
- Carpenter MA, Brown EW, MacDonald DW, O'Brien SJ: **Phylogeographic patterns of feline immunodeficiency virus genetic diversity in the domestic cat.** *Virology* 1998, **251(2)**:234-243.
- Carpenter MA, Brown EW, Culver M, Johnson WE, Pecon-Slattery J, Brousset D, O'Brien SJ: **Genetic and phylogenetic divergence of feline immunodeficiency virus in the puma (Puma concolor).** *J Virol* 1996, **70(10)**:6682-6693.
- Hofmann-Lehmann R, Fehr D, Grob M, Elgizoli M, Packer C, Martenson JS, O'Brien SJ, Lutz H: **Prevalence of antibodies to feline parvovirus, calicivirus, herpesvirus, coronavirus, and immunodeficiency virus and of feline leukemia virus antigen and the interrelationship of these viral infections in free-ranging lions in east Africa.** *Clin Diagn Lab Immunol* 1996, **3(5)**:554-562.
- Brown EW, Yuhki N, Packer C, O'Brien SJ: **A lion lentivirus related to feline immunodeficiency virus: epidemiologic and phylogenetic aspects.** *J Virol* 1994, **68(9)**:5953-5968.
- Carpenter MA, O'Brien SJ: **Coadaptation and immunodeficiency virus: lessons from the Felidae.** *Curr Opin Genet Dev* 1995, **5(6)**:739-745.
- Biek R, Rodrigo AG, Holley D, Drummond A, Anderson CR Jr, Ross HA, Poss M: **Epidemiology, genetic diversity, and evolution of endemic feline immunodeficiency virus in a population of wild cougars.** *J Virol* 2003, **77(17)**:9578-9589.
- IUCN: **IUCN Red List of Threatened Species.** 2007 [<http://www.iucnredlist.org>].
- Langley RJ, Hirsch VM, O'Brien SJ, Adger-Johnson D, Goeken RM, Olmsted RA: **Nucleotide sequence analysis of puma lentivirus (PLV-14): genomic organization and relationship to other lentiviruses.** *Virology* 1994, **202(2)**:853-864.
- Olmsted RA, Barnes AK, Yamamoto JK, Hirsch VM, Purcell RH, Johnson PR: **Molecular cloning of feline immunodeficiency virus.** *Proc Natl Acad Sci U S A* 1989, **86(7)**:2448-2452.
- Barr MC, Zou L, Long F, Hoose WA, Avery RJ: **Proviral organization and sequence analysis of feline immunodeficiency virus isolated from a Pallas' cat.** *Virology* 1997, **228(1)**:84-91.
- Burkala E, Poss M: **Evolution of feline immunodeficiency virus Gag proteins.** *Virus Genes* 2007.
- Troyer JL, VandeWoude S, Pecon-Slattery J, O'Brien SJ: **FIV cross-species transmission: an evolutionary perspective.** *Vet Immunol Immunopathol* in press.
- Bendinelli M, Pistello M, Lombardi S, Poli A, Garzelli C, Matteucci D, Ceccherini-Nelli L, Malvaldi G, Tozzini F: **Feline immunodeficiency virus: an interesting model for AIDS studies and an important cat pathogen.** *Clin Microbiol Rev* 1995, **8(1)**:87-112.
- Willett BJ, Flynn JN, Hosie MJ: **FIV infection of the domestic cat: an animal model for AIDS.** *Immunol Today* 1997, **18(4)**:182-189.
- Bull ME, Gebhard DG, Tompkins WA, Kennedy-Stoskopf S: **Poly-morphic expression in the CD8alpha chain surface receptor**

- of African lions (*Panthera leo*).** *Vet Immunol Immunopathol* 2002, **84(3-4)**:181-189.
22. Bull ME, Kennedy-Stoskopf S, Levine JF, Loomis M, Gebhard DG, Tompkins WA: **Evaluation of T lymphocytes in captive african lions (*Panthera leo*) infected with feline immunodeficiency virus.** *Am J Vet Res* 2003, **64(10)**:1293-1300.
  23. Roelke ME, Pecon-Slattery J, Taylor S, Citino S, Brown E, Packer C, Vandewoude S, O'Brien SJ: **T-lymphocyte profiles in FIV-infected wild lions and pumas reveal CD4 depletion.** *J Wildl Dis* 2006, **42(2)**:234-248.
  24. de Rozieres S, Mathiason CK, Rolston MR, Chatterji U, Hoover EA, Elder JH: **Characterization of a highly pathogenic molecular clone of feline immunodeficiency virus clade C.** *J Virol* 2004, **78(17)**:8971-8982.
  25. Talbott RL, Sparger EE, Lovelace KM, Fitch WM, Pedersen NC, Luciw PA, Elder JH: **Nucleotide sequence and genomic organization of feline immunodeficiency virus.** *Proc Natl Acad Sci U S A* 1989, **86(15)**:5743-5747.
  26. Sodora DL, Courcelle J, Brojatsch J, Berson A, Wang YC, Dow SW, Hoover EA, Ji M: **Analysis of a feline immunodeficiency virus provirus reveals patterns of gene sequence conservation distinct from human immunodeficiency virus type 1.** *AIDS Res Hum Retroviruses* 1995, **11(4)**:531-533.
  27. Poss M, Ross HA, Painter SL, Holley DC, Terwee JA, Vandewoude S, Rodrigo A: **Feline lentivirus evolution in cross-species infection reveals extensive G-to-A mutation and selection on key residues in the viral polymerase.** *J Virol* 2006, **80(6)**:2728-2737.
  28. MS GenomeNet 2007 [<http://motif.genome.jp>].
  29. Wernersson R, Pedersen AG: **RevTrans: Multiple alignment of coding DNA from aligned amino acid sequences.** *Nucleic Acids Res* 2003, **31(13)**:3537-3539.
  30. Biek R, Drummond AJ, Poss M: **A virus reveals population structure and recent demographic history of its carnivore host.** *Science* 2006, **311(5760)**:538-541.
  31. Poss M, Idoine A, Ross HA, Terwee JA, Vandewoude S, Rodrigo A: **Recombination in feline lentiviral genomes during experimental cross-species infection.** *Virology* 2007, **359(1)**:146-151.
  32. Vigne JD, Guilaine J, Debue K, Haye L, Gerard P: **Early taming of the cat in Cyprus.** *Science* 2004, **304(5668)**:259.
  33. Driscoll CA, Mullikan JC, Mennotti-Raymond M, Roca AL, Pontius JU, Stephens R, Johnson WE, Lindblad-Toh K, Smith D, Geffan E, Harley e, Godoy J, Pontier D, Kitchener A, Yamaguchi N, Macdonald DW, O'Brien SJ: **The domestication of cats—Origins and genomic landscape.** *Science* 2007, **317**:519-523.
  34. Johnson WE, Eizirik E, Pecon-Slattery J, Murphy WJ, Antunes A, Teeling E, O'Brien SJ: **The late Miocene radiation of modern Felidae: a genetic assessment.** *Science* 2006, **311(5757)**:73-77.
  35. Fromont E, Sager A, Leger F, Bourguemestre F, Jouquelet E, Stahl P, Pontier D, Artois M: **Prevalence and pathogenicity of retroviruses in wildcats in France.** *Vet Rec* 2000, **146(11)**:317-319.
  36. Antunes A, Troyer JL, Roelke ME, Pecon-Slattery J, Packer C, Winterbach C, Winterbach H, Hemson G, Frank L, Stander P, Siefert L, Driciru M, Funston PJ, Alexander KA, Prager KC, Mills G, Wildt DE, Bush M, O'Brien SJ, Johnson WE: **The evolutionary history of lions: Integrating host/pathogen molecular genetics.** in press.
  37. Pecon-Slattery J, Troyer JL, Johnson WE, O'Brien SJ: **Evolution Of Feline Immunodeficiency Virus In Felidae: Implications For Human Health And Wildlife Ecology.** *Vet Immunol Immunopathol* in press.
  38. Phillips TR, Lamont C, Konings DA, Shacklett BL, Hamson CA, Luciw PA, Elder JH: **Identification of the Rev transactivation and Responsive elements of feline immunodeficiency virus.** *J Virol* 1992, **66(9)**:5464-5471.
  39. Smirnova N, Troyer JL, Schissler J, Terwee J, Poss M, VandeWoude S: **Feline lentiviruses demonstrate differences in receptor repertoire and envelope structural elements.** *Virology* 2005, **342(1)**:60-76.
  40. Willett BJ, Hosie MJ, Neil JC, Turner JD, Hoxie JA: **Common mechanism of infection by lentiviruses.** *Nature* 1997, **385(6617)**:587.
  41. Lombardi S, Garzelli C, Pistello M, Massi C, Matteucci D, Baldinotti F, Cammarota G, da Prato L, Bandecchi P, Tozzini F, et al.: **A neutralizing antibody-inducing peptide of the V3 domain of feline immunodeficiency virus envelope glycoprotein does not induce protective immunity.** *J Virol* 1994, **68(12)**:8374-8379.
  42. Lombardi S, Massi C, Tozzini F, Zaccaro L, Bazzichi A, Bandecchi P, La Rosa C, Bendinelli M, Garzelli C: **Epitope mapping of the V3 domain of feline immunodeficiency virus envelope glycoprotein by monoclonal antibodies.** *J Gen Virol* 1995, **76(Pt 8)**:1893-1899.
  43. Tozzini F, Matteucci D, Bandecchi P, Baldinotti F, Siebelink K, Osterhaus A, Bendinelli M: **Neutralizing antibodies in cats infected with feline immunodeficiency virus.** *J Clin Microbiol* 1993, **31(6)**:1626-1629.
  44. Hohdatsu T, Hirabayashi H, Motokawa K, Koyama H: **Comparative study of the cell tropism of feline immunodeficiency virus isolates of subtypes A, B and D classified on the basis of the env gene V3-V5 sequence.** *J Gen Virol* 1996, **77(Pt 1)**:93-100.
  45. Vahlenkamp TW, Verschoor EJ, Schuurman NN, van Vliet AL, Horzinek MC, Egberink HF, de Ronde A: **A single amino acid substitution in the transmembrane envelope glycoprotein of feline immunodeficiency virus alters cellular tropism.** *J Virol* 1997, **71(9)**:7132-7135.
  46. Verschoor EJ, Boven LA, Blaak H, van Vliet AL, Horzinek MC, de Ronde A: **A single mutation within the V3 envelope neutralization domain of feline immunodeficiency virus determines its tropism for CRFK cells.** *J Virol* 1995, **69(8)**:4752-4757.
  47. Willett BJ, Hosie MJ: **The role of the chemokine receptor CXCR4 in infection with feline immunodeficiency virus.** *Mol Membr Biol* 1999, **16(1)**:67-72.
  48. HIV, sequence, database [<http://www.hiv.lanl.gov/content/sequence/HIV/CRFs/CRFs.html>]
  49. Sharp PM, Robertson DL, Hahn BH: **Cross-species transmission and recombination of 'AIDS' viruses.** *Philos Trans R Soc Lond B Biol Sci* 1995, **349(1327)**:41-47.
  50. Robertson DL, Hahn BH, Sharp PM: **Recombination in AIDS viruses.** *J Mol Evol* 1995, **40(3)**:249-259.
  51. Jin MJ, Hui H, Robertson DL, Muller MC, Barre-Sinoussi F, Hirsch VM, Allan JS, Shaw GM, Sharp PM, Hahn BH: **Mosaic genome structure of simian immunodeficiency virus from west African green monkeys.** *Embo J* 1994, **13(12)**:2935-2947.
  52. Beer BE, Bailes E, Dapolito G, Campbell BJ, Goeken RM, Axthelm MK, Markham PD, Bernard J, Zagury D, Franchini G, Sharp P, Hirsch VM: **Patterns of Genomic Sequence Diversity among Their Simian Immunodeficiency Viruses Suggest that L'Hoest Monkeys (*Cercopithecus lhoesti*) Are a Natural Lentivirus Reservoir.** *J Virol* 2000, **74**:3892-3898.
  53. Salemi M, De Oliveira T, Courgnaud V, Moulton V, Holland B, Cassol S, Switzer WM, Vandamme A-M: **Mosaic Genomes of the Six Major Primate Lentivirus Lineages Revealed by Phylogenetic Analyses.** *J Virol* 2003:7202-7213.
  54. Sharp PM, Shaw GM, Hahn BH: **Simian immunodeficiency virus infection of chimpanzees.** *J Virol* 2005, **79(7)**:3891-3902.
  55. Miyazawa T, Furuya T, Itagaki S, Tohya Y, Takahashi E, Mikami T: **Establishment of a feline T-lymphoblastoid cell line highly sensitive for replication of feline immunodeficiency virus.** *Arch Virol* 1989, **108(1-2)**:131-135.
  56. Snyder HW Jr, Hardy WD Jr, Zuckerman EE, Fleissner E: **Characterisation of a tumour-specific antigen on the surface of feline lymphosarcoma cells.** *Nature* 1978, **275(5681)**:656-658.
  57. Sodora DL, Courcelle J, Brojatsch J, Berson A, Wang YC, Dow SW, Hoover EA, Mullins JL: **Analysis of a feline immunodeficiency virus provirus reveals patterns of gene sequence conservation distinct from human immunodeficiency virus type 1.** *AIDS Res Hum Retroviruses* 1995, **11(4)**:531-533.
  58. Olmsted RA, Hirsch VM, Purcell RH, Johnson PR: **Nucleotide sequence analysis of feline immunodeficiency virus: genome organization and relationship to other lentiviruses.** *Proc Natl Acad Sci U S A* 1989, **86(20)**:8088-8092.
  59. BLAST [<http://www.ncbi.nlm.nih.gov>]
  60. Whitwam T, Peretz M, Poeschla E: **Identification of a central DNA flap in feline immunodeficiency virus.** *J Virol* 2001, **75(19)**:9407-9414.
  61. Thompson JD, Gibson TJ, Plewniak F, Jeanmougin F, Higgins DG: **The CLUSTAL\_X windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools.** *Nucleic Acids Res* 1997, **25(24)**:4876-4882.
  62. Rambaut A: **Se-AI: Sequence Alignment Editor.** Oxford, UK: University of Oxford 1996.
  63. Posada D, Crandall KA: **MODELTEST: testing the model of DNA substitution.** *Bioinformatics* 1998, **14(9)**:817-818.

64. Swofford D: **Phylogenetic Analysis Using Parsimony (\*and Other Methods)**. version 4 edition. Sunderland, Massachusetts: Sinauer Associates; 2002.
65. Kumar S, Tamura K, Nei M: **MEGA3: Integrated software for Molecular Evolutionary Genetics Analysis and sequence alignment**. *Brief Bioinform* 2004, **5(2)**:150-163.

Publish with **BioMed Central** and every scientist can read your work free of charge

*"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."*

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:  
[http://www.biomedcentral.com/info/publishing\\_adv.asp](http://www.biomedcentral.com/info/publishing_adv.asp)

