











Use of Poultry By-product and Plant Protein Sources in Diets of Redclaw Crayfish (*Cherax quadricarinatus*)

Orhan Tufan Eroldoğan^{1,*} , Mabrouk Elsabagh^{2,*} , Hüseyin Sevgili³ , Brett Glencross⁴ , Marina Paolucci⁵ , Metin Kumlu¹ , Enes Kinay¹ , Ece Evliyaoğlu¹ , Hatice Asuman Yılmaz¹ , Merve Sariipek⁶ 

¹Çukurova University, Faculty of Fisheries, Department of Aquaculture, 01330, Adana, Turkey

²Niğde Ömer Halisdemir University, Faculty of Agricultural Sciences and Technologies, Department of Animal Production and Technologies, 51240, Niğde, Turkey

³Isparta University of Applied Sciences, Eğirdir Faculty of Fisheries, Department of Aquaculture, Isparta, Turkey

⁴University of Stirling, Institute of Aquaculture, Stirling FK9 4LA, United Kingdom

⁵University of Sannio, Department of Science and Technologies, 82100 Benevento, Italy

⁶Sinop University, Faculty of Fisheries, Department of Aquaculture, 57000 Sinop, Turkey

How to cite

Eroldoğan, O.T., Elsabagh, M., Sevgili, H., Glencross, B., Paolucci, M., Kumlu, M., Kinay, E., Evliyaoğlu, E., Yılmaz, H.A., Sariipek, M. (2022). Use of Poultry By-product and Plant Protein Sources in Diets of Redclaw Crayfish (*Cherax quadricarinatus*) *Turkish Journal of Fisheries and Aquatic Sciences*, 22(8), TRJFAS21188. <http://doi.org/10.4194/TRJFAS21188>

Article History

Received 20 December 2021

Accepted 04 March 2022

First Online 07 March 2022

Corresponding Author

Tel.: +903223386084

E-mail: mtufan@cu.edu.tr

mabrouk.elsabagh@vet.kfs.edu.eg

Keywords

Poultry meal

Fish meal

Alternative sources

Fatty acid profile

Essential amino acid index

Abstract

A total of 300 juvenile crayfish (13.0±0.03 g) were randomly distributed among 5 dietary groups (n=60, 3 replicates) held within 15×500 L⁻¹ fiberglass tanks connected to a recirculation system (RAS), at 20 crayfish per tank. Each group was fed for 12 weeks one of five experimental diets where the main protein sources were: 1) control, fish-meal-based diet (FM, 48% of the diet); 2) 10% FM + 52.5% poultry by-product meal (PoM); 3) 34.5% soybean meal + 34.5% corn gluten meal (Pmix); 4) 34.5% PoM + 32.5% soybean/corn gluten meal mix (PoM/Pmix); and 5) 10% FM + 27.5% soybean + 27.5% corn gluten meal (FM/Pmix). The results demonstrated that there were no significant differences among diets in terms of growth and feed utilization efficiency. Muscle amino acid profile of redclaw crayfish fed the FM diet had the highest level of total essential amino acids, followed by FM/PMix, Pmix, PoM/Pmix, and PoM diets. Particularly, in all experimental groups, the highest essential amino acids (EAA) were lysine, arginine, and leucine. Based on these findings, we conclude that redclaw can perform well with FM-free vegetable diets and PoM-based diets although more research is needed into the total composition of EAA and FA in muscle.

Introduction

The freshwater crustacean redclaw crayfish, *Cherax quadricarinatus*, is characterized by a range of positive biological, ecological and commercial attributes for aquaculture (Jones, 1995; Saoud et al., 2012; Saoud et al., 2013). These attributes have contributed to the expansion of its aquaculture across the world beyond its native origin in North-Eastern Australia (Zeng et al., 2019). Given that redclaw crayfish is a highly sought after food resource for human consumption, being the second most globally cultured crayfish species after red swamp crayfish, *Procambarus clarkii* (FAO, 2020) The research has long been devoted to develop practical

diets enhancing a successful culture of such crayfish species in pond based production.

Since the cost of feed for crustacean production accounts for 70% of the total cost of production, it is important to identify options for low-cost feeds for commercial-scale production without sacrificing essential dietary nutrients (Thompson et al., 2005; Pavasovic, 2008). For decades, nutritionists have endeavoured to develop aquafeed formulations that support or enhance the growth of cultured redclaw crayfish while controlling costs (Pavasovic, 2008). Numerous researchers have collectively made great strides in addressing the many constraints associated with optimal feed formulation in crustaceans (Webster

et al., 2004; Thompson *et al.*, 2005; Thompson *et al.*, 2006; Saoud *et al.*, 2008, Venero *et al.*, 2008; Garza *et al.*, 2012; Saoud *et al.*, 2012). While the cultivation of omnivorous aquatic species easily switched to feeds containing minimal amounts of fish meal or oil (Turchini *et al.*, 2019). Until recently, replacing or minimizing the inclusion of fishmeal in commercial red swamp crayfish diets examined only the use of vegetable proteins (Tan *et al.*, 2018). However, more recently potential alternatives to fish meal including the by-products of animal-based product chains such as poultry, have been considered potentially useful sources of animal protein (Pavasovic *et al.*, 2007; Saoud *et al.*, 2008; Garza de Yta *et al.*, 2012).

To date, the substitution of fish meal with plant or animal by-product proteins has been mostly reported to produce cost-effective diet formulations without any detrimental effects on the production performance of crustaceans, particularly penaeids (Muzinic *et al.*, 2004; Thompson *et al.*, 2005; 2006; Saoud *et al.*, 2008; Fuertes *et al.*, 2012; 2013; 2014; Glencross *et al.*, 2014; Glencross *et al.*, 2018; Tan *et al.*, 2018). However, regarding crayfish, the efforts have been partially limited, compared to other commercially cultivated penaeid shrimps (Saoud *et al.*, 2008; Lunda Roy *et al.*, 2020).

Most recently, Qian *et al.* (2021) examined four plant protein sources (cottonseed, soybean, peanut and rapeseed meal) incorporated at 30% inclusion level in commercial diets for redclaw. However, there are limited available data for the growth performance in redclaw crayfish fed with 100% plant and/or animal by-product meal sources under intensive culture conditions such as in closed recirculating aquaculture systems. Rodríguez-Canto *et al.*, 2002) and García-Ulloa *et al.* (2012) examined the growth performance of redclaw crayfish fed with a commercial diet containing 35% protein at high stocking density. However, in the light of the previous findings, there is still a need to elucidate the plant and/or animal-based protein sources for optimum growth of redclaw crayfish cultured in intensive culture conditions. Given the importance of endogenous productivity to the nutrition of crustaceans farmed in pond systems (Gamboa-Delgado, 2014), we hypothesised that their nutrition would be subject to additional sensitivities under intensive culture conditions.

Several studies have attempted to assess the growth performance of the crayfish in either indoor or outdoor systems. In particular, distinct differences in growth performance were observed in these studies when the protein sources and level of the nutrients in the experimental diets were altered (Rodríguez-Canto *et al.*, 2002; García-Ulloa *et al.*, 2012; Saoud *et al.*, 2012). As carbon, nitrogen and phosphorous outputs are generally a greater concern in high stocking density in crayfish culture systems feed is the ultimate source of those elements a significant effort has been devoted to reducing their losses (Van Rijn, 2013; Schneider *et al.*,

2004). Despite these significant findings, no studies have been recently evaluated the suitability of various plant and animal by-product-based ingredients and their combination as potential substitutes for fish meal for redclaw feed formulation. Therefore, aim of this study was to to develop economic and sustainable diets for the production of redclaw crayfish. The impacts of dietary modifications on growth performances, proximate body composition, muscle percentage of essential amino acids and fatty acids as well as nutrient retentions are reported.

Materials and Methods

Crayfish Rearing System and Experimental Diets

The juvenile redclaw crayfish were produced at the Research and Development Center in the Faculty of Fisheries, Cukurova University, Adana, Turkey. The juveniles were collected and placed in an indoor nursery tank system for three weeks before the experiment. Individually selected and weighed crayfish were stocked into one of 15 cylindrical fiberglass tanks (500 L, 0.5 m² bottom area) connected to the RAS. Within the RAS system, water was re-circulated through a sand filter and a biofilter aerated by air stones distributed along with the experimental tanks. The recycling rate of water in the system was maintained at 3-4 L min⁻¹ throughout the experiment. At the beginning of the experiment, 300 acclimated crayfish juveniles (13.0±0.03 g) were randomly distributed into five treatment groups with triplicate fiberglass tanks, at 20 crayfish per tank (40 crayfish m⁻²). Each tank contained two pieces of 15 × 15 cm hollow bricks used for refuge of individuals and one air stone for aeration and water mixing. During the experimental period, water conditions were measured weekly with mean ± S.D. as follows: temperature 25.47±1.97°C, pH 7.85±0.41 (Hanna pH-meter), dissolved oxygen content 6.5-7.0 mg L⁻¹ (YSI Oxygen meter, USA), and total ammonia nitrogen content <0.05 mg L⁻¹, nitrite from 0 to 0.5 mg L⁻¹, nitrate from 0.25 to 100 mg L⁻¹. Nitrite, nitrate, ammonium, and phosphate concentrations were measured using Aquamerck® (Darmstadt, Germany) reagent kits. All water quality parameters were considered within optimum levels for redclaw crayfish.

Each tank of crayfish was fed for 12 weeks with one of five isolipidic (crude lipid, %=9.1±0.18) and isonitrogenous (crude protein, %=38.0±0.36) experimental diets. The protein contents of each diet were based on (Tables 1 and 2):

- 1) Fish meal only (FM, 48% of the diet);
- 2) 10% FM + 52.5% poultry by-product meal (PoM);
- 3) 34.5% soybean meal + 34.5% corn gluten meal (Pmix);
- 4) 10% FM + 27.5% soybean + 27.5% corn gluten meal (FM/Pmix);
- 5) 34.5% PoM + 32.4 SBM/CGM mix (PoM/Pmix).

All ingredients used as protein sources were ground to pass through a 100 – 150 μ size screen and were thoroughly homogenized. Other components, such as mineral and vitamin premixes, were mixed and added into the blended ingredients. Fish oil and water (150 ml) were added to the dry ingredients and completely mixed to form a homogeneous mixture. Experimental diets were pelleted with a diameter of 1.9 mm using a pellet machine (Pasfil Machinery, Istanbul, Turkey), then kept at 110°C and 0.5 bar for 30 min in a pressure vessel. This process increases the water stability of the experimental diets according to Kumlu (2018). All pellets were air-dried to approximately 10% moisture and stored at -20°C until used.

All groups of crayfish were fed three times daily at 09:00, 13:00 and 18:00 at 12% body weight per day in the first week and then gradually reduced to 3-5% body weight day⁻¹ by the end of the experiment. Uneaten feed and faecal waste were removed daily by siphoning before the first feeding. Collected uneaten feed was used to calculate feed intake after drying at 40°C for 6 hours. The experimental tanks were visually checked daily for the presence of exuvia as described by Barki *et al.* (1997) and the average number of the exuvia was recorded.

Sampling and Analysis

At the commencement of the trial, five individuals from a common pool crayfish were sampled randomly and stored at -20°C for subsequent determination of initial proximate composition. At the end of the trial, three crayfish per tank (n=3, N=9 per treatment) were sacrificed and frozen until analysis. Proximate body composition analysis was performed according to AOAC (1995) and the measurement of the fatty acid profile of the experimental diets and the whole body was carried out following the total lipid extraction (Folch *et al.*, 1957). The amino acid analyses of the experimental diets and muscle were determined using the Ultra-Fast Liquid Chromatography system with a UV detector (Gheshlaghi *et al.*, Douglas, 2008).

Calculations and Statistical Analysis

The growth and feed utilization parameters were calculated as follows: specific growth rate (SGR %day⁻¹)=100 (Ln final weight – Ln initial weight) t⁻¹, weight gain (WG, %)=(final weight-initial weight)/initial weight x 100, feed conversion ratio

(FCR)=dry feed intake (g)/weight gain (g).

Lipid, protein and energy retention (deposition) were calculated as follows: nutrient gain (g kg ABW day⁻¹)=(final body nutrient content (g) – initial body nutrient content (g))/kg ABW/days, nutrient gain (g kg ABW day⁻¹)=(final body nutrient content (g) – initial body nutrient content (g))/kg ABW/days, nutrient retention (%)=100 × (nutrient gain/nutrient intake). Average body weight (ABW)=(final weight (g) + initial weight (g))/2 (Akpınar *et al.*, 2012). Gross energy was calculated using conversion factors of 39.5, 23.6, and 17.2 kJ g⁻¹ for lipid, protein, and carbohydrate, respectively (NRC, 2011).

Essential amino acid index (EAAI), based on A/E ratios, was calculated according to Peñaflores (1989) and Gunasekera *et al.*, (2002). According to EAAI derivation, the following equations were applied:

$$A/E \text{ ratio} = (\text{essential amino acid} / \text{total essential amino acids}) \times 100$$

$$aa/AA = A/E \text{ ratio in feed} / A/E \text{ ratio in crayfish}$$

$$EAAI = \sqrt[n]{\frac{aa1}{AA1} \times \frac{aa2}{AA2} \times \dots \times \frac{aan}{AA_n}}$$

The numerical values of aa1/AA1, aa2/AA2... were set to 0.01 minimum and an EAAI closer to 1 is indicative of the degree of conformity of the EAA profiles of the ingredient used in the experimental crayfish diets and therefore, the potential of the ingredient to meet the essential amino acid requirements of the species.

Data were analyzed by one-way Analysis of Variance (ANOVA) at a significance level of 0.05%, after confirmation of normality and homogeneity of variance. After all variables were homogenous and normally distributed, significant differences were detected. Then the data were subjected to a Tukey-Kramer post hoc test for identifying homogeneous subsets. All statistical analyses were performed using SPSS 11.5 (SPSS Inc., Chicago, IL, USA). The significance level was set at P<0.05. The results are presented as means \pm standard deviations (n=3), except amino acid results (n=2).

Results

The experimental diets were observed to be reasonably water-stable and acceptable by the crayfish. The results of dietary amino acid profiles (Table 2) revealed that each diet was rich in one or more essential

Table 1. Proximate composition (%) of the test ingredients used in the trial

	Protein	Lipids	Dry matter	Ash	Carbohydrate*
Fish meal	64.9	9.7	91.2	12.1	4.5
Soybean meal	40.1	11.7	91.9	7.6	32.2
Corn gluten meal	56.5	3.4	89.3	4.5	25
Poultry by-product meal	55.5	12.5	89.8	10.7	9.5
Wheat meal	10.8	1.4	88.5	0.5	75.8

*Carbohydrate = Dry matter – (protein+lipid+ash)

Table 2. Formulation and proximate composition of the experimental diets

Ingredients	FM	PoM	Pmix	PoM/Pmix	FM/Pmix
Fish meal (Anchovy meal) ¹	48.0	10.0	-	-	10.0
Soybean meal ¹	-	-	34.5	16.2	27.5
Corn gluten meal ¹	-	-	34.5	16.2	27.5
Poultry by-product meal ²	-	52.5	-	34.5	-
Wheat meal ¹	42.5	24.3	16.1	18.5	21.5
Fish oil ¹	2.4	2.7	2.4	2.5	2.5
Soy lecithin (70%) ³	2.0	2.0	2.0	2.0	2.0
Gluten (wheat) ⁴	3.1	3.1	3.1	3.1	3.1
Dicalcium phosphate ⁴	-	1.1	1.8	1.8	1.0
CaCO ₃ ⁵	-	2.1	2.9	2.8	2.5
DL-methionine ¹	-	0.1	0.1	0.1	0.1
L-Lysine ¹	-	0.1	0.6	0.3	0.3
Vitamin premix ^{†,1}	0.8	0.8	0.8	0.8	0.8
Mineral premix ^{‡,1}	0.2	0.2	0.2	0.2	0.2
Vitamin C (Stay C) ¹	0.1	0.1	0.1	0.1	0.1
<i>Composition (% DM unless otherwise stated)</i>					
Protein	38.1±0.2	40.7±0.3	38.2±0.1	39.4±0.02	38.0±0.4
Lipids	9.1±0.28	11.3±0.16	9.2±0.27	10.9±0.14	9.3±0.28
Dry matter	89.6±0.2	89.6±0.3	90.1±0.4	89.8±0.2	90.0±0.3
Ash	6.6±0.2	11.1±0.4	9.0±0.3	10.8±0.3	8.3±0.2
Energy (kJ g ⁻¹)	18.7±0.9	18.6±0.2	18.6±0.4	18.5±0.6	18.5±0.4
<i>EAA (g/100 g)</i>					
Arginine	2.0±0.00	1.9±0.01	1.1±0.00	1.1±0.00	1.0±0.00
Histidine	1.0±0.03	1.0±0.02	0.8±0.01	0.8±0.01	1.0±0.01
Isoleucine	1.8±0.01	2.1±0.01	1.7±0.01	2.4±0.01	1.8±0.01
Leucine	3.4±0.01	3.5±0.01	5.1±0.04	5.0±0.02	5.1±0.01
Lysine	3.1±0.01	2.8±0.04	2.7±0.01	2.0±0.01	4.0±0.02
Methionine	0.9±0.01	0.8±0.00	0.6±0.00	0.8±0.00	0.9±0.01
Phenylalanine	1.9±0.01	2.2±0.01	2.5±0.01	3.2±0.01	2.7±0.00
Threonine	1.3±0.00	2.0±0.01	1.7±0.00	2.0±0.01	1.4±0.00
Valine	2.2±0.01	3.0±0.02	1.8±0.01	3.3±0.01	2.1±0.01
<i>NEAA (g/100 g)</i>					
Alanine	2.4±0.01	2.9±0.01	2.6±0.01	3.2±0.01	2.7±0.01
Tyrosine	1.4±0.03	1.7±0.01	1.8±0.01	1.9±0.00	1.9±0.01
Aspartate	2.5±0.01	1.5±0.00	1.1±0.01	0.8±0.01	1.2±0.01
Glutamate	6.7±0.03	5.1±0.02	7.2±0.02	3.3±0.04	6.8±0.03
Glycine	2.3±0.01	4.1±0.01	1.9±0.01	3.3±0.01	2.0±0.01
Serine	1.3±0.01	2.6±0.02	2.0±0.01	1.7±0.01	1.9±0.01
<i>CEAA (g/100g)</i>					
Proline	2.4±	4.8±	4.5±	7.4±	4.1±
Cystine	vest	vest	vest	ND	vest
Hydroxyproline	vest	vest	ND	ND	vest
ΣAA	36.6±	42.0±	39.1±	42.2±	40.6±
ΣEAA	16.4±	19.3±	18±	20.6±	20.0±
<i>Fatty acid (mole%)</i>					
Total SFA	30.1±0.40	24.9±0.09	19.5±0.01	23.4±0.46	23.4±0.22
Total MUFA	25.0±0.20	32.1±0.12	38.2±0.25	30.7±0.85	27.2±0.16
Total PUFA	43.8±0.59	42.3±0.18	41.9±0.05	45.4±0.38	48.7±0.41
n-6 LC PUFA	1.1±0.01	0.8±0.01	1.3±0.01	0.8±0.00	0.9±0.01
n-6 PUFA	18.2±0.03	31.4±0.06	33.3±0.04	37.4±0.65	35.9±0.34
n-3 LC PUFA	25.0±0.29	10.4±0.27	7.90±0.06	7.67±0.32	12.5±0.47
n-3 PUFA	25.6±0.59	10.9±0.23	8.53±0.02	8.08±0.31	12.8±0.74

¹Sibal Corporation, Sinop, Turkey (Anchovy meal-crude protein 70%).²Egehan Feed, (Poultry meal rendering and bone meal rendering-pet food grade) Turkey.³Tito Soy lecithin, Germany, Smart Chemical Trading and Consultancy Company Limited, Turkey.⁴Kimetsan Ltd., Turkey.⁵Sigma Aldrich Co Ltd, St. Louis, USA.[†]Vitamin Premix (mg/kg): retinyl acetate 2.58 mg, DL-cholecalciferol 0.037 mg, DL-α tocopheryl acetate 30 mg, menadione sodium bisulphite 2.5 mg, 4hiamine 7.5 mg, riboflavin 15 mg, pyridoxine 7.5 mg, nicotinic acid 87.5 mg, folic acid 2.5 mg, calcium pantothenate 2.5 mg, vitamin B12 0.025 mg, ascorbic acid 250 mg, inositol 500 mg, biotin 1.25 mg and choline chloride 500 mg.[‡]Mineral Premix (mg/kg): calcium carbonate (40% Ca) 2.15 g, magnesium hydroxide (60% Mg) 1.24 g, potassium chloride 0.9 g, ferric citrate 0.2 g, potassium iodine 4 mg, sodium chloride 0.4 g, calcium hydrogen phosphate 50 g, copper sulphate 0.3 g, zinc sulphate 40 g, cobalt sulphate 2 g, manganese sulphate 30 g, sodium selenite 0.3 g.EAA, essential amino acids; CEAA, conditionally essential amino acids; NEAA, nonessential amino acids; vest, vestigial amount of amino acid (< 0.01 mg g⁻¹); ND, not determined.

amino acids. The dietary total EAA was similar among the experimental diets. Of the EAA, leucine and lysine were the most abundant (ranged from 3.4 to 5.1 g/100 g and 2 to 4 g/100 g, respectively), while methionine was the least abundant (0.6-0.9 g/100 g) (Table 2). The FM diet was characterized by a high content of saturated fatty acid (SFA), n-3 polyunsaturated fatty acids (n-3 PUFA), n-3 long-chain polyunsaturated fatty acids (n-3 LC PUFA), including eicosapentaenoic acid (EPA, 20:5n-3) and docosahexaenoic acid (DHA, 22:6n-3). As expected, the Pmix was higher in monounsaturated fatty acids (MUFA) and n-6 long-chain polyunsaturated fatty acids (n-6 LC-PUFA) while the PoM/Pmix was higher in n-6 polyunsaturated fatty acids (n-6 PUFA) (Table 2).

Growth performance parameters, survival and molt cycle data of each treatment are presented in Table 3. The results indicated that there were no significant differences among diets in terms of growth performance, SGR, WG, FCR, survival, and exuviation number ($P>0.05$). Although not significant, the molt cycle was longer in crayfish fed PoM/Pmix while survival was higher in crayfish fed FM.

The moisture content ranged between the highest significant value of 79.7% (PoM) and the lowest significant value of 77.7% (FM). The protein content ranged between the highest significant value of 22.1% (FM/Pmix) and the lowest significant value of 17.5 (PoM) ($P<0.05$). The lipid content ranged between the highest significant value of 1.07% (PoM/Pmix) and the lowest significant value of 0.96% (PoM) ($P>0.05$). Ash content ranged between the highest significant value of 1.47 (FM) and the lowest significant value 1.27 (FM/Pmix) FM group was the lowest in moisture and the highest in ash. PoM group was the highest in moisture and the lowest in protein and lipids; Pmix group showed an average content of moisture, protein, lipids and ash; FM/Pmix group was as high in moisture as FM, the highest in protein and the lowest in ash ($P<0.05$) (Table 3).

While the muscle AA concentrations were identical (ranged from 79.8 to 84.2%); EAA, as proportion of the total AA, was ranged from 47.9 to 51.6% among the experimental diets, without any significant different ($P>0.05$) (Table 4). However, some specific amino acids were dominant across the experimental diets. In all groups, the highest EAA was lysine, followed by arginine, leucine (except than in PoM where leucine was higher than arginine), valine, isoleucine, phenylalanine, threonine, methionine (except than in PoM and PoM/Pmix where threonine was higher than methionine) and finally histidine ($P<0.05$). Crayfish fed FM were superior in arginine, lysine, methionine and total EAA contents. Crayfish fed PoM showed higher content of histidine, threonine, methionine and valine ($P<0.05$). The Pmix enhanced the isoleucine, phenylalanine and threonine contents in crayfish. Lysine and threonine were greater in crayfish fed PoM/Pmix. Overall, a correlation between the essential amino acid results of each experimental diet and the respective EAAI was 1.02 to 1.06 (Table 5).

Whole-body fatty acids composition is reported in Table 6. The SFAs were higher in PoM (23.4 mole %) and lower in FM (21.9 mole %) but there were no significant differences among Pmix, PoM/Pmix and FM/Pmix groups (22.1, 23.1 and 22.0 mole %, respectively). MUFA showed no significant differences among groups; however, the poultry meal-based diets (PoM and PoM/Mix) were lower in MUFA (26.6 and 26.8 mole %) than FM (28.0 mole %) and vegetable protein-based (Pmix and FM/Pmix, 28.2 and 27.0 mole %) groups. PoM/Pmix group showed the highest percentage of linoleic acid (18:2n-6, 16.8 mole %) and n-6 PUFA (19.4 mole %). The lowest levels of linoleic and n-6 PUFA were observed in the FM group (8.7 and 10.8 mole %, respectively). There were no differences in levels of linoleic and n-6 PUFA among PoM, Pmix and FM/Pmix groups ($P>0.05$). There were no significant differences among groups in n-6 LC-PUFA. Levels of n-3 LC PUFA and n-3 PUFA were higher in those crayfish fed the FM diet with respect to the other groups (38.6 and 39.1 mole %)

Table 3. Growth performance parameters and muscle compositions of redclaw crayfish, *Cherax quadricarinatus*, fed the five experimental diets for 12-weeks[†]

	Dietary treatments				
	FM	PoM	Pmix	PoM/Pmix	FM/Pmix
Initial W, g	13.0±2.2 ^a	12.8±0.0 ^a	13.2±0.0 ^a	13.0±0.0 ^a	12.9±0.0 ^a
Final W, g	20.7±0.42 ^a	20.2±0.98 ^a	19.6±0.99 ^a	20.2±0.95 ^a	20.8±0.99 ^a
SGR	0.50±0.024 ^a	0.54±0.058 ^a	0.47±0.09 ^a	0.52±0.04 ^a	0.56±0.05 ^a
WGR, %	59.3±3.21 ^a	57.8±7.70 ^a	48.5±4.89 ^a	55.4±7.37 ^a	61.±10.9 ^a
Feed intake, g	10.1±0.89 ^a	11.1±1.5 ^a	9.42±1.1 ^a	10.7±0.86 ^a	10.0±0.79 ^a
FCR	1.31±0.08 ^a	1.55±0.36 ^a	1.53±0.35 ^a	1.52±0.19 ^a	1.28±0.30 ^a
Survival %	86.7±10.4 ^a	76.7±11.5 ^a	81.7±2.89 ^a	75.0±2.89 ^a	81.7±0.0 ^a
Exuviation number	7.0±3.0 ^a	5.3±1.5 ^a	7.3±2.3 ^a	8.7±1.2 ^a	5.7±3.1 ^a
<i>Proximate composition (%)</i>					
Dry matter	22.3±0.49 ^a	20.3±1.25 ^a	21.0±0.63 ^a	21.8±1.01 ^a	20.4±1.04 ^a
Protein	20.1±0.35 ^b	17.5±0.20 ^d	18.6±0.60 ^c	19.9±0.00 ^b	22.1±0.35 ^a
Lipids	1.00±0.00 ^a	1.00±0.00 ^a	1.00±0.00 ^a	1.07±0.06 ^a	1.00±0.10 ^a
Ash	1.47±0.11 ^a	1.37±0.06 ^{ab}	1.33±0.6 ^{ab}	1.43±0.06 ^{ab}	1.27±0.06 ^b
Energy (kJ g ⁻¹)	5.20±0.05	4.63±0.16	4.86±0.07	5.12±0.02	5.60±0.09

[†]Data are expressed as mean ± SD. Values in the same row with different superscripts are significantly different by the Tukey post-hoc test ($P<0.05$).

Table 4. Essential amino acids in the muscle tissue (g/100 g) of redclaw crayfish fed the five experimental diets[‡]

	Dietary treatments				
	FM	PoM	Pmix	PoM/Pmix	FM/Pmix
<i>EAA</i>					
Arginine	7.37±0.27 ^a	6.31±0.00 ^d	6.61±0.07 ^{cd}	6.98±0.01 ^b	6.70±0.02 ^{bc}
Histidine	1.69±0.01 ^b	1.98±0.01 ^a	1.37±0.07 ^c	1.71±0.00 ^b	1.29±0.00 ^c
Isoleucine	3.77±0.02 ^b	3.81±0.02 ^{ab}	3.82±0.02 ^{ab}	3.46±0.02 ^c	3.85±0.00 ^a
Leucine	6.33±0.01 ^c	6.45±0.00 ^b	6.46±0.01 ^b	6.06±0.01 ^d	6.52±0.04 ^a
Lysine	12.21±0.29 ^a	9.22±0.09 ^c	11.48±0.15 ^b	11.86±0.05 ^{ab}	11.57±0.06 ^b
Methionine	1.77±0.02 ^{ab}	1.85±0.01 ^a	1.75±0.06 ^{bc}	1.68±0.01 ^c	1.84±0.02 ^a
Phenylalanine	3.53±0.01 ^b	3.49±0.01 ^c	3.54±0.02 ^{ab}	3.33±0.00 ^d	3.57±0.01 ^a
Threonine	2.25±0.13 ^b	2.46±0.00 ^a	2.31±0.08 ^{ab}	2.29±0.00 ^{ab}	2.23±0.00 ^b
Valine	4.23±0.03 ^b	4.30±0.01 ^a	4.17±0.03 ^{bc}	3.80±0.01 ^d	4.15±0.01 ^c
<i>NEAA</i>					
Alanine	4.53±0.06 ^b	5.10±0.00 ^a	4.60±0.00 ^b	4.20±0.00 ^c	4.60±0.00 ^b
Tyrosine	9.23±0.06 ^a	8.67±0.06 ^c	8.90±0.10 ^b	8.40±0.10 ^d	8.87±0.06 ^{bc}
Aspartate	3.20±0.00 ^b	3.30±0.00 ^a	3.17±0.06 ^b	2.87±0.06 ^c	3.20±0.00 ^b
Glutamate	4.50±0.00 ^c	5.20±0.00 ^a	4.67±0.06 ^b	4.53±0.06 ^c	4.77±0.06 ^b
Glycine	2.60±0.00 ^{ab}	2.70±0.00 ^a	2.60±0.10 ^{ab}	2.63±0.06 ^{ab}	2.50±0.00 ^b
Serine	13.67±0.15 ^b	15.20±0.10 ^a	13.27±0.25 ^c	12.93±0.06 ^c	13.00±0.00 ^c
<i>CEAA</i>					
Proline	3.30±0.00 ^a	3.27±0.06 ^{ab}	3.23±0.06 ^{ab}	3.10±0.00 ^c	3.20±0.08 ^b
Cystine	vest	vest	ND	ND	ND
Hydroxyproline	vest	vest	ND	ND	vest
ΣAA	84.2	83.2	81.9	79.8	81.9
ΣEAA	43.1	39.9	41.5	41.2	41.7
ΣAA/ΣAA (%)	51.2	47.9	50.7	51.6	51.0

[‡] Data are expressed as mean ± SD. Values in the same row with different superscripts are significantly different by the Tukey post-hoc test (P<0.05).

EAA, essential amino acids; CEAA, conditionally essential amino acids; NEAA, nonessential amino acids; vest, vestigial amount of amino acid (<0.01 mg g⁻¹); ND, not determined.

Table 5. Muscle essential amino acid indices (EAAI[†]) of the redclaw fed experimental diets[‡]

	Dietary treatments				
	FM	PoM	Pmix	PoM/Pmix	FM/Pmix
Arginine	0.75±0.01 ^b	0.78±0.00 ^a	0.62±0.00 ^c	0.55±0.00 ^d	0.56±0.00 ^d
Histidine	1.11±0.01 ^c	1.04±0.01 ^d	1.15±0.03 ^b	0.97±0.00 ^e	1.26±0.01 ^a
Isoleucine	1.01±0.00 ^c	1.06±0.00 ^b	1.01±0.00 ^c	1.19±0.01 ^a	1.00±0.00 ^c
Leucine	1.06±0.00 ^c	1.06±0.00 ^c	1.35±0.01 ^a	1.29±0.00 ^b	1.28±0.00 ^b
Lysine	1.02±0.01 ^a	0.79±0.00 ^c	0.74±0.03 ^d	0.58±0.00 ^e	0.85±0.00 ^b
Methionine	1.03±0.00 ^a	0.95±0.00 ^c	0.92±0.01 ^d	0.98±0.00 ^b	0.98±0.00 ^b
Phenylalanine	1.07±0.00 ^e	1.14±0.00 ^d	1.27±0.01 ^b	1.38±0.00 ^a	1.25±0.00 ^c
Threonine	1.12±0.03 ^b	1.29±0.00 ^a	1.29±0.04 ^a	1.31±0.00 ^a	1.16±0.00 ^b
Valine	1.03±0.00 ^c	1.20±0.00 ^b	1.01±0.02 ^d	1.32±0.01 ^a	1.02±0.00 ^{cd}
Average EAAI	1.02±0.01	1.04±0.00	1.04±0.02	1.04±0.00	1.06±0.01

[†] EAAI=Vaa/AA= V(A/E ratio in feed)/ (A/E ratio in crayfish)

[‡] Data are expressed as mean ± SD. Values in the same row with different superscripts are significantly different by the Tukey post-hoc test (P<0.05).

among which there were no significant differences (PoM, Pmix, PoM/Pmix and FM/Pmix, with 31.1, 30.8, 29.7, 32.8 mole % and 31.8, 31.8, 30.6 and 33.7 mole % for n-3 LC PUFA and n-3 PUFA, respectively (P>0.05). Alpha-linolenic acid (18:3n-3) content was significantly higher in PM/Pmix group (0.98 mole %) and lower in the FM (0.16 mole %) group. EPA (20:5-n3) content was higher in the FM group (18.3 mole %) and lower in the PoM (15.3 mole %) and PoM/Pmix (14.3 mole %) groups, while there were no significant differences between FM/Pmix (15.9 mole %) and Pmix (16.7 mole %) groups and between FM and Pmix groups. DHA (22:6-n3) was higher in FM/Pmix group (7.9 mole %) and lower in PoM/Pmix group (5.8 mole %), while there were no

differences among FM, PoM, and Pmix groups (6.6, 6.0, and 6.4 mole %, respectively).

Following the 12-week growth trial different lipid, protein and energy retention (deposition) responses were observed by the redclaw crayfish (Figure 1). Lipid retention (ranged from 10.1% to 13.1%) was similar across the diets (P>0.05). In contrast, nitrogen retention efficiency varied among treatments and was highest with the FM/Pmix diet (69.7%) and lowest with the PoM diet (34.4%) (P<0.05). In the present study, energy retention was significantly higher in the FM/Pmix, FM and PoM/Pmix diets compared with the PoM and Pmix diets that were comparable to each other (P>0.05).

Table 6. Muscle fatty acids composition (mole%) of redclaw fed the experimental diets[‡]

	Dietary treatments				
	FM	PoM	Pmix	PoM/Pmix	FM/Pmix
14:0	0.36±0.01 ^a	0.26±0.01 ^b	0.26±0.00 ^b	0.27±0.03 ^b	0.26±0.02 ^b
16:0	13.3±0.52 ^{ab}	13.6±0.16 ^{ab}	12.9±0.15 ^b	13.8±0.37 ^a	13.0±0.29 ^{ab}
18:0	8.0±0.25 ^c	9.4±0.05 ^a	8.8±0.05 ^b	8.9±0.26 ^{ab}	8.6±0.33 ^b
24:0	0.2±0.09 ^a	0.15±0.01 ^{ab}	0.13±0.01 ^b	0.14±0.01 ^b	0.15±0.01 ^{ab}
Total SFA ¹	21.9±0.67 ^b	23.4±0.24 ^a	22.1±0.21 ^{ab}	23.1±0.62 ^{ab}	22.0±0.62 ^{ab}
15:1	0.2±0.01 ^a	0.14±0.17 ^b	0.15±0.0 ^b	0.14±0.01 ^b	0.14±0.02 ^b
16:1n-7	1.7±0.05 ^a	0.77±0.02 ^c	0.89±0.07 ^b	0.74±0.03 ^c	0.98±0.10 ^b
17:1n-7	0.3±0.01 ^a	0.10±0.00 ^c	0.14±0.02 ^b	0.11±0.02 ^{bc}	0.13±0.00 ^{bc}
18:1n-7	2.4±0.11 ^{ab}	2.2±0.08 ^c	2.5±0.03 ^{ab}	2.3±0.07 ^{bc}	2.5±0.07 ^a
18:1n-9	22.6±1.24	22.9±0.36	23.7±0.06	22.8±0.59	22.3±0.66
20:1n-11	0.8±0.01 ^a	0.63±0.01 ^b	0.87±0.04 ^a	0.68±0.03 ^b	0.89±0.065 ^a
Total MUFA ²	28.0±1.43	26.6 ±0.42	28.2±0.14	26.8±0.73	27.0±0.78
18:2n-6	8.7±0.30 ^c	15.6±0.31 ^b	15.0±0.04 ^b	16.8±0.59 ^a	14.9±0.37 ^b
18:3n-6	0.2±0.01 ^b	0.16±0.01 ^c	0.18±0.00 ^a	0.16±0.01 ^c	0.17±0.01 ^b
20:2n-6	1.78±0.06	2.06±0.02	2.4±0.06	2.2±0.05	1.7±1.4
20:3n-6	0.09±0.01 ^d	0.16±0.01 ^a	0.12±0.01 ^c	0.15±0.01 ^b	0.11±0.0 ^c
20:4n-6	0.08±0.01	0.1±0.09	0.09±0.01	0.07±0.01	0.06±0.01
Total n-6 LC PUFA ³	1.9±0.07	2.3±0.11	2.6±0.11	2.5±0.06	1.9±1.43
Total n-6 PUFA ⁴	10.8±0.37 ^c	18.0±0.42 ^{ab}	17.7±0.07 ^{ab}	19.4±0.64 ^a	17.0±1.20 ^b
18:3n-3	0.16±0.02 ^d	0.80±0.01 ^c	0.96±0.04 ^{ab}	0.98±0.04 ^a	0.89±0.03 ^b
20:3n-3	3.2±0.26 ^c	5.1±0.15 ^a	2.7±0.09 ^c	4.24±0.30 ^b	2.7±0.33 ^c
20:5n-3	18.3±1.07 ^a	15.3±0.24 ^c	16.7±0.16 ^{ab}	14.3±0.06 ^c	15.9±0.83 ^b
22:5n-3	10.5±4.02 ^a	4.6±1.14 ^b	5.0±0.17 ^{ab}	5.3±1.87 ^{ab}	6.2±1.00 ^{ab}
22:6n-3	6.6±0.23 ^b	6.0±0.34 ^b	6.4±0.15 ^b	5.8±0.58 ^b	7.9±0.56 ^a
Total n-3 LC PUFA ⁵	38.6±2.49 ^a	31.1±1.12 ^b	30.8±0.16 ^b	29.7±2.04 ^b	32.8±0.49 ^b
Total n-3 PUFA ⁶	39.1±2.46 ^a	31.8±1.11 ^b	31.8±0.14 ^b	30.6±2.01 ^b	33.7±0.52 ^b
Total PUFA	49.9±2.10	49.8±0.69	49.5±0.20	50.0±1.37	50.7±1.40

[‡]Data are expressed as mean ± SD. Values in the same row with different superscripts are significantly different by the Tukey-Kramer test (P < 0.05).

¹SFA (saturated fatty acid) the sum of all fatty acids without double bounds; includes 12:0, 14:0, 15:0, 16:0, 17:0, 18:0, 20:0, 24:0.

²MUFA (monounsaturated fatty acid) the sum of all fatty acids with a single double bounds; includes 14:1, 15:1, 16:1n-7, 17:1n-7, 18:1n-7, 18:1n-9, 20:1n-11, 22:1n-9 and 24:1n-9

³n-6 LC PUFA; n-6 fatty acids with chain length ≥20 carbon atoms and ≥ 2double bounds

⁴n-6 PUFA; Sum of all n-6 fatty acids

⁵n-3 LC PUFA; n-3 fatty acids with chain length ≥20 carbon atoms and ≥2 double bounds

⁶n-3 PUFA; Sum of all n-3 fatty acids




Figure 1. Lipid, nitrogen and energy retention (%) in crayfish fed with experimental diets.

Discussion

In this trial, redclaw crayfish were fed a range of diets formulated from commercially available local ingredients in Turkey to investigate the potential for increasing the intensive production of crayfish. The crayfish's apparent growth and survival were similar among diets. Levels of moulting were not similar to previous studies on crayfish (4 times in 3 months; Jones, De Silva, & Mitchell, 1996). However, the growth rate of our crayfish in the RAS system was not correspondent to those of other studies based in pond or flow through systems. Noteworthy, our crayfish increased approximately 7 g within 84 days; a value lower than that observed in previous studies. Redclaw crayfish can reach 40-60 g from 12.8 g after six months of cultivation (Masser & Rouse, 1997; Jones & Ruscoe, 2001). García-Guerrero *et al.*, (2013) stocked 10 redclaw crayfish (0.8 g) into 60 L plastic tanks and recorded a final weight of 6.68 g after 90 days of feeding on a 35% CP diet. Crayfish appear to grow faster in open ponds systems implicating a pond-derived factor in their performance. For instance, Naranjo-Páramo *et al.*, (2004) recorded a body weight of 25 g after 80 days feeding trial in natural ponds, with an initial weight of 1.3 g at a stocking density of 11 crayfish m⁻², with once-daily feeding of a 35% CP diet. Crayfish (8 g initial weight) reared in earthen ponds at a very low stocking density of 1.2-2.4 m⁻² achieved a final body weight of 58 g after 70 days of feeding on a commercial marine shrimp diets containing 35% CP (Webster *et al.*, 2004). Garza de Yta *et al.*, (2012) observed that crayfish (0.125 g initial weight) stocked at 12.5 m⁻² and fed for 8 weeks diets with 35% CP containing SBM (~58% of diet) together with 10% of poultry by-product meal, fish meal, ground pea meal or distillers dried grains attained a final body weight of about 4.98 g. We hypothesize that the comparatively low growth rate observed in our study, relative to some of the literature data, is attributable to the higher stocking density that was 40 crayfish m⁻² and the higher initial weight that was 12-13 g. The specific growth rate of our crayfish was similar to that observed in the marron (*Cherax cainii*) fed for two months a 30% CP poultry by-product (~39% of diet) based-diet (Foysal *et al.*, 2019). Although our study demonstrates the potential of poultry by-product meal as a suitable replacer for fish meal in crayfish diets, it is important to note that other factors can affect the crayfish growth, such as the aggressive behaviour, temperature, farming system, the ingredient composition of the diets and photoperiod along with the stocking density. Previous studies report no adverse effects of replacing dietary fish meal with poultry by-product meals on crayfish growth performance (Saoud *et al.*, 2008, 2012; Garza de Yta *et al.*, 2012; Fuertes *et al.*, 2013; Foysal *et al.*, 2019), so much that poultry by-product meal has been routinely added up to 21% (Saoud *et al.*, 2008), 31.3% (Fuertes *et al.*, 2013), 39.0% (Foysal *et al.*, 2019) in crayfish diets. One possible explanation can reside in the

fact that the use of animal protein sources has not been shown to have an advantage over plant protein sources for crayfish's apparent growth and survival. Indeed, redclaw crayfish demonstrated higher digestibility for plant proteins than animal protein sources (Pavasovic *et al.*, 2007). Redclaw crayfish performed comparably with fish meal-free vegetable diets with 28-35% CP or 30% CP diet containing 15% anchovy FM (Thompson *et al.*, 2005; 2006). With a mainly omnivorous feeding habit, redclaw crayfish can grow well on plant proteins-based diets (Lawrence & Jones, 2002; Saoud *et al.*, 2012) most probably as it produces various digestive enzymes including cellulases (Xue *et al.*, 1999; Figueiredo *et al.*, 2001; Saoud *et al.*, 2012).

In the present study, all diets containing plant and animal resources were equally accepted by crayfish with concomitant comparable growth performance, suggesting a general acceptance of both dietary plant and animal proteins by this species. It has been reported that the feeding preferences and growth potential of this species are depended on the activity of digestive enzymes (Lu *et al.*, 2020; Xu *et al.*, 2021). Moreover, oversupply of dietary protein can mask the deficiencies of various protein sources when fed to fish (Glencross *et al.*, 2008). Previous studies propose that crayfish require relatively low levels of dietary protein (22-28%) (Thompson *et al.*, 2005; 2006). It has been reported that 26-33% CP and 8% lipid are optimum for redclaw crayfish growth and survival during juvenile and pre-adult stages (Cortés-Jacinto *et al.*, 2004; Cortés-Jacinto *et al.*, 2005), whilst a dietary protein level of 32% is optimal for reproductive performance (Rodríguez-González *et al.*, 2006; Rodríguez-González *et al.*, 2009). Diets with 30% CP were reported to be ideal for crayfish (Jones *et al.*, 1996; Foysal *et al.*, 2019). The relatively higher CP level in our diets compared with those reported in the literature may not be as valuable for redclaw crayfish in terms of increasing protein use for energy expenditure. With this respect, the highest nitrogen retention was in the FM/Pmix which, in turn, a remarkable decrease in solid and dissolved nitrogenous waste. This could be a result of the better availability of FM/Pmix protein and amino acid profiles, which could be favoured in high stocking density systems ie. RAS. Our findings are in accordance with numerous studies (Steffens, 1999; McGoogan & Gatlin, 2000). Furthermore, Cho & Bureau (2001) noted that diets with unbalanced amino acid profiles in relation to species requirements increase nitrogenous losses and increase protein use for energy production. However, it appears that high protein (as nitrogen) retention, in the present study, can be attributed through improving dietary amino acid balance, specifically when lysine was sufficient in the diet which is also evident in other studies (Fynn-Aikins *et al.*, 1995; Gaylord & Barrows, 2009). Likewise, the FM/Pmix group, and particularly the PoM/Pmix group, preserve nitrogen and energy for maximizing the feed efficiency and growth in the present study. In this context, our nitrogen retention

was high when compare to the normal ranged reported for fish (Peres & Oliva-Teres, 1999; Gaylord & Barrows, 2009; Meiler *et al.*, 2021). This was probably due to different biological response on protein retention efficiency (or the efficiency of retention of synthesised protein) in invertebrates, which has lower protein turnover than in fish and mammals (Mente Coutteau *et al.*, 2002; Carter & Mente, 2014). Further research, thus, is suggested on the nutritional requirement of crayfish reared under various stocking density and feeding practices with the consideration of ammonia production and energy utilization.

There was no consistent trend in the response of the muscle AA profile of crayfish to the experimental diet. Nevertheless, in the present study, the major part of the variation among the EAA is interpreted in concentrations of single amino acid rather than uniformly high or low concentrations in all amino acid pool (Carter *et al.*, 2000). Each diet induced an increase of one or more of EAA content in crayfish except lysine and arginine contents that were consistently higher in crayfish muscle than their respective values supplied in experimental diets. Irrespective of the dietary groups; arginine, leucine and lysine respectively accounted for 8.3, 7.7 and 13.7 of the total EAA even though EAA, as proportion of the total AA, was not significantly different across dietary groups. There is no clear explanation for this response since data on EAA digestibility are missing in our study. Overall, there was no great difference among diets in the EAA profile of crayfish in terms of absolute values. This is consistent with previous studies stating that a mix of plant protein ingredients or animal-plant protein mix could be satisfactory for optimal growth and performance of crayfish (Saoud *et al.*, 2012). Dietary protein quality based on EAAI is considered good when the index is above 0.90, sufficient when it is around 0.8 and inadequate for values below 0.70 (Peñaflorida, 1989). Considering these reference values, our EAAI data was in the range of 1.02 to 1.06 which suggested that overall EAAI was sufficient for the redclaw crayfish. The arginine EAAI data, in the present study, exhibited inadequate value (0.55 to 0.78) across the experimental diets. However, it is important to note that fulfilment of arginine requirement of crayfish calculated as FCR multiplied by arginine content of the experimental diets (as suggested by Lunda *et al.* (2020) showed redclaw crayfish in FM (2.6), PoM (2.9) and Pmix (1.8) group had sufficient arginine in these diets. Moreover, methionine and lysine are usually considered the most critical EAA for marine and crayfish (NRC, 2011; Lunda *et al.*, 2020). In the present study, these two important EAA were at or above standards of NRC (2011), which cover well for most of the EAA requirement in crayfish (Lunda *et al.*, 2020).

There is limited information about the fatty acid requirements of crayfish (Thomson *et al.*, 2010; Saoud *et al.*, 2012). Redclaw crayfish grew equally well when fed diets with various lipid sources including menhaden

oil, linseed oil, canola oil, corn oil and beef tallow, suggesting that crayfish may have an ability to synthesize long-chain fatty acids (20:5n-3, 22:6n-3 and 20:4n-6) from their 18 carbon precursors (18:3n-3 and 18:2n-6) (Thomson *et al.*, 2010), which was verified to a certain degree by Wu *et al.* (2018). In the present study, fish oil and soy lecithin were used as lipid and phospholipid sources in all diets but the incorporation of various protein sources created remarkable changes in dietary fatty acid profiles which resulted in comparable growth performance. However, the fatty acid profile of crayfish on the poultry by-product-based diets showed a lower level of LC-PUFA (particularly EPA) compared to other treatments, which could be a reflection of the high lipid content of poultry by-product meal (19.9%) and its high level of inclusion in the diets at 35 and 52%. In terms of health and growth performance, previous studies reported no adverse effects of replacing fish meal with poultry by-product meals in crayfish diets up to 21% (Saoud *et al.*, 2008), 31.3% (Fuertes *et al.*, 2013), and 39% (Foysal *et al.*, 2019). However, further research is needed regarding the effect of poultry by-product meal/oil on the fatty acid profile of LC-PUFA in crayfish.

In conclusion, our study shows that redclaw crayfish can perform well with low-level and fish meal-free vegetable diets (Pmix) and poultry by-product meal-based diets (PoM and PoM/Pmix). Furthermore, better nitrogen and energy retentions by crayfish fed the FM/Pmix diet (10% FM + 27.5% SBM + 27.5% CGM) compared with the 48% FM diet were observed. Increasing dietary terrestrial plant and animal protein sources while more closely considering diet amino acid profiles may be a strategy for improving the growth performance and nitrogen losses of redclaw crayfish reared under high stocking density. Thus, future studies should consider a more stringent focus on the EAA requirements and development of diets containing different plant, animal by-products and their mixtures in redclaw crayfish reared in RAS.

Ethical Statement

All fish handling procedures complied with Turkish Ethical guidelines for animal care (No. 28141) set by the Ministry of Food, Agriculture and Livestock. Tests were conducted according to the guidance of ethical systems of Cukurova University (CU). The care and use of experimental animal were permitted by the Ethics Committee of CU.

Funding Information

This study was supported by the Scientific Research Project Unit of Cukurova University, Turkey (Grant Number: FBA-2016-6642). The author Dr. Mabrouk Elsabagh received financial grant from the Scientific and Technologic Research Council of Turkey-TÜBİTAK (Grant Number: 1059B211601103).

Author Contribution

O.T.E. and M.E. Conceived and designed the research; H.E., M.K., E.K., E.E., H.A.Y. design the feed formulation and conducted a feeding trial analyses data; B.G. and M.P. interpreted the results, O.T.E. and M.E. contribute to the interpreted and presentation of the experimental full data set. All authors wrote, read and contributed to and approved the manuscript.

Conflict of Interest

No conflict of interest exists in the submission of this manuscript. All authors have seen the manuscript and approve it to submit to your journal. It is not being submitted to any other journal.

References

- AOAC (Association of Official Analytical Chemists). (1995). *Official methods of analysis of AOAC international* (Washington, DC: Association of Official Analytical Chemists).
- Akpınar, Z., Sevgili, H., Demir, A., Özgen, T., Emre, Y., & Tufan Eroldoğan, O. (2012). Effects of dietary lipid levels on growth, nutrient utilization, and nitrogen and carbon balances in shi drum (*Umbrina cirrosa* L.). *Aquaculture International*, 20(1), 131-143.
- Carter, C., & Mente, E. (2014). Protein synthesis in crustaceans: a review focused on feeding and nutrition. *Open Life Sciences*, 9(1), 1-10. <https://doi.org/10.2478/s11535-013-0134-0>
- Carter, C.G., Houlihan, D.F., & He, Z.Y. (2000). Changes in tissue free amino acid concentrations in Atlantic salmon, *Salmo salar* L., after consumption of a low ration. *Fish Physiology and Biochemistry*, 23(4), 295-306.
- Cho, C.Y., & Bureau, D.P. (2001). A review of diet formulation strategies and feeding systems to reduce excretory and feed wastes in aquaculture. *Aquaculture Research*, 32, 349-360. <https://doi.org/10.1046/j.1355-557x.2001.00027.x>
- Cortés-Jacinto, E., Villarreal-Colmenares, H., Civera-Cerecedo, R., & Naranjo-Páramo, J. (2004). Effect of dietary protein level on the growth and survival of pre-adult freshwater crayfish *Cherax quadricarinatus* (von Martens) in monosex culture. *Aquaculture Research*, 35(1), 71-79. <https://doi.org/10.1111/j.1365-2109.2004.00988.x>
- Cortés-Jacinto, E., Villarreal-Colmenares, H., Cruz-Suárez, L.E., Civera-Cerecedo, R., Nolasco-Soria, H., & Hernandez-Llamas, A. (2005). Effect of different dietary protein and lipid levels on growth and survival of juvenile Australian redclaw crayfish, *Cherax quadricarinatus* (von Martens). *Aquaculture Nutrition*, 11(4), 283-291. <https://doi.org/10.1111/j.1365-2095.2005.00353.x>
- FAO, 2020. (2021, November 10). The State of World Fisheries and Aquaculture—Sustainability in Action. Rome. 224 pp. <http://www.fao.org/3/ca9229en/ca9229en.pdf>
- Figueiredo, M.S.R.B., Krickler, J.A., & Anderson, A.J. (2001). Digestive enzyme activities in the alimentary tract of redclaw crayfish, *Cherax quadricarinatus* (Decapoda: Parastacidae). *Journal of Crustacean Biology*, 21(2), 334-344. <https://doi.org/10.1163/20021975-99990133>
- Folch, J., Lees, M., & Stanley, G.S. (1957). A simple method for the isolation and purification of total lipides from animal tissues. *Journal of Biological Chemistry*, 226(1), 497-509.
- Foyсал, M.J., Fotedar, R., Tay, C.Y., & Gupta, S.K. (2019). Dietary supplementation of black soldier fly (*Hermetica illucens*) meal modulates gut microbiota, innate immune response and health status of marron (*Cherax cainii*, Austin 2002) fed poultry-by-product and fishmeal based diets. *PeerJ*, 7, e6891. <https://doi.org/10.7717/peerj.6891>
- Fuertes, J.B., Celada, J.D., Carral, J.M., Sáez-Royuela, M., & González-Rodríguez, Á. (2014). Effects of fishmeal replacement by feather meal in practical diets for juvenile crayfish (*Pacifastacus leniusculus* Dana, Astacidae). *Aquaculture Nutrition*, 20(1), 36-43. <https://doi.org/10.1111/anu.12044>
- Fuertes, J.B., Celada, J.D., Carral, J.M., Saez-Royuela, M., & Gonzalez-Rodriguez, A. (2013). Replacement of fish meal with poultry by-product meal in practical diets for juvenile crayfish (*Pacifastacus leniusculus* Dana, Astacidae) from the onset of exogenous feeding. *Aquaculture*, 404, 22-27. <https://doi.org/10.1016/j.aquaculture.2013.04.019>
- Fuertes, J.B., Celada, J.D., Carral, J.M., Sáez-Royuela, M., & González-Rodríguez, Á. (2012). Effects of dietary protein and different levels of replacement of fish meal by soybean meal in practical diets for juvenile crayfish (*Pacifastacus leniusculus*, Astacidae) from the onset of exogenous feeding. *Aquaculture*, 364, 338-344. <https://doi.org/10.1016/j.aquaculture.2012.08.050>
- Fynn-Aikins, K., Hughes, S.G., & Vandenberg, G.W. (1995). Protein retention and liver aminotransferase activities in Atlantic salmon fed diets containing different energy sources. *Comparative Biochemistry and Physiology Part A: Physiology*, 111(1), 163-170. [https://doi.org/10.1016/0300-9629\(95\)98533-M](https://doi.org/10.1016/0300-9629(95)98533-M)
- Gamboa-Delgado, J. (2014). Nutritional role of natural productivity and formulated feed in semi-intensive shrimp farming as indicated by natural stable isotopes. *Reviews in Aquaculture*, 6(1), 36-47. <https://doi.org/10.1111/raq.12023>
- García-Guerrero, M., Hernández-Sandoval, P., Orduña-Rojas, J., & Cortés-Jacinto, E. (2013). Effect of temperature on weight increase, survival, and thermal preference of juvenile redclaw crayfish *Cherax quadricarinatus*. *Hidrobiológica*, 23(1), 73-81.
- García-Ulloa, G.M., Pérez-Moreno, M.R., Rodríguez-González, D., Gallo-García, M.C., Ponce-Palafox, J.T., Rodríguez-González, H., & Góngora-Gómez, A.M. (2012). Stocking density for nursery production of Redclaw Crayfish, *Cherax quadricarinatus*, in a recirculating system. *Journal of Applied Aquaculture*, 24(1), 8-15. <https://doi.org/10.1080/10454438.2012.650599>
- Garza de Yta, A., Davis, D.A., Rouse, D.B., Ghanawi, J., & Saoud, I.P. (2012). Evaluation of practical diets containing various terrestrial protein sources on survival and growth parameters of redclaw crayfish *Cherax quadricarinatus*. *Aquaculture Research*, 43(1), 84-90. <https://doi.org/10.1111/j.1365-2109.2011.02806.x>
- Gaylord, T.G., & Barrows, F.T. (2009). Multiple amino acid supplementations to reduce dietary protein in plant-based rainbow trout, *Oncorhynchus mykiss*, feeds. *Aquaculture*, 287(1-2), 180-184. <https://doi.org/10.1016/j.aquaculture.2008.10.037>
- Gheshlaghi, R., Schärer, J.M., Moo-Young, M., & Douglas, P.L. (2008). Application of statistical design for the

- optimization of amino acid separation by reverse-phase HPLC. *Analytical Biochemistry*, 383(1), 93-102. <https://doi.org/10.1016/j.ab.2008.07.032>
- Glencross, B., Blyth, D., Wade, N., & Arnold, S. (2018). Critical variability exists in the digestible value of raw materials fed to black tiger shrimp, *Penaeus monodon*: The characterisation and digestibility assessment of a series of research and commercial raw materials. *Aquaculture*, 495, 214-221. <https://doi.org/10.1016/j.aquaculture.2018.05.026>
- Glencross, B., Hawkins, W., Evans, D., Rutherford, N., McCafferty, P., Dods, K., & Sipsas, S. (2008). Assessing the implications of variability in the digestible protein and energy value of lupin kernel meals when fed to rainbow trout, *Oncorhynchus mykiss*. *Aquaculture*, 277(3-4), 251-262. <https://doi.org/10.1016/j.aquaculture.2008.02.012>
- Glencross, B., Irvin, S., Arnold, S., Blyth, D., Bourne, N., & Preston, N. (2014). Effective use of microbial biomass products to facilitate the complete replacement of fishery resources in diets for the black tiger shrimp, *Penaeus monodon*. *Aquaculture*, 431, 12-19. <https://doi.org/10.1016/j.aquaculture.2014.02.033>
- Gunasekera, R.M., Turoczy, N.J., De Silva, S.S., Gavine, F., & Gooley, G.J. (2002). An evaluation of the suitability of selected waste products in feeds for three fish species. *Journal of Aquatic Food Product Technology*, 11(1), 57-78. https://doi.org/10.1300/J030v11n01_06
- Jones, C.M. (1995). Production of juvenile redclaw crayfish, *Cherax quadricarinatus* (von Martens) (Decapoda, Parastacidae) II. Juvenile nutrition and habitat. *Aquaculture*, 138(1-4), 239-245. [https://doi.org/10.1016/0044-8486\(95\)00069-0](https://doi.org/10.1016/0044-8486(95)00069-0)
- Jones, C.M., & Ruscoe, I.M. (2001). Assessment of five shelter types in the production of redclaw crayfish *Cherax quadricarinatus* (Decapoda: Parastacidae) under earthen pond conditions. *Journal of the World Aquaculture Society*, 32(1), 41-52. <https://doi.org/10.1111/j.1749-7345.2001.tb00920.x>
- Jones, P.L., De Silva, S.S., & Mitchell, B.D. (1996). The effect of dietary protein source on growth and carcass composition in juvenile Australian freshwater crayfish. *Aquaculture International*, 4(4), 361-376.
- Kumlu, M. (2018). Developing an economic and sustainable shrimp production model for Turkey. Project Report (TÜBİTAK) 223 p.
- Lawrence, C., & Jones, C. (2002). *Cherax*. In: Holdich DM. (Ed), *Biology of freshwater crayfish*, (pp. 635-669).
- Lu, X., Peng, D., Chen, X., Wu, F., Jiang, M., Tian, J., Liu, W., Wen, H., & Wei, K. (2020). Effects of dietary protein levels on growth, muscle composition, digestive enzymes activities, hemolymph biochemical indices and ovary development of pre-adult red swamp crayfish (*Procambarus clarkii*). *Aquaculture Reports*, 18, 100542. <https://doi.org/10.1016/j.aqrep.2020.100542>
- Lunda, R., Roy, K., Dvorak, P., Kouba, A., & Mraz, J. (2020). Recycling biofloc waste as novel protein source for crayfish with special reference to crayfish nutritional standards and growth trajectory. *Scientific Reports*, 10(1), 1-10. <https://doi.org/10.1038/s41598-020-76692-0>
- Masser, M.P., & Rouse, D.B. (1997). Australian red claw crayfish (Report No. 244). Stoneville: Southern Regional Aquaculture Center.
- McGoogan, B.B., & Gatlin III, D.M. (2000). Dietary manipulations affecting growth and nitrogenous waste production of red drum, *Sciaenops ocellatus*: II. Effects of energy level and nutrient density at various feeding rates. *Aquaculture*, 182(3-4), 271-285. [https://doi.org/10.1016/S0044-8486\(99\)00260-4](https://doi.org/10.1016/S0044-8486(99)00260-4)
- Meiler, K.A., & Kumar, V. (2021). Organic and inorganic zinc in the diet of a commercial strain of diploid and triploid rainbow trout (*Oncorhynchus mykiss*): Effects on performance and mineral retention. *Aquaculture*, 545, 737126. <https://doi.org/10.1016/j.aquaculture.2021.737126>
- Mente, E., Coutteau, P., Houlihan, D., Davidson, I., & Sorgeloos, P. (2002). Protein turnover, amino acid profile and amino acid flux in juvenile shrimp *Litopenaeus vannamei*: effects of dietary protein source. *Journal of Experimental Biology*, 205(20), 3107-3122. <https://doi.org/10.1242/jeb.205.20.3107>
- Muzinic, L.A., Thompson, K.R., Morris, A., Webster, C.D., Rouse, D.B., & Manomaitis, L. (2004). Partial and total replacement of fish meal with soybean meal and brewer's grains with yeast in practical diets for Australian red claw crayfish *Cherax quadricarinatus*. *Aquaculture*, 230(1-4), 359-376. [https://doi.org/10.1016/S0044-8486\(03\)00420-4](https://doi.org/10.1016/S0044-8486(03)00420-4)
- Naranjo-Páramo, J., Hernandez-Llamas, A., & Villarreal, H. (2004). Effect of stocking density on growth, survival and yield of juvenile redclaw crayfish *Cherax quadricarinatus* (Decapoda: Parastacidae) in gravel-lined commercial nursery ponds. *Aquaculture*, 242(1-4), 197-206. <https://doi.org/10.1016/j.aquaculture.2004.05.017>
- NRC. (2011). Nutrient Requirements of fish and shrimp. National Research Council Washington, DC.
- Oser, B.L. (1959). An integrated essential amino acid index for predicting the biological value of proteins. In A.A. Albanese (Ed), *Protein and amino acid nutrition* (pp. 281). New York: Academic Press.
- Pavasovic, A., 2008. Evaluation of the nutritional requirements of redclaw crayfish. *Cherax quadricarinatus* (Doctoral dissertation. Queensland University of Technology). 165 p.
- Pavasovic, A., Anderson, A.J., Mather, P.B., & Richardson, N.A. (2007). Effect of a variety of animal, plant and single cell-based feed ingredients on diet digestibility and digestive enzyme activity in redclaw crayfish, *Cherax quadricarinatus* (Von Martens 1868). *Aquaculture*, 272(1-4), 564-572. <https://doi.org/10.1016/j.aquaculture.2007.08.027>
- Peñaflorida, V.D. (1989). An evaluation of indigenous protein sources as potential component in the diet formulation for tiger prawn, *Penaeus monodon*, using essential amino acid index (EAAI). *Aquaculture*, 83(3-4), 319-330. [https://doi.org/10.1016/0044-8486\(89\)90043-4](https://doi.org/10.1016/0044-8486(89)90043-4)
- Peres, H., & Oliva-Teles, A. (1999). Influence of temperature on protein utilization in juvenile European seabass (*Dicentrarchus labrax*). *Aquaculture*, 170(3-4), 337-348. [https://doi.org/10.1016/S0044-8486\(98\)00422-0](https://doi.org/10.1016/S0044-8486(98)00422-0)
- Qian, D., Yang, X., Xu, C., Chen, C., Jia, Y., Gu, Z., & Li, E. (2021). Growth and health status of the red claw crayfish, *Cherax quadricarinatus*, fed diets with four typical plant protein sources as a replacement for fish meal. *Aquaculture Nutrition*, 27(3), 795-806. <https://doi.org/10.1111/anu.13224>
- Rodríguez-Canto, A., Arredondo-Figueroa, J.L., Ponce-Palafox, J.T., & Rouse, D.B. (2002). Growth characteristics of the Australian redclaw crayfish, *Cherax quadricarinatus*,

- cultured in an indoor recirculating system. *Journal of Applied Aquaculture*, 12(3), 59-64.
https://doi.org/10.1300/J028v12n03_06
- Rodríguez-González, H., García-Ulloa, M., Hernández-Llamas, A., & Villarreal, H. (2006). Effect of dietary protein level on spawning and egg quality of redclaw crayfish *Cherax quadricarinatus*. *Aquaculture*, 257(1-4), 412-419.
<https://doi.org/10.1016/j.aquaculture.2006.01.020>
- Rodríguez-González, H., Villarreal, H., García-Ulloa, M., & Hernández-Llamas, A. (2009). Evaluation of practical diets containing different protein levels on gonad development of female redclaw crayfish *Cherax quadricarinatus*. *Aquaculture Nutrition*, 15(4), 347-355.
<https://doi.org/10.1111/j.1365-2095.2008.00599.x>
- Saoud, I.P., Ghanawi, J., Thompson, K.R., & Webster, C.D. (2013). A review of the culture and diseases of redclaw crayfish *Cherax quadricarinatus* (von Martens 1868). *Journal of the World Aquaculture Society*, 44(1), 1-29.
<https://doi.org/10.1111/jwas.12011>
- Saoud, I.P., Rodgers, L.J., Davis, D.A., & Rouse, D.B. (2008). Replacement of fish meal with poultry by-product meal in practical diets for redclaw crayfish (*Cherax quadricarinatus*). *Aquaculture Nutrition*, 14(2), 139-142.
<https://doi.org/10.1111/j.1365-2095.2007.00513.x>
- Saoud, I.P., Garza De Yta, A., & Ghanawi, J. (2012). A review of nutritional biology and dietary requirements of redclaw crayfish *Cherax quadricarinatus* (von Martens 1868). *Aquaculture Nutrition*, 18(4), 349-368.
<https://doi.org/10.1111/j.1365-2095.2011.00925.x>
- Schneider, O., Amirkolaie, A.K., Vera-Cartas, J., Eding, E.H., Schrama, J.W., & Verreth, J.A. (2004). Digestibility, faeces recovery, and related carbon, nitrogen and phosphorus balances of five feed ingredients evaluated as fishmeal alternatives in Nile tilapia, *Oreochromis niloticus* L. *Aquaculture Research*, 35(14), 1370-1379.
<https://doi.org/10.1111/j.1365-2109.2004.01179.x>
- Steffens, W. (1999). Effect of two lipid levels on growth, feed utilization, body composition and biochemical parameters of rainbow trout, *Oncorhynchus mykiss* (Walbaum 1792). *Journal of Applied Ichthyology*, 15, 159-164.
- Tan, Q., Song, D., Chen, X., Xie, S., & Shu, X. (2018). Replacing fish meal with vegetable protein sources in feed for juvenile red swamp crayfish, *Procambarus clarkii*: Effects of amino acids supplementation on growth and feed utilization. *Aquaculture Nutrition*, 24(2), 858-864.
<https://doi.org/10.1111/anu.12621>
- Thompson, K.R., Bailey, T.J., Metts, L.S., Brady, Y.J., & Webster, C.D. (2010). Growth response and fatty acid composition of juvenile red claw crayfish (*Cherax quadricarinatus*) fed different sources of dietary lipid. *Aquaculture Nutrition*, 16(6), 604-615.
<https://doi.org/10.1111/j.1365-2095.2009.00697.x>
- Thompson, K.R., Metts, L.S., Muzinic, L.A., Dasgupta, S., & Webster, C.D. (2006). Effects of feeding practical diets containing different protein levels, with or without fish meal, on growth, survival, body composition and processing traits of male and female Australian red claw crayfish (*Cherax quadricarinatus*) grown in ponds. *Aquaculture Nutrition*, 12(3), 227-238.
<https://doi.org/10.1111/j.1365-2095.2006.00407.x>
- Thompson, K.R., Muzinic, L.A., Engler, L.S., & Webster, C.D. (2005). Evaluation of practical diets containing different protein levels, with or without fish meal, for juvenile Australian redclaw crayfish (*Cherax quadricarinatus*). *Aquaculture*, 244(1-4), 241-249.
<https://doi.org/10.1016/j.aquaculture.2004.11.018>
- Turchini, G.M., Trushenski, J.T., & Glencross, B.D. (2019). Thoughts for the future of aquaculture nutrition: realigning perspectives to reflect contemporary issues related to judicious use of marine resources in aquafeeds. *North American Journal of Aquaculture*, 81(1), 13-39. <https://doi.org/10.1002/naaq.10067>
- Van Rijn, J. (2013). Waste treatment in recirculating aquaculture systems. *Aquacultural Engineering*, 53, 49-56. <https://doi.org/10.1016/j.aquaeng.2012.11.010>
- Venero, J.A., Davis, D.A., & Lim, C.E. (2008). Use of plant protein sources in crustacean diets. *Alternative protein sources in aquaculture diets* (pp 163-203). New York, USA, The Haworth press.
- Webster, C.D., Thompson, K.R., Muzinic, L.A., Yancey, D.H., Dasgupta, S., Xiong, Y.L., & Manomaitis, L. (2004). A preliminary assessment of growth, survival, yield, and economic return of Australian redclaw crayfish, *Cherax quadricarinatus*, stocked at three densities in earthen ponds in a cool, temperate climate. *Journal of Applied Aquaculture*, 15(3-4), 37-50.
https://doi.org/10.1300/J028v15n03_03
- Wu, D.L., Huang, Y.H., Liu, Z.Q., Yu, P., Gu, P.H., Fan, B., & Zhao, Y.L. (2018). Molecular cloning, tissue expression and regulation of nutrition and temperature on $\Delta 6$ fatty acyl desaturase-like gene in the redclaw crayfish (*Cherax quadricarinatus*). *Comparative Biochemistry and Physiology Part B: Biochemistry and Molecular Biology*, 225, 58-66. <https://doi.org/10.1016/j.cbpb.2018.07.003>
- Xu, L., Yuan, J., Chen, X., Zhang, S., Xie, M., Chen, C., & Wu, Z. (2021). Screening of intestinal probiotics and the effects of feeding probiotics on the digestive enzyme activity, immune, intestinal flora and WSSV resistance of *Procambarus clarkii*. *Aquaculture*, 540, 736748. <https://doi.org/10.1016/j.aquaculture.2021.736748>
- Xue, X.M., Anderson, A.J., Richardson, N.A., Anderson, A.J., Xue, G.P., & Mather, P.B. (1999). Characterisation of cellulase activity in the digestive system of the redclaw crayfish (*Cherax quadricarinatus*). *Aquaculture*, 180(3-4), 373-386.
[https://doi.org/10.1016/S0044-8486\(99\)00213-6](https://doi.org/10.1016/S0044-8486(99)00213-6)
- Zeng, Y., Shakir, K.K., & Yeo, D.C. (2019). Competition between a native freshwater crab and an invasive crayfish in tropical Southeast Asia. *Biological Invasions*, 21(8), 2653-2663. <https://doi.org/10.1007/s10530-019-02009-6>