

Sección Agrícola / Agriculture

Artículos de revisión / Revisional paper

Management of *Aceria litchii* (Acari: Eriophyidae) on *Litchi chinensis*Manejo del ácaro *Aceria litchii* (Acari: Eriophyidae) en *Litchi chinensis*BÁRBARA MONTEIRO DE CASTRO E CASTRO¹, ANGELICA PLATA-RUEDA², WIANE MELONI SILVA³,
CLAUBERT WAGNER GUIMARÃES DE MENEZES⁴, CARLOS FREDERICO WILCKEN⁵
and JOSÉ COLA ZANUNCIO⁶

Abstract: *Litchi chinensis* fruits are very popular in the international market. The erinose mite, *Aceria litchii* (Eriophyidae), a major pest of this crop, feeds on the leaves and on all parts of the plants forming erinea. The purpose of this review is to present the most common control methods and to propose integrated management techniques for *L. chinensis* production systems. *A. litchii* infestations were detected in Brazilian orchards in 2008. Chemical pesticides are currently the main control measure for this pest; however, the combination of biological, chemical and cultural methods are more promising in the long term. Controlling and preventing the erinose mite are key factors for reducing its spread and establishment.

Key words: Fruit, pest, pesticides, Sapindaceae.

Resumen: El cultivo del litchi, *Litchi chinensis*, tiene gran popularidad en el mercado internacional. El ácaro del litchi, *Aceria litchii* (Eriophyidae), es una de las principales plagas de este cultivo, se alimenta de hojas, sobre todo en la parte inferior de las plantas que forman erineos. El objetivo de esta revisión es presentar los principales métodos de control y proponer técnicas de manejo en el sistema de producción integrado del cultivo de *L. chinensis*. Se detectaron infestaciones de *A. litchii* en los huertos de Brasil en 2008 y el uso de insecticidas químicos es el principal método de control de esta plaga. La combinación de métodos de cultivo biológico y químico hace que el sistema sea más satisfactorio en el largo plazo. El control y la prevención del ácaro del litchi son clave para reducir su propagación y establecimiento.

Palabras clave: Fruta, plaga, pesticidas, Sapindaceae.

Introduction

The litchi, *Litchi chinensis* Sonn. (Sapindaceae) is a tropical plant native to Southeast Asia that produces climacteric fruits (Cabral *et al.* 2014) and is cultivated in several countries (Gontier *et al.* 2008; Jiang *et al.* 2013). It is widely sought after on the international market (Yang *et al.* 2016). China and India are the largest producers (Ranjan and Ran 2015). In Brazil, the litchi was introduced in 1810 and by 1970 commercial production began in São Paulo (Yamanishi *et al.* 2010). Currently, it is one of the most popular exotic fruits in the country (Alves *et al.* 2016), with the “Bengal” variety being the most consumed (Suguino 2006). Harvesting is carried out between November and January (Martins *et al.* 2001; Yamanishi *et al.* 2001), when the demand for fruit is high and there is no competition from other countries (Lins *et al.* 2015). The fruit has a red epicarp surrounding the mesocarp consisting of a white layer, with tasty pulp and high nutritional value (Bhoopat *et*

al. 2011). The fruits of this plant can be eaten fresh and used to manufacture juices, vinegar, marmalades and fermented alcoholic beverages (Alves *et al.* 2011; Saxena *et al.* 2011). Bioactive compounds such as lignans and flavonoids are present in large quantities in *L. chinensis* leaves and epicarp (Wen *et al.* 2014). In addition, consumption of lychee fruits and seeds can prevent cancer cell growth (Bhat and Al-Daihan 2014; Wen *et al.* 2015).

Among the main *L. chinensis* pests, the mite *Aceria litchii* (Keifer) (Acari: Eriophyidae) stands out (Lall and Rahman 1975; Hameed *et al.* 1992; Huang 2008), having been reported in Australia, Brazil, China, Hawaii, India and Pakistan (Jeppson *et al.* 1975; Sabelis and Bruin 1996; Hong *et al.* 2006, Huang 2008). The lesions caused by *A. litchii* were detected on litchi plants in São Paulo state, Brazil in January 2008, being the first report of this pest in South America (Raga *et al.* 2010). Espírito Santo was the second Brazilian state where infestation was confirmed (Fornazier *et al.* 2014). *Aceria litchii* is located and feeds

¹ M. Sc. Estudante de doctorado, Departamento de Fitotecnia, Universidade Federal de Viçosa, 36571-900, Viçosa, Minas Gerais, Brazil, *barbaramcastro@hotmail.com*. ² Pós-Doc. Pós-Doctor, Departamento de Agronomia, Campus de Rio Paranaíba, Universidade Federal de Viçosa, 38810-000, Rio Paranaíba, Minas Gerais, Brazil, *angelicaplata@yahoo.com.mx*. ³ M. Sc. Estudante de doctorado, Departamento de Ciência Florestal, Universidade Federal de Viçosa, 36570-900, Viçosa, Minas Gerais, Brazil, *wianems@yahoo.com.br*. ⁴ D. Sc. Professor del Departamento de Ciências Agrárias, Instituto Federal do Norte de Minas Gerais, 39480-000, Januária, Minas Gerais, Brazil, *claubertmenezes@yahoo.com.br*. ⁵ D. Sc. Professor del Departamento de Proteção Vegetal, Universidade Estadual Paulista Júlio de Mesquita Filho (UNESP), 18610-307, Botucatu, São Paulo, Brazil, *cwilcken@fca.unesp.br*. ⁶ Ph. D. Professor del Departamento de Entomologia/BIOAGRO, Universidade Federal de Viçosa, 36570-900, Viçosa, Minas Gerais, Brazil, *zanuncio@ufv.br*. Corresponding author: Bárbara Monteiro de Castro e Castro. M. Sc. Estudante de doctorado, Departamento de Fitotecnia, Universidade Federal de Viçosa, 36571-900, Viçosa, Minas Gerais, Brazil, *barbaramcastro@hotmail.com*.

on the leaves, flowers and underside of young *L. chinensis* fruits (Nishida and Holdaway 1955; Butani 1977; Sharma *et al.* 1986). Abiotic factors such as temperature, relative humidity, wind speed and precipitation affect the population growth of this mite (Singh *et al.* 1987). The symbiotic relationship between *A. litchii* and the algae *Cephaleuros virescens* Kunze (Tentropoliaceae) favors the formation of erineos (trichomes developed abnormally) (Saha *et al.* 1996). The erineo generates an adequate environment for mites, protecting them from precipitation, temperature, wind (Sharma 1984; Sharma *et al.* 1986; Thakur and Sharma 1990) and the action of chemical products (Jeppson *et al.* 1975; Westphal and Manson 1996). Colonization by these algae causes detrimental effects to the plant through stomatal blockage and hinders photosynthesis (Alam and Wadud 1963). Initially, the erineo is silvery white, changing to light brown, dark reddish brown and black as the infestation advances (Waite 2005). The leaves affected turn thick and wavy, wither and finally fall. Damaged shoots do not produce flowers or fruits (Lall and Rahman 1975).

Practices for litchi mite control should be adopted throughout most of the year (Azevedo *et al.* 2014) mainly in the development of the inflorescence and the expansion of new leaves when the mites migrate to young organs to take refuge, feed themselves and multiply by establishing new erineos (Arthur and Machi 2016). Several chemical products are widely used to control the mite (Azevedo *et al.* 2013), but the adoption of other control methods is desirable.

This review aims to present the main control methods, their advances and propose management techniques for the litchi mite, *A. litchii*.

Sampling

The behavior and small size of *A. litchii* hampers the verification of population density. This mite lives in erineos, which impedes direct observation (Azevedo *et al.* 2014). Methods to estimate lychee mite populations and other insects that form erineos have been adapted from one used to evaluate nematode densities in plant roots (Coolen and D'Herde 1972) consisting of blending litchi infested leaves in a blender for 1 min in 250 ml distilled water and screening the material with a set of sieves with openings of 2, 0.2 and 0.037 mm, respectively, from top to bottom. Larger particles of leaves are retained in the two larger aperture screens, while smaller ones and mites are retained in the screen with a smaller aperture. The mite passage through the upper screens is driven by a jet of tap water for a few seconds. The material retained in the lowest sieve is transferred to a 25 ml cylinder, using sucrose solution (density of 1.15 g/ml, corresponding to 400 g of commercial sucrose dissolved in 750 ml of water). The material should be shaken several times and left to rest for 15 hours allowing the mites to concentrate near the solution surface. The supernatant is poured through a 0.037 mm opening screen and the material retained (mainly mites) is washed with distilled water to remove excess sucrose. The mites should be transferred to a bottle with 15 ml of 70 % ethanol. Mites are counted in Peters sheets, commonly used in nematode counts. The estimate is made by extrapolating the count of three aliquots of 1 ml. This procedure is progressive with higher precision as the *A. litchii* infestation level increases (Azevedo *et al.* 2014). The highest *A. litchii* density on plants in Brazil is recorded at the beginning of the

rainy season (October) and in the middle of the dry season (May-June) (Azevedo *et al.* 2014). *Aceria litchii* density can also be determined by stirring 2 cm² leaf pieces with erineos in a 0.5 % neutral detergent solution for five seconds to dislodge this mite from the erineos. The counting of mites present in the solution is done with a stereoscopic microscope (Picoli 2010).

Chemical control

Pesticides must be applied before and/or during the emergence of the inflorescence and leaf expansion of *L. chinensis* (Picoli 2010). Spraying outside this range has unsatisfactory results (Waite 2005). The main problem with Eriophyidae mite control is its hidden lifecycle. Pesticide application should occur during the stages when the mites temporarily leave the erineos. Control is better provided with acaricides with long residual effect, since the transfer (erine exit) of mites to new leaves extends over several days or weeks (Van Leeuwen *et al.* 2010).

Three preventative dimethoate or sulfur applications in soluble powder at two to three weeks intervals are recommended in Thailand (Waite and Hwang 2002). Dichlorvos, chlorpyrifos, dimethoate and isocarbophos are effective and used in China (Waite 2005). The fenpyroxime, sulfur, abamectin and hexythiazox application cause high *A. litchii* mortality. However, these insecticides were considered harmful to the predator *Phytoseius intermedius* Evans & MacFarlane (Acari: Phytoseiidae) (Azevedo *et al.* 2013). The complete elimination of *A. litchii* was achieved with a double application of espiromesifeno at a concentration of 0.144 g a.i. L⁻¹ in Thailand (Schulte *et al.* 2007).

When effectively used, sulfur presents low toxicity to mammals, but has an impact on natural enemies (Prischmann *et al.* 2005) such as *P. intermedius* (Azevedo *et al.* 2013); *Galendromus occidentalis* (Nesbitt, 1951) (Beers *et al.* 2009); *Euseius victoriensis* (Womersley, 1954) (Bernard *et al.* 2010) and *Typhlodromus pyri* (Scheuten, 1857) (Acari: Phytoseiidae) (Gadino *et al.* 2011). Sulfur is inorganic and ineffective against *Tetranychus urticae* (Koch, 1836) (Acari: Tetranychidae) (Auger *et al.* 2003).

The development of resistance by Eriophyidae mites to pesticides has been reviewed (Messing and Croft 1996). Resistance to organophosphate by *Aculus cornutus* and *A. lycopersici* (Acari: Eriophyidae) was confirmed in the laboratory (Abou-Awad and El-Banhawy 1995), but products with different action mechanisms can reduce the development of resistance in mite pests (Azevedo *et al.* 2013).

Azadirachtin causes low *A. litchii* mortality but has low effect on the predator *P. intermedius* (Azevedo *et al.* 2013). This insecticide is toxic to phytophagous and predatory mites (Castagnoli *et al.* 2000; Brito *et al.* 2006; Duso *et al.* 2008).

In Australia, the management of *A. litchii* consists of three acaricide applications during shoot formation at intervals of 10 to 14 days if erineos on most plants. Satisfactory control has been achieved with successive sprays alternating sulfur and dimethoate, the first performed at the beginning of shoot formation, if the mite is present during *L. chinensis* flowering (Waite 2011). However, there is no pesticide registered for litchi cultivation in Brazil (Brasil 2016).

The chemical control of Eriophyidae mites was reviewed (Childers *et al.* 1996), but the efficiency of other chemical

compounds to control these mites requires further study. The main reason for lacking information on the toxicity of new compounds is the lower economic importance of these mites compared to others such as Tetranychidae mites (Van Leeuwen *et al.* 2010).

Biological control

Predatory mites. *Aceria litchii* can be transported by bees from the flowers of infested plants (Waite and Mcalpine 1992; Waite 1999) and can be preyed upon when migrating to new leaves, before forming the erineo (Azevedo *et al.* 2013). Predators attack *A. litchii* when the mite leaves the erineo (Azevedo *et al.* 2014). The main predators of phytophagous mites are Phytoseiidae species (Gerson *et al.* 2003; McMurtry *et al.* 2015). The natural presence and periodic releases of predatory mites has been used to control pests (Moraes and Lima 1983; Momen and Hussein 1999). Phytoseiids are associated with *A. litchii* in India (Thakur and Sharma 1989), Australia, China (Waite and Gerson 1994) and Brazil (Picoli 2010; Azevedo *et al.* 2013). The predator *Amblyseius compositus* (Denmark and Muma, 1973) (Acari: Phytoseiidae) (42.6 %) predominated in Casa Blanca, São Paulo state, Brazilian plantations followed by the other predatory mite *P. intermedius* (31.2 %) (Picoli 2010), but in Limeira, São Paulo, *A. compositus* corresponded to 10 % of phytoseiids observed (Azevedo *et al.* 2014). The highest predator population densities generally coincide with the presence of *A. litchii* between October and December (Picoli 2010; Azevedo *et al.* 2014).

Aceria litchii is suitable prey for *P. intermedius* (Azevedo *et al.* 2016). This predator was found on plants with leaves covered with trichomes, such as *Helicteres brevispira* Saint-Hilaire and *Helicteres lhotzkyana* (Schott & Endlincher) (Malvaceae), *Guazuma ulmifolia* Lamarck (Sterculiaceae), *Miconia* sp. (Melastomataceae) and *Cordia sellowiana* Chamisso (Boraginaceae) in São Paulo state, Brazil (Demite *et al.* 2008). However, the erineos on *A. litchii* on litchi leaves reduces the action of the predatory mites. Eight phytoseiid species are associated with *A. litchii* in India (Thakur and Sharma 1990) and *A. compositus*, *Euseius concordis* (Chant, 1959) and *Iphiseiodes zuluagai* (Denmark & Muma, 1972) (Acari: Phytoseiidae) are positively correlated and promising for the biological control of *A. litchii* (Picoli 2010).

Entomopathogenic fungi. Entomopathogenic fungi can regulate arthropod populations by penetrating their cuticle and destroying internal tissues (Kurtti and Keyhani 2008; Rossoni *et al.* 2014; Costa *et al.* 2015). The ease of dispersion of these microorganisms in the field justifies research to investigate their potential for biological control (Meyling *et al.* 2009; Costa *et al.* 2015).

Entomopathogenic fungi (Acari: Eriophyidae) have been reported on mites (Tanzini *et al.* 2000; Demite and Feres 2008); and the erineos conserving the humidity inside the gills providing a favorable microclimate for growth of these microorganisms (Picoli and Vieira 2013). *Hirsutella thompsonii* (Fischer) (Ascomycota: Ophiocordycipitaceae) is the most commonly found fungus on Eriophyidae (McCoy 1996). This fungus is an alternative to control *A. litchii* (Picoli and Vieira 2013). The *H. thompsonii* infestation was higher during rainy periods with a positive correlation with rainfall and relative humidity (Demite and Feres 2008)

simultaneously with the population density growth of *A. litchii* on litchi plants in Brazil (Azevedo *et al.* 2014).

Entomopathogenic fungi are recommended for the biological control of other mites (Sreerema Kumar and Singh 2002, 2008; Alves *et al.* 2005; Fernando *et al.* 2007; Paz *et al.* 2007; Edgington *et al.* 2008; Gerson *et al.* 2008). Entomopathogenic fungi presence and impacts on Eriophyidae mites were reviewed (McCoy 1996; Van der Geest *et al.* 2000; Balazy *et al.* 2008) and they can contribute to *A. litchii* management programs.

Cultural control

Branch and leaf pruning of erineos and resultant damage is an effective measure to manage mites in litchi plantations. *A. litchii* populations are higher in the lower third of the plants and pruning of symptomatic branches in this region should be complete (Raga *et al.* 2011). However, *A. litchii* management with pruning and acaricide spraying increases production costs for this crop, therefore, litchi plants should be inspected in nurseries to prevent or reduce the spread of this mite (Raga *et al.* 2010).

Final considerations

Aceria litchii is the most important litchi pest in the world. Chemicals are the main control measure for this mite. However, adverse effects of pesticides justify the search for ecologically sustainable pest control strategies (Khederi *et al.* 2014). The combination of biological, cultural and chemical methods (Timprasert *et al.* 2014) make the production system more promising over the long term and can reduce pesticide dependence (Hashemi *et al.* 2008). The natural biological control of Eriophyidae mites is based mainly on predators and, to a lesser degree, on pathogens. However, the implementation of this control method presents difficulties. Predatory thrips are commercialized to manage Tetranychidae mites (Gerson *et al.* 2003; de Faria and Wraight 2007). They are also natural enemies of Eriophyidae mites but none of them is sold specifically to control these mites (Van Leeuwen *et al.* 2010). Few studies have investigated the efficiency of new chemical compounds to control *A. litchii*. Therefore, research into control methods, with special attention on the tritrophic complex (cultural, natural enemies and pests) is fundamental to achieve integrated control and reduce the possibilities of *A. litchii* propagation and establishment.

Acknowledgements

To “Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES), Fundação de Amparo à Pesquisa do Estado de Minas Gerais (FAPEMIG)” and “Programa Cooperativo sobre Proteção Florestal/PROTEF of the Instituto de Pesquisas e Estudos Florestais/IPEF”. Dr. Phillip John Villani (University of Melbourne, Australia) revised and corrected the English language used in this manuscript.

Literature cited

ABOU-AWAD, B. A.; EL-BANHAWY, E. M. 1995. Susceptibility of the tomato russet mite, *Aculops lycopersici* in Egypt to

- methamidiphos, pyridaphenthion, cypermethrin, dicofol and fenarimol. *Experimental and Applied Acarology* 1 (1): 11-15.
- ALAM, M. Z.; WADUD, M. A. 1963. On the biology of litchi mite, *Aceria litchi* Keifer (Eriophyidae, Acarina) in east Pakistan. *Pakistan Journal of Science* 15 (5): 232-240.
- ALVES, S. B.; TAMAI, M.S.; ROSSI, L. S.; CASTIGLIONE, E. 2005. *Beauveria bassiana* pathogenicity to citrus rust mite *Phyllocoptruta oleivora*. *Experimental and Applied Acarology* 37 (1-2): 117-122.
- ALVES, J. A.; DE OLIVEIRA, L. L. C.; NUNES, C. A.; DIAS, D. R.; SCHWAN, R. F. 2011. Chemical, physical-chemical, and sensory characteristics of lychee (*Litchi chinensis* Sonn.) wines. *Journal of Food Science* 76 (5): 330-336.
- ALVES, E. C.; GUIMARÃES, J. E. R.; FRANCO, C. K. B.; MARTINS, A. B. G. 2016. Number of leaflets on rooting of lychee herbaceous cuttings. *Ciência Rural* 46 (6): 1003-1006.
- ARTHUR, V.; MACHI, A. R. 2016. Development of phytosanitary irradiation against *Aceria litchii* (Trombidiformes: Eriophyidae) on lychee. *Florida Entomologist* 99 (2): 143-149.
- AUGER, P.; GUICHOU, S.; KREITER, S. 2003. Variations in acaricidal effect of wettable sulfur on *Tetranychus urticae* (Acari: Tetranychidae): effect of temperature, humidity and life stage. *Pest Management Science* 59 (5): 559-565.
- AZEVEDO, L. H.; DE MORAES, G. J.; YAMAMOTO, P. T.; ZANARDI, O. Z. 2013. Development of a methodology and evaluation of pesticides against *Aceria litchii* and its predator *Phytoseius intermedius* (Acari: Eriophyidae, Phytoseiidae). *Journal of Economic Entomology* 106 (5): 2183-2189.
- AZEVEDO, L. H.; MAEDA, E. Y.; INOMOTO, M. M.; MORAES, G. J. 2014. A method to estimate the population level of *Aceria litchi* (Prostigmata: Eriophyidae) and a study of the population dynamics of this species and its predators on litchi trees in southern Brazil. *Journal of Economic Entomology* 107 (1): 361-367.
- AZEVEDO, L. H.; CASTILHO, R. D.; DE MORAES, G. J. 2016. Suitability of the litchi erineum mite, *Aceria litchii* (Keifer), as prey for the mite *Phytoseius intermedius* Evans & MacFarlane (Acari: Eriophyidae, Phytoseiidae). *Systematic and Applied Acarology* 21 (3): 270-278.
- BALAZY, S.; MIETKIEWSKI, R.; THACZUK, C.; WEGENSTEINER, R.; WRZOSEK, M. 2008. Diversity of acaropathogenic fungi in Poland and other European countries. *Experimental and Applied Acarology* 46 (1-4): 53-70.
- BEERS, E. H.; MARTINEZ-ROCHA, L.; TALLEY, R. R.; DUNLEY, J. E. 2009. Lethal, sublethal and behavioral effects of sulfur-containing products in bioassays of three species of orchard mites. *Journal of Economic Entomology* 102 (1): 324-335.
- BERNARD, M. B.; COLE, P.; KOBELT, A.; HORNE, P. A.; ALTMANN, J.; WRATTEN, S. D.; YEN, A. L. 2010. Reducing the impact of pesticides on biological control in Australian vineyards: pesticide mortality and fecundity effects on an indicator species, the predatory mite *Euseius victoriensis* (Acari: Phytoseiidae). *Journal of Economic Entomology* 103 (6): 2061-2071.
- BHAT, R. S.; AL-DAIHAN, S. 2014. Antimicrobial activity of *Litchi chinensis* and *Nephelium lappaceum* aqueous seed extracts against some pathogenic bacterial strains. *Journal of King Saud University-Science* 26 (1): 79-82.
- BHOOPAT, T.; BHOOPAT, L.; SRICHAIRATANAKOOL, S.; KANJANAPOTHI, D.; TAESOTIKUL, T.; THANANCHAI, H. 2011. Hepatoprotective effects of lychee (*Litchi chinensis* Sonn.): A combination of antioxidant and anti-apoptotic activities. *Journal of Ethnopharmacology* 136 (1): 55-66.
- BRASIL. 2016. Ministério da Agricultura, Pecuária e Abastecimento. Instrução Normativa 52. Dez 2007. Available in: <http://extranet.agricultura.gov.br/sislegis-consulta/consultarLegislacao.do?operacao=visualizar&id=18316>. [Review date: 13 April 2018].
- BRITO, H. M.; GONDIM J. R., M. G. C.; OLIVEIRA, J. V.; CÂMARA, C. A. G. 2006. Toxicity of the natuneem over the *Tetranychus urticae* Koch (Acari: Tetranychidae) and mites predators from the family Phytoseiidae. *Ciência e Agrotecnologia* 30 (4): 685-691.
- BUTANI, D. K. 1977. Pests of litchi in India and their control. *Fruits* 32 (1): 269-270.
- CABRAL, T. A.; CARDOSO, L. M.; PINHEIRO-SANT'ANA, H. M. 2014. Chemical composition, vitamins and minerals of a new cultivar of lychee (*Litchi chinensis* cv. Tailandes) grown in Brazil. *Fruits* 69 (6): 425-435.
- CASTAGNOLI, M.; SIMONI, S.; GOGGIOLI, D. 2000. Attività biologica di sostanze vegetali nei confronti di *Tetranychus urticae* Koch (Acari: Tetranychidae) e del suo predatore *Neoseiulus californicus* (McGregor) (Acari: Phytoseiidae). *Redia* 83 (1): 141-150.
- CHILDERS, C. C.; EASTBROOK, M. A.; SOLOMON, M. G. 1996. Chemical control of eryophyoid mites. pp. 695-726. In: Lindquist, E. E.; Sabelis, M. W.; Bruin, J. (Eds.). *Eryophyoid mites - their biology, natural enemies and control*. Vol. 6. Elsevier Publishers, Amsterdam, The Netherlands, 690 p.
- COOLEN, W. A.; D'HERDE, C. J. 1972. A method for the quantitative extraction of nematodes from plant tissue. State Agriculture Research Center. Ghent, Bélgica. 77 p.
- COSTA, V. H. D.; SOARES, M. A.; RODRÍGUEZ, F. A. D.; ZANUNCIO, J. C.; SILVA, I. M.; VALICENTE, F. H. 2015. *Nomuraea rileyi* (Hypocreales: Clavicipitaceae) in *Helicoverpa armigera* (Lepidoptera: Noctuidae) larvae in Brazil. *Florida Entomologist* 98 (2): 796-798.
- DE FARIA, M. R.; WRAIGHT, S. P. 2007. Mycoacaricides an mycoacaricides: a comprehensive list with worldwide coverage and international classification of formulation types. *Biological Control* 43 (3): 237-256.
- DEMITE, P. R.; FERES, R. J. F. 2008. Seasonality of pathogenic fungi in mites of rubber tree plantations adjacent to fragments of Cerrado. *Brazilian Journal of Biology* 68 (3): 535-538.
- DEMITE, P. R.; LOFEGO, A. C.; FERES, R. J. F. 2008. Three new species of *Phytoseius ribaga* (Acari: Phytoseiidae), and a new record from Brazil. *Zootaxa* 1909: 16-26.
- DUSO, C.; MALAGNINI, V.; POZZEBON, A.; CASTAGNOLI, M.; LIGUORI, M.; SIMONI, S. 2008. Comparative toxicity of botanical and reduced-risk insecticides to Mediterranean populations of *Tetranychus urticae* and *Phytoseiulus persimilis* (Acari: Tetranychidae, Phytoseiidae). *Biological Control* 47 (1): 16-21.
- EDGINGTON, S.; FERNANDO, L. C. P.; JONES, K. 2008. Natural incidence and environmental profiling of the mitopathogenic fungus *Hirsutella thompsonii* Fisher for control of the coconut mite in Sri Lanka. *International Journal of Pest Management* 54 (2): 123-127.
- FERNANDO, L. C. P.; MANOJ, P.; HAPUARACHCHI, D. C. L.; EDGINGTON, S. 2007. Evaluation of four isolates of *Hirsutella thompsonii* against coconut mite (*Aceria guerreronis*) in Sri Lanka. *Crop Protection* 26 (7): 1062-1066.
- FORNAZIER, M. J.; MARTINS, D. S.; FORNAZIER, D. L.; AZEVEDO, L. H.; ZANUNCIO J. R., J. S.; ZANUNCIO, J. C. 2014. Range expansion of the litchi erineum mite *Aceria litchii* (Acari: Eriophyidae) in Brazil. *Florida Entomologist* 97 (2): 846-848.
- GADINO, A. N.; WALTON, V. M.; DREVES, A. J. 2011. Impact of vineyard pesticides on a beneficial arthropod, *Typhlodromus yri* (Acari: Phytoseiidae), in laboratory bioassays. *Journal of Economic Entomology* 104 (3): 970-977.
- GERSON, U.; SMILEY, R. L.; OCHOA, R. 2003. Mites (Acari) for pest control. Oxford, Blackwell Sciences. 539 p.
- GERSON, U.; GAFNI, A.; PAZ, Z.; SZTEJNBERG, A. 2008. A tale of three acaropathogenic fungi in Israel: *Hirsutella*, *Meira* and *Acaromyces*. *Experimental and Applied Acarology* 46 (1-4): 183-194.
- GONTIER, E.; BOUSSOUEL, N.; TERRASSE, C.; JANNOYER, M.; MÉNARD, M.; HUANG, K. 2008. *Aceria* (Acarina:

- Eriophyoidea) in Taiwan: five new species and plant abnormalities caused by sixteen species. *Zootaxa* 1829: 1-30.
- HAMEED, S. F.; SHARMA, D. D.; AGRAWAL, M. L. 1992. Integrated pest management in litchi. Proceedings of National Seminar on Recent Development in Litchi production held in 1992 at RAU, Pusa. 38 p.
- HASHEMI, S. M.; MOKHTARNIA, M.; ERBAUGH, J. M.; ASADI, A. 2008. Potential of extension workshops to change farmers' knowledge and awareness of IPM. *Science of the Total Environment* 407 (1): 84-88.
- HONG, X. Y.; WANG, D. S.; ZHANG, Z. Q. 2006. Distribution and damage of recent invasive eriophyoid mites (Acari: Eriophyoidea) in mainland China. *International Journal of Acarology* 32 (3): 227-240.
- HUANG, K. W. 2008. *Aceria* (Acarina: Eriophyoidea) in Taiwan: five new species and plant abnormalities caused by sixteen species. *Zootaxa* 1829: 1-30.
- JEPPSON, L. R.; KEIFER, H. H.; BAKER, E. W. 1975. Mites injurious to economic plants. Berkeley, University of California. 641 p.
- JIANG, G.; LIN, S.; WEN, L.; JIANG, Y.; ZHAO, M.; CHEN, F.; PRASAD, K. N.; DUAN, X.; YANG, B. 2013. Identification of a novel phenolic compound in litchi (*Litchi chinensis* Sonn.) pericarp and bioactivity evaluation. *Food Chemistry* 136 (2): 563-568.
- KHEDERI, S. J.; LILLO, E.; KHANJANI, M.; GHOLAMI, M. 2014. Resistance of grapevine to the erineum strain of *Colomerus vitis* (Acari: Eriophyidae) in western Iran and its correlation with plant features. *Experimental and Applied Acarology* 63 (1): 15-35.
- KURTTI, T. J.; KEYHANI, N. O. 2008. Intracellular infection of tick cell lines by the entomopathogenic fungus *Metarhizium anisopliae*. *Microbiology* 154 (6): 1700-1709.
- LALL, B. S.; RAHMAN, M. F. 1975. Studies on the bionomics and control of Erionose mite, *Eriophyes litchii* Keifer, Acarina: Eriophyidae. *Pesticides* 9 (11): 49-54.
- LINS, L. C. R.; SALOMAO, L. C. C.; CECON, P. R.; SIQUEIRA, D. L. 2015. The lychee tree propagation by layering. *Revista Brasileira de Fruticultura* 37 (2): 480-487.
- MARTINS, A. B. G.; BASTOS, D. C.; SCALOPPI JUNIOR, E. J. 2001. Lichieira (*Litchi chinensis* Sonn.). Sociedade Brasileira de Fruticultura, Jaboticabal, Brasil. 48 p.
- MCCOY, C. W. 1996. Pathogens of Eriophyoids. pp. 481-490. In: Lindquist, E. E.; Sabelis, M. W.; Bruin, J. (Eds.). *Eriophyoid mites - their biology, natural enemies and control*. Vol. 6. Elsevier Publishers, Amsterdam, The Netherlands, 690 p.
- McMURTRY, J. A.; SOURASSOU, N. F.; DEMITE, P. R. 2015. The Phytoseiidae (Acari: Mesostigmata) as biological control agents. pp. 133-150. In: Carrillo, D.; Moraes, G. J.; Peña, J. E. (Eds.). *Prospects for biological control of plant feeding mites and other harmful organisms*. Suíza, Springer.
- MESSING, R. H.; CROFT, B. A. 1996. Pesticide resistance in eriophyoid mites, their competitors and predators. pp. 689-694. In: Lindquist, E. E.; Sabelis, M. W.; Bruin, J. (Eds.). Vol. 6. Elsevier Publishers, Amsterdam, The Netherlands, 690 p.
- MEYLING, N. V.; LÜBECK, M.; BUCKLEY, E. P.; EILENBERG, J.; REHNER, S. A. 2009. Community composition, host range and genetic structure of the fungal entomopathogen *Beauveria* in adjoining agricultural and seminatural habitats. *Molecular Ecology Resources* 18 (6): 1282-1293.
- MOMEN, F. M.; HUSSEIN, H. 1999. Relationships between food substances, developmental success and reproduction in *Typhlodromus transvaalensis* (Acari: Phytoseiidae). *Acarologia* 40 (2): 108-111.
- MORAES, G. J.; LIMA, H. C. 1983. Biology of *Euseius concordis* (Chant) (Acarina: Phytoseiidae) a predator of the tomato russet mite. *Acarologia* 24 (3): 251-255.
- NISHIDA, T.; HOLDAWAY, F. G. 1955. The erineose mite of lychee. Agricultural Experiment Station, Hawaii. 48 p.
- PAZ, Z.; GERSON, U.; SZTEJNBERG, A. 2007. Assaying three new fungi against citrus mites in the laboratory, and a field trial. *Biocontrol* 52 (6): 855-862.
- PICOLI, P. R. F. 2010. *Aceria litchii* (Keifer) em lichia: ocorrência sazonal, danos provocados e identificação de possíveis agentes de controle biológico. Universidade Estadual Paulista, Faculdade de Engenharia de Ilha Solteira (Dissertação), Ilha Solteira, Brasil. 75 p.
- PICOLI, P. R. F.; VIEIRA, M. R. 2013. First report of pathogenic activity of *Hirsutella thompsonii* (Fischer) on the litchi erineum mite *Aceria litchii* (Keifer). *Semina: Ciências Agrárias* 34 (1): 187-190.
- PRISCHMANN, D. A.; JAMES, D. G.; WRIGHT, L. C.; TENEYCK, R. D.; SNYDER, W. E. 2005. Effects of chlorpyrifos and sulfur on spider mites (Acari: Tetranychidae) and their natural enemies. *Biological Control* 33 (3): 324-334.
- RAGA, A.; MINEIRO, J. L. C.; SATO, M. E.; MORAES, G. J.; FLECHTMANN, C. H. W. 2010. Primeiro relato de *Aceria litchii* (Keifer) (Prostigmata: Eriophyidae) em plantas de lichia no Brasil. *Revista Brasileira de Fruticultura* 32 (2): 628-629.
- RAGA, A.; MINEIRO, J. L. C.; SILOTO, R. C.; BERTON, L. H. C. 2011. Registros do ácaro-da-erinoze *Aceria litchii* (Keifer) (acari: Trombidiformes) no Estado de São Paulo. In: Simpósio Brasileiro de Acarologia, 3. Campinas. Anais. Campinas: Instituto Biológico.
- RANJAN, R.; RAY, R. 2015. Effect of abiotic factors on the incidence of litchi mite, *Aceria litchii* Keifer. *Pest Management in Horticultural Ecosystems* 21 (2): 225-227.
- ROSSONI, C.; KASSAB, S. O.; LOUREIRO, E. S.; PEREIRA, F. F.; COSTA, D. P.; BARBOSA, R. H.; ZANUNCIO, J. C. 2014. *Metarhizium anisopliae* and *Beauveria bassiana* (Hypocreales: Clavicipitaceae) are compatible with *Cotesia flavipes* (Hymenoptera: Braconidae). *Florida Entomologist* 97 (4): 1794-1804.
- SABELIS, M. W.; BRUIN, J. 1996. Evolutionary ecology: life history patterns, food plant choice and dispersal. pp. 329-365. In: Lindquist, E. E.; Sabelis, M. W.; Bruin, J. (Eds.). *World crop pests: Eriophyoid mites their biology, natural enemies and control*. Vol. 6. Elsevier Publishers, Amsterdam, The Netherlands, 690 p.
- SAHA, K.; SOMCHOUDHURY, A. K.; SARKAR, P. K. 1996. Structure and ecology of *Cephaleuros virescens* Kunze and its relationship with *Aceria litchii* Keifer (Prostigmata: Acari) in forming litchi erineum. *Journal of Mycopathological Research* 34 (2): 159-171.
- SAXENA, S.; HAJARE, S. N.; MORE, V.; KUMAR, S.; WADHAWAN, S.; MISHRA, B. B.; PARTE, M. N.; GAUTAM, S.; SHARMA, A. 2011. Antioxidant and radioprotective properties of commercially grown litchi (*Litchi chinensis*) from India. *Food Chemistry* 126 (1): 39-45.
- SCHULTE, M. J.; MARTIN, K.; SAUERBORN, J. 2007. Efficacy of spiromesifen on *Aceria litchii* (Keifer) in relation to *Cephaleuros virescens* Kunze colonization on leaves of litchi (*Litchi chinensis* Sonn.). *Journal of Plant Diseases and Protection* 114 (3): 133-137.
- SHARMA, D. D. 1984. Control of litchi mite. *Indian Horticulture* 29 (1): 27-28.
- SHARMA, D. D.; SINGH, S. P.; AKHAURI, R. K. 1986. Relationship between the population of *Aceria litchii* Keifer on litchi and weather factors. *Indian Journal of Agricultural Sciences* 56 (1): 59-63.
- SINGH, P.; SOMCHOUDHARY, A. K.; MUKHERJEE, A. B.; SARKAR, P. K. 1987. Susceptibility of different varieties of litchi to *Aceria litchii* Keifer (Acarina: Eriophyidae). Proceedings of 1st National Seminar on Agril. Acarology, held at Kalyani, pp. 35.
- SREEREMA KUMAR, P. S.; SINGH, L. 2002. Development of 'Mycohit', the first mycoacaricide based exclusively on *Hirsutella thompsonii*, for suppressing the coconut mite in

- India. pp. 209-213. In: Reddy, S. M.; Redy, S. R.; Singarachary, M. A.; Girisham, S. (Eds.). Proceedings, national symposium on bioinoculants for sustainable agriculture and forestry. India.
- SREEREMA KUMAR, P. S.; SINGH, L. 2008. Enabling mycelia application of *Hirsutella thompsonii* for managing the coconut mite. *Experimental and Applied Acarology* 46 (1-4): 169-182.
- SUGUINO, E. 2006. Influencia dos substratos no desenvolvimento de mudas de plantas frutíferas. Escola Superior de Agricultura "Luiz de Queiroz", Universidade de São Paulo (Tese), Piracicaba, Brasil.
- TANZINI, M. R.; ALVES, S. B.; TAMAI, M. A.; MORAES, G. J.; FERLA, N. J. 2000. An epizootic of *Calacarus heveae* (Acari: Eriophyidae) caused by *Hirsutella thompsonii* on rubber trees. *Experimental and Applied Acarology* 24 (2): 141-144.
- THAKUR, A. P.; SHARMA, D. D. 1989. New records of predatory mites on mite pests of economic crops in Bihar. *Biojournal* 1 (1): 155-156.
- THAKUR, A. P.; SHARMA, D. D. 1990. Influence of weather factors and predators on the populations of *Aceria litchii* Keifer. *Indian Journal of Plant Protection* 18 (1): 109-112.
- TIMPRASERT, S.; DATTA, A. A.; RANAMUKHAARACHCHI, S. L. 2014. Factors determining adoption of integrated pest management by vegetable growers in Nakhon Ratchasima Province, Thailand. *Crop Protection* 62 (1): 32-39.
- VAN DER GEEST, L. P. S.; ELLIOT, S. L.; BREEUWER, J. A. J.; BEERLING, E. A. M. 2000. Diseases of mites. *Experimental and Applied Acarology* 24 (7): 497-560.
- VAN LEEUWEN, T.; WITTERS, J.; NAUEN, R.; DUSO, C.; TIRRY, L. 2010. The control of eriophyoid mites: state of the art and future challenges. *Experimental and Applied Acarology* 51 (1-3): 205-224.
- WAITE, G. K. 1999. New evidence further incriminates honeybees as vectors of lychee erinose mite *Aceria litchii* (Acari: Eriophyiidae). *Experimental and Applied Acarology* 23 (2): 145-147.
- WAITE, G. K. 2005. Pests. pp. 237-259. In: Menzel, C. M.; Waite, G. K. (Eds.). *Litchi and Longan, Botany, Production and Uses*, CAB International, Wallingford, Reino Unido.
- WAITE, G. 2011. Lychee erinose mite in lychees. Queensland: Department of Plant Industries and Fisheries. Available in: <http://www2.dpi.qld.gov.au/horticulture/5432.ht>. [Review date: 13 April 2018].
- WAITE, G. K.; GERSON, U. 1994. The predator guild associated with *Aceria litchii* (Acari: Eriophyidae) in Australia and China. *Entomophaga* 39 (3): 275-280.
- WAITE, G. K.; HWANG, J. S. 2002. Pests of litchi and longan. pp. 331-359. In: Peña, J. E.; Sharp, J. L.; Wysoki, M. (Eds.). *Tropical fruit pests and pollinators: biology economic importance, natural enemies and control*. Wallingford: CABI.
- WAITE, G. K.; McALPINE, J. D. 1992. Honey bees as carriers of lychee erinose mite *Eriophyes litchii* (Acari: Eriophyiidae). *Experimental and Applied Acarology* 15 (4): 299-302.
- WEN, L.; WUA, D.; JIANGA, Y.; PRASADD, K. N.; LINA, S.; JIANGA, G.; HEA, J.; ZHAOB, M.; LUOB, W.; YANGA, B. 2014. Identification of flavonoids in litchi (*Litchi chinensis* Sonn.) leaf and evaluation of anticancer activities. *Journal of Functional Foods* 6 (1): 555-563.
- WEN, L.; YOU, L.; YANG, X.; YANG, J.; CHEN, F.; JIANG, Y.; YANG, B. 2015. Identification of phenolics in litchi and evaluation of anticancer cell proliferation activity and intracellular antioxidant activity. *Free Radical Biology and Medicine* 84 (1): 171-184.
- WESTPHAL, E.; MANSON, D. C. M. 1996. Feeding effects on host plants: gall formation and other distortations. pp. 231-242. In: Lindquist, E. E.; Sabelis, M. W.; Bruin, J. (Eds.). *World crop pests: Eriophyoid mites - their biology, natural enemies and control*. Vol. 6. Elsevier Publishers, Amsterdam, The Netherlands, 690 p.
- YAMANISHI, O. K.; MACHADO, J. A.; KAWATI, R. 2001. Overview of litchi production in São Paulo state Brasil. *Acta Horticulturae* 558 (1): 59-62.
- YAMANISHI, O. K.; PIRES, M. C.; ALMEIDA, L. F. P. 2010. The Brazilian lychee industry - present and future. *Acta Horticulturae* 863 (1): 59-65.
- YANG, B.; PRASAD, K. N.; JIANG, Y. 2016. Structure identification of a polysaccharide purified from litchi (*Litchi chinensis* Sonn.) pulp. *Carbohydrate Polymers* 137 (1): 570-575.

Received: 11-Oct-2016 • Accepted: 16-Feb-2018

Suggested citation:

CASTRO, B. M. C.; PLATA-RUEDA, A.; SILVA, W. M.; MENEZES, C. W. G.; WILCKEN, C. F.; ZANUNCIO, J. C. 2018. Management of *Aceria litchii* (Acari: Eriophyidae) on *Litchi chinensis*. *Revista Colombiana de Entomología* 44 (1): 2-7. Enero - Junio 2018.