

BIROn - Birkbeck Institutional Research Online

Nguyen, H.H. and Carter, Andrew and Hoang, L.V. and Fox, M. and Pham, S.N. and Vinh, H.B. (2022) Evolution of the continental margin of south to central Vietnam and its relationship to opening of the South China Sea (East Vietnam Sea). *Tectonics* 41 (2), ISSN 0278-7407.

Downloaded from: <https://eprints.bbk.ac.uk/id/eprint/47538/>

Usage Guidelines:

Please refer to usage guidelines at <https://eprints.bbk.ac.uk/policies.html>
contact lib-eprints@bbk.ac.uk.

or alternatively

1 **Evolution of the continental margin of south to central Vietnam and its relationship**
2 **to opening of the South China Sea (East Vietnam Sea)**

3 **Hiep Huu Nguyen¹, Andrew Carter², Long Van Hoang³, Matt Fox⁴ Sang Nhu Pham¹, Hau**
4 **Bui Vinh¹**

5 ¹Hanoi University of Mining and Geology, 18 Vien Street, Duc Thang ward, Bac Tu Liem
6 District, Hanoi, Vietnam

7 ²Department of Earth and Planetary Sciences, Birkbeck, University of London, Malet Street,
8 London WC1E 7HX, UK

9 ³Petroleum Exploration & Production Center Vietnam Petroleum Institute, VPI Tower, 167
10 Trung Kinh Street, Yen Hoa Ward, Cau Giay District, Hanoi, Vietnam

11 ⁴Department of Earth Sciences, UCL, Gower Street, London WC1E 6BT, UK

12 Corresponding author: Hiep Huu Nguyen (nguyenhuuhiep@humg.edu.vn)

13

14 **Key Points:**

- 15 • Apatite thermochronometry record South Vietnam margin exhumation history
- 16 • Thermal models are used to test different causes of thermal history
- 17 • Regional enhanced cooling and rock uplift between 37-30 Ma record early stage rifting

18 **Abstract**

19 The continental margin of south to central Vietnam is notable for its high elevation plateaus
20 many of which are covered by late Cenozoic basalt flows. It forms the westernmost margin of a
21 wide continental rift of the South China Sea (East Vietnam Sea), and uplift has been considered a
22 result of either rifting or younger intraplate basalt magmatism. To investigate margin
23 development apatite thermochronometry was applied to a dense array of samples collected from
24 across and along the margin of south to central Vietnam. Results, including thermal history
25 models, identified a distinct regional episode of fast cooling between c. 37 and 30 Ma after
26 which cooling rates remained low. The fast cooling coincides with a period of fast extension

27 across the South China Sea (East Sea) region that preceded continental break-up recorded by
28 Oligocene grabens onshore. A thermal model is used test different processes that might influence
29 the inferred cooling including a distinct pulse of exhumation; a decrease in exhumation followed
30 by an associated transient decrease in geothermal gradients and, underplating coincident with
31 rifting. Thermal relaxation following Mesozoic arc magmatism is ruled out as geotherms
32 returned to background rates within 20 Myrs of emplacement, well before the onset of fast
33 cooling. Models support fast cooling attributed to accelerated erosion during early stages of
34 rifting. Some additional heating from either underplating, and/or hot mantle upwellings is also
35 possible. No evidence was found to support regional uplift associated with the intraplate
36 magmatism, enhanced monsoon-driven erosion or seafloor spreading dynamics.

37

38 **Plain Language Summary**

39 This study used apatite thermochronometry to examine the uplift history of South-Central
40 Vietnam to better understand how the elevated landscape formed, and to see if it is connected to
41 the eruption of the widespread basalt flows that cover many of the high elevation plateaus.
42 Results showed a match in the timing between a period of rapid cooling between 37-30 million
43 years ago that affected the entire region and a period of fast rifting and extension that ended with
44 continental breakup. To fully understand the cause of cooling thermal models tested different
45 scenarios. Thermal relaxation after magmatism was discounted as it took place well before the
46 period of rapid cooling. Models support fast cooling attributed to accelerated erosion during
47 early stages of rifting. Magmatic underplating associated with rifting might also have produced
48 additional heating but no evidence was found to support regional uplift associated with the basalt
49 magmatism.

50

51 **1. Introduction**

52 The continental margin of Indochina represents the westernmost margin of the South China Sea,
53 a wide continental rift that in the west is mostly submerged. In the region of south to central
54 Vietnam, the location of this study (**Fig. 1**), the continental margin extends offshore for > 100
55 km before oceanic crust is reached. The extension that led to continental break-up has yet to be
56 fully explained but is known to be related to the subduction of the paleo-Pacific plate, which at

57 the time formed a wide Cretaceous magmatic arc with widespread calc-alkaline magmatism
58 across south to central Vietnam and southern China. Subduction was directed to the northeast
59 beneath southern China and westwards below Vietnam, the eastern margin of the Indochina
60 block (Zhu et al., 2004; Hall, 2012; Hennig-Breitfeld et al., 2021).

61
62 Extension is imprinted on the landscape and geology of Vietnam but to what extent is not yet
63 clear. It also remains uncertain as to when surface uplift took place. Was uplift staggered over a
64 long time or did it occur within a short episode? Stratigraphic evidence provides some
65 constraints. In south-central Vietnam many of the high plateaus, with typical elevations between
66 500-1500m a.s.l. (Fig. 1), are covered by basalt flows (mainly tholeiites) the majority of which
67 were erupted between the late Miocene to Quaternary (Hoang et al., 2013; An et al., 2017).
68 Eruptions were often from extensional faults or small axial rifts that formed previously during
69 extrusion and opening of the South China Sea (Flower et al., 1998; Hoang and Flower, 1998).
70 Rangin et al., (1995) recognized that many of the normal faults associated with these basalts
71 were reactivated N 160°E to N-S right-lateral faults. However, the timing of extension across
72 this margin is not well defined. A common view is that extension involved uplift of the rift
73 shoulders and that this drove widespread erosion from the latest Cretaceous-early Paleocene
74 (Matthews et al., 1997; Roques et al., 1997; Lee and Watkins, 1998; Lee et al., 2001; Andersen
75 et al., 2005; Pubellier et al., 2005). However, onshore, there are no geological constraints at
76 outcrop to confirm this. Basalt flows can be seen to drape over Precambrian to Proterozoic
77 crystalline and metamorphic basement rocks and in some places Jurassic to Cretaceous and late
78 Neogene sedimentary and magmatic rocks, but the temporal gap between the youngest rocks is
79 large, e.g., exhumed Cretaceous granites, and their basalt cover spans 80-90 Myrs. Offshore
80 constrains are little better. In southern Vietnam the Cuu Long and Nam Con Son rift basins
81 record a c. 50 Myr gap between the Eocene clastic sediments that were deposited on exhumed
82 Cretaceous granites and older rocks (Fyhn et al., 2009a).

83
84 As the deformation and tectonic evolution of the continental margin of south-central Vietnam
85 and its relationship to the opening of the South China Sea is poorly resolved we conducted a
86 thermochronometry study based on apatite U-Th-He and fission track (FT) data using rock
87 samples collected from across the continental margin. The utility of this approach is

88 demonstrated by previous work on the Kontum region (**Fig. 1**) of central Vietnam (Carter et al.,
89 2000) that identified a late Miocene increase in exhumation rates across the central margin
90 contemporaneous with increased sedimentation in the adjacent Phu Khan marine basin. It was
91 unclear if this phase of exhumation was caused by surface uplift of the continental margin and
92 base level change associated with breakup of the southwest sub-basin of the South China Sea
93 (Fyhn et al., 2009; Savva et al., 2014), and/or switching between rifting and left-lateral
94 transtension along the East Vietnam Boundary Fault Zone (**Fig. 1**). An alternative explanation is
95 that late Cenozoic magmatism had a local influence on rock uplift (Carter et al., 2000). Finally,
96 work by Clift et al., (2015) proposed that basin subsidence histories may have been affected by
97 increased loading associated with monsoon intensification driving faster sediment delivery to
98 offshore basins and therefore enhanced exhumation could be due to erosional unloading related
99 to changes in monsoon intensity.

100

101 **2. Geological Background**

102 During the Cretaceous eastern Indochina was part of a wide Andean type continental margin.
103 Calc-alkaline magmatism dates to between 118 – 87 Ma (Thuy et al., 2004; Shellnutt et al., 2013;
104 Hennig-Breitfield et al., 2021). The end of magmatism coincided with docking of the Luconia-
105 Dangerous Ground block with Indochina, and South China moving to the southeast to become
106 part of the South Asia margin (Fyhn et al., 2009a; Hall, 2012). These changes likely drove the
107 early rifting of the Proto-South China Sea in either a back-arc setting on the East Asian margin
108 or slab-pull induced microcontinent detachment from subduction along Northern Borneo (Doust
109 and Sumner, 2007).

110

111 Spreading in the South China Sea, that started at c. 32 Ma (Barckhausen et al., 2014) affected the
112 continental margins of both Indochina and SE China. Pre-spreading extension of the continental
113 lithosphere produced more or less E-W oriented normal faults across SE China and South
114 Vietnam margins although in the Cu Long basin the orientation is closer to SW-NE (Schmidt et
115 al., 2019). With the onset of ocean spreading, the orientation of new faults moved to NE-SW
116 reflecting the progression of spreading to the southwest (Savva et al., 2014). A ridge jump at 25
117 Ma relocated spreading to a southwestern sub-basin until spreading ended at 15 Ma (Li et al.,

118 2014), when the subduction zone became blocked by collision with continental fragments
119 (northern part of Palawan and/or a part of the Dangerous Grounds).

120

121 It is widely assumed that the Late Cretaceous to Early Paleocene early extension drove uplift of
122 the rift shoulders followed by widespread erosion and peneplanation (Taylor and Hayes, 1983),
123 but this is based on offshore seismic interpretations of poorly dated early syn-rift sediments. The
124 imprint of extension and rifting on the south-central Vietnam continental margin is different
125 from the SE China margin due to a combination of re-activation of inherited structures (mainly
126 Indosinian) and the local influence of tectonic extrusion driven by India-Eurasia collision (Fyhn
127 et al, 2009a). Although the amount of left-lateral extrusion is debated (Searle et al., 2010) the
128 position and orientation of Indochina must have changed throughout the period of South China
129 Sea opening, and any change would have affected the regional stress field. Four phases of late
130 Mesozoic–Cenozoic deformation are recognised in Vietnam; 1) E-W compression, N-S
131 extension, 2) NNW-SSE compression, ENE-WSW Extension, 3) N-S compression, E-W
132 extension and 4) NNE-SSW compression, ESE-WNW extension (Nguyen and Hoai, 2019).

133

134 **2.1. Records from offshore basins**

135 Extension and rifting led to the formation of the Phu Khanh and Cuu Long basins (**Fig. 1**). The
136 Phu Khanh Basin was controlled by the N-S striking East Vietnam Boundary Fault, considered a
137 continuation of the Red River Shear Zone, as well as NE-SW directed normal faults (Fyhn et al.,
138 2009a). Further south the Cuu Long Basin is bounded by NE-SW to E-W striking normal faults
139 (Schmidt et al., 2019). The sedimentary archives of these basins reflect onshore erosion.

140

141 **2.2. Phu Khanh Basin**

142 This mainly deep-water basin is situated at the base of the continental slope off central Vietnam
143 and is separated from the mainland by the East Vietnam Boundary Fault (**Fig. 1**) (Fyhn et al.,
144 2009b). The basin history records two major rifting events (Lee, and Watkins, 1998; Savva et al.,
145 2014). Syn-Rift I in the Palaeogene was accompanied by the deposition of clastic and lacustrine
146 sediments. Syn-Rift II is associated with the opening of the South China Sea and from the end of
147 the Oligocene involved mainly marine sedimentation. A distinct unconformity marks the
148 Paleogene–Neogene boundary (Fyhn et al., 2009b) across which rifting decreased significantly.

149 Post-rift sediments include upper Miocene–Pliocene turbidite fans interpreted as the product of
150 high rates of onshore erosion (**Fig. 2**).

151

152 **2.3. Cuu Long Basin**

153 This basin is located offshore southeast Vietnam (**Fig. 1**) across the continental shelf and was
154 formed by rifting during the Late Eocene–Early Oligocene (Schmidt et al., 2019). The basin
155 contains up to 8 km of mainly clastic sediments deposited between the Eocene to Quaternary
156 (Morley et al., 2019). A narrow valley developed during the first (syn-rift) phase of extension,
157 and in the basin axial zone rapid subsidence took place in the late Oligocene. Whilst extensional
158 deformation extended into the Miocene, a distinct unconformity at ~ 27 Ma marks inversion and
159 a transition to post-rift sag. The main phase of post-rift thermal sag took place after 23 Ma
160 (Morley et al., 2019).

161

162 In the Early Miocene, a new phase of seafloor spreading was accompanied by a period of sea
163 level rise. This caused a marine transgression in all of the basins along the western edge of the
164 South China Sea (Lee, and Watkins, 2001) leading to the formation of carbonate and coral reefs
165 (Fyhn et al., 2009a). Later post rift subsidence formed a broader shallow sag basin with clastic
166 sedimentation. Our low-temperature thermochronometry results should show which parts of the
167 continental margin experienced the enhanced erosion that drove the clastic sedimentation.

168

169 **3. Methodology**

170 The partial retention zone (c. 80–40 °C) of the (U-Th-Sm)/He (AHe) and partial annealing zone
171 (c. 130–60 °C) of the apatite fission track (AFT) methods are well placed to provide constraints
172 on timing, rate, and magnitude of bedrock exhumation in the uppermost crust (Lisker et al.,
173 2009). For this study a primary goal is to use apatite thermochronometry to detect when rock
174 uplift and exhumation accelerated in response to either surface uplift and/or denudation.

175 Mapping the pattern of exhumation in relation to the extended continental margin will allow
176 assessment of the impact of extension on the continental margin of southern Vietnam. This aim
177 guided our bedrock sampling strategy that comprised a series of north to south, coast to interior
178 transects across the study area, spanning the local relief. Studies of rift and passive margin
179 erosion patterns based on apatite fission track data (Wildman et al., 2019) have shown that the

180 amount of erosion is generally greatest along the coastal areas and decreases inland. This is
181 mainly due to isostatic unloading: As a rift escarpment migrates inland, rock uplift rates remain
182 high close to the margin leading to higher exhumation rates and younger ages. Variations in
183 erosion caused by local geomorphic conditions and geology such as reactivation of inherited
184 faults and magmatic underplating add noise to this dominant trend.

185

186 In total 67 samples were collected of which 42 (**Fig. 3**) produced good quality apatite suitable for
187 analysis. Each sample weighed between 2–5 kg and were mainly granites as early sampling
188 found that other rock types contained little, if any, apatite. Supplementary table S1 details
189 sample locations. All thermochronometry analyses were conducted at the London
190 Geochronology Centre. Fission track analyses used conventional neutron irradiation procedures
191 based on the external detector method and zeta calibration approach (Hurford and Green, 1983).
192 Apatites were mounted, polished and etched in 5N nitric acid 20 °C for 20 seconds. Track length
193 measurements were used to constrain sample cooling rate, and grain bulk composition was
194 monitored using etch pit length (DPAR) measured parallel to the crystallographic c-axis
195 (Donelick 1993). (U-Th-Sm)/He analyses typically involved between 4 and 6 replicates. Apatite
196 grains were placed in platinum tubes and outgassed using a 25 W, 808 nm diode laser and ⁴He
197 measured on a Balzers quadrupole mass spectrometer. Following apatite dissolution and spiking
198 U, Th, and Sm concentrations were measured on an Agilent 7700x ICP-MS. Further details on
199 the analytical protocols can be found in the supplementary section.

200

201 **4. Results**

202 **4.1. Low-Temperature Thermochronometry**

203 Apatite (U-Th-Sm)/He analysis was performed on 18 representative samples from across the
204 elevation range (Table S2). These typically show age dispersion between individual replicates,
205 some of which can be explained by variations in grain size, accumulated radiation damage, and
206 apatite chemical composition (Fig. S1). Such factors can affect helium diffusion kinetics and
207 produce a range of (U-Th)/He dates from a single rock. However, spurious ages can also arise
208 from alpha particle implantation from external sources, U-Th rich inclusions, inaccurate grain

209 size measurement and alpha ejection correction, problematic outgassing, incomplete grain
210 dissolution and sample handling. Outlier ages were only rejected where there was clear evidence
211 of experimental issues.

212 AFT analysis was carried out on 47 samples of which 44 yielded good quality age data (Table
213 S3. By contract, track length data was affected by low spontaneous track densities and only 11
214 samples yielded sufficient track lengths for modelling. Sample central ages range from 70 ± 5 to
215 22 ± 3 Ma, with a weighted average of 41 Ma. Apatite etch pit lengths measured parallel to the
216 grain c-axis (DPAR), a proxy for grain bulk composition, ranges from 1.7 to 3.8 μm . If cooling
217 rates were low DPAR and AFT age would be expected to correlate, due to a compositional
218 influence on FT annealing. As there is no correlation this implies cooling rates were not slow,
219 borne out by FT track length data (samples with > 50 measurements) that show unimodal
220 distributions and long mean lengths between 13.5 μm and 14.2 μm ($n = 16$). Furthermore, a
221 comparison between AHe ages and effective uranium (eU) values and grain radius, based on a
222 spherical geometry, revealed no trends to support slow cooling (Fig. S1). If this were the case
223 single AHe ages should correlate with eU (proxy for radiation damage) (Flowers et al., 2009)
224 Likewise, larger grains, with a slightly higher closure temperature, would be appreciably older
225 than smaller grains if cooling was slow.

226
227 Figure 4 plots all replicate AHe ages and AFT central ages against sample elevation. Most ages
228 fall between 20-60 Ma, irrespective of sample elevation, that points to a phase of regional uplift.
229 There is no well-developed age trend between coastal areas and furthest points inland as would
230 be expected from a conventional passive margin whereby the youngest ages would be confined
231 to the coastal areas and oldest ages restricted to the highest elevations inland. Within the dataset,
232 AHe ages tend to increase with elevation but this is less well developed in the AFT data. Clear
233 evidence is seen in **Fig. 5** that shows ages plotted for a suite of samples collected across a 1421m
234 elevation range within a 92 Ma granite, located close to Nha Trang on the coast. Proximity to the
235 coast means that the data are more likely to record a primary signal of rock uplift and erosion:
236 points further from the coast will be less sensitive to a geomorphic response to rock uplift. In
237 other words, any delay between uplift and erosion will depend on geomorphic processes, which
238 may take longer to respond to changes in tectonics further from the baselevel (sea-level). The

239 plot shows exhumation rates accelerated after 40-50 Ma based on the changing slopes of the age-
240 elevation relationship.

241

242 **4.2. Thermal history models.**

243 To constrain rock uplift history, the thermal histories of representative samples, with sufficient
244 track length measurements, were obtained using the QTQt software of Gallagher (2012). This is
245 based on a Bayesian transdimensional approach to data inversion to extract probable thermal
246 histories. Model outputs are accepted thermal history models that can be combined to give a
247 mean thermal history model weighted by the posterior probability of each individual thermal
248 history with 95% credible intervals that provide a measure of uncertainty. Model runs allowed
249 the temperature offset to vary over time and data were predicted using the annealing and
250 diffusion models Ketcham et al (2007) and Gautheron et al., (2009). Granite emplacement age
251 was the only time-temperature constraint used in the inversion based on our own zircon U-Pb
252 analyses or published ages (Shellnutt et al., 2013; Hennig-Breitfield 2021) .

253

254 Results from the multi-sample vertical profile inversion shown in **Fig. 5** are presented in **Fig. 6**.
255 Models produce a reasonable fit to the data although some AHe ages are underpredicted. For the
256 low elevation data underprediction, could be due to a recent increase in cooling such that the
257 amount of rock uplift and exhumation is insufficient to provide a clear signal in the AHe data.
258 However, thermal history models of low elevation samples with the youngest AHe ages from
259 across the study area (S2) show no significant departure from the post 30 Ma cooling path in
260 figure 6. Likewise models of high elevation samples from across the study area are consistent
261 (residence at crustal temperatures < 60°C since 50-60 Ma) with the thermal history model in **Fig.**
262 **6**. The models suggest the region experienced a multi-stage thermal history that involved
263 accelerated cooling between 60-50 Ma, minor reheating between 50-35 Ma, and accelerated
264 cooling between ~ 37-30 Ma followed by constant cooling to the present. The early phases of
265 cooling are not well constrained as oldest tracks do not extend to the middle Cretaceous hence
266 the thermal history between 90 Ma and about 50 Ma are largely driven by the requirement of
267 granite emplacement. Resolution increases through time as shown by the difference between the
268 credible intervals for specific samples. The reheating stage of the history from 50 to 37 Ma is
269 relatively well resolved. From 37 Ma, however, resolution increases and the accelerated cooling

270 between 37-30 Ma is well constrained and consistent with the regional age-elevation data plotted
271 in **Fig. 5**. These two stages require average offset temperatures (differences between lowest and
272 highest elevation samples) to rise to up to 80°C equivalent to a geothermal gradient of
273 ~57°C/km, but then temperature offsets decrease through time. This decrease in geothermal
274 gradient through time is partly required by the fact that both high and low elevation samples
275 must reach similar temperatures at the surface. However, the ages are sensitive to times as recent
276 as 20 Ma and therefore, the change in geothermal gradient is likely to be robust and not an
277 artifact.

278

279 **5. Discussion**

280 Dense sampling across the study area, extending from the coast up to 75 km inland yielded 42
281 AFT ages and 77 AHe replicate single grain ages distributed across an elevation range of 1524
282 m. Ages do not increase from the coast to inland as would be expected from a typical rifted
283 margin where there is limited erosion landward of an escarpment (Gallagher and Brown, 1997).
284 The majority of sample ages cluster between 20-50 Ma, independent of elevation, consistent with
285 regional block uplift. Although some of the highest elevation data show a relationship between
286 age and elevation reflecting an earlier exhumation history, most of the data below 800 m record
287 similar ages indicative of faster cooling. This contrast in cooling rates is best seen in data from a
288 suite of samples collected across a 1.4 km vertical section of a 92 Ma granite (**Fig. 5**) that
289 captures the regional thermal history. QTQt thermal history models (**Fig. 6**) show a multi-stage
290 history with reheating between ~ 50-37 Ma, and well-resolved accelerated cooling between 37-
291 30 Ma followed by uninterrupted slower cooling to the present. Differences in temperature
292 between the lowermost and uppermost samples between 60-30 Ma require a high geothermal
293 gradient (c. 57°C/km) that when applied to the accelerated cooling between ~37-30 Ma would
294 crudely translate to about 1500 m of rock uplift and exhumation. This regional thermal history
295 raises several questions: 1) What caused the interval of reheating between 50-37 Ma? 2) What
296 caused the interval of accelerated cooling between 37-30 Ma and was this period related to the
297 generation of present-day topography? 3) Are there any connections between changes in cooling
298 rate, the opening of the South China Sea (East Vietnam Sea) and sedimentation in the marine
299 basins?

300

301 In order to address these questions, we require a thermal model that relates cooling histories of
302 rocks to changes in exhumation rate and the thermal structure of the crust. Using this thermal
303 model we can test different factors that might influence the inferred cooling. In the next section,
304 we first highlight that the elevated heating from arc magmatism and emplacement of the sampled
305 granites is a transient feature that does not persist beyond about 20 Myrs. Therefore our resolved
306 changes in cooling since about 50 Ma are related to other processes. We highlight three different
307 processes that predict similar cooling histories: a distinct pulse of exhumation; a decrease in
308 exhumation followed by an associated transient decrease in geothermal gradients; and, recent
309 underplating coincident with rifting but unrelated to arc magmatism and recent basaltic
310 magmatism.

311
312 We use a simple 1D transient thermal model to explore these processes. The model solves the
313 heat transport equation with Dirichlet boundary conditions at surface and lower crustal
314 temperatures, tracking material points through the evolving thermal field to provide time-
315 temperature paths. Numerically, the model uses finite differences in space and the Crank-
316 Nicolson method for temporal integration (see Fox and Carter, 2020). The thermal model is 35
317 km thick, used boundary conditions set at 15 °C at the surface and 800 °C at the base to give an
318 unperturbed geothermal gradient of 22 °C/km as this is about the average geothermal gradient
319 predicted by the QTQt models over the last 40 Ma. A thermal diffusivity of 25 km²/Ma is used
320 for all models. The timestep length is set to 1x10⁻³ Myr and the vertical resolution is 0.2318 km.
321 The initial condition is modified to reflect the emplacement of magmas at different depths.
322 Exhumation rate histories are also modified to explore the different scenarios.

323

324 **5.1. Relationship between cooling and exhumation**

325

326 **The influence of magmatic heating during emplacement**

327 As a former magmatic arc the study area would have experienced increased heating and elevated
328 geothermal gradients followed by post-magmatic thermal relaxation. A fundamental question is
329 whether this impacted on the low-temperature thermochronology data. Murray et al., (2018)
330 investigated this question using 1D and 3D models in which heat diffused from a magmatic
331 emplacement and rocks were advected towards the surface. They were able to show that large

332 midcrustal plutons, emplaced at depths between 10–15 km, can reset low-temperature
333 thermochronometers in the upper crust. This is not surprising, but of importance to this study is
334 the question of how long such thermal effects persisted following magma emplacement? There is
335 at least a 40 Myr gap between granite ages and the earliest cooling constrained by the apatite
336 data. Could this early stage cooling be a consequence of post magmatic thermal relaxation? To
337 explore this we used the same approach as Murray et al., (2018) to assess the effects of heating at
338 depth on the time-temperature paths experienced by thermochronometry samples. The cartoon in
339 Figure 7 shows the basic evolution of the model. Heating can lead to increased geothermal
340 gradients at all depths. Samples that are exhuming will experience this heating and subsequent
341 cooling. The exact heating and cooling rates will depend on the duration of heating, the
342 exhumation rate, the depth of heating and the samples and the thermal diffusivity of the crust
343 (Murray et al., 2018).

344

345 Figure 8 shows the results of a numerical model designed to simulate high temperature at the
346 start of the model. Here, temperatures are high during the first 2 Myrs of the model run from 8-
347 12 km depth. This heats a large section of the crust. After this initialization period, heat is lost
348 through diffusion and temperatures rapidly return to a steady state solution. It is important to
349 note that the steady state solution shows slightly higher geothermal gradients than the initial
350 condition due to the advection of heat driven by erosion which is constant at 0.1 km/Myr. The
351 rocks that are at the surface at the end of the model are tracked along the red curve. This is
352 highlighted Figure 8B and shows the rapid cooling following emplacement. For reference, a
353 thermal history is shown in which there is no heating at the start of the model. Only the high
354 temperature $^{40}\text{Ar}/^{39}\text{Ar}$ in hornblende and muscovite thermochronometric systems would be
355 influenced by this heating event. The other thermochronometers would not be sensitive to this
356 thermal event and therefore, we can begin the subsequent models at 40 Ma.

357

358 Next, we consider two other processes that may produce similar, high rates of cooling over the
359 same sort of time interval: 1) relaxation of geothermal gradient following the cessation of
360 exhumation and, 2) cooling following a pulse of heating associated with emplacement of
361 underplating material below the lower crust associated with the rifting.

362

363 **Changes in exhumation rate**

364 We use a reference scenario in which the cooling is interpreted in terms of exhumation (Fig. 9).
365 In this scenario, a pulse of exhumation with rates of 0.6 km/Ma is simulated between 37 and 30
366 Ma and the background rate is set to 0.02 km/Ma (consistent with Quaternary cosmogenic
367 erosion rates measured in a river catchment to the north of the study area by Jonell et al., 2017).
368 This leads to relatively fast cooling over this time interval recreating the accelerated cooling
369 between 37 and 30 Ma. However, the peak in cooling rate is at 30 Ma and there is just a small
370 change in the geothermal gradient. This is in contrast with the results of the QTQt model that
371 highlights large changes in the geothermal gradient through time. It is important to note that we
372 are highlighting the geothermal gradient at the changing depth of the synthetic sample as it is
373 exhumed. This is to be consistent with the temperature offset resolved by QTQt. In addition, it is
374 not clear how exhumation rates might change so abruptly given the gradual changes in surface
375 processes expected following changes in rock uplift.

376
377 A second scenario is simulated in which a pulse of exhumation that began at 40 Ma at rates of
378 0.75 km/Ma and then decreased exponentially from 37 Ma to the present to a background rate of
379 0.01 km/Ma (**Fig. 9**). This exhumation rate history approximates a geomorphic response to rock
380 uplift in which erosion rates are high during rock uplift and then decay after the cessation of
381 active rock uplift. High rock uplift rates elevate the geothermal gradient due to the advection of
382 heat. As exhumation rates decrease, the geothermal gradients relax but rocks continue to
383 exhume. Importantly, the peak in cooling rate actually postdates the phase of active rock uplift
384 and therefore the phase of rapid cooling resolved by the thermochronometric data may be the
385 result of an earlier phase of active rock uplift. Although this scenario modifies the geothermal
386 gradient due to heat advection, the magnitude of the change in geothermal gradient is not as large
387 as inferred using QTQt. As with the first scenario, geothermal gradients decrease following the
388 phase of fast exhumation.

389

390 **Changes in boundary conditions**

391 A third scenario simulates the emplacement of underplating material below the lower crust (**Fig**
392 **9**). In seismic sections further to the north in the South China Sea, lower-crust high-velocity
393 anomalies have been observed and related to magmatic underplating at depths of about 25-30 km

394 (Li et al., 2020). In this model a pulse of heating is set to 1500 °C between depths of 25 and 35
395 km from 40-38 Ma and this simulates a change in temperature at the base of the crust associated
396 with rifting. Such depths and temperatures are not unrealistic as the geochemistry of local
397 Cenozoic calc alkali magmatism indicates a hotter than normal shallow asthenosphere (Hoang et
398 al., 2013) and the lithosphere in this study area is some 20-30 km thinner than northern Vietnam
399 (Vu et al., 2021), presumably in part due to extension and rifting. Following this heating event,
400 the lower boundary is returned to 800 °C and temperatures diffuse back to the normal conditions
401 over time. We acknowledge that it is unrealistic that temperature would return to a fixed value
402 immediately after this heating event, however, this is sufficient to highlight this process. The
403 background exhumation rate is set to 0.075 km/Ma for the duration of the model. This value is
404 slightly higher than the 0.02 km/Ma inferred using cosmogenic nuclide concentration
405 measurements (Jonell et al., 2017) but we hope to highlight that a constant exhumation rate
406 through time can produce the observed cooling history, and 0.02 km/Ma for the duration of the
407 model would require impossibly high geothermal gradients to exhume rocks from high enough
408 temperatures. This scenario produces a reheating signal as temperatures increase, then as
409 temperatures decrease and rocks continue to exhume, a pulse of cooling is predicted. There is a
410 hint of reheating in the QTQt models. However it is not very strong and is poorly resolved.
411 Cooling rates during the cooling pulse are lower than in the other two scenarios. However, the
412 geothermal gradient is much higher and decreases through time, as required by the data.

413

414 These three scenarios highlight the range of geological models that produce similar time-
415 temperature paths. Other geological models could be envisioned including changing thermal
416 diffusivities through time, different boundary conditions or many other factors. The large
417 changes in geothermal gradient through time predicted by QTQt are hard to reproduce with a
418 simple advection-diffusion equation and require changes in boundary conditions through time or
419 the emplacement of hot material at the base of the model, as simulated in scenario 3. However,
420 the rate of cooling following this is too slow in scenario 3 and this suggests that there is
421 additional exhumation during rifting. The data are, therefore, likely providing a record of
422 accelerated erosion during rifting and additional heating possibly related to magmatic
423 underplating in the lower crust.

424

425 **5.2. Enhanced denudation associated with monsoon climate?**

426 Climate is a key driver of denudation. The study area is affected by the East Asian monsoon and
427 therefore the changes in cooling rate detected by the thermochronometry might reflect enhanced
428 rates of erosion and exhumation driven by monsoon climate change. Reconstructions of the East
429 Asian monsoon show a relatively stable wet environment throughout the late Eocene and
430 Oligocene that strengthened after 23 Ma, reaching a maximum between 18-10 Ma (Clift et al.,
431 2014, Farnsworth et al., 2019). These timings do not coincide with the changes seen in the
432 apatite thermal histories. Notably, cooling rates declined at a time when the monsoon intensified.
433 Hence there is no obvious connection between changes in monsoon intensity and regional
434 thermal history.

435
436 In the offshore basins lacustrine/fluvial environments dominated the Eocene, and sediment
437 accumulation rates do not appear to change, although this could be due to limited
438 accommodation space and poor development of regional drainage systems. Between the
439 Oligocene and early Miocene there was a switch from siliciclastic sedimentation to carbonates
440 associated with shallow marine lagoonal and reefal environments. Only in the latest part of the
441 Miocene do offshore basins record a marked rise in siliciclastic sedimentation, largely associated
442 with shoreface facies. At this time accumulation rates were sufficiently high to cause
443 progradation, especially in the Phu Khanh Basin (Fyhn et al., 2009; Lee and Watkins, 1998).
444 Paradoxically, this occurred at a time when the East Asian monsoon began weakening (Clift et
445 al., 2014). Likewise, ¹⁰Be Cosmogenic derived denudation rates for the Song Gianh monsoon-
446 dominated river in northern central Vietnam (Jonell et al., 2017), show lower rates for the strong
447 monsoon of the Early Holocene compared to longer-term erosion rates based on local apatite
448 thermochronometry data. Based on this evidence past changes in monsoon climate intensity do
449 not explain the thermochronometry data.

450

451 **5.3. Timing of surface uplift**

452 By the early Cenozoic, it is widely held that a peneplain or low relief surface extended across
453 much of Indochina (Tran et al., 2011) hence surface uplift must post-date this. Tighter
454 constraints are provided by the onshore geology. Onshore South Vietnam, the c. 300 km long
455 Song Ba Rift formed within a major NW–SE-trending strike-slip fault zone (**Fig 1**) that

456 continues offshore and separates the Phu Khanh Basin from the Cuu Long Basin. Structural
457 observations show the faults cross-cut Cretaceous rocks and were therefore active in the
458 Cenozoic when deformation reactivated the strike-slip faults as extensional faults (Nielsen et al.,
459 2007). The Song Ba rift contains up to 500 m of syn-rift Oligocene sediments but their thermal
460 maturity (vitrinite reflectance % Ro c. 0.4) indicates that the grabens were originally deeper and
461 contained more fill, between 1-2 km. This means that substantial uplift and denudation occurred
462 after the Oligocene but before eruption of the Miocene basalts that drape the uplift unconformity
463 (Nielsen et al., 2007). Some basalt flows are also seen to infill pre-existing valleys. Hoang and
464 Flowers (2013) provide examples of 6–8 Ma lava flows (their Fig. 2) infilling over 300m of local
465 relief. Such evidence favours a close relationship between topographic growth and rifting
466 although it is possible surface uplift continued after active rifting had ended.

467

468 **5.4. Paleogene rifting**

469 The 37–30 Ma interval of accelerated cooling recorded by the thermochronometry data overlaps
470 active rifting in both onshore and offshore basins across central and south Vietnam. Onshore
471 evidence for active erosion during this time can be found within Oligocene clastic deposits of the
472 Di Linh Formation of the Da Lat zone. Detrital zircon U-Pb data from these rocks are dominated
473 by Cretaceous magmatic ages, consistent with erosion of the Cretaceous igneous rocks on which
474 they sit (Hennig et al., 2018). The Di Linh Formation is possibly an extension of the offshore
475 Cuu Long Basin where sediments were first deposited during the latest middle Eocene. Similar to
476 onshore, Paleogene sediments rest on an older basement that includes exhumed Cretaceous
477 granites. Active rifting and inversion took place during the late Eocene to early Oligocene and
478 had almost ceased by the end of the Lower Oligocene, recorded by a basin-wide unconformity
479 dated to c. 28 Ma (Morley, et al., 2019). After this the basin entered a short period of
480 compression due to the Mekong Delta Fault Zone switching from left-lateral to right-lateral
481 strike-slip (Schmidt et al., 2019). This regime ended circa 25 Ma when strike-slip motion on the
482 Mekong Delta Fault Zone stopped, and the basin then transitioned into post rift thermal
483 subsidence.

484

485 The influence of a large fault zone is also seen in the Phu Khan Basin where left-lateral motion
486 along the East Vietnam Boundary Fault Zone (**Fig. 1**) affected early rifting (Fyhn et al., 2009a).

487 Easternmost parts of the basin also show evidence for contemporaneous NW to SE extension
488 suggesting some of the extension was due to other factors, such as slab pull from subduction of a
489 proto-South China Sea (Vu et al., 2017). An unconformity at the Oligocene-Miocene boundary
490 marks the end of rifting in this basin, and a switch to regional post-rift subsidence led to a
491 transgression.

492

493 The 37–30 Ma interval of accelerated cooling recorded by the thermochronometry data also
494 spanned regional changes in the stress regime associated with rifting and the onset of ocean
495 spreading c. 32 Ma (**Fig. 10**). Ocean Discovery Program expeditions 367/368, at the northern
496 South China Sea margin, identified a major period of fast extension during the late Eocene to
497 early Oligocene (Larsen et al., 2018), roughly 40–30 Ma. This relatively short rifting event
498 spanned the rift-to-igneous crustal accretion transition and was associated with rapid upwelling
499 of the asthenosphere, seeding the seafloor spreading that took place soon after. This would have
500 affected all surrounding margins, and it culminated in the formation of a regional breakup
501 unconformity at circa 33–28 Ma after which there was a regional reduction in extensional
502 activity (Morley et al., 2016; 2019). This timing is shared by basins offshore the study area.
503 During this time Indochina also underwent extrusion along the Ailao Shan-Red River Fault Zone
504 (ARRFZ). Left-lateral shearing along the Day Nui Con Voi shear zone, the southern extension of
505 the ARRFZ, took place between ~35 Ma to ~20 Ma (Jolivet et al., 2001; Leloup et al., 2001 Liu
506 et al., 2020; Searle et al., 2010). However, it seems unlikely that this had a major influence on
507 deformation across the study area since the duration of extrusion extended well beyond the
508 episode of fast cooling.

509

510 **5.5. Relationship to ocean spreading.**

511 Onset of seafloor spreading at magnetic anomaly 11, c. 32 Ma (Barckhausen et al., 2014; Sibouet
512 et al., 2016) began in the East South China Sea sub-basin opening in a roughly N-S direction.

513 But, at 23 Ma (anomaly 6a) a ridge jump changed the position and direction of spreading by
514 rotating about 15–20° anticlockwise to form the South West sub-basin (Sibouet et al., 2016).

515 This shift in orientation caused a change in regional stress and coincides with unconformities in
516 the Cuu Long and Phu Khanh basins (**Figs. 1&8**) (Fyhn et al., 2009a; Schmidt et al., 2019).

517 Changes in the regional stress regime are also indicated by changes in rates of the propagation of

518 spreading. Between 32–23 Ma rates of propagation in the East South China Sea sub-basin were
519 slow but increased in both sub-basins after the ridge jump (Le Pourhiet et al., 2017).

520

521 Were these changes sufficient to explain the fast cooling? During spreading, as the asthenosphere
522 rises and spreads along the rift axis there will be accompanying changes in the force balance
523 between far-field stresses, resulting from margin composition, structure and topography, and
524 local buoyancy from thinning of the lithosphere. Margin uplift would be expected where the
525 rising asthenosphere induces gradients of gravitational potential energy by juxtaposing denser
526 asthenospheric material against thicker crustal material (Rey, 2001; Mondy et al., 2018)
527 however, a large topographic load from the surrounding continental margin can transmit
528 significant compressive stresses that oppose buoyancy forces. In the case of the South China Sea
529 (East Vietnam Sea) 3D numerical simulations have demonstrated how a small amount of
530 compression, and/or extension, acting normal to the direction of propagation can influence the
531 rate of breakup propagation (Le Pourhiet et al., 2017). The slow propagation stage of rifting
532 within the East South China Sea sub-basin can be explained by resistive, compressive stresses
533 from the west-to-east topographic load of Indochina. The change following the ridge jump at 23
534 Ma, caused the direction of spreading propagation to become oblique to Indochina and reduced
535 much of the resisting out-of plane compression allowing the propagation of spreading to the SW
536 to accelerate. Whilst the slow propagation stage overlaps with the fast cooling seen in the QTQt
537 models (**Fig. 6**), this is not the case for the timing of the switch to fast rift propagation since it
538 significantly post-dates the slowdown in cooling rates. Thus, whilst it may be appealing to link
539 the onshore thermal history to changes in spreading and regional stresses there is no strong
540 evidence to support this.

541

542 **5.6. Relationships to mantle structure**

543 Between ~17–0.2 Ma widespread intraplate volcanism took place across south-central Vietnam
544 (Hoang et al., 2013) and is effectively ongoing. A common view is that the late Miocene
545 increased siliciclastic sedimentation in basins along the margins of south-central Vietnam was a
546 response to margin surface uplift (Fyhn et al., 2009) and that this may have been linked to the
547 widespread basaltic magmatism (Carter et al., 2000). Hoang et al., (2013) observed that the large
548 magmatic centers in Vietnam formed in pull apart basins and that the pattern of volcanism had

549 evolved from SW–NE transtension to E–W extension linked to changes in the lithospheric stress
550 field due to either collision-related lithospheric response or an effect of asthenospheric flow.
551 Using a 1-D shear velocity model for the Indochina block, Yang et al. (2015) identified a low- V_s
552 anomaly below depths of 100 km along South-Central Vietnam and considered it likely reflects
553 active mantle flow in the asthenosphere beneath the region due to the stretching and thinning of
554 the lithosphere. The nature of flow is unclear, however, but could be related to edge-driven
555 convection as this has been used to explain intraplate volcanism and high topography in areas
556 where there is a significant gradient in lithosphere thickness (Kaislaniemi and Hunen, 2014).
557 Whilst the patterns and consequences of edge-driven convection on the continental side of a
558 passive margin has yet to be fully investigated, it is noted that local conditions for such flow are
559 suitable. The study area has an average crustal thickness between 29–33km (Bai et al., 2010; Vu
560 et al., 2021), thinner than global average values (~36–41 km; Szwillus et al., 2019) and the study
561 area lithospheric thickness is lower, between 110–120 km, compared to > 130km for northern
562 Vietnam (Vu et al., 2021). In relation to the apatite thermochronometry results early flow
563 associated with initial thinning and breakup would explain the interval of rapid cooling and
564 account for some of the margin uplift.

565
566 An alternative explanation to the above is related to low velocity anomalies detected in the lower
567 mantle below the study area (spanning depths between ~700–2889 km). Using multiscale global
568 tomography Zhao et al., (2021) interpreted the low velocity anomalies below southern Indochina
569 as hot mantle upwellings associated with a cluster of dying or already dead plumes that existed in
570 the Cenozoic. Basalt compositions across the study area are Ocean Island Basalt (OIB) mantle or
571 Enriched Mantle types (EM). An et al., (2016) argued that whilst there is large-ion lithophile
572 element enrichment without high field strength element depletion, typical of OIB, NiO -olivine
573 plots show compositions that fall within the range of Hawaiian and Hainan basalt olivines. As
574 these may be produced by partial melting of silica-poor eclogite and peridotite An et al., (2016)
575 favoured a Hainan plume source from recycled eclogitic oceanic crust. An alternative
576 explanation for basalt compositions is provided by Hoang et al., (2003) who highlighted the role
577 of active mantle flow in the asthenosphere by arguing that the late Cenozoic alkalic magmatism
578 in East Asia originated in the shallow ductile asthenosphere from adiabatic decompression
579 caused by pronounced changes in stress regimes linked to India - Eurasia collision and

580 subduction of the Pacific Plate. India - Eurasia collision and subduction clearly impacted on
581 mantle structure beneath Indochina as shown by a regional-scale receiver function study of the
582 mantle transition zone discontinuities (Yu et al., 2017). This work highlighted the potential role
583 of the subducted Indian slab and how broken segments sinking into the mantle transition zone
584 could have influenced mantle structure and dynamics beneath the Indochina Peninsula. Clearly
585 further work is needed to fully understand the nature of the magmatism and its relationship to
586 mantle structure.

587

588

589 **6. Conclusions**

590 The dense array of apatite thermochronometry results from across and along the margin of south-
591 central Vietnam revealed a regional episode of fast cooling between 37-30 Ma. This signal is
592 present within the coastal margins as well as inland. Consideration of the multiple factors that
593 affected the regional stress field at this time highlighted the coincidence between the episode of
594 fast cooling and the period of fast extension across the South China Sea (East Sea) region. This
595 shows a close relationship to rifting, however the episode of cooling might also be due to
596 transient changes in geothermal gradient. Thermal models explored the relationship between
597 cooling and exhumation. Models rule out thermal relaxation following Mesozoic arc magmatism
598 as geotherms returned to background rates 40 to 30 Myrs before the onset of fast cooling.
599 Instead, models suggest fast cooling could be attributed to accelerated erosion during early stages
600 of rifting, possibly with some additional heating from either underplating, and/or hot mantle
601 upwellings although the timing does not coincide with known periods of magmatism. No
602 evidence was found to connect regional uplift with the Miocene to Quaternary intraplate
603 magmatism. If this did occur the magnitude of associated surface uplift is beyond the resolution
604 of this study.

605

606 **Acknowledgements**

607 This work has been financially supported by the National Foundation for Science and
608 Technology Development of Vietnam – NAFOSTED (Grant No.: 105.99-2019.302). We thank
609 Birkbeck College, University of London, the London Geochronology Centre and Hanoi
610 University of Mining and Geology for access to their analytical and experimental facilities.

611 Our appreciation goes to Michael Fyhn and Lynne Elkins for their constructive reviews and
612 suggestions.

613

614 **Open Research**

615 **Data Availability Statement**

616 Data sets for this research are provided in Supporting Information Tables S1 to S3. Data are also
617 available in Earth and Space Science Open Archive doi.org/10.1002/essoar.10507473.1

618

619 **References**

- 620 An, A-R., Hi, S., Choi, S.H., Yu, Y., Lee, D-C. (2017) Petrogenesis of Late Cenozoic basaltic
621 rocks from southern Vietnam. *Lithos*, 272–273, 192–204.
- 622 Andersen, C., Mathiesen, A., Nielsen, L.H., Tiem, P.V., Petersen, H.I., Diem, P.T., (2005).
623 Evaluation of petroleum systems in the northern part of the Cainozoic Song Hong Basin (Gulf
624 of Tonkin), Vietnam. *Journal of Petroleum Geology* 28, 167–184
- 625 Bai, L., Iidaka, T., Kawakatsu, H., Morita, Y., Dzung, N.Q., (2009). Upper mantle anisotropy
626 beneath Indochina block and adjacent regions from shear-wave splitting analysis of Vietnam
627 broadband seismograph array data. *Phys. Earth Planet. Inter.* 176, 33–43.
- 628 Barckhausen, U., Engels, M., Franke, D., Ladage, S., Pubellier, M., (2014). Evolution of the
629 South China Sea: revised ages for breakup and seafloor spreading. *Mar. Pet. Geol.* 58, 599–
630 611. <http://dx.doi.org/10.1016/j.marpetgeo.2014.02.022>.
- 631 Carter, A., Roques, D. & Bristow, C. S. (2000). Denudation history of onshore Central Vietnam:
632 Constraints on the Cenozoic evolution of the western margin of the South China Sea.
633 *Tectonophysics* 322, 265-277.
- 634 Clift, P.D., Brune, S., Quinteros, J., (2015). Climate changes control offshore crustal structure at
635 South China Sea continental margin. *Earth Planet Sci. Lett.* 420, 66–72.
- 636 Clift, P.D., Wan, S. and Bluszrajn, J., (2014). Reconstructing chemical weathering, physical
637 erosion and monsoon intensity since 25 Ma in the northern South China Sea: a review of
638 competing proxies. *Earth-Science Reviews*, 130, 86–102.
- 639 Donelick R (1993) Apatite etching characteristics versus chemical composition. *Nucl Tracks*
640 *Radiat Meas* 21:604.

- 641 Doust, H and Sumner, S.H. (2007) Petroleum systems in rift basins – a collective approach in
642 Southeast Asian basins. *Petroleum Geoscience*, 13, 127 – 144.
- 643 Farnsworth, A., Lunt, D. J., Robinson, S. A., Valdes, P. J., Roberts, W. H. G., Clift, P. D.,
644 Markwick, P., Su, T., Wrobel, N., Bragg, F., Kelland, S-J., Pancost, R. D., (2019). Past East
645 Asian monsoon evolution controlled by paleogeography, not CO₂. *Science Advances*, 5,
646 eaax1697.
- 647 Fitzgerald P.G., Malusà M.G. (2019) Concept of the Exhumed Partial Annealing (Retention)
648 Zone and Age-Elevation Profiles in Thermochronology. In: Malusà M., Fitzgerald P. (eds)
649 Fission-Track Thermochronology and its Application to Geology. Springer Textbooks in
650 Earth Sciences, Geography and Environment. Springer, Cham. [https://doi.org/10.1007/978-3-](https://doi.org/10.1007/978-3-319-89421-8_9)
651 [319-89421-8_9](https://doi.org/10.1007/978-3-319-89421-8_9)
- 652 Flowers, R. M. R. A. Ketcham, D. L. Shuster, and K. A. Farley, (2009). Apatite (U–Th)/He
653 thermochronometry using a radiation damage accumulation and annealing model. *Geochim.*
654 *Cosmochim. Acta*, 73(8), 2347–2365.
- 655 Flower, M., Tamaki, K., Hoang, N., (1998). Mantle extrusion: a model for dispersed volcanism
656 and DUPAL-like asthenosphere in East Asia and the western Pacific. In: Flower, M., Chung,
657 S.-L., Lo, C.-H., Lee, T.Y. (Eds.), *Mantle Dynamics and Plate Interactions in East Asia*.
658 American Geophysical Union, Geophysical Monograph Series, 27, 67–88.
- 659 Fox, M. and Carter, A., (2020). Heated topics in Thermochronology and paths towards
660 resolution. *Geosciences*, 10(9), p.375.
- 661 Fyhn, M.B.W., Boldreel, L.O., Nielsen, L.H., (2009a). Geological development of the central
662 and south Vietnamese margin: implications for the establishment of the South China Sea,
663 indochinese escape tectonics and cenozoic volcanism. *Tectonophysics* 478, 184-214.
- 664 Fyhn, M. B. W., et al. (2009b), Geological evolution, regional perspectives and hydrocarbon
665 potential of the northwest Phu Khanh Basin, offshore Central Vietnam, *Mar. Pet. Geol.*, 26,
666 1–24, doi:10.1016/j.marpetgeo.2007.07.014
- 667 Fyhn, M.B.W., Pedersen, S.A.S., Boldreel, L.O., Nielsen, L.H., Green, P.F., Dien, P.T., Huyen,
668 L.T. and Frei, D., (2010) Palaeocene–early Eocene inversion of the Phuquoc–Kampot Som
669 Basin: SE Asian deformation associated with the suturing of Luconia. *J. Geol Soc. London*,
670 167, 281-296.

- 671 Gallagher, K., & Brown, R. (1997). The onshore record of passive margin evolution. *Journal of*
672 *the Geological Society*, 154(3), 451–457.
- 673 Gallagher, K. (2012). Transdimensional inverse thermal history modelling for quantitative
674 thermochronology. *Journal of Geophysical Research: Solid Earth*, 117(B2).
- 675 Hall, R., (2012). Late Jurassic-Cenozoic reconstructions of the Indonesian region and the Indian
676 Ocean. *Tectonophysics* 570, 1–41.
- 677 Hall, R., Spakman, W., (2015). Mantle Structure and Tectonic History of SE Asia.
678 *Tectonophysics*, 658: 14–45. <https://doi.org/10.1016/j.tecto.2015.07.003>.
- 679 Hennig-Breitfeld, J., Hennig-Breitfeld, H.T., Sang, D.Q., Vinh, M.K., Long, T.V., Thirwell, M.,
680 Cuong, T.X. (2021). Ages and character of igneous rocks of the Da Lat Zone in SE Vietnam
681 and adjacent offshore regions (Cuu Long and Nam Con Son basins). *J. Asian Earth Sciences*,
682 218, 104878.
- 683 Hiep, N. H., , Carter., A, Hoang, L. V., Fox, M., Pham, S.N., Bui, H.V. (2021) Evolution of the
684 continental margin of south to central Vietnam and its relationship to opening of the South
685 China Sea (East Vietnam Sea). *J Earth and Space Science Open Archive*, 24,
686 doi:10.1002/essoar.10507473.1
- 687 Hoang, N., Flower, M.F.J., 1998. Petrogenesis of Cenozoic basalts from Vietnam: implication
688 for origins of a ‘diffuse igneous province’. *Journal of Petrology* 39, 369–395.
- 689 Hoang, N., Flower, M.F.J., Chi, C.T., Xuan, P.T., Quy, H.V., Son, T.T., (2013). Collision-
690 induced basalt eruptions at Pleiku and Buon Me Thuot, south-central Viet Nam. *Journal of*
691 *Geodynamics* 69, 65–83.
- 692 Hurford, A.J., and Green, P.F., (1983). The zeta age calibration of fission track dating. *Chemical*
693 *Geology (Isotope Geoscience)*, 1, 285-317.
- 694 Jonell, T.N., Clift, P.D. Hoang, L.V., Hoang, T., Carter, A., Wittmann, H., Böning, P., and
695 Rittenour, T. (2017). Controls on Erosion Patterns and Sediment Transport in a Monsoonal,
696 Tectonically Quiescent Drainage, Song Gianh, Central Vietnam. *Basin Research*, 29, 659–
697 683.
- 698 Kaislaniemi, L., and J. van Hunen (2014), Dynamics of lithospheric thinning and mantle melting
699 by edge driven convection: Application to Moroccan Atlas mountains, *Geochem. Geophys.*
700 *Geosyst.*, 15, 3175–3189, doi:10.1002/2014GC005414.

- 701 Kincaid, C., Schilling, J. G. and Gable, C., (1996). The dynamics of off-axis plume-ridge
702 interaction in the uppermost mantle. *Earth and Planetary Science Letters* 137, 29–43.
- 703 Jolivet, L., Beyssac, O., Goffé, B., Avigad, D., Lepvrier, C., Maluski, H., & Thang, T. T. (2001).
704 Oligo-miocene midcrustal subhorizontal shear zone in Indochina. *Tectonics*, 20, 46–57.
- 705 Larsen et al., (2018). Rapid transition from continental breakup to igneous oceanic crust in the
706 South China Sea. *Nature Geoscience*, 11, 782-789.
- 707 Lee, G.H., Watkins, J.S., (1998). Seismic stratigraphy and hydrocarbon potential of the Phu
708 Khan Basin, offshore central Vietnam, south China Sea. *AAPG Bull.* 82, 1711-1735.
- 709 Lee, G.H., Lee, K., Watkins, J.S., (2001). Geological evolution of the Cuu Long and Nam Con
710 Son Basins, offshore southern Vietnam, South China Sea. *AAPG Bulletin* 85, 1055–1082.
- 711 Leloup, P. H., Arnaud, N., Lacassin, R., Kienast, J. R., Harrison, T. M., Phan Trong, T. T., et al.
712 (2001). New constraints on the structure, thermochronology, and timing of the Ailao Shan-
713 Red River shear zone, SE Asia. *Journal of Geophysical Research*, 106, 6683–6732.
- 714 Le Pourhiet, Laetitia, Chamot-Rooke, N., Delescluse, M., May, D. A., Watremez, L., Pubellier,
715 M. (2018). Continental break-up of the South China Sea stalled by far-field compression.
716 *Nature Geoscience*, 11, 605-609.
- 717 Li, C.-F., et al. (2014), Ages and magnetic structures of the South China Sea constrained by deep
718 tow magnetic surveys and IODP Expedition 349, *Geochem. Geophys. Geosyst.*, 15, 4958–
719 4983, doi:[10.1002/2014GC00556](https://doi.org/10.1002/2014GC00556)
- 720 Li, Y., Abbas, A., Li, C.F., Sun, T., Zlotnik, S., Song, T., Zhang, L., Yao, Z. and Yao, Y.,
721 (2020). Numerical modeling of failed rifts in the northern South China Sea margin:
722 Implications for continental rifting and breakup. *Journal of Asian Earth Sciences*, 199,
723 p.104402.
- 724 Lisker, F., Ventura, B., and Glasmacher, U.A. (2009) Apatite thermochronology in modern
725 geology. *Geological Society, London, Special Publications*, 324, 1-23,
726 <https://doi.org/10.1144/SP324.1>
- 727 Liu, J. L., Chen, X. Y., Tang, Y., Song, Z. J., & Wang, W. (2020). The Ailao Shan-Red River
728 shear zone revisited: Timing and tectonic implications. *GSA Bulletin*, 132(5–6), 1165–1182.
- 729 Matthews, S.J., Fraser, A.J., Lowe, S., Todd, S.P., Peel, F.J., (1997). Structure, stratigraphy and
730 petroleum geology of the SE Nam Con Son, offshore Vietnam. In: Fraser, A.J., Matthews,

- 731 S.J., Murphy, R.W. (Eds.), *Petroleum Geology of Southeast Asia*. Geological Society,
732 London, Special Publication, 126, pp. 89–106.
- 733 Mondy, L.S., Rey, P.F., Ducleaz, G., and Moresi, L. 2018. The role of asthenospheric flow
734 during rift propagation and breakup. *Geology*, 46, 103-106.
- 735 Morley, C.K., (2016). Major unconformities/termination of extension events and associated
736 surfaces in the South China Seas: review and implication for tectonic development. *J. Asian*
737 *Earth Sci.* 120, 62–86.
- 738 Morley, R.J., Dung, Bui Viet, Tung, Nguyen Thanh, Kullman, R.J., Bird, R., Van Kieu, Nguyen,
739 Chung, Nguyen Hoai, (2019). High-resolution Palaeogene sequence stratigraphic framework
740 for the Cuu Long Basin, offshore Vietnam, driven by climate change and tectonics,
741 established from sequence biostratigraphy. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 530,
742 113-135.
- 743 Murray, K. E., Braun, J., & Reiners, P. W. (2018). Toward robust interpretation of low-
744 temperature thermochronometers in magmatic terranes. *Geochemistry, Geophysics,*
745 *Geosystems*, 19, 3739–3763. <https://doi.org/10.1029/2018GC007595>.
- 746 Nguyen, V.V. and Hoai, L.T.T. (2019). Cenozoic paleostress evolution in south central Vietnam:
747 Implication for changing dynamics of faulting along the eastern Indochina continental margin.
748 *Journal of Asian Earth Sciences* 185, 104006.
- 749 Nguyen, D.H., Le, V.H., (2004). Hydrocarbon geology of Cuu Long basin, offshore Vietnam. In:
750 *Proceedings of AAPG International Conference and Exhibition*. Search and Discovery Article
751 #10062.
- 752 Nielsen, L.H., Petersen, H.I., Thai, N.D., Duc, N.A., Fyhn, M.B.W., Boldreel, L.O., Tuan, H.A.,
753 Lindstrom, S., Hien, L.V. (2007) A Middle–Upper Miocene fluvial–lacustrine rift sequence in
754 the Song Ba Rift, Vietnam: an analogue to oil-prone, small-scale continental rift basins.
755 *Petroleum Geosciences*, 13, 145-168.
- 756 Pham, V.N., (2003). *Bien Dong Monograph, Vol. II—Meteorology*. Hanoi National University
757 Publisher, Hanoi, 565 pp. (in Vietnamese).
- 758 Pubellier, M., Rangin, C., Le Pichon, X., Dotsea Working group, (2005). Dotsea: a synthesis of
759 deep marine data in Southeast Asia. *Memoires de la Société Géologique de France*, vol. 176,
760 pp. 1–32.

- 761 Rangin, C., Huchon, P., Le Pichon, X., Bellon, H., Lepvrier, C., Roques, D., Hoe, N.D., Quynh,
762 P.V., (1995). Cenozoic deformation of Central and South Vietnam. *Tectonophysics* 235, 179–
763 196.
- 764 Rey, P., (2001), From lithospheric thickening and divergent collapse to active continental rifting,
765 in Miller, J.A., et al., eds., *Continental Reactivation and Reworking: Geological Society of*
766 *London Special Publication* 184, 77–88.
- 767 Roques, D., Matthews, S.J., Rangin, C., (1997). Constraints on strike-slip motion from seismic
768 and gravity data along the Vietnam margin offshore Da Nang: implications for hydrocarbon
769 prospectivity and opening of the East Vietnam Sea. In: Fraser, A.J., Matthews, S.J., Murphy,
770 R.W. (Eds.), *Petroleum geology of Southeast Asia*, Geological Society, London, Special
771 Publication, 126, pp. 341–353.
- 772 Savva, D., Pubellier, M., Franke, D., Chamot-Rooke, N., Meresse, F., Steuer, S., Auxietre, J.L.,
773 (2014). Different Expressions of Rifting on the South China Sea Margins *Mar. Pet. Geol.* 58.
774 pp. 579e598d.
- 775 Schmidt, W.J., Bui, H.H., Handschy, J.W., Vu, T.H., Trinh, X.C., Nguyen, T.T., (2019).
776 Tectonic evolution and regional setting of the Cuu Long Basin, Vietnam. *Tectonophysics* 757,
777 36–57.
- 778 Searle, M.P. (2006). Role of the Red River Shear zone, Yunnan and Vietnam, in the continental
779 extrusion of SE Asia: *Journal of the Geological Society*, 163, 1025–1036, doi: 10.1144/0016-
780 76492005-144.
- 781 Searle, M. P., Yeh, M. W., Lin, T. H., & Chung, S. L. (2010). Structural constraints on the
782 timing of left-lateral shear along the Red River shear zone in the Ailao Shan and Diancang
783 Shan Ranges, Yunnan, SW China. *Geosphere*, 6, 316–338.
- 784 Shellnutt, J.G., Lan, C-Y., Long, T. V., Usuki, T., Yang, H-J., Mertzman, S.A., Lizuka, Y.,
785 Chung, S-L., Wang, K-L., Huse, W-Y., (2013). Formation of Cretaceous Cordilleran and
786 post-orogenic granites and their microgranular enclaves from the Dalat zone, southern
787 Vietnam: Tectonic implications for the evolution of Southeast Asia. *Lithos*, 182-183, 229-
788 241.
- 789 Sibouet, J-C., Yeh, Y-C., Lee, C-S. (2016). Geodynamics of the South China Sea.
790 *Tectonophysics*, 692, 98-119.

- 791 Szwillus, W., Afonso, J. C. C., Ebbing, J., & Mooney, W. D. (2019). Global crustal thickness
792 and velocity structure from geostatistical analysis of seismic data. *Journal of Geophysical*
793 *Research: Solid Earth*, 124, 1626–1652. <https://doi.org/10.1029/2018JB016593>
- 794 Taylor, B., Hayes, D.E., (1983). Origin and history of the South China Sea Basin. In: Hayes,
795 D.E. (Ed.), *The Tectonic and Geologic Evolution of Southeast Asian Seas and Islands. Part 2.*
796 *Geophysical Monograph*, 27. AGU, Washington, D.C., pp. 23–56
- 797 Thuy, N.T.B., Muharren, S., Wolfgang, S., Fukun, C, (2004). Granitoids in the Dalat zone,
798 Southern Vietnam: age constraints on magmatism and regional geological implications.
799 *International Journal of Earth Science* 93, 329-340.
- 800 Tran V.T., and Vu, K., (Eds.), (2011). *Geology and Earth Resources of Vietnam: General*
801 *Department. of Geology, and Minerals of Vietnam.* Hanoi, Publishing House for Science and
802 *Technology*, 634 pp.
- 803 Turcotte, D. L., & Schubert, G. (2002). *Geodynamics* (2nd ed.). New York: Cambridge
804 *University Press.* [https://doi.org/10.1017/](https://doi.org/10.1017/CBO9780511807442)
805 [CBO9780511807442](https://doi.org/10.1017/CBO9780511807442)
- 806 Vu, A.T., Fyhn, M.B.W., Xuan, C.T., Nguyen, T.T., Hoag, D.N., Pham, L.T., Van, H.N. (2017),
807 Cenozoic tectonic and stratigraphic development of the Central Vietnamese continental
808 margin. *Marine and Petroleum Geology*, 86, 386-410.
- 809 Vue, D.T., Bonvalot, S., Bruinsma, S., Bui, L.K. (2021), A local lithospheric structure model for
810 Vietnam derived from a high-resolution gravimetric geoid. *Earth, Planets and Space*, 73, 92.
811 [doi:10.1186/s40623-021-01415-2.](https://doi.org/10.1186/s40623-021-01415-2)
- 812 Wildman M., Cogné N., Beucher R. (2019), Fission-Track Thermochronology Applied to the
813 Evolution of Passive Continental Margins. In: Malusà M., Fitzgerald P. (eds) *Fission-Track*
814 *Thermochronology and its Application to Geology.* Springer Textbooks in Earth Sciences,
815 *Geography and Environment.* Springer, Cham. https://doi.org/10.1007/978-3-319-89421-8_20
- 816 Yan, Q., Shi, X., Metcalfe, I. *et al.* (2018). Hainan mantle plume produced late Cenozoic basaltic
817 rocks in Thailand, Southeast Asia. *Sci Rep* 8, 2640. [https://doi.org/10.1038/s41598-018-](https://doi.org/10.1038/s41598-018-20712-7)
818 [20712-7.](https://doi.org/10.1038/s41598-018-20712-7)
- 819 Yu, Y., S. S. Gao, K. H. Liu, T. Yang, M. Xue, and K. Phon Le (2017), Mantle transition zone
820 discontinuities beneath the Indochina Peninsula: Implications for slab subduction and mantle
821 upwelling, *Geophys. Res. Lett.*, 44, 7159–7167, [doi:10.1002/2017GL073528](https://doi.org/10.1002/2017GL073528)

822 Zhao, D., Toyokuni, G., and Kurata, K., (2021). Deep mantle structure and origin of Cenozoic
823 intraplate volcanoes in Indochina, Hainan and South China Sea. *Geophys. J. Int.* 225, 572–
824 588.

825 Zhu, B.-Q., Wang, H.-F., Chen, Y.-W., Chang, X.-Y., Hu, Y.-G., Xie, J. (2004),
826 Geochronological and geochemical constraint on the Cenozoic extension of Cathaysian
827 lithosphere and tectonic evolution of the border sea basins in East Asia. *J. Asian Earth Sci.* 24
828 (2), 163-175.

829

830

831

832 **Figure Captions**

833

834 **Fig. 1.** Location of the study area that compliments an earlier study of the Kontum region (Carter
835 et al., 2000).

836

837 **Fig. 2.** Seismic profile across the continental margin adjacent to the study area (Vietnam
838 Petroleum Institute). A distinct Oligocene unconformity records basin inversion and a mid
839 Miocene unconformity records the end of rifting after which sediment supply increased leading
840 to progradation and slope mass transport deposits.

841

842 **Fig. 3.** Sample locations and regional geology modified from Tran, et al., (1988).

843

844 **Fig. 4.** Plot comparing AFT and AHe (corrected) replicate ages against sample elevation.
845 Because the samples are from a range of locations with different exhumation rate histories, the
846 slope of this relationship will not approximate the exhumation rate. In addition, elevation varies
847 with distance from the coast and so this variability also masks obvious trends. Error bars are 1s.

848

849 **Fig. 5.** AFT and AHe (corrected) ages for a suite of samples collected across a 1421m elevation
850 range within a 92 Ma granite. Ages above 800m record slower exhumation than samples at lower
851 elevation. Key to geological units given in **Fig. 3.**

852

853 **Fig. 6.** Thermal history model of the vertical profile shown in Fig. 5. Dark column denotes the
854 circa 37-30 Ma onset of fast cooling recorded by the best-fit models. Solid lines are best fit
855 models. The credible intervals are shown (dashed lines) are shown only for the lowermost and
856 uppermost samples. Plot 6b shows how well the models fit the measured data by plotting the
857 predicted and observed values as a function of elevation. Error bars include the mean 95%
858 credible range for the predictions from all thermal history models accepted during the post-
859 burnin MCMC sampling.

860

861 **Fig. 7.** A simple 1D model is used to assess the effects of heating at depth on the time-
862 temperature paths experienced by thermochronometry samples. In this cartoon, the initial
863 condition is shown by the dotted line. Here temperature increases linearly with depth and then
864 there is an increase in temperature associated with the magma (1). Temperatures remain high to
865 simulate an active magma chamber. This leads to heating above the magma chamber (2). After a
866 set amount of time, temperatures at the base of the model return to a temperature defined by the
867 initial geothermal gradient (3). Temperatures remain high across the upper part of the model (4).
868 Over time temperature return to a steady state thermal model determined by the boundary
869 conditions and exhumation rate. The inset shows the evolution of temperature a material point
870 experiences as it is exhumed towards the surface. Initially there is heating to the elevation
871 geotherm (EG) and then temperatures decrease to the steady state geotherm (SSG).

872

873 **Fig. 8.** The duration of heating associated with magma intrusion. A magma intrusion is simulated
874 between 8-12 km depth. Temperatures remain high for 2 million years before the basal boundary
875 condition at 35 km is forced to 800C. The exhumation rate remains constant at 0.1 km/Ma. The
876 grey curves show the evolving thermal model over the first 10 million years of the simulated
877 history. After approximately 20 million years the steady state solution is reached and
878 temperature as a function of depth remains constant. The red line shows the temperature-depth
879 paths rocks that reach the surface took. The circles show the closure temperatures and depths
880 evaluated using Dodson's approximation. B) This same path can be plotted as a function of
881 temperature and age. This highlights the rapid cooling from temperatures of 1500°C to
882 temperatures of about 250°C over the first 20 million years of the simulated history. For

883 reference, a time-temperature path from a thermal model with no heating is shown by the blue
884 dashed line.

885

886 **Fig. 9:** Thermal models for three different thermokinematic scenarios used to predict the
887 evolving thermal field: Dotted line (scenario 1) is where the cooling was interpreted in terms of
888 exhumation, with rates of 0.6 km/Ma between 37.5 and 30 Ma and a background rate of 0.02
889 km/Ma. The solid line (scenario 2) approximates a geomorphic response to rock uplift where
890 exhumation began at 40 Ma at rates of 0.75 km/Ma but then decreased exponentially from 37 Ma
891 to the present background rate of 0.01 km/Ma. The dashed line (scenario 3) represents a pulse of
892 heating associated with emplacement of underplating material below the lower crust and
893 highlights how a pulse of heating within the lower crust could also produce a pulse of cooling.
894 Blue shaded areas show the interval of rapid cooling indicated by QTQt models (**Fig. 6**). All
895 curves show increased cooling rates between about 35 and 25 Ma. Scenario 3 also shows a
896 reheating event.

897

898 **Fig. 10.** Principal features of the marine basins adjacent to the study area (Fyhn et al., 2009;
899 Schmidt et al., 2019; Vu et al., 2017; Morely et al., 2019) compared to the South China Sea
900 spreading history (Larsen et al., 2018; Le Pourhiet et al., 2017), exhumation data and volcanism
901 across the Indochina margin (Hoang et al., 2013).

902

















