



Contents lists available at ScienceDirect

## Contact Lens and Anterior Eye

journal homepage: [www.elsevier.com/locate/clae](http://www.elsevier.com/locate/clae)

## International multi-centre study of potential benefits of ultraviolet radiation protection using contact lenses

James S Wolffsohn<sup>a,\*</sup>, Sandeep Dhallu<sup>a</sup>, Maana Aujla<sup>a</sup>, Debbie Laughton<sup>a</sup>, Keith Tempny<sup>b</sup>, Daniel Powell<sup>c,d</sup>, Kate Gifford<sup>e</sup>, Paul Gifford<sup>f</sup>, Kin Wan<sup>g</sup>, Pauline Cho<sup>g</sup>, Ulrike Stahl<sup>h</sup>, Jill Woods<sup>h</sup>

<sup>a</sup> Aston University, Birmingham, UK

<sup>b</sup> Leightons and Tempny, Poole, UK

<sup>c</sup> University of Houston, Texas, USA

<sup>d</sup> Southern College of Optometry, Memphis, TN, USA

<sup>e</sup> Queensland University of Technology, Brisbane, Australia

<sup>f</sup> University of New South Wales, Sydney, Australia

<sup>g</sup> Hong Kong Polytechnic University, Kowloon, Hong Kong

<sup>h</sup> University of Waterloo, Ontario, Canada

## ARTICLE INFO

## Keywords:

Ultraviolet radiation  
Contact lenses  
Macular degeneration  
Presbyopia  
Accommodation  
Ocular surface

## ABSTRACT

**Purpose:** To examine the effects of long-term ultraviolet radiation (UVR) blocking wearing contact lenses on ocular surface health, eye focus and macular pigment.

**Method:** 210 pre-presbyopic patients were recruited from Birmingham UK, Brisbane Australia, Hong Kong China, Houston USA and Waterloo Canada (n = 42 at each site). All patients had worn contact lenses for ≥ 5 years, half (test group) of a material incorporating a UVR-blocking filter. Ocular health was assessed using slit-lamp biomicroscopy and UV autofluorescence. Accommodation was measured subjectively with a push-up test and overcoming lens-induced defocus. Objective stimulus response and dynamic measures of the accommodative response were quantified with an open-field aberrometer. Macular pigment optical density (MPOD) was assessed using heterochromatic flicker photometry (MPS II).

**Results:** The two groups of participants were matched for age, sex, race, body-mass-index, diet, lifestyle, UVR exposure, refractive error and visual acuity. Limbal (p = 0.035), but not bulbar conjunctival redness (p = 0.903) was lower in eyes that had worn UVR-blocking contact lenses compared to controls. The subjective (8.0 ± 3.7D vs 7.3 ± 3.3D; p = 0.125) and objective (F = 1.255, p = 0.285) accommodative response was higher in the test group, but the differences did not reach significance. However, the accommodative latency was shorter in eyes that had worn UVR-blocking contact lenses (p = 0.003). There was no significant difference in MPOD with UVR filtration (p = 0.869).

**Conclusions:** Blocking the transmission of UVR is beneficial in maintaining the eye's ability to focus, suggesting that presbyopia maybe delayed in long-term UVR-blocking contact lenses wearers. These lenses also provide protection to the critical limbal region.

### 1. Introduction

The risk of damage to the skin from ultraviolet radiation (UVR) exposure is well known and there is strong evidence that blocking is beneficial [1,2]. However, there is comparatively little evidence on the ocular benefits of UVR shielding. While exposure to UVR in small quantities can be beneficial to human health in some instances, such as

in synthesis of vitamin D [3], there are many associated health risks from both acute and chronic exposure. While the anterior ocular structures absorb most UVR [4], research into the penetration to internal ocular structures is limited. Pterygium, a wing shaped thickening of the conjunctiva and cornea, has been shown to be related to prolonged UVR exposure [5,6]. Ascorbate is an antioxidant in the aqueous humour which has protective effects on the crystalline lens. Levels of ascorbate

\* Corresponding author at: School of Optometry, Aston University, Aston Triangle, Birmingham B47ET, UK.

E-mail address: [j.s.w.wolffsohn@aston.ac.uk](mailto:j.s.w.wolffsohn@aston.ac.uk) (J.S. Wolffsohn).

<https://doi.org/10.1016/j.clae.2022.101593>

Received 23 December 2021; Received in revised form 25 March 2022; Accepted 29 March 2022

1367-0484/© 2022 The Author(s). Published by Elsevier Ltd on behalf of British Contact Lens Association. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

can be affected by extended UVR exposure and may have implications in cataractogenesis [7]. There is some controversy regarding the link between sun exposure and age-related macular degeneration (AMD). Although it has been suggested that UVR is a risk factor in AMD, epidemiological studies have not shown a significant association [8-12].

Soft contact lenses with a UVR filter can offer protection to the cornea, limbus, some of the bulbar conjunctiva and internal ocular structures because of their large diameter, movement with the eye and position against the ocular surface. Incorporating a UVR-blocker into a soft contact lens has been shown to help protect the eye from the peripheral light focusing effect at the nasal limbus, with protection given from all angles of light incidence. Hence, UVR-blocking soft lenses may help reduce the prevalence of ocular conditions, such as pterygium and early cortical cataract [13]. Stem cells are important in the maintenance of tissue integrity and as a result tend to be found in protected areas in the body. Following injury to the corneal epithelium, limbal stem cell division is upregulated in order to replace lost cells [14]. It is thought that the limbal shielding afforded by contact lenses can help to protect the repositories of limbal stem cells from UVR damage, especially from tangential and temporal sources of UVR rays [15]. In a recent study, UVR-blocking contact lenses were found to have a protective effect in preventing short-term UV-B-induced limbal stem cell damage and inflammation compared to lenses without a UVR-blocker, and it was suggested that these lenses may be particularly useful in certain situations, such as after limbal stem cell transplantation or following pterygium surgery [16].

Exposure to UV-B causes cell death in the cornea and the use of UVR-blocking contact lenses have been shown to reduce levels of cell death compared to a minimal UVR-blocking contact lenses [7]. In addition, changes in epithelial, stromal and endothelial cells have been observed between eyes wearing standard hydrogel lenses and those wearing UVR-blocking lenses. [17] Trauma to keratocytes and endothelial cells can result in permanent damage to the cornea [18,19] thus UVR protection is imperative.

The level of protection given by a soft contact lens is dependent upon the transmittance characteristics of the lens material, with UVR absorption levels varying between brands [20]. Contact lens light transmittance is reported to change with contact lens wear and this is most likely due to the formation of biofilms on the lens surface, however this was not found to affect the UVR-blocking properties of the lens [21]. The level of UVR protection provided by a contact lens has been shown to be dependent upon its power, and therefore centre thickness [22]. Class I lenses must block at least 90% of UV-A and at least 99% of UV-B, and class II must block at least 50% of UV-A and at least 95% of UV-B [23].

Although there is strong evidence that shielding the skin from UVR is beneficial, there is little direct evidence for the eyes. This multi-site study examined the effects of wearing UVR-blocking contact lenses for at least five years on eye health, ocular accommodation, and macular pigment across five developed countries.

## 2. Methods

The study was approved by the Research Ethics Committees in each site location and conformed to the tenets of the Declaration of Helsinki. The variability (95% confidence interval) of the push up test is  $\pm 1.43$  D [24], so to detect a 0.6 D change (80% power, 0.05 significance level) a minimum of 20 participants in each study arm for non-parametric comparison (G\*Power v3.1.9.4) was required. The variability of defocus curves is not known, but the variability of subjective acuity measurement at each level of defocus is  $\pm 0.08$  logMAR [25,26], so to detect a 2 letter (0.04 logMAR) change would require at least 14 participants. The variability of Aston aberrometer measures of accommodation has been shown to be 0.2 D [27], so to detect a 0.10 D difference required at least 14 participants. The variability of MPS 900 macular pigment densitometry has been shown to be 0.04 [28] so to detect a 0.02 MPOD change would require 14 participants. Therefore 21 participants were

enrolled for each group at each site to allow for some invalid data. Once written informed consent had been obtained after explanation of the nature and possible consequences of the study, 42 participants aged 18 to 50 years were assessed at each of the five sites across the world (Birmingham (United Kingdom), Houston (USA), Brisbane (Australia), Waterloo (Canada) and Hong Kong SAR (China);  $n = 210$  in total). Recruitment was stratified with 7 participants in each of the 18–28 year old, 29–39 year old and 40–50 year old age ranges in both the UVR and minimal UVR-blocking contact lens groups.

The average temperature and UVR indices of each region are represented in Table 1. The UVR index is a standardised way of measuring the strength of UVR reaching the earth's surface at any given time and location. It was first released in 1995, with an updated version published in 2002 [28]. The index is based on the thickness of the ozone layer in the upper atmosphere, the angle of the sun and the amount of cloud cover. The scale ranges from lows of zero to highs of 11 or more in tropical countries, and can be grouped into five levels of risk: low (0–2), moderate (3–5), high (6–7), very high (8–10) and extreme (11 +) [29].

Participants were required to have worn contact lenses for a minimum of five days per week for the previous five years or more and to have been a resident of that country during the same period. Additionally, participants were required to have a best corrected visual acuity of 6/7.5 or better in each eye, have no previous history of eye surgery and not to be on any medication known to affect accommodation. Twenty-one participants from each site ( $n = 105$  in total) had worn contact lens materials with UVR-blocking for at least five years and the 21 controls ( $n = 105$  in total) had worn a contact lens material with minimal UVR-blocking properties for the same period. Participants reported the brand of their contact lenses; these were checked against clinical records where available and the UVR-blocking properties against published spectral profiles. Investigators were masked to the participant's contact lens history.

Participant demographics of age, sex, ethnicity, height and weight (to calculate body mass index = height  $\times$  weight (BMI)), general health (including smoking) and medication were recorded. Bulbar conjunctival and limbal redness were graded through a slit lamp biomicroscope using an Efron scale to one decimal place from 0.0 to 4.0, and the presence of any pinguecula and pterygium noted.

All research data was collected by the same researchers at all sites, on the same equipment. Ocular accommodation was assessed subjectively whilst wearing prescription optimized contact lenses by averaging three repeats of the push-up test [30], as well as from minus-to-blur testing: increasingly minus powered lenses were held in front of the participants distance corrected eye monocularly, as they viewed a distance letter chart. Participants attempted to minimise the resulting defocus by accommodating, reading the smallest letters they could see (randomized between lenses), so that the dioptric power at which they could no longer hold their best visual acuity line in clear focus (allowing for lens minification) could be identified [31,32]. Objective stimulus response and dynamic measures of the accommodative response were quantified with an open-field aberrometer [27]. The stimulus response curve for accommodation and pupil size was measured using a high contrast Maltese cross at 0.0D, 0.5D, 1.0D, 2.0D, 3.0D, 4.0D and 5.0D of accommodative demand (in a Badal optical system). The dynamics of accommodation were assessed by rotating a Maltese cross at 3.0D accommodative stimulus into and out of the line of sight of a second Maltese cross positioned at optical infinity, at 5 s intervals (servo motor transition time < 0.05 s). The 25 Hz dynamic recording of the Aston aberrometer [27] was captured by a personal computer which also controlled the stepper motor rotating the Maltese cross, so the speed of accommodation, response amplitude, latency and speed of disaccommodation could be evaluated from three repeated accommodation and relaxation cycles.

Macular pigment optical density (MPOD) was determined using the MPS II 9000 (Elektron eye technology, Cambridge, UK) that employs the psychophysical technique of heterochromatic flicker photometry [33].

**Table 1** Average maximum summer month temperature, monthly Ultraviolet Radiation index and sunshine hours by data collection region. <https://www.weather-atlas.com> accessed March 2022.

City, Country	Average maximum summer temperature (°C)	Ultraviolet Radiation Index												
		AVERAGE	January	February	March	April	May	June	July	August	September	October	November	December
Birmingham, UK	21.8	3.0	2	2	3	3	4	4	4	4	3	3	2	2
Houston, USA	34.4	5.5	3.9	3.6	5.4	6.8	6.7	7.2	6.9	6.4	4.1	4.1	3.7	4.1
Brisbane, Australia	29.1	7.7	5.3	5.8	6.8	8.3	10.4	10.4	10.5	8.6	6.9	6.9	5.6	4.8
Waterloo, Canada	21.0	3.1	1	1	2	3	5	5	5	5	6	6	6	6
Hong Kong SAR, China	31.4	10.1	2.5	3.0	5.2	6.9	8.7	9.4	9.7	8.3	4.5	4.5	4.1	2.6
		5.0	4.6	3.4	2.9	3.4	4.5	4.9	6.8	5.7	6.3	6.0	6.0	5.6

The test eye was fully corrected with full aperture trial lenses position in a trial frame to avoid any bias from the light transmission of the contact lens, with the fellow eye occluded to avoid distraction.

UVR autofluorescence images were captured of the nasal and temporal conjunctiva of the right eye using a custom-made attachment mounted to an iPhone camera (4032 × 3024 pixels resolution) incorporating a macro-lens to give 6x magnification. This bespoke device uses a 365 nm wavelength diode positioned 3 cm from the eye that produces non-collimated light to highly damaged cells on the ocular surface [34]. The edges of regions with conjunctival fluorescence were subjectively outlined using ImageJ software (<http://rsbweb.nih.gov/ij/>) to give an area in pixels. An object of known size was also imaged using the same camera system to obtain a pixel to mm conversion factor from which the area in square millimetres was ascertained.

Participants were asked to complete a lifestyle questionnaire related to their demographics (age and sex), refractive correction wear, lifestyle (self-reported as ‘sun-avoider’, ‘average sun exposure’ or ‘sun-worshipper’, use of sunglasses – worn most of the time outdoors, worn only when sunny, worn sometimes or never worn) and contact lens history (number of years lenses worn, current brand and years worn) [34]. Additionally, participants completed a validated food frequency questionnaire which collected information about average frequency (checking 1 of 9 frequency categories, ranging from “never or less than once per month” to “two or more times per day) and serving sizes (small, medium, or large) of consumption over the previous year. The selected frequency categories and serving sizes for each food item were converted into an average daily intake of medium servings and the foods items combined into food groups to reduce inter-person variation [35].

**3. Statistics**

Due to the association between right and left eyes, only data from right eyes were included in the analysis in order to avoid statistical bias. Amplitude of accommodation, minus lens to blur, accommodative dynamics and macular pigment were not significantly different from a normal distribution (one-sample Kolmogorov–Smirnov test  $p > 0.05$ ) and therefore analysis of variance (with Tukey post-hoc comparison) was conducted for site and material UVR comparison were used. For redness grading and autofluorescence area, independent sample Mann-Whitney U for lens comparison and Kruskal-Wallis for site and material UVR comparison were used. Only two autofluorescence images of the 420 captured (0.5%) could not be analysed due to low quality. Statistical significance was set as  $p \leq 0.05$ .

**4. Results**

The cohorts of participants who had worn UVR-blocking or minimal UVR-blocking contact lenses were matched for age, sex, race, body-mass-index, diet, lifestyle, refractive error and visual acuity (Table 2). Reported sun exposure differed between sites (all were ‘average’, except Hong Kong, where the median was ‘sun avoider’;  $p < 0.001$ ). Use of sunglasses was slightly lower in the minimal UVR-blocking cohort ( $p = 0.050$ ) and differed with site (all were ‘only worn when sunny’, except Hong Kong, where the median was ‘never worn’;  $p < 0.001$ ).

There was no statistically significant difference in bulbar conjunctival redness ( $p = 0.903$ ) in eyes that had worn UVR-blocking contact lenses (1.2 median, 0.5–2.9 range) compared to the controls (1.3 median; 0.2–2.3 range), but the median of the USA and Hong Kong cohorts was double that of Canada ( $p < 0.001$ ; Table 3). Limbal redness was statistically significantly lower ( $p = 0.035$ ) in eyes that had worn UVR-blocking contact lenses compared to the controls; the median was higher in the UK, USA and Hong Kong compared to Canada and Australia ( $p < 0.001$ ; Table 3). Pingueculae (UVR: 19 eyes versus minimal UVR-blocking contact lenses: 20 eyes;  $p = 0.859$ ) or pterygia (UVR: 4 eyes versus minimal UVR-blocking contact lenses: 5 eyes;  $p = 0.733$ ) did not differ in proportion between eyes that had worn UVR-blocking contact

**Table 2**

Comparison (mean  $\pm$  standard deviation) of age, sex, body mass index, mean spherical equivalent refraction, best corrected visual acuity, servings of fruit/vegetables, smokers, use of sunglasses and typical sun exposure between the UVR-blocking (n = 105) and minimal UVR-blocking (n = 105) cohorts. \*Independent-sample Mann-Whitney-U test. <sup>†</sup>Chi-squared test.

	UVR-blocking lens wearers	Minimal UVR-blocking lens wearers	Significance
Age (years)	31.9 $\pm$ 9.1	31.6 $\pm$ 9.2	p = 0.899
Sex (male/female)	23 / 82	26 / 79	p = 0.625 <sup>†</sup>
Body Mass Index	23.6 $\pm$ 5.0	23.3 $\pm$ 5.6	p = 0.746
Mean Spherical Equivalent Refraction (D)	-0.28 $\pm$ 0.85	-0.37 $\pm$ 0.89	p = 0.479
Best Corrected Visual Acuity (logMAR)	-0.04 $\pm$ 0.09	-0.02 $\pm$ 0.09	p = 0.182
Servings of fruit/vegetables	6.3 $\pm$ 7.0	5.3 $\pm$ 3.4	p = 0.082
Smokers	2	4	p = 0.407 <sup>†</sup>
Use of sunglasses (1 = most of time, 4 = never)	2.2 $\pm$ 1.0	2.5 $\pm$ 1.1	p = 0.050*
Typical sun exposure (1 = sun avoider, 3 = sun worshipper)	1.9 $\pm$ 0.5	1.8 $\pm$ 0.6	p = 0.270*

lenses compared to the controls.

Amplitude of accommodation push-up test results were higher, but not statistically significantly, in eyes that had worn UVR-blocking contact lenses (8.0  $\pm$  3.7D) versus those that had worn minimal UVR-blocking contact lenses (7.3  $\pm$  3.3D; p = 0.125); it was lower in Hong Kong compared to the other sites (p < 0.001; Table 3). The maximum negative powered lens that distance targets could be resolved through without debilitating blur was lower than the push-up amplitude of accommodation, but followed a similar pattern, with no significant difference with the UVR-blocking of the contact lenses worn (p = 0.758), but again was lower in Hong Kong compared to other sites (p = 0.050; Table 3).

Accommodative response increased with accommodative demand (F = 230.289, p < 0.001)(Fig. 1). There was no statistically significant difference in the stimulus response between UVR and minimal UVR-blocking lenses (F = 1.255, p = 0.285) or countries (F = 1.524, p = 0.066).

Accommodative latency was shorter in eyes that had worn UVR-blocking contact lenses compared to eye that had worn lenses with minimal UVR-blocking (p = 0.003). Accommodative latency was shorter in eyes from the USA than all the other countries except Australia (p < 0.001). The speed of accommodation and step size was higher in eyes from the USA than the UK (p = 0.014) and the step size was lower in Australia than the USA or Canada (both p = 0.002, Table 3).

There was no significant difference in UV autofluorescence (nasal: p = 0.786; temporal: p=0.639) and MPOD (p = 0.869) between eyes that had worn UVR-blocking contact lenses compared to those that had worn contact lenses with minimal UVR-blocking contact lenses (Table 3). However, there was a significant difference among countries (p = 0.031), with wearers in the UK having a lower macular pigment density than those in Hong Kong (p = 0.050).

## 5. Discussion

There are many associated health risks to both the skin and the eye from exposure to UVR. Soft contact lenses with a UVR filter can offer protection to the eye and its structures because of their relatively large diameters and the fact that they move with the eye, ensuring continued protection. There is a strong evidence base which supports shielding as a form of protection from UVR for the skin, however there is comparatively little evidence for the eyes. This multi-site study examined the retrospective effects of wearing UVR-blocking contact lenses on ocular health, accommodation, and macular pigment across five developed countries, with varying temperatures and levels of UVR exposure. The

cohorts of participants who had worn UVR-blocking or minimal UVR-blocking contact lenses for at least the past five years were matched for age, sex, race, body-mass-index, diet, lifestyle, refractive error and visual acuity. Interestingly in Hong Kong SAR, China, individuals tend to report being a 'sun avoider' (culturally fair coloured skin is preferred) and perhaps as a result, were more likely to report sunglasses are 'never worn'. However, in other sites, even those with high UVR indices such as Brisbane Australia and Houston USA (Table 1), with most individuals reporting 'average' sun exposure, sunglasses are typically worn 'only worn when sunny', risking damage from UVR penetration through clouds.

Soft contact lenses cover the cornea and limbus, with minimal coverage of the conjunctiva. However, compared to light filtering spectacles or sunglasses, they are in contact with the ocular surface and move with the eyes and can therefore block peripheral light rays entering through the cornea. Hence, the finding that they reduced limbal, but not conjunctival redness was not surprising. Likewise the higher levels of bulbar and limbal conjunctival redness in Houston USA and Hong Kong SAR China cohorts could reflect the higher temperatures and UVR indices in these regions, but this did not account for the higher level of limbal redness in the UK cohort that was less exposed to these environmental conditions, nor the low levels of redness in Brisbane Australia, where more harsh environmental conditions are common. UV autofluorescence of the conjunctiva was also not found to differ with UVR blocking of contact lenses in this study, perhaps due to contact lenses not offering much coverage of this area, although protection has been found in other studies [36]. Other factors such as public health messaging around sun exposure in some regions are likely to play a role in mitigating environmental effects.

Although UVR is recognized as a risk factor for cataract, very little is known about its effect on presbyopia [37]. The comparison of amplitude of accommodation and stimulus response curve did not show a statistically significant difference between participants who had worn UVR-blocking compared to those wearing minimal UVR-blocking contact lenses for the past five years or more. The effect size may have been reduced by the age range of the participants, with some older individuals demonstrating minimal objective accommodation. However, the additional amplitude of those who had worn UVR-blocking lenses was + 0.25D measured objectively at 5.0 D of accommodative demand and + 0.7D measured subjectively at the average amplitude of accommodation of around 7.5D of demand. As around 80% of the subjective amplitude can be utilised in a sustained task [38], this difference could be considered clinically meaningful in delaying the onset of presbyopia. The amplitude of accommodation measured by push-up or the maximum negative powered lens that distance targets could be resolved through without debilitating blur was lower in the Hong Kong cohort. As the region with the highest UVR exposure of the cohorts examined in this study, this concurs with the high incidence of presbyopia occurring at younger ages that has been reported in countries with high levels of UVR [39,40].


Measurement of the dynamics of accommodation, such as speed of accommodation/ disaccommodation and latency, can be used as an indicator of the state of the accommodative system [41,42] and the development of presbyopia [43-45], in addition to assessment of the amplitude of accommodation [41]. Accommodative latency was found to be shorter in those wearing UVR-blocking contact lenses versus those wearing minimal UVR-blocking lenses, suggesting that UVR exposure can have an impact on when presbyopia occurs. The cohort from Houston USA had a shorter latency, faster speed and higher step size than some of the other regions, which cannot be explained by the other data collected in this study, so warrants further investigation.

There was no difference in MPOD between eyes that had worn UVR-blocking contact lenses compared to those that had worn contact lenses with minimal UVR-blocking contact lenses. This suggests that any epidemiological association between macular degeneration and UVR is unlikely to be through the protection afforded by the macular pigment



**Table 3**  
Comparison of participants wearing UVR and minimal UVR-blocking contact lenses across five international sites. N = 21 in each cohort. Average ± S.D. or Median [range].

	Australia		Canada		Hong Kong		USA		UK		Overall	
	UVR-Blocking	Minimal UVR-Blocking	UVR-Blocking	Minimal UVR-Blocking	UVR-Blocking	Minimal UVR-Blocking	UVR-Blocking	Minimal UVR-Blocking	UVR-Blocking	Minimal UVR-Blocking	UVR-Blocking	Minimal UVR-Blocking
Redness	1.2[0.9-1.3]	1.2[1.0-2.3]	0.8[0.5-1.1]	0.9[0.5-1.6]	1.6[1.2-2.9]	1.6[0.2-2.5]	1.6[1.2-2.8]	1.8[1.3-3.0]	1.2[0.8-2.9]	1.2[0.8-2.5]	1.2[0.5-2.9]	1.3[0.2-3.0]
Bulbar conjunctiva												
Limbal	0.8[0.8-1.2]	0.8[0.5-1.1]	0.5[0.2-0.8]	0.5[0.2-1.3]	1.3[1.0-2.7]	1.4[0.7-2.5]	1.4[1.0-2.8]	1.5[1.0-2.8]	1.1[0.2-2.4]	1.2[0.2-2.5]	1.0[0.2-2.8]	1.1[0.2-2.8]
Accommodation Push-up (D)	12.5±4.5	8.4±2.3	9.7±2.6	9.1±2.8	4.6±2.2	4.9±2.0	6.0±1.9	5.4±2.5	8.9±3.5	8.6±3.4	8.0±3.7	7.3±3.3
Minus Lens to blur (D)	-6.8±1.3	-5.7±1.8	-5.6±2.6	-5.5±2.1	-3.7±2.5	-4.3±2.8	-6.3±3.6	-5.5±3.2	-5.0±2.8	-5.0±2.4	-5.2±2.8	-5.1±2.5
Accommodative Stimulus / Response	0.15±0.17	0.20±0.42	0.19±0.28	0.23±0.22	0.16±0.27	0.13±0.28	0.29±0.27	0.16±0.27	0.20±0.28	0.17±0.44	0.20±0.26	0.18±0.32
1.0D	0.46±0.20	0.36±0.49	0.19±0.98	0.37±0.59	0.37±0.41	0.32±0.44	0.37±0.53	0.28±1.12	0.25±0.50	0.38±0.65	0.32±0.59	0.34±0.53
2.0D	1.10±0.47	0.84±0.63	0.87±1.00	1.26±0.63	1.16±0.64	1.10±0.62	1.43±0.42	1.12±0.80	0.77±0.85	1.02±0.66	1.05±0.75	1.08±0.67
3.0D	1.71±0.70	1.47±0.83	2.05±1.05	2.28±0.71	1.92±0.91	1.85±1.75	2.45±0.63	1.88±0.96	1.51±1.12	1.77±0.86	1.92±0.96	1.87±1.10
4.0D	2.36±0.89	2.12±1.04	2.78±1.21	3.21±0.78	2.43±1.29	2.74±1.12	3.22±0.77	2.64±1.32	2.42±1.21	2.46±0.99	2.65±1.12	2.66±1.10
5.0D	2.74±1.13	2.59±1.16	3.40±1.39	3.37±1.38	3.29±1.49	3.14±1.14	3.75±0.97	3.12±1.45	3.23±1.38	2.92±1.29	3.30±1.31	3.05±1.30
Latency (ms)	0.24±0.23	0.68±0.98	0.46±0.35	0.75±0.44	0.68±0.34	0.91±0.52	0.18±0.24	0.16±0.16	0.46±0.36	1.26±0.88	0.39±0.35	0.73±0.71
Speed (D/s)	1.60±0.70	1.12±0.82	1.95±0.92	1.81±0.86	1.72±0.86	2.21±1.25	2.23±0.96	2.31±1.24	1.75±0.87	1.57±0.63	1.86±0.88	1.89±1.02
Step size (D)	1.72±0.69	1.48±0.61	2.07±0.86	1.98±0.50	1.91±0.82	1.85±0.49	2.05±0.53	1.95±0.58	1.44±0.54	1.65±0.62	1.84±0.73	1.81±0.57
Nasal Autofluorescence												
Temporal (mm <sup>2</sup> )	2.37±2.86	2.89±3.75	1.23±2.50	1.31±2.25	1.53±3.73	1.77±5.59	0.62±1.50	0.68±1.62	1.58±3.33	2.00±3.34	1.43±2.89	1.68±3.60
Macular Pigment	0.51±0.17	0.53±0.17	0.47±0.11	0.43±0.11	0.54±0.18	0.52±0.18	0.50±0.21	0.41±0.20	0.38±0.17	0.40±0.18	0.46±0.18	0.45±0.18



**Fig. 1.** Change in accommodative response between Ultraviolet Radiation blocking and minimally Ultraviolet Radiation blocking contact lens wearers.

filter. Several studies have shown a higher macular pigment level in south Asians compared to Caucasians eyes [46-49], supporting the observation of the highest levels in the Hong Kong cohort in this study, although other cohorts included native Asian participants.

The limitations of the study were its retrospective nature of eye protection and the confounding effects of lifestyle such as indoor UVR exposure, beyond those that were matched for, that may have influenced UVR exposure between individuals. However, the same masked observers collected the clinical metrics across all sites, using the same instrumentation, allowing the first standardised comparison of the clinical effects of UVR-blocking filters embedded in contact lenses. More protective effects may have been observed for a longer period of ocular UVR protection.

In conclusion, blocking the transmission of UVR through a contact lens seems beneficial in maintaining the eye's ability to focus, suggesting that presbyopia may be delayed in long-term UVR-blocking contact lens wearers. There is also evidence they provide protection to the critical limbal region.

**Financial disclosures**

**J.S. Wolffsohn:** British Contact Lens Association (S); Alcon, Bausch and Lomb, Coopervision, Johnson and Johnson Vision (F,C); **S. Dhalla:** None **M. Aujla:** None **D. Laughton:** None **K. Tempany:** None **D. Powell:** None **K. Gifford:** Alcon, Coopervision, Menicon, Johnson and Johnson Vision (C) **P. Gifford:** None **K. Wan:** None **P. Cho:** None **U. Stahl:** None **J. Woods:** None.

**Funding/Support**

This research was funded by an investigator-initiated grant by Johnson and Johnson Vision AII5-501.

**Declaration of Competing Interest**

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

**References**

[1] Berneburg M, Gattermann N, Stege H, Grewe M, Vogelsang K, Ruzicka T, et al. Chronically Ultraviolet-exposed Human Skin Shows a Higher Mutation Frequency of Mitochondrial DNA as Compared to Unexposed Skin and the Hematopoietic System. *Photochem Photobiol* 1997;66(2):271-5.

[2] Runger TM. How different wavelengths of the ultraviolet spectrum contribute to skin carcinogenesis: the role of cellular damage responses. *J Invest Dermatol* 2007; 127(9):2103-5. <https://doi.org/10.1038/sj.jid.5700988>.

- [3] Young AR. Acute effects of UVR on human eyes and skin. *Prog Biophys Mol Biol* 2006;92(1):80–5.
- [4] Boettner EA, Reimer WJ. Transmission of the ocular media. *Investig Ophthalmol Visual Sci* 1962;1:776–83.
- [5] Moran DJ, Hollands FC. Pterygium and ultraviolet radiation: a positive correlation. *Br J Ophthalmol* 1984;68(5):343–6.
- [6] Saw SM, Tan D. Pterygium: prevalence, demography and risk factors. *Ophthalmic Epidemiol* 1999;6(3):219–28. <https://www.ncbi.nlm.nih.gov/pubmed/10487976>.
- [7] Chandler HL, Reuter KS, Sinnott LT, Nichols JJ. Prevention of UV-Induced Damage to the Anterior Segment Using Class I UV-Absorbing Hydrogel Contact Lenses. *Invest Ophthalmol Vis Sci* 2010;51(1):172–8. <https://doi.org/10.1167/iov.09-3996>.
- [8] Darzins P, Mitchell P, Heller RF. Sun Exposure and Age-related Macular Degeneration: An Australian Case—Control Study. *Ophthalmology* 1997;104(5):770–6.
- [9] Khan J, Shahid H, Thurlby D, Bradley M, Clayton D, Moore A, et al. Age related macular degeneration and sun exposure, iris colour, and skin sensitivity to sunlight. *Br J Ophthalmol* 2006;90(1):29–32.
- [10] Margrain TH, Boulton M, Marshall J, Sliney DH. Do blue light filters confer protection against age-related macular degeneration? *Prog Retin Eye Res* 2004;23(5):523–31. <https://doi.org/10.1016/j.preteyeres.2004.05.001>.
- [11] Taylor HR. The Long-term Effects of Visible Light on the Eye. *Arch Ophthalmol* 1992;110(1):99.
- [12] West SK. Exposure to Sunlight and Other Risk Factors for Age-Related Macular Degeneration. *Arch Ophthalmol* 1989;107(6):875.
- [13] Kwok LS, Kuznetsov VA, Ho A, Coroneo MT. Prevention of the adverse photic effects of peripheral light-focusing using UV-blocking contact lenses. *Invest Ophthalmol Vis Sci* 2003;44(4):1501–7. <https://www.ncbi.nlm.nih.gov/pubmed/12657585>.
- [14] Daniels JT, Dart JK, Tuft SJ, Khaw PT. Corneal stem cells in review. *Wound Repair Regenerat* 2001;9(6):483–94.
- [15] Walsh JE, Bergmanson JP. Does the eye benefit from wearing ultraviolet-blocking contact lenses? *Eye Contact Lens* 2011;37(4):267–72. <https://doi.org/10.1097/ICL.0b013e3182235777>.
- [16] Notara M, Behboudifard S, Kluth MA, Maßlo C, Ganss C, Frank MH, et al. UV light-blocking contact lenses protect against short-term UVB-induced limbal stem cell niche damage and inflammation. *Sci Rep* 2018;8(1). <https://doi.org/10.1038/s41598-018-30021-8>.
- [17] Bergmanson JP, Pitts DG, Chu LW. Protection from harmful UV radiation by contact lenses. *J Am Optom Assoc* 1988;59(3):178–82. <https://www.ncbi.nlm.nih.gov/pubmed/3351185>.
- [18] Dohlman CH, Gasset AR, Rose J. The effect of the absence of corneal epithelium or endothelium on the stromal keratocytes. *Invest Ophthalmol Vis Sci* 1968;7(5):520–34.
- [19] Wilson SE, Kim W-J. Keratocyte apoptosis: implications on corneal wound healing, tissue organization, and disease. *Invest Ophthalmol Vis Sci* 1998;39(2):220–6.
- [20] Moore L, Ferreira JT. Ultraviolet (UV) transmittance characteristics of daily disposable and silicone hydrogel contact lenses. *Contact Lens Anterior Eye J Br Contact Lens Associat* 2006;29(3):115–22. <https://doi.org/10.1016/j.clae.2006.03.002>.
- [21] Lira M, Dos Santos Castanheira EM, Santos L, Azeredo J, Yebra-Pimentel E, Real Oliveira ME. Changes in UV-visible transmittance of silicone-hydrogel contact lenses induced by wear. *Optomet Vis Sci Off Publicat Am Acad Optomet* 2009;86(4):332–9. <https://doi.org/10.1097/OPX.0b013e318198d047>.
- [22] Quesnel NM, Perron MJ, Giasson CJ. Effect of back vertex power on transmittance of contact lenses with UV-protection. *San Diego: AAO; 2005*.
- [23] International Organization for Standardization, & ISO 18369–1: 2017. (2010). *Ophthalmic optics—contact lenses—part 1: vocabulary, classification system and recommendations for labelling specifications*.
- [24] Rosenfield M, Cohen AS. Repeatability of clinical measurements of the amplitude of accommodation. *Ophthalmic Physiol Opt* 1996;16(3):247–9. <https://www.ncbi.nlm.nih.gov/pubmed/8977892>.
- [25] Raasch TW, Bailey IL, Bullimore MA. Repeatability of visual acuity measurement. *Optomet Vis Sci Off Publicat Am Acad Optomet* 1998;75(5):342–8. <https://doi.org/10.1097/00006324-199805000-00024>.
- [26] Lovie-Kitchin JE, Brown B. Repeatability and intercorrelations of standard vision tests as a function of age. *Optomet Vis Sci Off Publicat Am Acad Optomet* 2000;77(8):412–20. <https://doi.org/10.1097/00006324-200008000-00008>.
- [27] Bhatt UK, Sheppard AL, Shah S, Dua HS, Mihashi T, Yamaguchi T, et al. Design and validity of a miniaturized open-field aberrometer. *J Cataract Refract Surg* 2013;39(1):36–40. [S0886-3350\(12\)01305-3 \[pii\]](https://doi.org/10.1088/175137571235033).
- [28] Bartlett H, Stainer L, Singh S, Eperjesi F, Howells O. Clinical evaluation of the MPS 9000 Macular Pigment Screener. *Br J Ophthalmol* 2010;94(6):753–6. <https://doi.org/10.1136/bjo.2009.175901>.
- [29] Protection WHOaCoN-IR. Global solar UV index: a practical guide. *no. WHO/SDE/OEH/02.2 World Health Organization; 2002:28*.
- [30] Gies P, van Deventer E, Green AC, C S, Tinker R. Review of the global solar UV index 2015 workshop report Health physics 2018;114(1):84.
- [31] Chen A-H, O'Leary APDJ. Validity and repeatability of the modified push-up method for measuring the amplitude of accommodation. *Clin Exp Optomet* 1998;81(2):63–71. <https://doi.org/10.1111/j.1444-0938.1998.tb06628.x>.
- [32] Ostrin LA, Glasser A. Accommodation measurements in a presbyopic and presbyopic population. *J Cataract Refract Surg* 2004;30(7):1435–44. <https://doi.org/10.1016/j.jcrs.2003.12.045>.
- [33] Vargas V, Radner W, Allan BD, Reinstein DZ, Burkhard Dick H, Alio JL, et al. Methods for the study of near, intermediate vision, and accommodation: an overview of subjective and objective approaches. *Surv Ophthalmol* 2019;64(1):90–100. <https://doi.org/10.1016/j.survophthal.2018.08.003>.
- [34] van der Veen RL, Berendschot TT, Hendrikse F, Carden D, Makridaki M, Murray LJ. A new desktop instrument for measuring macular pigment optical density based on a novel technique for setting flicker thresholds. *Ophthalmic Physiol Opt* 2009;29(2):127–37. <https://doi.org/10.1111/j.1475-1313.2008.00618.x>.
- [35] Wolffsohn JS, Drew T, Sulley A. Conjunctival UV autofluorescence—Prevalence and risk factors. *Contact Len Anter Eye* 2014;37(6):427–30.
- [36] Wolffsohn JS, Drew T, Sulley A. Conjunctival UV autofluorescence—prevalence and risk factors. *Contact Lens Anterior Eye*. 2014;37(6):427–30.
- [37] Chiu C-J, Chang M-L, Zhang FF, Li T, Gensler G, Schleicher M, et al. The relationship of major American dietary patterns to age-related macular degeneration. *Am J Ophthalmol* 2014;158(1):118–127.e1.
- [38] Presbyopia TRJ. Emerging from a blur towards an understanding of the molecular basis for this most common eye condition. *Exp Eye Res* 2009;88(2):241–7. <https://doi.org/10.1016/j.exer.2008.07.003>.
- [39] Wolffsohn JS, Sheppard AL, Vakani S, Davies LN. Accommodative amplitude required for sustained near work. *Ophthalm Physiol Opt* 2011;31(5):480–6. <https://doi.org/10.1111/j.1475-1313.2011.00847.x>.
- [40] Stevens MA, Bergmanson JP. Does sunlight cause premature aging of the crystalline lens? *J Am Optom Assoc* 1989;60(9):660–3. <https://www.ncbi.nlm.nih.gov/pubmed/2677104>.
- [41] Nwosu SN. Ocular problems of young adults in rural Nigeria. *Int Ophthalmol* 1998;22(5):259–63. <https://www.ncbi.nlm.nih.gov/pubmed/10826540>.
- [42] Del Águila-Carrasco AJ, Esteve-Taboada JJ, Papadatou E, Ferrer-Blasco T, Montés-Micó R. Amplitude, Latency, and Peak Velocity in Accommodation and Disaccommodation Dynamics. *Biomed Res Int* 2017;2017:1–8.
- [43] Bharadwaj SR, Schor CM. Dynamic control of ocular disaccommodation: first and second-order dynamics. *Vision Res* 2006;46(6–7):1019–37.
- [44] Kasthurirangan S, Glasser A. Age related changes in accommodative dynamics in humans. *Vision Res* 2006;46(8–9):1507–19.
- [45] Mordí JA, Ciuffreda KJ. Dynamic aspects of accommodation: age and presbyopia. *Vision Res* 2004;44(6):591–601.
- [46] Taylor HR, West S, Muñoz B, Rosenthal FS, Bressler SB, Bressler NM. The long-term effects of visible light on the eye. *Arch Ophthalmol* 1992;110(1):99–104.
- [47] Davey PG, Lievens C, Ammono-Monney S. Differences in macular pigment optical density across four ethnicities: a comparative study. *Ther Adv Ophthalmol* 2020;12: 2515841420924167. [10.1177/2515841420924167](https://doi.org/10.1177/2515841420924167).
- [48] Huntjens B, Asaria TS, Dhanani S, Konstantakopoulou E, Ctori I. Macular pigment spatial profiles in South Asian and white subjects. *Invest Ophthalmol Vis Sci* 2014;55(3):1440–6. <https://doi.org/10.1167/iov.13-13204>.
- [49] Howells O, Eperjesi F, Bartlett H. Macular pigment optical density in young adults of South Asian origin. *Invest Ophthalmol Vis Sci* 2013;54(4):2711–9. <https://doi.org/10.1167/iov.12-10957>.